Title: A cross-sectional study of psychosocial factors and sickness presenteeism in Japanese workers during the COVID-19 pandemic

Masashi Masuda, M.D.¹, Tomohiro Ishimaru, M.D., M.P.H., Ph.D.¹, Ayako Hino, M.D., Ph.D.², Hajime Ando, M.D., M.O.H., Ph.D.³, Seiichiro Tateishi, M.D., Ph.D.⁴, Tomohisa Nagata, M.D., Ph.D.⁵, Mayumi Tsuji, M.D., Ph.D.⁶, Shinya Matsuda, M.D., Ph.D.⁷, and Yoshihisa Fujino, M.D., M.P.H., Ph.D.¹, for the CORoNaWork Project

¹ Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

² Department of Mental Health, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

³ Department of Work Systems and Health, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

⁴ Department of Occupational Medicine, School of Medicine, University of Occupational and Environmental Health, Japan

⁵ Department of Occupational Health Practice and Management, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

6 Department of Environmental Health, School of Medicine, University of Occupational
and Environmental Health, Japan

7 Department of Preventive Medicine and Community Health, School of Medicine,
University of Occupational and Environmental Health, Japan

Author contributions:

MM, Writing the manuscript; TI, AH and HA, creating the questionnaire, review of the
manuscript, and advice on interpretation; ST, TN, MT and SM, review of the manuscript,
advice on interpretation, and funding for research; and YF, Overall survey planning,
creating the questionnaire, analysis, and drafting the manuscript

Correspondence to: Yoshihisa Fujino¹, M.D., M.P.H., Ph.D.

Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,
University of Occupational and Environmental Health, Japan

1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

Tel: +81-93-691-7401

Email: zenq@med.uoeh-u.ac.jp

Funding Sources

This study was supported and partly funded by the University of Occupational and Environmental Health, Japan; General Incorporated Foundation (Anshin Zaidan); The Development of Educational Materials on Mental Health Measures for Managers at Small-sized Enterprises; Health, Labour and Welfare Sciences Research Grants; Comprehensive Research for Women's Healthcare (H30-josei-ippan-002); Research for the Establishment of an Occupational Health System in Times of Disaster (H30-roudou-ippan-007); and scholarship donations from Chugai Pharmaceutical Co., Ltd., the Collabo-Health Study Group, and Hitachi Systems, Ltd.

Conflict of Interest: The authors declare no conflicts of interest associated with this manuscript.

Acknowledgements

The current members of the CORoNaWork Project, in alphabetical order, are as follows: Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira Ogami,

Dr. Arisa Harada, Dr. Ayako Hino, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr. Kazunori Ikegami, Dr. Kei Tokutsu, Dr. Keiji Muramatsu, Dr. Koji Mori, Dr. Kosuke Mafune, Dr. Kyoko Kitagawa, Dr. Masako Nagata, Dr. Mayumi Tsuji, Ms. Ning Liu, Dr. Rie Tanaka, Dr. Ryutaro Matsugaki, Dr. Seiichiro Tateishi, Dr. Shinya Matsuda, Dr. Tomohiro Ishimaru, and Dr. Tomohisa Nagata. All members are affiliated with the University of Occupational and Environmental Health, Japan.

Ethical approval: This study was approved by the ethics committee of the University of Occupational and Environmental Health, Japan (reference No. R2-079 and R3-006).

Running title: Socioeconomic status and sickness presenteeism in Japanese workers

A cross-sectional study of psychosocial factors and sickness presenteeism in Japanese workers during the COVID-19 pandemic

Abstract:

Background: We examined the association between socioeconomic and health status, and lifestyle and sickness presenteeism among Japanese workers during the COVID-19 epidemic.

Methods: A cross-sectional study using an Internet-monitor survey was conducted in December, 2020 in Japan. Of 33,302 survey participants, we analyzed 27,036 participants (13,814 men and 13,222 women) who reported experience with sickness presenteeism.

Results: The odds ratio (OR) of sickness presenteeism associated with unmarried versus married status was 1.15. Respective figures for other variables were 1.11 for manual laboring work compared to desk work; 1.79 and 2.29 for loss of employment at the time the pandemic began and continuation of unemployment compared to maintaining employment during the pandemic; and 3.34 for a feeling of financial instability compared to stability.

Conclusion: The issue of sickness presenteeism has become more prominent under the COVID-19 epidemic.

Keywords: COVID-19; Japan; Socioeconomic Factors; Sickness Presenteeism

Introduction

Sickness presenteeism has rapidly attracted attention in occupational health. It has been defined as "the phenomenon of people, despite complaints and ill health that should prompt rest and absence from work, still turning up at their jobs"¹. Sickness presenteeism has been reported in a number of countries²⁻⁶. It is known that workers with sickness presenteeism are aware of suboptimal general health and at increased risk of developing coronary artery disease and depression⁷⁻⁹. Known workplace impacts of sickness presenteeism include increased long-term leave, reduced work capacity, and the spread of infectious diseases¹⁰⁻¹².

Vulnerable socioeconomic conditions, insecure employment status, as well as individual health views, work attitudes, company leave systems and culture influence workers' experience of sickness presenteeism¹³. Workers who are worried about their job security are reluctant to take time off work for treatment. Workers with insecure employment are more likely to engage in sickness presenteeism because they fear that complaining about their health condition will be detrimental to the status of their employment contract. The shortage of employees and inadequate leave system also lead to an increase in sickness presenteeism^{14,15}. Sick leave rates are low in areas where unemployment rate is high¹⁶, and it is thought that the fear of losing employment and poverty have led to situations in which people feel they have no choice but to work even if they are in poor physical condition¹⁷. In addition, worker behaviors, such as a positive job attitude, feelings of strong obligation, and considering absence a less legitimate option, are also known to result in sickness presenteeism¹⁸.

We hypothesized that the coronavirus disease 2019 (COVID-19) epidemic would affect worker experience of sickness presenteeism via an effect on their socioeconomic

status and employment instability, as well as on their health behaviors. Shortly after the initial confirmation of COVID-19 in China, the infection spread globally, and its presence in Japan was confirmed in January 2020. Since then, COVID-19 has continued to have a profound impact on healthcare systems, economic activities and people's lives around the world. Under such circumstances, the impact of the COVID-19 epidemic on sickness presenteeism is considered to be an occupational hygiene issue.

During a COVID-19 epidemic, workers may be more hesitant to report their health condition to the company when they are not feeling well. Reports worldwide have also noted the interruption of treatment for previously controlled diseases during the COVID-19 outbreak^{19,20}, which also acts to increase sickness presenteeism. Furthermore, the management of sickness presenteeism is important in terms of preventing the spread of infection. Among various measures taken to halt the spread of COVID-19 in the workplace, self-reporting of physical condition and restriction of attendance by employees who have fever or other health problems are particularly important - the fact that some employees still come to work despite feeling ill is now an issue.

Some studies have considered the assumption that the COVID-19 epidemic affects the experience of workers' sickness presenteeism^{21,22}. An analysis that combined survey data from 20,974 employees collected in Europe with Eurostat's regional unemployment data shows that high unemployment rates enhance the presenteeism of disadvantaged workers²¹. The COVID-19 pandemic and consequent increase in labor market instability can increase the behavior of presenteeism. An Internet survey conducted in Belgium and the Netherlands after 8 weeks of coronavirus lockdown showed that the respondents experienced considerable levels of stress and concern about their financial situation, and also cancelled/delayed general healthcare and productivity loss during the COVID-19

70 pandemic²². The study also estimated presenteeism during the 8 weeks, but did not show
71 an association between presenteeism and the factors experienced by respondents.
72 However, there are few reports on the characteristics of workers experiencing sickness
73 presenteeism during the COVID-19 epidemic. We examined the association between
74 socioeconomic and health status, and lifestyle and sickness presenteeism among Japanese
75 workers during the COVID-19 epidemic.

Methods

Study Design and Participants

A cross-sectional study was conducted from December 22 to 26, 2020 at the time of the third wave of COVID-19 infection in Japan. The study was conducted as an Internet-monitor survey. Details of the study protocol have been described elsewhere²³. Briefly, the data were obtained from workers currently under an employment contract at the time of the survey, and categorized by prefecture, type of job, and sex. Of 33,302 survey participants, 215 respondents were excluded because they were deemed to have provided fraudulent responses by the surveying company (Cross Marketing Inc.,). Subsequently, 6,051 respondents determined to have provided invalid or erroneous responses were excluded (See reference 23 for details). Finally, 27,036 people were included in the study. The present analysis included all participants who responded that they had attended work despite needing to recuperate at home.

This study was approved by the ethics committee of the University of Occupational and Environmental Health, Japan (R2-079, February 5, 2021). Informed consent was obtained via a form on the survey website.

Assessment of sickness presenteeism

Sickness presenteeism was determined using the following single-item question: “How many days have you worked (including at home) in the last 30 days under conditions in which you would really like to take a day off?” To evaluate the sickness presenteeism during the third wave of COVID-19 in Japan, we set the target period to 30 days. Respondents were asked to respond with a specific number of days; based on the distribution of days with sickness presenteeism, we set the cut-off to 3 days.

Survey of participants socioeconomic conditions, health status, and lifestyle factors

The questionnaire was conducted via the Internet and enquired about the participant's socioeconomic conditions. Specifically, the survey enquired about age, sex, marital status (married; unmarried; widowed/divorced), occupation (job mainly involves desk work; interpersonal communication; labor), educational background (Junior high school; High school; Vocational school/College; University; Graduate school), equivalised income (household income divided by the square root of household size), job change or unemployment after April 2020 (no resignation or job change; transfer to another company; resignation and immediate start at a new job; unemployment for a period, but presently employed; retired and started a business), and perception of financial situation (very difficult; somewhat difficult; slightly difficult; comfortable).

With regard to health status and psychological factors, we asked about self-rated health, psychological distress, loneliness, presence of supportive friends, and having a health condition that requires company consideration to allow work. Psychological distress was addressed using the Kessler Psychological Distress Scale (K6)²⁴, whose validity in Japanese has been confirmed²⁵. Participants with a K6 score of 5 or higher were considered to have mild psychological distress. Loneliness was examined with the question: “Have you ever felt alone in the past month?”, with response options of: “never,” “a little,” “sometimes,” “usually,” and “always.” This method follows previous studies that assessed loneliness with a single question²⁶. The presence of a health condition that required the employer’s consideration to permit work was surveyed using the question: “Do you require consideration or support from your company to continue working in your current health condition?”, with the three response options of: “no,” “yes, but I have not

received support;” and “yes, and I have received support.”

The questionnaire also contained a variety of questions on lifestyle and work-related factors, including smoking (never; quit more than one year ago; quit within the past year; started to smoke < 1 year ago; have smoked for > 1 year), drinking habit (6–7 days/week; 4–5 days/week; 2–3 days/week; < 1 day/week; almost never), exercise habit, breakfast routine, time spent in one-way commuting and number of overtime hours worked/day. With regard to exercise, participants indicated the number of days each week on which they exercised for 30 minutes or more. Regarding breakfast, they indicated the number of days each week on which they had breakfast.

Statistical analysis

Age- and sex-adjusted odds ratios (ORs) and multivariate adjusted ORs were analyzed using a multilevel logistic model with nesting by residential prefecture²³. All analyses included the incidence rate of COVID-19 since declared pandemic status was an area-level variable. In the analyses of socioeconomic conditions, the model included sex, age, marital status, equivalised income (divided into tertiles), job type, employment status, educational background, and level of comfort with financial condition. The model for analysis of health-related factors included age, sex, psychological distress, self-rated health, feeling of loneliness, having a friend who can provide support, and having a health condition which requires the employer’s support to allow work to occur. The model used to analyse lifestyle and work-related factors incorporated age, sex, drinking habit, smoking habit (classified into three groups, "non-smoker"; "ex-smoker"; "current smoker"), exercise, breakfast routine, overtime hours worked/day and time required for one-way commuting.

148 A *P* value of less than .05 (two-tailed) was considered to indicate statistical
149 significance. Stata Statistical Software version 16 (StataCorp LLC, TX, USA) was used
150 for all analyses.

151

152

153

Results

Basic participant characteristics are shown in Table 1. Among all 27,036 participants, approximately 19 percent of participants had experienced sickness presenteeism. On comparison, participants who had experienced sickness presenteeism were more likely than those who had not to have lower level of subjective health, greater psychological distress, greater loneliness, and greater need for employment consideration from their employer.

ORs of socioeconomic status associated with sickness presenteeism are shown in Table 2. On multivariate analysis, the odds ratio (OR) of sickness presenteeism associated with not being married versus being married was 1.15 (95% confidence interval (95%CI): 1.04-1.28, p=0.009). Respective values were 1.11 (95%CI: 1.03-1.20, p=0.009) for manual labor versus desk work; 1.79 (95%CI: 1.47-2.19, p<0.001) and 2.29 (95%CI: 1.79-2.92, p<0.001) for lost employment when the COVID-19 pandemic started and continued unemployment compared with employment across the entire pandemic period; and 3.34 (95%CI: 2.94-3.81, p<0.001) for a feeling of financial instability versus stability.

The ORs of health status and psychological distress associated with sickness presenteeism is shown in Table 3. On multivariate analysis, the OR of sickness presenteeism associated with poor self-rated health was 11.21 (95%CI: 10.24-12.28, p<0.001); experience of psychological distress was 6.11 (95%CI: 5.70-6.56, p<0.001); feeling always alone was 8.30 (95%CI: 7.34-9.39, p<0.001); and the need for company consideration to allow work but not receiving it was 9.57 (95%CI: 8.87-10.32, p<0.001).

The ORs of lifestyle and occupational factors associated with sickness presenteeism is shown in Table 4. Smoking, breakfast routine, commute time, and hours worked overtime were associated with sickness presenteeism. In contrast, we saw no association

between drinking habit (except less than 1 day a week) and sickness presenteeism. On multivariate analysis, current smokers were at higher likelihood of experiencing sickness presenteeism (OR=1.20, 95%CI: 1.12-1.29, $p<0.001$), as were those who rarely ate breakfast (OR=1.49, 95%CI: 1.38-1.62, $p<0.001$), spent longer than 2 hours in one-way commuting (OR=1.45, 95%CI: 1.18-1.77, $p<0.001$), or worked more than 2 hours overtime (OR=2.53, 95%CI: 2.53-2.80, $p<0.001$).

Discussion

This study found that 19% of workers in Japan had experienced sickness presenteeism during the COVID-19 pandemic. A survey of Japanese workers conducted before the COVID-19 pandemic found that 27.6% of workers had experienced sickness presenteeism due to mental health problems 2 to 5 times in the last 12 months²⁷. It is not possible to make a simple comparison between this previous and the present study because the conditions of the two surveys are different. However, considering that the target period of this survey was only one month, it can be said that this result shows that COVID-19 is sufficiently associated with sickness presenteeism. Risk of experiencing sickness presenteeism was higher in workers with a lower socioeconomic status and those in poor health. Further, risk was also higher in workers with an unfavorable lifestyle and working conditions.

Socioeconomic status is known to be an important factor in a worker's experience of sickness presenteeism^{14,28}. Further, emergencies such as disasters impair access to health care for people with a disadvantaged socioeconomic status^{29,30}. As a result, opportunities for workers to experience sickness presenteeism can be expected to increase. The present study reveals that socioeconomically disadvantaged workers were more likely to experience sickness presenteeism during the COVID-19 pandemic. In particular, risk of sickness presenteeism was increased in subjects with low income, unemployment experience, and economic insecurity, a finding consistent with previous studies of the impact of COVID-19 on sickness presenteeism^{21,22}. Economic deprivation is a direct reason for sickness presenteeism. Workers with unstable employment or a low income have little choice but to work hard for their livelihoods, even if they are not feeling well¹⁴. Conversely, workers with sickness presenteeism are more likely to exacerbate their

illness¹⁰. As a result, sickness presenteeism will work in the direction of lower income and employment instability. Workers with unstable employment may not be entitled to sick leave or may be reluctant to take it^{31,32}, rendering them prone to sickness presenteeism. Moreover, experience of sickness presenteeism has also been shown to exacerbate health status, resulting in turn in employment instability as well as economic disadvantage.

Our present results revealed that those with a poor health condition had a greater likelihood of experiencing sickness presenteeism: those whose self-rated health was poor, or with psychological distress, loneliness, and workplace support were at greater risk of experiencing sickness presenteeism. It is clear from the definition that sickness presenteeism is more likely in poor health¹. Conversely, workers with sickness presenteeism are known to be more likely to exacerbate their illness¹⁰. In our present study, we found an association between greater psychological distress and a higher likelihood of sickness presenteeism. A number of studies have noted that the fear of infection when visiting a hospital is a major reason for treatment interruption³³⁻³⁵. Anxiety concerning infection may affect treatment discontinuation, resulting in continued working in poor physical condition. In addition, participants without company support were more likely to have sickness presenteeism. This result is consistent with reports that there is less sickness presenteeism with the support of workplace managers³⁶. For those who tend to feel lonely, it is thought that work efficiency is likely to decline due to the infectious disease countermeasures (eg Stay Home and Remote Work) that have become widespread during the COVID-19 epidemic, and the resulting decline in workplace communication.

Our present results also revealed the association of non-preferable lifestyle and work-related factors with sickness presenteeism. Previous studies have shown a

relationship between lifestyle and sickness presenteeism³⁷⁻³⁹. It is thought that stress has likely to have increased due to environmental changes caused by infectious disease countermeasures (eg, Stay Home and Remote Work) that have become widespread during the COVID-19 epidemic. Sickness presenteeism will increase due to a deterioration in lifestyle, such as the stress caused by countermeasures against COVID-19 and work at home, where the lack of restrictions on smoking leads to an increase in the number of cigarettes smoked⁴⁰. Long commute times put some physical burden on workers. Also, longer commute times reduce leisure and sleep time in daily life. Returning home late at night and having a supper also contribute to the worsening of lifestyle-related diseases. Therefore, long commute times can lead to sickness presenteeism⁴¹. Countermeasures against COVID-19 recommended in Japan such as staggered commuting and ventilation inside the train by opening windows require workers to go to work quite early in the morning and to travel in a train that is not adjusted to a comfortable temperature or humidity. This can also put a physical and mental burden on workers and increase sickness presenteeism. This is considered to be especially noticeable for workers who commute for long hours.

Sickness presenteeism is a major occupational health issue and public health challenge. The COVID-19 pandemic has brought a particular focus to it. Our present study emphasizes that sickness presenteeism is associated with socioeconomic status. Regardless of causation, individuals with a lower socioeconomic or health status were at greater risk of sickness presenteeism. It is therefore critical to recognize not only the clinical importance of COVID-19, but in addition the deterioration in labor productivity and the impact on the lives and health of workers through the socioeconomic environment. In a survey of Japanese workers, 62% of those with symptoms such as fever went to work

within 7 days after symptom onset. Being hired by a company and not being able to work remotely were associated with going to work within 7 days of the onset of symptoms⁴².

It is important to foster the social consensus that workers should take the day off from work when they are feeling unwell. It is known that companies with a leave system that is easy to use when workers are not feeling well have a low incidence of infectious diseases¹⁵. An easy-to-use leave system is effective not only in reducing sickness presenteeism but also with regard to COVID-19 countermeasures.

There are several limitations in this study. First, the reasons for sickness presenteeism are not known. Many possible explanations may explain why workers experience sickness presenteeism, including financial hardship which prevents medical visits, unstable employment, or unavailability of a sick leave entitlement. Second, we did not investigate the types of illnesses and health problems experienced by people with sickness presenteeism. Third, we did not determine the chronological relationship between sickness presenteeism and socioeconomic status and lifestyle.

In this study, we found that around 19% of Japanese workers experienced sickness presenteeism during the period of rapid spread of COVID-19 infection. Those who were socioeconomically disadvantaged, in poor health, or with an unfavorable lifestyle were more likely to report sickness presenteeism. Accordingly, the issue of sickness presenteeism has become more prominent under the COVID-19 epidemic. The increase in sickness presenteeism may not only worsen the health status of individuals, but also have long-term effects on society, such as reduced productivity and increased social security burden due to employment instability. These findings indicate the need for efforts to decrease sickness presenteeism in workers who need to recuperate at home.

References

1. Aronsson G, Gustafsson K, Dallner M. Sick but yet at work. An empirical study of sickness presenteeism. *J Epidemiol Community Health*. 2000;54:502-509.
2. Eurofound. Health and well-being at work: a report based on the fifth european working conditions survey. Dublin: 2012.
3. Webster RK, Liu R, Karimullina K, et al. A systematic review of infectious illness Presenteeism: prevalence, reasons and risk factors. *BMC Public Health*. 2019 Jun 21;19:799.
4. Shan G, Wang S, Wang W, et al. Presenteeism in Nurses: Prevalence, Consequences, and Causes From the Perspectives of Nurses and Chief Nurses. *Front Psychiatry*. 2021 Jan 8;11:584040.
5. Park JW, Cho SS, Lee J, et al. Association between employment status and sickness presenteeism among Korean employees: a cross-sectional study. *Ann Occup Environ Med*. 2020 Jun 12;32:e17.
6. Oshio T, Tsutsumi A, Inoue A, et al. The reciprocal relationship between sickness presenteeism and psychological distress in response to job stressors: evidence from a three-wave cohort study. *J Occup Health*. 2017 Nov 25;59:552-561.
7. Taloyan M, Aronsson G, Leineweber C, et al. Sickness presenteeism predicts suboptimal self-rated health and sickness absence: A nationally representative study of the Swedish working population. *PLoS One*. 2012;7:e44721.
8. Kivimäki M, Head J, Ferrie JE, et al. Working while ill as a risk factor for serious coronary events: the Whitehall II study. *Am J Public Health*. 2005;95:98-102.
9. Conway PM, Hough A, Rugulies R, et al. Is sickness presenteeism a risk factor for depression? A Danish 2-year follow-up study. *J Occup Environ Med*. 2014;56:595-603.

- 305 10. Hansen CD, Andersen JH. Sick at work — a risk factor for long-term sickness absence
306 at a later date? *J Epidemiol Community Health*. 2009; 63:397-402.
- 307 11. Dellve L, Hadzibajramovic E, Ahlborg G Jr. Work attendance among healthcare
308 workers: prevalence, incentives, and long-term consequences for health and
309 performance. *J Adv Nurs*. 2011;67:1918-1929.
- 310 12. Rosvold EO, Bjertness E. Physicians who do not take sick leave: hazardous heroes?
311 *Scand J Public Health*. 2001;29:71-75.
- 312 13. Johns G. Presenteeism in the workplace: A review and research agenda. *Journal of*
313 *Organizational Behavior* 2010;31:519-542.
- 314 14. Aronsson G, Gustafsson K. Sickness presenteeism: Prevalence, attendance-pressure
315 factors, and an outline of a model for research. *J Occup Environ Med*. 2005;47:958-
316 966.
- 317 15. Böckerman P, Laukkanen E. What makes you work while you are sick? Evidence
318 from a survey of workers. *Eur J Public Health*. 2010;20:43-46.
- 319 16. Allebeck P, Mastekaasa A. Chapter 5. Risk factors for sick leave - general studies.
320 *Scand J Public Health* 2004;32:49–108.
- 321 17. Sverke M, Hellgren J, Näswall K. No security: a meta-analysis and review of job
322 insecurity and its consequences. *J Occup Health Psychol* 2002;7:242–64.
- 323 18. Miraglia M, Johns G. Going to work ill: A meta-analysis of the correlates of
324 presenteeism and a dual-path model. *J Occup Health Psychol*. 2016 Jul;21:261-283.
- 325 19. Bong CL, Brasher C, Chikumba E, et al. The COVID-19 Pandemic: Effects on Low-
326 and Middle-Income Countries. *Anesth Analg*. 2020 Jul; 131: 86-92.
- 327 20. Das AK, Mishra DK, Gopalan SS. Reduced access to care among older American
328 adults during CoVID-19 pandemic: results from a prospective cohort study. *medRxiv*.

329 2020: 2020.11.29.20240317.

330 21. Reuter M, Dragano N, Wahrendorf M. Working while sick in context of regional
331 unemployment: a Europe-wide cross-sectional study. *J Epidemiol Community Health*.
332 2020 Nov 13;jech-2020-214888.

333 22. van Ballegooijen H, Goossens L, Bruin RH, et al. Concerns, quality of life, access to
334 care and productivity of the general population during the first 8 weeks of the
335 coronavirus lockdown in Belgium and the Netherlands. *BMC Health Serv Res*. 2021
336 Mar 12;21:227.

337 23. Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a nationwide Internet-based health
338 survey in workers during the COVID-19 pandemic in 2020. *medRxiv*. Published online
339 February 5, 2021:2021.02.02.21249309.

340 24. Kessler RC, Andrews G, Colpe LJ, et al. Short screening scales to monitor population
341 prevalences and trends in non-specific psychological distress. *Psychol Med*.
342 2002;32:959-976.

343 25. Furukawa TA, Kawakami N, Saitoh M, et al. The performance of the Japanese version
344 of the K6 and K10 in the World Mental Health Survey Japan. *Int J Methods Psychiatr*
345 *Res*. 2008;17:152-158.

346 26. Courtin E, Knapp M. Social isolation, loneliness and health in old age: a scoping
347 review. *Health Soc Care Community*. 2017 May;25:799-812.

348 27. Doki S, Sasahara S, Suzuki S, et al. Relationship between sickness presenteeism and
349 awareness and presence or absence of systems for return to work among workers with
350 mental health problems in Japan: an Internet-based cross-sectional study. *J Occup*
351 Health. 2015;57:532-539.

352 28. Kim JY, Lee J, Muntaner C, et al. Who is working while sick? Nonstandard

employment and its association with absenteeism and presenteeism in South Korea. *Int Arch Occup Environ Health*. 2016;89:1095-1101.

29. Feinstein JS. The Relationship between Socioeconomic Status and Health: A Review of the Literature. *M 323 ilbank Q*. 1993;71:279-322.

30. Swain GR. How does economic and social disadvantage affect health. *Focus*. 2016;33:1-6.

31. Kinman G, Grant C. Presenteeism during the COVID-19 pandemic: risks and solutions. *Occup Med* . Published online November 18, 2020. doi:10.1093/occmed/kqaa193

32. Skagen K, Collins AM. The consequences of sickness presenteeism on health and wellbeing over time: A systematic review. *Soc Sci Med*. 2016;161:169-177.

33. Czeisler MÉ, Marynak K, Clarke KEN, et al. Delay or Avoidance of Medical Care Because of COVID-19-Related Concerns - United States, June 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:1250-1257.

34. Chudasama YV, Gillies CL, Zaccardi F, et al. Impact of COVID-19 on routine care for chronic diseases: A global survey of views from healthcare professionals. *Diabetes Metab Syndr*. 2020;14:965-967.

35. Erol MK. Treatment Delays and In-Hospital Outcomes In Acute Myocardial Infarction During The Covid-19 Pandemic: A Nationwide Study. *The Anatolian Journal of Cardiology*. Published online 2020. doi:10.14744/anatoljcardiol.2020.98607

36. Caverley N, Cunningham JB, MacGregor JN. Sickness presenteeism, sickness absenteeism, and health following restructuring in a public service organization. *J Manag Studies*. 2007;44:304-319.

37. Robroek SJ, van den Berg TI, Plat JF, et al. The role of obesity and lifestyle behaviours

377 in a productive workforce. *Occup Environ Med.* 2011;68:134-139.

378 38. Kirkham HS, Clark BL, Bolas CA, et al. Which modifiable health risks are associated
379 with changes in productivity costs? *Popul Health Manag.* 2015;18:30-38.

380 39. Merchant JA, Kelly KM, Burmeister LF, et al. Employment status matters: a statewide
381 survey of quality-of-life, prevention behaviors, and absenteeism and presenteeism. *J*
382 *Occup Environ Med.* 2014; 56:686-698.

383 40. Siegel A, Korbman M, Erbllich J. Direct and Indirect Effects of Psychological Distress
384 on Stress-Induced Smoking. *J Stud Alcohol Drugs.* 2017 Nov;78:930-937.

385 41. Min D, Lee SJ. Factors Associated With the Presenteeism of Single-Person Household
386 Employees in Korea: The 5th Korean Working Conditions Survey (KWCS). *Occup*
387 *Environ Med.* 2021 Sep 1;63:808-812.

388 42. Machida M, Nakamura I, Saito R, et al. The actual implementation status of self-
389 isolation among Japanese workers during the COVID-19 outbreak. *Trop Med Health.*
390 2020 Aug 3;48:63. doi: 10.1186/s41182-020-00250-7. eCollection 2020.

Table 1. Basic characteristics of the study participants

	Sickness presenteeism <3day (n=21997)	Sickness presenteeism ≥3day (n=5039)
Sex		
Women	10580 (48.1%)	2642 (52.4%)
Men	11417 (51.9%)	2397 (47.6%)
Marital status		
Married	12503 (56.8%)	2526 (50.1%)
Unmarried	7212 (32.8%)	1952 (38.7%)
Divorced/widowed	2282 (10.4%)	561 (11.1%)
Job type		
Mainly desk work	11126 (50.6%)	2342 (46.5%)
Jobs mainly involving interpersonal communication	5556 (25.3%)	1371 (27.2%)
Mainly labor	5315 (24.2%)	1326 (26.3%)
Annual equivalent household income (JPY)		
500000-2650000	7075 (32.2%)	1919 (38.1%)
2650000-4500000	7139 (32.5%)	1543 (30.6%)
>4500000	7783 (35.4%)	1577 (31.3%)
Self-rated health, very good/good	12305 (55.9%)	1151 (22.8%)
Psychological distress (K6≥5)	7054 (32.1%)	3763 (74.7%)
Have you ever felt alone? usually or always	1535 (7.0%)	1215 (24.1%)
Do you require consideration or support from your company to continue working in your current health condition?		
No	18122 (82.4%)	2139 (42.4%)
Yes, but I have not received support	2001 (9.1%)	2297 (45.6%)
Yes and I have received support	1874 (8.5%)	603 (12.0%)
Smoking status		
Non-smoker	11897 (54.1%)	2690 (53.4%)
Ex-smoker	4096 (18.6%)	841 (16.7%)
Current smoker	6004 (27.3%)	1508 (29.9%)
Alcohol consumption, 6–7 days a week	4690 (21.3%)	984 (19.5%)
How often do you eat breakfast? less than 1 day a week	3697 (16.8%)	1095 (21.7%)
Time spent on one-way commute (excluding time spent at home), more than 2 hours	434 (2.0%)	151 (3.0%)
Overtime work, more than 2 hours	1732 (7.9%)	721 (14.3%)

JPY, Japanese yen; K6, Kessler Psychological Distress Scale.

Table 2. Association between socioeconomic status and sickness presenteeism

	Age-sex adjusted				Multivariate*			
	OR	95%CI	p		OR	95%CI	p	
Education								
Junior high school	1.32	1.03	1.69	0.027	1.17	0.91	1.50	0.212
High school	1.02	0.95	1.10	0.538	0.97	0.90	1.05	0.464
Vocational school/college, university, graduate school	reference				reference			
Marital status								
Married	reference				reference			
Unmarried	1.25	1.13	1.39	<0.001	1.15	1.04	1.28	0.009
Divorced/bereavement	1.19	1.11	1.28	<0.001	1.13	1.05	1.21	0.001
Job type								
Mainly desk work	reference				reference			
Jobs mainly involving interpersonal communication	1.13	1.05	1.22	<0.001	1.11	1.03	1.20	0.005
Mainly labor	1.16	1.08	1.26	<0.001	1.11	1.03	1.20	0.009
Equivalized income (JPY)								
500000-2650000	1.29	1.19	1.38	<0.001	1.19	1.10	1.29	<0.001
2650000-4500000	1.03	0.95	1.11	0.474	1.01	0.94	1.10	0.721
>4500000	reference				reference			
Have you resigned or changed jobs since April 2020?								
Did not resign or change jobs	reference				reference			
Transferred to another company	2.66	1.92	3.69	<0.001	2.69	1.94	3.73	<0.001
Resigned and entered into a new job immediately	1.40	1.18	1.65	<0.001	1.36	1.15	1.61	<0.001
Unemployed for a period, but currently working	1.90	1.56	2.32	<0.001	1.79	1.47	2.19	<0.001
Retired and started a business (e.g., running a company, personal business, or self-employment)	2.42	1.89	3.09	<0.001	2.29	1.79	2.92	<0.001
How do you feel about your current financial situation?								
Very difficult	3.41	3.01	3.88	<0.001	3.34	2.94	3.81	<0.001
Slightly difficult	1.95	1.73	2.19	<0.001	1.94	1.72	2.18	<0.001
Fairly difficult	1.13	1.01	1.27	0.035	1.13	1.01	1.27	0.033
Comfortable	reference				reference			

OR, odds ratio; 95%CI, 95% confidence interval; JPY, Japanese yen.

Table 3. Association between health status, psychological distress and sickness presenteeism

	Age-sex adjusted				Multivariate*			
	OR	95%CI		p	OR	95%CI		p
Self-rated health								
Very good/good	reference				reference			
Neither good nor bad	2.81	2.60	3.04	<0.001	2.81	2.60	3.04	<0.001
Bad	11.21	10.25	12.27	<0.001	11.21	10.24	12.28	<0.001
Psychological distress (K6≥5)	6.14	5.73	6.59	<0.001	6.11	5.70	6.56	<0.001
Have you ever felt alone?								
Never	reference				reference			
A little	2.51	2.30	2.73	<0.001	2.52	2.32	2.74	<0.001
Sometimes	4.30	3.93	4.69	<0.001	4.32	3.95	4.73	<0.001
Usually	6.13	5.45	6.90	<0.001	6.21	5.51	7.00	<0.001
Always	8.17	7.24	9.23	<0.001	8.30	7.34	9.39	<0.001
Do you require consideration or support from your company to continue working in your current health condition?								
No	reference				reference			
Yes, but I have not received support	9.67	8.97	10.42	<0.001	9.57	8.87	10.32	<0.001
Yes and I have received support	2.64	2.38	2.92	<0.001	2.64	2.38	2.92	<0.001

OR, odds ratio; 95%CI, 95% confidence interval; K6, Kessler Psychological Distress Scale.

Table 4. Association between lifestyle, occupational factors and sickness presenteeism

	Age-sex adjusted				Multivariate*			
	OR	95%CI	p		OR	95%CI	p	
Smoking status								
Non-smoker	reference				reference			
Ex-smoker	1.06	0.97	1.16	0.189	1.06	0.97	1.16	0.207
Current smoker	1.26	1.17	1.36	<0.001	1.25	1.16	1.35	<0.001
Drinking habit								
6–7 days a week	0.96	0.88	1.05	0.337	0.98	0.90	1.07	0.674
4–5 days a week	0.91	0.80	1.03	0.142	0.94	0.83	1.06	0.301
2–3 days a week	0.96	0.87	1.06	0.385	0.98	0.89	1.08	0.677
Less than 1 day a week	0.88	0.80	0.96	0.004	0.89	0.81	0.97	0.008
Hardly ever	reference				reference			
How often do you eat breakfast?								
6–7 days a week	reference				reference			
4–5 days a week	1.43	1.29	1.59	<0.001	1.41	1.27	1.57	<0.001
2–3 days a week	1.71	1.52	1.93	<0.001	1.68	1.49	1.90	<0.001
Less than 1 day a week	1.72	1.50	1.97	<0.001	1.70	1.49	1.95	<0.001
Hardly ever	1.54	1.42	1.66	<0.001	1.49	1.38	1.62	<0.001
Time spent on one-way commute (excluding time spent at home)								
More than 2 hours	1.35	1.11	1.66	0.003	1.45	1.18	1.77	<0.001
More than 1 hour	1.06	0.95	1.19	0.272	1.13	1.01	1.27	0.029
More than 30 minutes	0.92	0.83	1.01	0.086	0.96	0.87	1.06	0.386
Less than 30 minutes	0.79	0.73	0.87	<0.001	0.81	0.73	0.88	<0.001
Almost none	reference				reference			
Overtime work								
More than 2 hours	2.39	2.16	2.64	<0.001	2.53	2.28	2.80	<0.001
More than 1 hour	1.67	1.53	1.83	<0.001	1.75	1.60	1.91	<0.001
More than 30 minutes	1.45	1.32	1.58	<0.001	1.50	1.37	1.64	<0.001
Less than 30 minutes	1.15	1.05	1.28	0.005	1.19	1.07	1.31	0.001
Almost none	reference				reference			

OR, odds ratio; 95%CI, 95% confidence interval.