TITLE

COVID-19 risk factors amongst 14,786 care home residents: An observational longitudinal analysis including daily community positive test rates of COVID-19, hospital stays, and vaccination status in Wales (UK) between 1st September 2020 and 1st May 2021.

AUTHORS

Joe Hollinghurst – Swansea University

Robyn Hollinghurst – Open University, Swansea University

Laura North – Swansea University

Amy Mizen – Swansea University

Ashley Akbari – Swansea University

Sara Long – Swansea University

Ronan A Lyons - Population Data Science, Swansea University

Rich Fry – Swansea University

AUTHOR CONTRIBUTIONS

JH – Conceived the study, determined the research questions, created the analysis datasets, analysed the data, and wrote the manuscript.

RH – Performed a literature review, interpreted the results, and edited the manuscript.

LN – Contributed to data preparation and reviewed the manuscript.

AM – Contributed to the design of the study and edited the manuscript.

SL – Contributed to the design of the study and edited the manuscript.

RAL – Secured funding, contributed to the design and edited the manuscript.

RF – Conceived the study, determined the research questions, edited the manuscript.

Corresponding author: Joe Hollinghurst, j.p.hollinghurst@swansea.ac.uk

The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence on a worldwide basis to the BMJ Publishing Group Ltd to permit this article to be published in BMJ editions and any other BMJ PGL products and sublicences such use and exploit all subsidiary rights, as set out in our licence.

All authors have completed the Unified Competing Interest form (available on request from the corresponding author) and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years, no other relationships or activities that could appear to have influenced the submitted work.

The lead author (the manuscript's guarantor) affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned have been explained.

SUMMARY BOXES

Section 1: What is already known on this topic

- Care home residents are at a high risk of COVID-19 infection, but existing literature has mainly focussed on excess mortality rather than infection risk.
- In our study we were able to investigate associations between COVID-19 infections and the community positive test rate of COVID-19, the vaccination status of care home residents, hospital admissions, and frailty.

Section 2: What this study adds

- Our study suggests an increased community positive test rate and hospital inpatients had an increased likelihood of a positive SARS-CoV-2 polymerase chain reaction test, whilst one or two doses of vaccination indicated a decreased chance of a positive test.
- Our findings suggest care providers need to stay vigilant despite the vaccination rollout, and extra precautions should be taken when caring for the most vulnerable, especially in a hospital setting.

ABSTRACT

Objectives: Determine individual level risk factors for care home residents testing positive for SARS-CoV-2.

Study Design: Longitudinal observational cohort study using individual-level linked data.

Setting: Care home residents in Wales (United Kingdom) between 1st September 2020 and 1st May 2021.

Participants: 14,786 older care home residents (aged 65+). Our dataset consisted of 2,613,341 individual-level daily observations within 697 care homes.

Methods: We estimated odds ratios (ORs [95% confidence interval]) using multilevel logistic regression models. Our outcome of interest was a positive SARS-CoV-2 polymerase chain reaction (PCR) test. We included time dependent covariates for the estimated community positive test rate of COVID-19, hospital admissions, and vaccination status. Additional covariates were included for age, positive PCR tests prior to the study, sex, frailty (using the hospital frailty risk score), and specialist care home services.

Results: The multivariable logistic regression model indicated an increase in age (OR 1.01 [1.00,1.01] per year of age), community positive test rate (OR 1.13 [1.12,1.13] per percent increase in positive test rate), hospital inpatients (OR 7.40 [6.54,8.36]), and residents in care homes with non-specialist dementia care (OR 1.42 [1.01,1.99]) had an increased odds of a positive test. Having a positive test prior to the observation period (OR 0.58 [0.49,0.68]) and either one or two doses of a vaccine (0.21 [0.17,0.25] and 0.05 [0.02,0.09] respectively) were associated with a decreased odds of a positive test.

Conclusions: Our findings suggest care providers need to stay vigilant despite the vaccination rollout, and extra precautions should be taken when caring for the most vulnerable. Furthermore,

minimising potential COVID-19 infection for care home residents admitted to hospital should be prioritised.

INTRODUCTION

Care homes are a keystone of adult social care. They provide accommodation and care for those needing substantial help with personal care, but more than that, they are people's homes (1,2). In 2016, there were 11,300 care homes in the UK, with a total of 410,000 residents (3). Within care homes people live in proximity, and may live with frailty and many different health conditions, making them susceptible to outbreaks of infectious disease (1). COVID-19 is described by Lithande et al, as '...a dynamic, specific and real threat to the health and well-being of older people' (2020,p.10) (4). The impacts of COVID-19 on this sub-population have been reported widely in both international and UK media, and in a growing peer reviewed literature.

Vaccinations for SARS-CoV-2 in the UK have been prioritised for those identified at higher risk including to older people living in care homes (5,6). However, very few vaccination trials have recruited older people or older people with frailty (7). Most studies involving care homes have used data sources at an aggregated residential-level, rather than individual-level. This has included the risk of outbreaks of COVID-19 following hospital discharges (8), along with increased risk of SARS-CoV-2 infection due to differing levels of community prevalence (9). Individual-level analyses have focussed on mortality due to COVID-19 in care homes, rather than the risk factors for individuals being infected (10–12).

This is the first study we are aware of investigating this vulnerable sub-population at an individual-level with the inclusion of the community positive test rate of COVID-19, hospital admissions, and vaccination status. Furthermore, we included information on previous positive SARS-CoV-2 polymerase chain reaction (PCR) tests, age, sex, and frailty. As suggested in (13) we were able to do this using up-to-date linked data on care homes from the Secure Anonymised Information Linkage (SAIL) Databank (14–16).

Objectives

We aimed to identify individual level risk factors for SARS-CoV-2 infection with the inclusion of community positive test rate of COVID-19, hospital admissions, and vaccination status.

METHODS

Study design

Longitudinal observational cohort study using anonymised linked data from the Secure Anonymised Information Linkage (SAIL) Databank.

Participants and Setting

Our cohort was 14,786 older care home residents (aged 65+) living in Wales between 1st September 2020 and 1st May 2021. Our dataset consisted of 2,613,341 individual level daily observations for the same period within 697 care homes. Residents were included if they lived in a care home at any period during the observation window. Only residents with at least one PCR test (positive or negative) were included within the dataset. Residents were censored if they moved out of the care home or died. All data were collected retrospectively and linked anonymously within the SAIL Databank.

Data sources

We used linked longitudinal data from the SAIL Databank to create our dataset (14–16). We used the Welsh Demographic Service Dataset (WDS) to determine care home residents and residency dates by linking an anonymised residential linkage field to an anonymised care home registry derived from Care Inspectorate Wales (CIW). The WDS also contains the Lower-layer Super Output Area (LSOA) for each address, which is an area containing approximately 1,500 people, as well as demographic information (age and sex). We used the COVID Vaccine Dataset (CVVD) to identify when individuals had received their vaccinations. The Pathology COVID-19 Daily (PATD) data was used to identify dates of positive and negative SARS-CoV-2 PCR tests. We linked to the Patient Episode Database for Wales (PEDW) to include an indicator for hospital admissions and to calculate the hospital frailty risk score. We used a combination of the Office for National Statistics (ONS) Annual District Death Extract (ADDE), WDS, and Consolidated Death Data Source (CDDS) for mortality information for censoring.

Variables

Outcome - positive SARS-CoV-2 PCR test

Our outcome of interest was a positive SARS-CoV-2 PCR test. The date of the positive test was recorded as the date when the specimen was collected. We used a binary variable to indicate the positive test dates for individuals.

Exposures

Exposure variables were daily COVID-19 community positive test rate estimates and a daily indicator for if a resident was a hospital inpatient. Daily inpatient status was identified using PEDW, residents were identified as being a hospital inpatient for all dates from the admission to the discharge date (inclusive) of a hospital spell.

The estimated COVID-19 community positive test rate was calculated by removing all tests for care home residents and creating a geospatial model for each daily observation. The model includes a spatial correlation term that decays with distance. In other words, areas that are close in proximity are likely to have a more similar estimate than those further away. This reduces the impact of artificial boundaries introduced by statistical geographies and provides a more realistic spatial distribution estimate of positive test rates in the communities surrounding a care home. The community test rate estimates were calculated for each LSOA using a 14-day lookback window for each date. The testing strategy in Wales in the observation period included asymptomatic testing of those caring for vulnerable people (e.g. care home workers and healthcare workers), and symptomatic testing for those in the community (17). Our community positive test rate estimate included both symptomatic and asymptomatic testing. The geospatial model used is an extension of the logistic regression model for binomial (numerator/denominator) data, in which the log-odds of the probability, $P(x)$, of at least one positive PCR test is the unobserved realisation of a spatially correlated stochastic process. Specifically, we used the total number of positive PCR tests in the 14-day window as the numerator, and total tests in the 14-day window as the denominator. The model has three parameters that determine the mean and variance of $P(x)$ and the rate at which the correlation between the values of $P(x)$ at two different locations decays with increasing distance between them, for more detail on the methodology see (18).

Predictors and Confounders

We included the number of doses of a vaccine and a binary indicator for if an individual had a positive PCR test prior to the observation period as predictor variables. The number of vaccine doses received was time varying and was recorded as a categorical variable on each date (0 doses, 1 dose, 2 doses). Additional variables included were specialist services provided by the care home where individuals were resident during the study period. This included: nursing care (yes/no), learning disability (yes/no), mental health (yes/no), dementia (no or unknown, non-specialist, specialist). Age (continuous), sex (male/female) and hospital frailty risk score (no score, low, intermediate, high) were included at the individual level. The Hospital Frailty Risk Score (HFRS) was developed using Hospital Episode Statistics (HES), a database containing details of all admissions, Emergency Department attendances and outpatient appointments at NHS hospitals in England, and validated on over 1-million older people using hospitals in 2014/15 (19). The HFRS uses the International Classification of Disease version 10 (20) (ICD-10) codes to search for specific conditions from secondary care. A weight is then applied to the conditions and a cumulative sum is used to determine a frailty status of: Low, Intermediate or High. We additionally included a HFRS score of 'No score' for people who had not been admitted to hospital in the look back period. We calculated the HFRS using PEDW, the Welsh counterpart to HES, on each daily observation, with a 2-year look back of all hospital admissions recorded in Wales on each date.

Longitudinal dataset design

In the longitudinal dataset created for the analysis individuals have multiple daily observations. Each daily observation includes updated time-dependent covariates; the estimated community COVID-19 positive test rate, whether an individual was in hospital, and the number of vaccine doses received. All other variables were fixed at the value of entry into the study. Additionally, the dataset has anonymous individual and care home identifiers used to cluster the observations. For an example of the dataset design see Figure S1 and Table S1.

Bias

To help minimise selection bias we only included individuals who had at least one SARS-CoV-2 PCR test. Similarly, to minimise time interval bias we used a large observation period where we believe testing of care home residents was more consistent than at the start of the pandemic. We also included anonymous individual and care home level identifiers to account for correlation amongst repeated measures.

Statistical methods

Descriptive statistics included the demographic information for individuals at the start of their residency and stratifications for those who did and did not have a positive PCR test within the observation period. We produced time plots for the daily estimated community positive test rate, positive test rate of residents, numbers of PCR tests for residents, and number of positive PCR tests for residents. For our analysis we used multilevel logistic regression models with a random intercept term for each care home. The regression was applied to the individual-level daily dataset. For sensitivity analysis we calculated null (intercept only) models with random effects at the individual level, care home level, and both (see Table S1). Observations with a missing LSOA were removed, and individuals were censored if they moved residence or died.

Patient and public involvement

No patients or members of the public were involved in setting the research question, study design, outcome measures, or the conduct of the study.

RESULTS

Participants and descriptive data

We analysed 2,613,341 daily observations, consisting of 14,786 care home residents within 697 care homes. The reasons for exclusion from the analysis dataset are included in Figure 1. Descriptive statistics taken from the first observation for each individual are included in

Table 1. We observed a high mean age of 85 (standard deviation 8.2), a large proportion of females (69%), and a high proportion of individuals with frailty (61.8% with low, intermediate, or high HFRS). To provide an indication of demographic differences we also stratified the variables by those who did and did not have a positive SARS-CoV-2 PCR test at any point within the observation period.

Cross tabulations for the observations with and without positive tests with hospital inpatients and the number of vaccine doses is displayed in Table 2. Results indicated a high proportion of observations with a positive PCR test had not been vaccinated (96%), and of those with a positive test who were unvaccinated a significant proportion were hospital inpatients (19%). A significantly larger proportion of observations with a positive PCR test were identified for hospital inpatients (0.93%) compared to not (0.08%).

Figure 2 shows the daily estimated community positive test rate of COVID-19 along with the positive test rate, number of tests taken, and number of positive tests for care home residents. The estimated community positive test rate of COVID-19 was largely correlated with the positive test rate amongst care home residents, with peaks in November and January. There was a large decrease in testing and positive tests amongst care home residents after February.

Figure 1. Flow diagram for the number of observations, individuals, and care homes in the study.

Table 1. Descriptive statistics for care home residents at the start of the observation period (1st September 2020, or first recorded observation).

	All		Negative		Positive	
Individuals (N)	14,786		12,629		2,157	
Individuals with a positive test during the observation period	2,157	14.6%				
Care homes (N)	697		684		289	
Mean age (S.D)	85 (8.2)		85 (8.2)		84.9 (8)	
Gender						
Female	10,190	68.9%	8,720	69.0%	1,470	68.2%
Male	4,596	31.1%	3,909	31.0%	687	31.8%
Hospital Frailty Risk Score						
No score	5,651	38.2%	4,756	37.7%	895	41.5%
Low	1,209	30.4%	1,029	30.9%	180	27.6%
Intermediate	3,430	23.2%	2,944	23.3%	486	22.5%
High	4,496	8.2%	3,900	8.1%	596	8.3%
Specialist services						
Nursing - No	7,132	48.2%	6,057	48.0%	1,075	49.8%
Nursing - Yes	7,654	51.8%	6,572	52.0%	1,082	50.2%
Learning disabilities - No	12,495	84.5%	10,760	85.2%	1,735	80.4%
Learning disabilities - Yes	2,291	15.5%	1,869	14.8%	422	19.6%
Mental Health - No	8,275	56.0%	7,307	57.9%	968	44.9%
Mental Health - Yes	6,511	44.0%	5,322	42.1%	1,189	55.1%
Dementia care - No or Unknown	2,790	18.9%	2,469	19.6%	321	14.9%
Dementia care - Non-specialist	8,708	58.9%	7,349	58.2%	1,359	63.0%
Dementia care - Specialist	3,288	22.2%	2,811	22.3%	477	22.1%

Table 2. Time varying cross tabulation for daily observations dependent on hospital status (inpatient) and number of vaccine doses received (0,1, or 2). The number of positive tests exceeds the total number of individuals testing positive as individuals could test positive on more than one date.

		Observations with a Positive PCR test			
		Yes	N = 2,618	No	N = 2,610,723
Hospital inpatient	Vaccine doses	N	%	N	%
No	0	2,018	77.08%	1,544,048	59.14%
No	1	74	2.83%	581,521	22.27%
No	2	≤6	<0.23%	428,945	16.43%
Yes	0	495	18.91%	39,410	1.51%
Yes	1	24	0.92%	11,665	0.45%
Yes	2	≤6	<0.23%	5,134	0.20%
Total		2,618	100%	2,610,723	100%

Figure 2. Daily estimated community positive test rate of COVID-19 (top left), positive test rate of care home residents (top right), number of tests taken for care home residents (bottom left), and the number of positive tests for care home residents (bottom right). For privacy protection the daily total of care home residents with positive tests has been masked (removed) where there were less than 5 positive tests on a particular date.

Logistic Regression for Positive SARS-CoV-2 PCR tests

The univariable and multivariable multilevel logistic regression results for positive SARS-CoV-2 PCR tests are presented in Table 3. The multivariable model indicated an increase in age (OR 1.01 [1.00,1.01] per year of age), community positive test rate (OR 1.13 [1.12,1.13] per percentage increase in the estimated community positive test rate of COVID-19), hospital inpatients (OR 7.40 [6.54,8.36] compared to residents not in hospital), and residents in care homes with non-specialist dementia care (OR 1.42 [1.01,1.99] compared to those without specialist dementia care) had an increased odds of a positive test. Having a positive test prior to the observation period (OR 0.58 [0.49,0.68]) and either one or two doses of a vaccine (0.21 [0.17,0.25] and 0.05 [0.02,0.09] respectively compared to no vaccine) were associated with a decreased odds of a positive test. The univariable model indicated an increased odds of a positive test was also associated with sex (male compared to female OR 1.20 [1.10,1.31]), care homes with specialist mental health services (1.41 [1.06,1.86]), and increased frailty severity (ORs of 1.27 [1.09,1.47], 1.25 [1.13,1.39], and 1.48 [1.34,1.63] for low, intermediate and high hospital frailty risk scores respectively compared to no score).

The random effects term indicated significant variance at the care home level in the multivariable model (intercept variance of 2.51 with standard error 0.178). We compared intercept only (null) models with random effects terms at the individual, care home, and individual and care home level and found most of the variability was accounted for at the care home level, see Table S2.

Table 3. Univariable and multivariable multilevel logistic regression results for positive SARS-CoV-2 PCR testing amongst care home residents. Results are displayed as Odds Ratios with 95% confidence intervals.

Odds Ratios (95% CI)	Univariable	Multivariable
Fixed effects		
Age (continuous)	1.005 (1.000,1.010)	1.010 (1.004,1.015)
Sex (baseline: female)		
Male	1.197 (1.099,1.305)	1.082 (0.987,1.187)
Predicted COVID-19 positive test rate % (continuous)	1.160 (1.153,1.167)	1.125 (1.118,1.133)
Positive PCR test before September (baseline: no)		
Positive test - yes	0.651 (0.556,0.762)	0.579 (0.490,0.682)
Hospital inpatient (baseline: no)		
Hospital inpatient - yes	9.333 (8.357,10.422)	7.398 (6.546,8.362)
Vaccine doses (baseline: no dose)		
1 dose	0.117 (0.096,0.142)	0.206 (0.170,0.251)
2 doses	0.012 (0.006,0.024)	0.045 (0.023,0.090)
Specialist services (baseline for each category: no)		
Nursing care - yes	1.084 (0.814,1.444)	1.182 (0.885,1.579)
Learning disability - yes	1.218 (0.881,1.683)	1.109 (0.767,1.603)
Mental health - yes	1.406 (1.062,1.861)	1.252 (0.924,1.694)
Dementia - non specialist care	1.469 (1.064,2.028)	1.418 (1.014,1.985)
Dementia - specialist care	1.295 (0.859,1.952)	1.132 (0.739,1.732)
Hospital Frailty Risk Score (baseline: no score)		
Low	1.267 (1.090,1.472)	1.131 (0.969,1.321)
Intermediate	1.254 (1.128,1.394)	1.054 (0.943,1.179)
High	1.481 (1.343,1.633)	1.000 (0.896,1.115)
	<i>Null model</i>	
Intercept	0.001 (0.001,0.001)	0 (0,0)
Random effects		
Variance - intercept	2.670	2.506

Standard error	0.177	0.176
----------------	-------	-------

DISCUSSION

An increase in the community positive test rate of COVID-19 led to an increase in the odds of care home residents testing positive, with an OR of 1.13 (1.12,1.13) per percentage increase of community positive test rate. As the community positive test rate estimates ranged between 0-20% in the observed time period this suggests a potential 10-fold increase in the odds of a positive test in care home residents at the peak of community positive test rate. The association between community positive test rate and care home residents having a COVID-19 infection warrants further research with routes of potential ingress needing further exploration to provide evidence and develop robust policies. Overall this finding highlights the importance of promoting strategies that maintain lower levels of community positive test rate which help reduce the odds of infection amongst vulnerable populations.

We found care home residents who were in hospital had a large increased odds of testing positive for SARS-CoV-2 with an adjusted OR of 7.40 (6.54,8.36). This could be due to the type of hospital care home residents are admitted to, with an estimated 61.9% of COVID-19 infection being hospital acquired in residential community care hospitals, compared to 9.7% in hospitals provided acute and general care (21). Residents entering a hospital environment may be more likely to be exposed to COVID-19 due to an increased contact with healthcare workers and close proximity to other patients in wards (22). We also postulate that the increased odds of testing positive could be associated with an increased probability of being PCR tested whilst in hospital following a lateral flow test in a care home.

Increased age, sex (male), and increased frailty severity were associated with an increased odds of a positive test in the univariable models. These factors are individual level characteristics that would possibly require an increased level of care, and therefore increased daily personal contact, potentially increasing the chance of transmission of COVID-19. In the multivariable model sex and frailty were statistically insignificant, suggesting a reduced importance compared to community positive test rate and hospital stays.

Those testing positive for SARS-CoV-2 prior to the observation period and individuals with one and two doses of a vaccine were at a reduced odds of a positive PCR test, with adjusted odds ratios of 0.58 (0.49,0.68), 0.21 (0.17,0.25) and 0.05 (0.02,0.09) respectively. Those who had previously tested positive may have built up immunity to COVID-19 in a similar way to receiving a vaccine (23). The cross tabulated results and interaction models indicated the odds of a positive test were elevated for those in hospital with a vaccine compared to those not in hospital. This suggests although the vaccine may be effective, there is still an increased risk of a positive test for hospital inpatients.

Strengths

We were able to create a large longitudinal dataset of care home residents and include linked individual-level data on hospital admissions, COVID-19 community positive test rate, vaccinations, and demographic information. This was all possible using existing data from the SAIL Databank.

Limitations

We observed a change in the testing regime from February, where the total number of tests was reduced. We did not explicitly investigate the impact of the change in testing amongst care home residents but found the care home positive test rate remained consistent in shape and magnitude

compared with the estimated community positive test rate. We were unable to link all care homes in Wales in the SAIL databank (91%, 948 of 1048). We were unable to include details on care home staff who may have tested positive for SARS-CoV-2 as data on care home staff is currently cannot be linked to specific care homes.

Conclusion

Our findings indicate that measures taken to prevent COVID-19 from spreading to the most vulnerable in society are not completely effective. Whilst vaccination profoundly decreased the odds of testing positive for SARS-CoV-2, there was still an increased risk of infection for vaccinated individuals admitted to hospital. We also found that an increased community positive test rate of COVID-19 was associated with an increased odds of infection in care home residents. This may reflect higher risk from visitors or care staff living locally. Achieving and maintaining very high rates of vaccination in residents, health and care staff, and visitors is essential. We suggest that care providers need to stay vigilant despite the vaccination rollout, and extra precautions should be taken when caring for the most vulnerable. Follow up research is needed to evaluate the impact of vaccine waning and changing prevalence of viral strains.

ADDITIONAL INFORMATION

Funding

This work was supported by the Medical Research Council [MR/V028367/1]; Health and Care Research Wales [Project: SCF-18-1504]; Health Data Research UK [HDR-9006] which receives its funding from the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation (BHF) and the Wellcome Trust; and Administrative Data Research UK which is funded by the Economic and Social Research Council [grant ES/S007393/1].

Ethics

The data used in this study are available in the SAIL Databank at Swansea University, Swansea, UK. All proposals to use SAIL data are subject to review by an independent Information Governance Review Panel (IGRP). Before any data can be accessed, approval must be given by the IGRP. The IGRP gives careful consideration to each project to ensure proper and appropriate use of SAIL data. When access has been approved, it is gained through a privacy-protecting safe haven and remote access system referred to as the SAIL Gateway. SAIL has established an application process to be followed by anyone who would like to access data via SAIL <https://www.saildatabank.com/application-process>.

Acknowledgements

This work uses data provided by patients and collected by the NHS as part of their care and support. We would also like to acknowledge all data providers who make anonymised data available for research.

We wish to acknowledge the collaborative partnership that enabled acquisition and access to the de-identified data, which led to this output. The collaboration was led by the Swansea University Health Data Research UK team under the direction of the Welsh Government Technical Advisory Cell (TAC) and includes the following groups and organisations: the Secure Anonymised Information Linkage (SAIL) Databank, Administrative Data Research (ADR) Wales, NHS Wales Informatics Service (NWIS), Public Health Wales, NHS Shared Services and the Welsh Ambulance Service Trust (WAST). All research conducted has been completed under the permission and approval of the SAIL independent Information Governance Review Panel (IGRP) project number 0911.

We used the STROBE checklist to create this manuscript, an Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

REFERENCES

1. Challis D, Mozley CG, Sutcliffe C, Bagley H, Price L, Burns A, et al. Dependency in older people recently admitted to care homes. *Age Ageing*. 2000;29(3):255–60.
2. Gordon AL, Goodman C, Achterberg W, Barker RO, Burns E, Hanratty B, et al. Commentary: COVID in Care Homes—Challenges and Dilemmas in Healthcare Delivery. *Age Ageing*. 2020;
3. Competitions and Markets Authority. (2017). Care homes market study. Final report. Competitions and Markets Authority, UK.
4. Lithander FE, Neumann S, Tenison E, Lloyd K, Welsh TJ, Rodrigues JCL, et al. COVID-19 in Older People: A Rapid Clinical Review. *Age Ageing*. 2020;
5. Public Health Wales. Vaccination Priority List - Wales [Internet]. 2020 [cited 2021 Feb 1]. Available from: <https://phw.nhs.wales/topics/immunisation-and-vaccines/covid-19-vaccination-information/eligibility-for-the-vaccine/>
6. Public Health England. Vaccination Priority List - England [Internet]. 2020 [cited 2021 Feb 1]. Available from: <https://www.gov.uk/government/publications/covid-19-vaccination-care-home-and-healthcare-settings-posters/covid-19-vaccination-first-phase-priority-groups>
7. Soiza RL, Scicluna C, Thomson EC. Efficacy and safety of COVID-19 vaccines in older people. *Age Ageing* [Internet]. 2020 Dec 2; Available from: <https://doi.org/10.1093/ageing/afaa274>
8. Emmerson C, Adamson JP, Turner D, Gravenor MB, Salmon J, Cottrell S, et al. Risk factors for outbreaks of COVID-19 in care homes following hospital discharge: A national cohort analysis. *Influenza Other Respi Viruses* [Internet]. 2021 May 1;15(3):371–80. Available from: <https://doi.org/10.1111/irv.12831>
9. Burton JK, McMinn M, Vaughan JE, Fleuriot J, Guthrie B. Care-home outbreaks of COVID-19 in Scotland March to May 2020: National linked data cohort analysis. *Age Ageing* [Internet]. 2021 May 8; Available from: <https://doi.org/10.1093/ageing/afab099>
10. Hollinghurst J, Lyons J, Fry R, Akbari A, Gravenor M, Watkins A, et al. The impact of COVID-19 on adjusted mortality risk in care homes for older adults in Wales, UK: a retrospective population-based cohort study for mortality in 2016–2020. *Age Ageing*. 2020;
11. Burton JK, Bayne G, Evans C, Garbe F, Gorman D, Honhold N, et al. Evolution and effects of COVID-19 outbreaks in care homes: a population analysis in 189 care homes in one geographical region of the UK. *Lancet Heal Longev* [Internet]. 2020;1(1):e21–31. Available from: <https://www.sciencedirect.com/science/article/pii/S266675682030012X>
12. Comas-Herrera A, Zalakain J. Mortality associated with COVID-19 outbreaks in care homes: early international evidence. *Resour to Support community institutional Long-Term Care responses to COVID-19*. 2020;
13. Hanratty B, Burton JK, Goodman C, Gordon AL, Spilsbury K. Covid-19 and lack of linked datasets for care homes. *BMJ* [Internet]. 2020;369. Available from: <https://www.bmj.com/content/369/bmj.m2463>
14. Ford D V, Jones KH, Verplancke J-P, Lyons RA, John G, Brown G, et al. The SAIL Databank: building a national architecture for e-health research and evaluation. *BMC Health Serv Res* [Internet]. 2009;9(1):157. Available from: <http://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-9-157>
15. Lyons RA, Jones KH, John G, Brooks CJ, Verplancke J-P, Ford D V, et al. The SAIL databank: linking multiple health and social care datasets. *BMC Med Inform Decis Mak* [Internet]. 2009;9(1):3. Available from: <http://bmcmmedinformdecismak.biomedcentral.com/articles/10.1186/1472-6947-9-3>
16. Jones KH, Ford D V., Jones C, Dsilva R, Thompson S, Brooks CJ, et al. A case study of the secure anonymous information linkage (SAIL) gateway: A privacy-protecting remote access system for health-related research and evaluation. *J Biomed Inform*. 2014;
17. Welsh Government. COVID-19 testing strategy [Internet]. Testing Strategy for Wales. 2021 [cited 2021 Sep 23]. Available from: <https://gov.wales/covid-19-testing-strategy>
18. Fry R, Hollinghurst J, Stagg HR, Thompson DA, Fronterre C, Orton C, et al. Real-time spatial health surveillance: mapping the UK COVID-19 epidemic. *Int J Med Inform* [Internet]. 2021;104400. Available from: <https://www.sciencedirect.com/science/article/pii/S1386505621000265>
19. Gilbert T, Neuburger J, Kraindler J, Keeble E, Smith P, Ariti C, et al. Development and validation of a Hospital Frailty Risk Score focusing on older people in acute care settings using electronic hospital records: an observational study. *Lancet*. 2018;

20. Organization WH, others. ICD-10: international statistical classification of diseases and related health problems: tenth revision. 2004;
21. Read JM, Green CA, Harrison EM, Docherty AB, Funk S, Harrison J, et al. Hospital-acquired SARS-CoV-2 infection in the UK's first COVID-19 pandemic wave. *Lancet* [Internet]. 2021 Sep 18;398(10305):1037–8. Available from: [https://doi.org/10.1016/S0140-6736\(21\)01786-4](https://doi.org/10.1016/S0140-6736(21)01786-4)
22. Jewkes S V, Zhang Y, Nicholl DJ. Nosocomial spread of COVID-19: lessons learned from an audit on a stroke/neurology ward in a UK district general hospital. *Clin Med* [Internet]. 2020/07/27. 2020 Sep;20(5):e173–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/32719035>
23. Block J. Vaccinating people who have had covid-19: why doesn't natural immunity count in the US? *BMJ* [Internet]. 2021 Sep 13;374:n2101. Available from: <http://www.bmj.com/content/374/bmj.n2101.abstract>