

Supplementary Table 1: Study population sample by age and sex, numbers and sampling fractions from Swedish total population

Age group, years	Men, sampled n	Men, sampled proportion	Women, sampled n	Women, sampled proportion	Total, sampled n	Total, sampled proportion
0-4	25000	0.081	25000	0.086	50000	0.083
5-9	25000	0.078	25000	0.082	50000	0.080
10-14	25000	0.080	25000	0.085	50000	0.082
15-19	25000	0.084	25000	0.092	50000	0.088
20-24	25000	0.080	25000	0.090	50000	0.085
25-29	25000	0.066	25000	0.070	50000	0.068
30-34	25000	0.068	25000	0.072	50000	0.070
35-39	25000	0.075	25000	0.079	50000	0.077
40-44	25000	0.078	25000	0.081	50000	0.079
45-49	25000	0.074	25000	0.076	50000	0.075
50-54	25000	0.073	25000	0.075	50000	0.074
55-59	25000	0.080	25000	0.081	50000	0.081
60-64	25000	0.088	25000	0.088	50000	0.088
65-69	25000	0.094	25000	0.092	50000	0.093
70-74	25000	0.091	25000	0.087	50000	0.089
75-79	25000	0.122	25000	0.112	50000	0.117
80-84	25000	0.206	25000	0.165	50000	0.183
85-89	25000	0.391	25000	0.251	50000	0.306
90-94	24740	1	25000	0.482	49740	0.649
95+	5580	1	17403	1	22983	1
	480320		492403		972723	

Supplementary Table 2. Definitions of adverse events of interest (AESIs)

Outcome	ICD-10 codes
Aseptic meningitis	A87
	B261
	G030
Febrile seizure	R560
Kawasaki's syndrome	M303
Multisystem inflammatory syndrome in children (MISC)	U109
Postinfectious arthritis	M015
	M018
	M02
	M03
Arthritis, broad (all types)	M015
	M018
	M02
	M03
	M05
	M06
	M07
	M08
	M09
M13	
Myocarditis	I012
	I090
	I40
	I41
	I514
Acute respiratory distress syndrome (ARDS)	J80
	P220
Myocardial infarction, broad	I21
	I22
	I23

Stroke	F010
	F011
	F013
	I63
	P101
	P524
	P526
	I60
	I61
	I62
	I64
Ischemic stroke	F010
	F011
	F013
	I63
Haemorrhagic stroke	P101
	P524
	P526
	I60
	I61
	I62
Venous thromboembolism	I822
	I823
	I828
	I829
Pulmonary embolism	I26
Acute kidney injury	N17
	O904
Acute Liver failure	K720
	K729
Erythema multiforme	L51
Disseminated intravascular coagulation	D65
	P60
Autoimmune thyroiditis	E063
Appendicitis	K35
	K36
	K37

