1   **Is Human Bocavirus Infection Associated with Gastroenteritis in Children? An** 2   **Updated Systematic Review and Meta-analysis**

3   Minyi Zhang¹, Minyi Liang¹, Qiushuang Li¹, Juxian Xian¹, Fei Wu¹, Liting Zheng¹,
4   Qing Chen^{1*}

5   ¹ Guangdong Provincial Key Laboratory of Tropical Disease Research, Department of
6   Epidemiology, School of Public Health, Southern Medical University, Guangzhou
7   510515, China

8   ***Corresponding authors**

9   Qing Chen, Guangdong Provincial Key Laboratory of Tropical Disease Research,
10   Department of Epidemiology, School of Public Health, Southern Medical University,
11   Guangzhou 510515, China.
12   Telephone and fax: +86-020-61648312; email: qch.2009@163.com

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background: Human bocavirus (HBoV) figures as an increased risk factor of respiratory and gastrointestinal tract infections among children. A great deal of data is available to support the pathogenic role of HBoV in acute respiratory diseases. However, the association between HBoV infection and gastroenteritis remains controversial due to the ambiguous results. The present work aims to clarify the role of HBoV as a cause of gastroenteritis in children.

Methodology/Principal findings: A systematic search of the literature was carried out from 1 January 2016 to 29 August 2021 in Embase, PubMed, Scopus, Web of Science, and the Chinese bibliographic database of biomedicine (CBM). Data from included studies were analyzed by use of a random-effects model. The pooled estimates of HBoV prevalence among all cases of gastroenteritis were generated and stratified by potential confounders. The pooled odds ratio (OR) and 95% confidence interval (CI) were computed for HBoV infection in relation to the risk of gastroenteritis. The overall prevalence of HBoV in children with gastroenteritis (9.1%, 95% CI: 6.7-11.8%) was considerably higher than that detected in children without gastroenteritis (4.0%, 95% CI: 1.1-8.5%). HBoV prevalence tended to be higher in cases of gastroenteritis under five years of age (12.1%, 95% CI: 6.8-18.7%). The highest frequency of HBoV was found in Egypt (57.8%, 95% CI: 47.7-67.6%). The predominant genotypes of HBoV circulating in children with gastroenteritis were genotype 1 (HBoV1, 3.8%, 95% CI: 2.7-5.2%) and genotype 2 (HBoV2, 2.4%, 95% CI: 1.3-3.7%). HBoV infection was significantly associated with an increased risk of gastroenteritis in children (OR 1.620,

36 95% CI: 1.023-2.566).

37 **Conclusion:** The HBoV prevalence in pediatric cases of gastroenteritis is higher than
38 that in children without gastroenteritis, demonstrating an increasing global burden of
39 gastroenteritis in children caused by HBoV infection. Targeted intervention to reduce
40 the HBoV burden should be established.

Author summary

In addition to the known gastroenteritis-associated pathogens (rotavirus and norovirus et al.), several novel viruses that might be caused agents of diarrhea have been gradually determined, such as HBoV. The impact of HBoV infection on the development of childhood gastroenteritis remains ambiguous and in need of verifications. This work clarifies the association between HBoV infection and the risk of gastroenteritis in children based on the review literature. The estimated HBoV prevalence in all cases of gastroenteritis was categorized into different subgroups following the potential confounders, including age, gender, country, and HBoV genotypes. The main finding of this work was the significant association between HBoV infection and the development of childhood gastroenteritis. Our systematic review highlighted that HBoV infection contributes to the increased global burden of gastroenteritis in children.

Introduction

Childhood Diarrheal disease is a leading cause of malnutrition and death in children younger than five years, which accounted for roughly 1.7 billion novel cases and greater than 500 thousand deaths each year throughout the world (1). Infection diarrhea is one of the most frequent clinical manifestations of gastrointestinal tract infection resulting from either bacteria, viruses, parasites, or fungi (2). As being preventable, fully understanding its causative agents would be of vital benefit from substantially decreasing the burden of diarrheal disease in children. Of these, viruses turn out to be the primary causative agents responsible for around 70% of gastroenteritis cases in children (3, 4). For decades, rotavirus was the primary pathogen of gastroenteritis-related hospitalizations and other severe diseases in the first five years of life (5). However, the recent wide application of rotavirus vaccines is an essential contributor to the marked decrease in rotavirus-related illnesses (6, 7). Since then, other novel detected viral pathogens have become the potential risk factors of childhood gastroenteritis, such as human bocavirus (HBoV).

HBoV consists of ~5kb linear and small non-enveloped single-stranded DNA genome organizing in three open reading frames that encode two forms of non-structural protein 1 (NS1), a nuclear phosphoprotein (NP1), and two viral capsid proteins (VP1 and VP2) (8-10). Following the genomic analysis, HBoV belongs to the genus *Bocavirus* within the family *Parvoviridae* and the subfamily *Parvovirinae*, with four currently identified genotypes. HBoV genotype 1 (HBoV1) was firstly discovered in nasopharyngeal aspirates from children with respiratory tract infections (5). With its

77 initial discovery in 2005 by Tobias Allander and colleagues (5), numerous studies
 78 regarding the epidemiology, pathogenesis, diagnosis, and clinical features of this virus
 79 have been carried out in many countries. Subsequently, three additional genotypes were
 80 detected in feces and named HBoV2, 3, and 4 (11, 12). A great deal of data is available
 81 to support the association between HBoV1 infection and acute respiratory diseases,
 82 such as asthma, pneumonia, bronchiolitis, and acute wheezing (13-17). On the contrary,
 83 the remaining genotypes are primarily identified in human specimens. Previous studies
 84 have suggested a comparable prevalence of HBoV in patients with gastroenteritis
 85 compared to healthy or non-diarrheal individuals (11, 18). For instance, a prior study
 86 involving 632 children with gastroenteritis and 162 healthy controls has shown a
 87 statistically significant association between HBoV infection and childhood
 88 gastroenteritis (18). It should be noted that, in the same study, the majority of HBoV
 89 infections were found in children less than 12 months of age (18). Most recently,
 90 Neveen et al. concluded that HBoV could be the significant etiologic agent of
 91 gastroenteritis due to the presence of HBoV in some children suffering from diarrheal
 92 disease without any association of any other etiological agents (19). However, a
 93 controversial issue that arises in this domain is the actual pathogen role of HBoV in
 94 gastrointestinal symptoms due to the growing appears of HBoV detection in
 95 asymptomatic patients (20, 21).

96 A systematic review and meta-analysis have been published to assess whether HBoV
 97 infection is a risk factor for gastroenteritis in children under five years (22). This review
 98 included 36 unique studies from 18 countries between Oct. 2005 and Oct. 2016 and

estimated that 6.90% of all cases of gastroenteritis in children younger than five years were correlated with HBoV infection. However, this review failed to identify a statistically significant association of HBoV infection with the risk of gastroenteritis. Also, the prevalence and distribution of HBoV3 and HBoV4 are needed to be established due to the lack of pooled estimations.

With the increasing use of molecular diagnostics and much attention of scientists, the data on different genotypes of HBoV associated with gastroenteritis in children has increased in the last five years and accordingly prompted us to update previous estimates of HBoV prevalence and genotype distribution in pediatric cases of gastroenteritis up to 29 August 2021. Additionally, the second aim of this work was to determine the association between HBoV infection and gastroenteritis in children through a meta-analytic method.

Methods

Data sources

Following the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) statement (23), a comprehensive literature search was conducted in Embase, PubMed, Scopus, Web of Science, and CBM (Chinese bibliographic database of biomedicine) databases for identifying eligible records from 1 January 2016 to 29 August 2021. The Medical Subject Heading (MeSH) terms “human bocavirus”, “gastroenteritis”, “diarrhea”, “child”, and related terms were used in all possible combinations. More detailed search strategies in each database are showed in S1 Table. Since fewer studies in a case-control design were identified during the present time

horizon of the literature search, we additionally included the case-control studies from a previous systematic by Ri De and colleagues (22) and extracted data for other variables as required.

Study selection

Studies that met the following criteria were deemed eligible for inclusion: 1) enrolled pediatric patients with gastroenteritis or healthy children; 2) applied PCR-based diagnostics for HBoV on fecal samples; 3) original cross-sectional or case-control documents of HBoV infection. The exclusion criteria were the following: 1) duplicate records; 2) HBoV prevalence in adults; 3) samples other than stools; 4) HBoV infection employing antigen assays and studies that reported seroprevalence of HBoV antibodies; 5) lack of detailed data; 6) animal studies, conference abstracts, reviews, letters, case reports, or obviously irrelevant.

Data extraction

Two researchers independently reviewed and selected all potentially relevant articles following the inclusion and exclusion criteria for further assessment. Suppose any conflicts exist between the investigators, a third evaluator will make a final decision. The following information was extracted from the included publications by two researchers: first author, year of publication, country, study period, study design, used diagnostic method, age range or mean, gender, sample size, number of total cases, number of HBoV-positive cases, number of total controls, number of HBoV-positive controls, and genotypes of HBoV.

Assessment of Study Quality

Two investigators independently assessed the quality of each retrieved study in a case-control design through the Newcastle-Ottawa Scale (NOS) (24). A study was awarded a maximum of one star for each numbered item within the selection and exposure sections. A maximum of two stars was awarded regarding the comparability category. Studies that received seven or more out of a maximum of 9 stars were judged as being of high quality. Meanwhile, medium-quality and poor-quality studies were considered those with four to six stars and those with fewer than four stars, respectively.

Statistical analysis

Using the HBoV prevalence as the primary outcome, we firstly generated pooled estimated prevalence of HBoV infection in all pediatric patients with gastroenteritis. Further, a meta-analysis of risk effect estimates was carried out regarding the HBoV exposure and gastroenteritis in children by the DerSimonian and Laird random-effects model (25). Pooled estimates of odds ratios (ORs) and 95% confidence interval (CIs) were computed. The I^2 statistic was utilized to measure the proportion of the total variability explained by heterogeneity across studies. I^2 values $< 25\%$ indicate a low level of heterogeneity, whereas I^2 values $\geq 75\%$ is regarded as a high degree of heterogeneity (26). Cochrane's Q statistic was applied to examine the statistical significance of the heterogeneity. We stratified the pediatric patients based on age, gender, country, and HBoV genotype and subsequently calculated the summary risk of gastroenteritis for HBoV infection. A Pearson's chi-square test was employed to identify the significant difference between the subgroups. Additionally, we addressed whether the distribution of HBoV infection varied between pediatric cases of

gastroenteritis and healthy children.

Publication bias was assessed by combining the funnel plot, Begg's adjusted rank correlation test, and Egger's regression asymmetry test (27, 28). All statistical analyses were conducted in Stata 16. A probability level less than 0.05 was considered statistically significant except the heterogeneity test set at less than 0.1, and all statistical tests were two-sided.

Results

Literature search

Based on the search strategy, seven hundred and seventy-three relevant articles were initially retrieved from Embase, PubMed, Scopus, Web of Science, and CBM, of which 266 were finally identified between 2016 and 2021. One hundred and twenty-seven documents were subsequently removed due to duplication, and a total of 222 records were excluded after manual screening of titles and abstracts. The remaining 44 full-text articles were further obtained for eligibility, of which 29 unique studies were included after applying the inclusion and exclusion criteria in this systematic review and meta-analysis. Case-control studies originally identified by Ri De and colleagues were appended to our search. However, two of the ten studies were excluded due to reporting HBoV prevalence either in serum samples or in adults. Also, one study not published in original research was removed, leaving seven additional studies in a case-control design. Consequently, the present updated review consisted of 36 studies. A PRISMA-based flowchart of the literature search and study selection is depicted in Fig 1.

Fig 1. Flowchart of the literature search and study selection

Study characteristics

The final dataset included data on children from 20 countries (Russia, United States, India, China, Brazil, Albania, Korea, Japan, Sudan, Gabon, Latvia, Iraq, Iran, South Africa, Bangladesh, Australia, Finland, United Kingdom, Thailand, and Egypt). Of the

192 36 included documents, 11 were case-control (11, 18, 29-37) and 25 were cross-
193 sectional in study design (20, 38-61). Amidst the cross-sectional studies, 23 were case-
194 only, and the remaining 2 presented data for healthy individuals. The largest study
195 investigated 5250 pediatric patients with gastroenteritis in Russia, and the smallest one
196 comprised 83 cases. This review consisted of 15 and 29 studies reporting gender and
197 genotype distribution of HBoV infection, respectively. The majority of these data
198 represented children less than five years of age (n=20). Most studies were carried out
199 in China (n=8), followed by Brazil (n=4) and Bangladesh (n=3). The characteristics of
200 cross-sectional and case-control studies are summarized in Table 1 and 2.

201 **Table 1 Summary of characteristics of cross-sectional studies**

First author	Year	Country	Study period	Age range/ mean	Number of HBoV positive	Total
With gastroenteritis						
Gubio Soares Campos	2016	Brazil	Jan. 2012-July 2012	≤ 5 years	44	105
G. La Rosa	2016	Albania	Oct. 2013-Jan. 2015	2 months-7 years	13	142
Eun Jin Lee	2016	Korea	Feb. 2015-May 2015	≤ 6 years	10	155
Min Zhao	2016	China	Oct. 2010-Dec. 2012	NA	25	1121
Neha Lasure (1)	2017	India	Jan. 2009-Dec. 2011	≤ 5 years	24	418
Naoki Shigemoto	2017	Japan	Jan. 2006-Dec. 2013	0-14 years	10	312
Tao Zhou	2017	China	Jan. 2012-Dec. 2013	≤ 6 years	60	346
Mosab A. Adam	2018	Sudan	April-Dec. 2014	≤ 5 years	5	437
Sonia Etenna Lekana-Douki	2018	Gabon	March 2010-June 2011	≤ 5 years	7	317
Zaiga Nora-Krukle	2018	Latvia	Nov. 2013-April 2017	≤ 5 years	21	83
Luana S. Soares	2019	Brazil	Nov. 2011-Nov. 2012	≤ 10 years	54	225
Ghasaq Mundher Ali	2020	Iraq	July 2019-March 2020	≤ 5 years	9	100
Gyu Ri Kim	2020	Korea	July 2016-July 2017	< 5 years	2	740
Marco Yung-Cheng Lin	2020	China	Sep. 2013-June 2015	NA	15	176
Fábio Correia Malta	2020	Brazil	Jan. 2016-Dec. 2017	≤ 2 years	110	886
Mehrdad Mohammadi	2020	Iran	2017-2018	≤ 3 years	72	500
Rembuluwani Netshikweta	2020	South Africa	April 2009-April 2015	≤ 5 years	212	3765

Mpumelelo Casper Rikhotso	2020	South Africa	June 2017-July 2018	≤ 5 years	8	141
Nadim Sharif (1)	2020	Bangladesh	Jan. 2014-Jan. 2019	≤ 15 years	19	387
Nadim Sharif (2)	2020	Bangladesh	Jan. 2015-Jan. 2019	≤ 15 years	21	290
Nattika Nantachit	2021	Tailand	Jan. 2012-Dec. 2018	≤ 5 years	141	2727
Neveen M. Rizk	2021	Egypt	NA	≤ 5 years	59	102
JI Yan-li	2017	China	Jan.-Dec. 2016	< 60 months	26	354
Without gastroenteritis						
Teresinha Teixeira de Sousa	2017	Brazil	2014-2015	≤ 5 years	1	119
Shoko Okitsu	2020	Bangladesh	June-Oct. 2016	73 ± 4 months	1	227

202 NA, Not available

204 **Table 2 Summary of case-control studies**

First author	Year	Country	Study period	Sample size	Age range/ mean	Number of HBoV positive	Total	Age range/ mean	Number of HBoV positive	Total
Alexander Tymentsev	2016	Russia	Feb 2010-Dec 2012	5502	≤ 5 years	62	5250	≤ 5 years	1	252
Rimma Melamed	2017	Unite States	Nov 2003-Nov 2005	356	44.8 (11-60) months	22	178	43.8 (11-59) months	14	178
Neha Lasure (2)	2019	India	2007-2011	985	≤ 5 years	46	778	≤ 5 years	0	207
Ri De	2018	China	May-Dec 2016	480	≤ 5 years	18	240	≤ 5 years	12	240
Jane L. Arthur ^a	2009	Australia	Jan-Dec 2001	372	≤ 18 years	54	186	≤ 18 years	30	186
Minna Risku ^a	2012	Finland	Aug 2006- Aug 2008	990	≤ 15 years	85	878	≤ 15 years	6	112
Sameena Nawaz ^a	2012	United Kingdom	1993-1996	1772	≤ 19 years	106	851	≤ 19 years	163	921
Thaweesak Chieochansin ^a	2008	Thailand	Nov 2005- Sep 2007	427	4 months-4 years	2	225	2 months-5 years	0	202
Wei-xia Cheng ^a	2008	China	July 2006-Sep 2007	512	≤ 5 years	14	397	≤ 5 years	4	115
Yu Jin ^a	2011	China	July 2006- June 2008	794	≤ 5 years	162	632	≤ 5 years	24	162
Zeng Mei ^a	2010	China	Oct-Dec 2008	420	≤ 5 years	6	220	≤ 5 years	4	200

205 ^aStudies originally from Ri De and colleagues.

HBoV prevalence in children with gastroenteritis

The overall pooled prevalence of HBoV infection among 23,664 pediatric cases of gastroenteritis was estimated at 9.1% (95% CI: 6.7-11.8%; $I^2 = 97.7\%$, $P < 0.001$ for heterogeneity test) (Fig 2). Subgroup analysis demonstrated the prevalence of HBoV exposure significantly differed across age category, country, and genotype (Table 3). By age, HBoV prevalence was significantly higher in cases of gastroenteritis younger than 5 years (12.1%, 95% CI: 6.8-18.7%), and the lowest prevalence was found in children between 5 and 10 years (3.4%, 95% CI: 0.2-8.6) ($P < 0.001$). By country, the highest prevalence of HBoV infection occurred in Egypt at 57.8% (95% CI: 47.7-67.6%), followed by Australia (29.0%, 95% CI: 22.6-36.1%) and Latvia (25.3%, 95% CI: 16.4-36.0%), and the lowest one was observed in Korea (0.7%, 95% CI: 0.2-1.4%) ($P < 0.001$). HBoV prevalence was slightly greater in male patients (9.4%, 95% CI: 5.8-13.8%) compared with female patients (8.7%, 95% CI: 5.7-12.2%) ($P = 0.785$). The estimated pooled HBoV prevalence did not markedly change when meta-analysis was limited to the studies that lasted at least 1 year (8.1%, 95% CI: 5.7-10.8, $P = 0.221$, $I^2 = 97.7\%$, $P < 0.001$ for heterogeneity test, $n=27$) (S1 Fig).

Whether the distribution of HBoV infection differed across pediatric cases of gastroenteritis and healthy children was further explored. In the 13 studies that included 3121 children without gastroenteritis, the pooled HBoV prevalence was detected in 4.0% (95% CI: 1.1-8.5%) (S2 Fig), which was significantly lower than that in cases of gastroenteritis ($P < 0.001$).

Fig 2. Overall pooled prevalence of HBoV infection in children with gastroenteritis based on case-control and cross-sectional studies. HBoV, human bocavirus

231 **Table 3** Estimated HBoV prevalence in children with gastroenteritis by age, gender,
232 country, and genotype.

Category	Subgroup	Number of studies	Pooled prevalence (% , 95% CI)	Heterogeneity test (I ² %, <i>P</i> value)	Chi-square test (<i>P</i> value)
Age	≤ 1 year	8	7.3 (3.6-12.2)	97.142%, < 0.001	<i>P</i> < 0.001
	> 1-5 years	10	5.2 (1.8-9.9)	96.373%, < 0.001	
	≤ 5 years	13	12.1 (6.8-18.7)	97.921%, < 0.001	
	> 5-10 years	2	3.4 (0.2-8.6)	< 0.001, < 0.001	
	Mixed	12	8.3 (6.2-10.7)	96.949%, < 0.001	
Genger	Male	14	9.4 (5.8-13.8)	95.789%, < 0.001	<i>P</i> = 0.785
	Female	14	8.7 (5.7-12.2)	92.220%, < 0.001	
Country	Russia	1	1.2 (0.9-1.5)	NA	<i>P</i> < 0.001
	Unite States	1	12.4 (7.9-18.1)	NA	
	India	2	5.8 (4.6-7.3)	< 0.001, <0.001	
	China	8	8.2 (3.1-15.2)	97.572%, <0.001	
	Brazil	3	24.7 (10.8-42.0)	< 0.001, <0.001	
	Albania	1	9.2 (5.0-15.1)	NA	
	Korea	2	0.7 (0.2-1.4)	< 0.001, <0.001	
	Japan	1	3.2 (1.5-5.8)	NA	
	Sudan	1	1.1 (0.4-2.6)	NA	
	Gabon	1	2.2 (0.9-4.5)	NA	
	Latvia	1	25.3 (16.4-36.0)	NA	
	Iraq	1	9.0 (4.2-16.4)	NA	
	Iran	1	14.4 (11.4-17.8)	NA	
	South Africa	2	5.5 (4.8-6.2)	< 0.001, <0.001	
	Bangladesh	2	5.9 (4.2-7.8)	< 0.001, <0.001	
	Australia	1	29.0 (22.6-36.1)	NA	
	Finland	1	9.7 (7.8-11.8)	NA	
	United Kingdom	1	12.5 (10.3-14.9)	NA	
	Tailand	2	4.7 (3.9-5.5)	< 0.001, <0.001	
	Egypt	1	57.8 (47.7-67.6)	NA	
Genotype	HBoV1	28	3.8 (2.7-5.2)	94.878%, < 0.001	<i>P</i> < 0.001
	HBoV2	24	2.4 (1.3-3.7)	96.249%, < 0.001	
	HBoV3	21	0.9 (0.4-1.6)	93.467%, < 0.001	
	HBoV4	16	0	14.799, 0.284	

Prevalence and distribution of HBoV genotypes

All four genotypes of HBoV were identified among pediatric cases of gastroenteritis (Table 3). The predominant genotype of HBoV circulating in children with gastroenteritis belonged to genotype 1 (HBoV1, 3.8%, 95% CI: 2.7-5.2%) and genotype 2 (HBoV2, 2.4%, 95% CI: 1.3-3.7%). Sixteen countries had reported the genotype distribution of HBoV. At the global level, HBoV1 dominated (50.5%), followed by HBoV2 (36.6%) and HBoV3 (12.0%). The estimated genotype distribution by country is depicted in Fig 3.

Fig 3. Genotype distribution by country

Association between HBoV infection and the risk of gastroenteritis in children

A quality assessment was performed for each study with a case-control design utilizing the NOS recommended by the Cochrane Collaboration (62). All eleven included case-control studies met the criteria for NOS and none of the poor quality (S2 Table). We

conducted the second analysis based on case-control studies to investigate the relationship between HBoV infection and the risk of gastroenteritis in children, with the pooled OR value of 1.620 (95% CI: 1.023-2.566; $I^2 = 73.6\%$, $P < 0.001$ for heterogeneity test) being reported (Fig 4). In regard to HBoV genotypes, only HBoV2 showed a statistically significant correlation with gastroenteritis in children (OR 1.75, 95% CI: 1.30-2.35). Among different countries, the significant association of HBoV infection with gastroenteritis was observed in India, China, and Australia. Detailed associations between HBoV infection and gastroenteritis for subgroups were displayed in Fig 5. Using a visual inspection of the funnel plot and statistical tests, no evidence of publication bias was recognized ($P = 0.876$ for Begg's test and $P = 0.145$ for Egger's test) (Fig 6).

Fig 4. Forest plot of the association between HBoV infection and the risk of

263 **gastroenteritis in children.** HBoV, human bocavirus; OR, odds ratio; CI, confidence
264 interval

265 Odds ratio

266 **Fig 5. Forest plot of the association between HBoV infection and the risk of**
267 **gastroenteritis in children by genotype and country.** HBoV, human bocavirus; OR,
268 odds ratio; CI, confidence interval

Fig 6. Funnel plot for assessment of publication bias

Discussion

HBoV has been considered a major pathogen that contributed to respiratory and gastrointestinal infections in particular younger children (63, 64). Following its first identification in Sweden, HBoV has been frequently detected in patients with various diseases throughout the Europe (65-67), North (68, 69) and South America (70, 71), Australia (72, 73), Asia (74, 75), and Africa (76, 77). In addition to respiratory and feces specimens, HBoV has also been identified in blood (78), urine (79), saliva (80), cerebrospinal fluid (81), and even environmental samples (82, 83). The current evidence clearly illustrates the potential risk of HBoV exposure for human health. In this review and meta-analysis, we summarized the results of 36 studies, including a total of 23,664 pediatric cases of gastroenteritis. The updated overall estimate suggested

that HBoV infection is associated with 9.1% of all gastroenteritis cases in children, which confirms and extends the earlier results of a systematic review and meta-analysis of HBoV conducted by Ri De (22). This growing estimate emphasized that HBoV infection contributes to the increased global burden of gastroenteritis in children.

In comparison to the previous meta-analysis (22) that only involved children aged under five years, the age distribution of HBoV-associated gastroenteritis in children younger than 18 years was available in the present work. These data revealed that children under five years of age were the most susceptible to develop gastroenteritis when infected with HBoV, with the highest prevalence of HBoV-associated gastroenteritis cases (12.1%). From the results of more specific age groups, the relatively high frequencies of HBoV were also found in the age groups less than one year old (7.3%) and between 1 and 5 years old (5.2%). These findings tie well with previous reports wherein younger children might undergo a higher occurrence of HBoV-associated gastroenteritis. For example, a recent large-scale survey conducted in South Africa with 3765 fecal samples collected from hospitalized children for gastroenteritis illustrated that more than 90% of HBoV-associated cases were children younger than two years (84). A case-control study in the UK concluded that although HBoV is relatively common across all age groups, the risk of HBoV-associated gastroenteritis derived from preschool-age children was expected to be higher compared with adults (37). Furthermore, Mehrdad et al. reported that the significant age category in HBoV-positive samples was between 1 and 2 years old and have concluded that the prevalence of gastrointestinal tract infection with HBoV occurs

precisely in the age range of digestive illnesses resulted from rotavirus (85). Our findings also matched several reviews focused on global norovirus (NoV) prevalence, demonstrating that most pediatric NoV-associated gastroenteritis occurred in children under two years of age (86, 87). Taken together, childhood gastroenteritis caused by HBoV and other known gastroenteritis-related viruses occurs precisely within the same age range. Thus, HBoV is much likely to figure as a potential etiological agent of gastroenteritis.

Comparisons across different countries showed that the highest HBoV prevalence in children with gastroenteritis was observed in Egypt (57.8%), followed by Australia (29.0%), Latvia (25.3%), and Brazil (24.7%). The lowest HBoV prevalence occurred in pediatric patients from Korea (0.7%). In line with the previous demonstration, these results confirmed that HBoV had been identified worldwide without any regional or geographic restrictions (63). However, these findings go beyond previous reports, which indicated a trend of decreasing prevalence from the developing countries to the developed countries (88, 89). For illustration, a case-control assessment conducted in the United States revealed that HBoV infection was related to 12.4% of all cases of acute childhood diarrhea (29). Nonetheless, the pooled HBoV prevalence was estimated at 5.8% among 1196 pediatric cases in India. It remains unclear to what complicated reasons attributed to the wide variation of HBoV prevalence across the world in addition to ethnicity, nutritional status, hygiene conditions, healthcare settings, and the various sample size of the study. More comprehensive studies should take into account the concrete reasons for HBoV prevalence in multiple geographies.

Consistent with previous evidence indicating less susceptibility to viral infections in females (90), our data supported that males are more susceptible than females to develop HBoV-associated gastroenteritis, with 9.4% and 8.7% of HBoV prevalence in male patients and female patients, respectively. However, no significant differences were observed in HBoV prevalence based on gender. Further research is needed to address this information lacuna.

Four HBoV genotypes have been described based on the genomic analyses of VP1 and VP2 sequences. Globally, HBoV1 accounted for more than half of all infections in children with gastroenteritis, which also dominated in most countries. It was also found that only one genotype was presented in Korea (HBoV1), Gabon (HBoV1), and Bangladesh (HBoV2). The estimated genotype distribution by country displayed considerable regional variations. As already mentioned, HBoV1 is commonly prevalent in clinical specimens of the respiratory tract, whereas HBoV2-4 are more frequently detected in patients with gastrointestinal infections and symptoms (64). Additionally, a recent relevant study in Taiwan demonstrated that HBoV1 and HBoV2 are generally the primary species in childhood gastrointestinal infections worldwide (43). The prevalence of HBoV genotypes has been investigated in children with gastroenteritis in the previous meta-analysis (22). The researchers showed that HBoV1 and HBoV2 were detected in 3.49% and 8.59% of pediatric cases with gastroenteritis. These previous results from Ri De et al. seem to be inappropriate for comparison because of which estimated data based on only ten case-control studies. In our updated review, HBoV1 (3.8%) figured as the predominant genotype circulating in children with gastroenteritis,

and HBoV2, with a prevalence of 2.4% in the next rank. Even though HBoV1 can also be found in fecal samples from pediatric patients suffering from gastroenteritis, this novel finding suggested the prevalent HBoV variant changed related to childhood gastroenteritis during the past years. Likewise, during a 3-year observation in Russia, all four HBoV genotypes with differing detection rates have been identified in children hospitalized with diarrhea (91). Hence, we believe that continuous surveillance of this virus would be vital in pointing out how the prevalent HBoV genetic variants change with time. With regard to other genotypes, we identified the relatively low prevalence of HBoV3 in children with gastroenteritis (0.9%), while HBoV4 was only determined in four included studies. Since the lower prevalence and frequency of the other genotypes, it remains to be resolved whether other genotypes display a pathogenic role in childhood gastroenteritis.

Using data from 11 case-control studies, a meta-analysis provided new evidence to construct pooled risk effects of HBoV-associated gastroenteritis. It was found that children infected with HBoV have approximately 1.6 times increased chance of developing gastrointestinal symptoms compared with uninfected controls, indicating that HBoV exposure is significantly associated with a higher risk of gastroenteritis in children (OR 1.620, 95% CI: 1.023-2.566). Additionally, the presence specifically of all four HBoV genotypes was examined for the purpose of determining any possible associations between specific HBoV genotypes and gastroenteritis in children. The analytical results indicated that HBoV2 was significantly correlated with gastroenteritis in children (OR 1.746, 95% CI: 1.296-2.351), whereas other genotypes have not yet

disease association. This supports Arthur and coworkers who have proven the association between HBoV2 infection and gastroenteritis using multivariate conditional logistic regression analysis (11). However, the pathogenic role of HBoV2 has been challenged by Ong et al. (92), demonstrating that HBoV in stool is probably an innocent bystander rather than a true pathogen due to high coinfection with other pathogens. It has also been questioned owing to the fact that HBoV was frequently detected in healthy or diarrhea-free individuals (17). In this updated review, the pooled prevalence of HBoV infection was estimated at 4.0% in children without gastroenteritis, which significantly lower than that in pediatric patients ($P < 0.001$). The most likely explanation for the presence of HBoV in feces from asymptomatic individuals can be linked to the long-term viral shedding following symptomatic infection (93, 94). This regard does not repudiate the role of HBoV as the etiological agent of gastrointestinal symptoms in children. Furthermore, the significantly positive association of HBoV infection with childhood gastroenteritis were observed in India (OR 26.34, 95% CI: 1.62-429.32), China (OR 1.70, 95% CI: 1.18-2.43), and Australia (OR 2.13, 95% CI: 1.29-3.52), while it was shown inverse association in the UK (OR 0.66, 95% CI: 0.51-0.86). The findings varied across countries might be due to the regional differences in viral epidemiology, geographic location, and seasonality (18, 95, 96).

Several limitations of the previous study (22) were addressed in the present work. However, it remains some limitations such as the lack of a large number of eligible research data and not considering other factors that might affect the results. Furthermore, the present work only included children diagnosed with gastroenteritis or diarrhea as

cases, making it probably underestimate the burden of HBoV infection on the development of childhood gastroenteritis.

Conclusion

In summary, the pooled prevalence of HBoV infection in children suffering from gastroenteritis was estimated at 9.1% between 2016 and 2021, showing a higher global burden of gastroenteritis in children caused by HBoV than in the last decade. The considerably higher HBoV prevalence was found in pediatric patients less than five years of age and in Egypt. Notably, the correlation of HBoV infection with gastroenteritis in children was established in this updated review. Our findings also add to the growing evidence that HBoV2 would be a causative agent of gastrointestinal symptoms. However, this conclusion should be treated with caution as the known gastroenteritis-associated viruses are not excluded. The actual pathogenic role of HBoV in leading to childhood gastroenteritis were needed to be determined by large-scale continuous surveillance, serological investigation, and appropriate animal models.

References

1. World Health Organization. Diarrhoeal Disease. 2020 [Available from: <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease>.
2. Harb A, Abraham S, Rusdi B, Laird T, O'Dea M, Habib I. Molecular Detection and Epidemiological Features of Selected Bacterial, Viral, and Parasitic Enteropathogens in Stool Specimens from Children with Acute Diarrhea in Thi-Qar Governorate, Iraq. *Int J Environ Res Public Health*. 2019;16(9).
3. Reymão TK, Hernandez JD, Costa ST, Sousa MS, Oliveira DS, Silva LD, et al. Sapoviruses in children with acute gastroenteritis from Manaus, Amazon region, Brazil, 2010-2011. *Rev Inst Med Trop Sao Paulo*. 2016;58:81.
4. Chow CM, Leung AK, Hon KL. Acute gastroenteritis: from guidelines to real life. *Clin Exp Gastroenterol*. 2010;3:97-112.
5. Allander T, Tammi MT, Eriksson M, Bjerkner A, Tiveljung-Lindell A, Andersson B. Cloning of a human parvovirus by molecular screening of respiratory tract samples. *Proceedings of the National Academy of Sciences*. 2005;102(36):12891-6.
6. Riera-Montes M, O'Ryan M, Verstraeten T. Norovirus and Rotavirus Disease Severity in Children: Systematic Review and Meta-analysis. *Pediatr Infect Dis J*. 2018;37(6):501-5.
7. Aliabadi N, Antoni S, Mwenda JM, Weldegebriel G, Biey JNM, Cheikh D, et al. Global impact of rotavirus vaccine introduction on rotavirus hospitalisations among children under 5 years of age, 2008-16: findings from the Global Rotavirus Surveillance Network. *Lancet Glob Health*. 2019;7(7):e893-e903.
8. Schildgen O, Qiu J, Söderlund-Venermo M. Genomic features of the human bocaviruses. *Future Virol*. 2012;7(1):31-9.
9. Brieu N, Gay B, Segondy M, Foulongne V. Electron microscopy observation of human bocavirus (HBoV) in nasopharyngeal samples from HBoV-infected children. *J Clin Microbiol*. 2007;45(10):3419-20.
10. Gurda BL, Parent KN, Bladek H, Sinkovits RS, DiMattia MA, Rence C, et al. Human bocavirus capsid structure: insights into the structural repertoire of the

436 parvoviridae. J Virol. 2010;84(12):5880-9.

437 11. Arthur JL, Higgins GD, Davidson GP, Givney RC, Ratcliff RM. A novel bocavirus
438 associated with acute gastroenteritis in Australian children. PLoS Pathog.
439 2009;5(4):e1000391.

440 12. Kapoor A, Simmonds P, Slikas E, Li L, Bodhidatta L, Sethabutr O, et al. Human
441 bocaviruses are highly diverse, dispersed, recombination prone, and prevalent in enteric
442 infections. J Infect Dis. 2010;201(11):1633-43.

443 13. Allander T, Jartti T, Gupta S, Niesters HG, Lehtinen P, Osterback R, et al. Human
444 bocavirus and acute wheezing in children. Clin Infect Dis. 2007;44(7):904-10.

445 14. Fry AM, Lu X, Chittaganpitch M, Peret T, Fischer J, Dowell SF, et al. Human
446 bocavirus: a novel parvovirus epidemiologically associated with pneumonia requiring
447 hospitalization in Thailand. J Infect Dis. 2007;195(7):1038-45.

448 15. Moriyama Y, Hamada H, Okada M, Tsuchiya N, Maru H, Shirato Y, et al.
449 Distinctive clinical features of human bocavirus in children younger than 2 years. Eur
450 J Pediatr. 2010;169(9):1087-92.

451 16. Falahi S, Sayyadi H, Abdoli A, Kenarkoochi A, Mohammadi S. The prevalence of
452 human bocavirus in <2-year-old children with acute bronchiolitis. New Microbes New
453 Infect. 2020;37:100736.

454 17. Jartti T, Hedman K, Jartti L, Ruuskanen O, Allander T, Söderlund-Venermo M.
455 Human bocavirus-the first 5 years. Rev Med Virol. 2012;22(1):46-64.

456 18. Jin Y, Cheng WX, Xu ZQ, Liu N, Yu JM, Li HY, et al. High prevalence of human
457 bocavirus 2 and its role in childhood acute gastroenteritis in China. J Clin Virol.
458 2011;52(3):251-3.

459 19. Rizk NM, Abd-Elmaksoud S, Farid TM, Abohashish MMA, Al-Herrawy AZ,
460 Hamza IA. Etiology of diarrheal disease among children under 5 years in Egypt: a high
461 incidence of human bocavirus. J Egypt Public Health Assoc. 2021;96(1):24.

462 20. Sousa TT, Almeida TNV, Fiaccadori FS, Souza M, Badr KR, Cardoso D.
463 Identification of Human Bocavirus type 4 in a child asymptomatic for respiratory tract
464 infection and acute gastroenteritis - Goiânia, Goiás, Brazil. Braz J Infect Dis.
465 2017;21(4):472-6.

21. Okitsu S, Khamrin P, Takanashi S, Thongprachum A, Hoque SA, Takeuchi H, et al. Molecular detection of enteric viruses in the stool samples of children without diarrhea in Bangladesh. *Infect Genet Evol.* 2020;77:104055.
22. De R, Liu L, Qian Y, Zhu R, Deng J, Wang F, et al. Risk of acute gastroenteritis associated with human bocavirus infection in children: A systematic review and meta-analysis. *PLoS One.* 2017;12(9):e0184833.
23. Knobloch K, Yoon U, Vogt PM. Preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement and publication bias. *J Craniomaxillofac Surg.* 2011;39(2):91-2.
24. Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa: Ottawa Hospital Research Institute. 2011:1-12.
25. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials.* 1986;7(3):177-88.
26. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Bmj.* 2003;327(7414):557-60.
27. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics.* 1994;50(4):1088-101.
28. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Bmj.* 1997;315(7109):629-34.
29. Melamed R, Storch GA, Holtz LR, Klein EJ, Herrin B, Tarr PI, et al. Case-control assessment of the roles of noroviruses, human bocaviruses 2, 3, and 4, and novel polyomaviruses and astroviruses in acute childhood diarrhea. *Journal of the Pediatric Infectious Diseases Society.* 2017;6(3):e49-e54.
30. Lasure N, Gopalkrishna V. Changing pattern of genotypic circulation of human bocavirus variants associated with acute gastroenteritis in Pune, Western India: A 5-year retrospective study. *J Med Virol.* 2019;91(10):1887-90.
31. De R, Liu LY, Qian Y, Jia LP, Dong HJ, Zhu RN, et al. [Correlation between human bocavirus 2 infection in children and acute diarrhea in Beijing]. *Zhonghua Er Ke Za Zhi.* 2018;56(12):945-9.

32. Tymentsev A, Tikunov A, Zhirakovskaia E, Kurilshikov A, Babkin I, Klemesheva V, et al. Human bocavirus in hospitalized children with acute gastroenteritis in Russia from 2010 to 2012. *Infection, Genetics and Evolution*. 2016;37:143-9.
33. Zeng M, Wang XH, Yu H, Zhu QR. [Clinical relevance of human bocavirus with acute respiratory tract infection and diarrhea in children: a prospective case-control study]. *Zhonghua Er Ke Za Zhi*. 2010;48(8):580-4.
34. Cheng WX, Jin Y, Duan ZJ, Xu ZQ, Qi HM, Zhang Q, et al. Human bocavirus in children hospitalized for acute gastroenteritis: a case-control study. *Clin Infect Dis*. 2008;47(2):161-7.
35. Chieochansin T, Thongmee C, Vimolket L, Theamboonlers A, Poovorawan Y. Human bocavirus infection in children with acute gastroenteritis and healthy controls. *Jpn J Infect Dis*. 2008;61(6):479-81.
36. Risku M, Ktk M, Lappalainen S, Rsnen S, Vesikari T. Human bocavirus types 1, 2 and 3 in acute gastroenteritis of childhood. *Acta Paediatr*. 2012;101(9):e405-10.
37. Nawaz S, Allen DJ, Aladin F, Gallimore C, Iturriza-Gmara M. Human bocaviruses are not significantly associated with gastroenteritis: results of retesting archive DNA from a case control study in the UK. *PLoS One*. 2012;7(7):e41346.
38. Ji Y, Wang Y, Yang Q, Cui H, Jin B. Epidemiological and phylogenetic analysis of human bocavirus from children with acute gastroenteritis in Beijing in 2016. *Journal of Pathogen Biology*. 2017;12(9).
39. Ali GM, Hussein AA, Kadim JI, Aufi IM. Detecting and Genotyping of Human Bocavirus among Children with Gastroenteritis in Diyala Governorate. *Annals of Tropical Medicine and Public Health*. 2020;23(14).
40. Shigemoto N, Hisatsune Y, Toukubo Y, Tanizawa Y, Shimazu Y, Takao S, et al. Detection of Gastroenteritis Viruses Among Pediatric Patients in Hiroshima Prefecture, Japan, Between 2006 and 2013 Using Multiplex Reverse Transcription PCR-Based Assays Involving Fluorescent Dye-Labeled Primers. *Journal of Medical Virology*. 2017;89(5):791-800.
41. Lekana-Douki SE, Behillil S, Enouf V, Leroy EM, Berthet N. Detection of human bocavirus-1 in both nasal and stool specimens from children under 5 years old with

influenza-like illnesses or diarrhea in Gabon. BMC Research Notes. 2018;11(1).

42. Rizk NM, Abd-Elmaksoud S, Farid TM, Abohashish MMA, Al-Herrawy AZ, Hamza IA. Etiology of diarrheal disease among children under 5 years in Egypt: a high incidence of human bocavirus. Journal of the Egyptian Public Health Association. 2021;96(1).

43. Lin MY, Chan HC, Chi H, Chiu SC, Nora-Krukke Z, Rasa-Dzelzkaleja S, et al. Genetic Diversity and Phylogenetic Analysis of Human Bocavirus 2 in Pediatric Patients with Acute Gastroenteritis in Taiwan. Int J Environ Res Public Health. 2020;17(3).

44. Nantachit N, Kochjan P, Khamrin P, Kumthip K, Maneekarn N. Human bocavirus genotypes 1, 2, and 3 circulating in pediatric patients with acute gastroenteritis in Chiang Mai, Thailand, 2012-2018. J Infect Public Health. 2021;14(2):179-86.

45. Campos GS, Silva Sampaio ML, Menezes AD, Tigre DM, Moura Costa LF, Chinalia FA, et al. Human bocavirus in acute gastroenteritis in children in Brazil. J Med Virol. 2016;88(1):166-70.

46. Malta FC, Varella RB, Guimaraes M, Miagostovich MP, Fumian TM. Human Bocavirus in Brazil: Molecular Epidemiology, Viral Load and Co-Infections. Pathogens. 2020;9(8).

47. La Rosa G, Della Libera S, Iaconelli M, Donia D, Cenko F, Xhelilaj G, et al. Human bocavirus in children with acute gastroenteritis in Albania. Journal of Medical Virology. 2016;88(5):906-10.

48. Lee EJ, Kim HS, Kim HS, Kim JS, Song W, Kim M, et al. Human Bocavirus in Korean Children with Gastroenteritis and Respiratory Tract Infections. BioMed Research International. 2016;2016.

49. Nora-Krukke Z, Vilmane A, Xu M, Rasa S, Ziemele I, Silina E, et al. Human Bocavirus Infection Markers in Peripheral Blood and Stool Samples of Children with Acute Gastroenteritis. Viruses. 2018;10(11).

50. Mohammadi M, Armin S, Yazdanpour Z. Human bocavirus infections and co-infections with respiratory syncytial virus and Rotavirus in children with acute respiratory or gastrointestinal disease. Brazilian Journal of Microbiology.

2020;51(1):45-51.

51. Sharif N, Nobel NU, Sakib N, Liza SM, Khan ST, Billah B, et al. Molecular and epidemiologic analysis of diarrheal pathogens in children with acute gastroenteritis in Bangladesh during 2014-2019. *Pediatric Infectious Disease Journal*. 2020:580-5.

52. Sharif N, Parvez AK, Haque A, Talukder AA, Ushijima H, Dey SK. Molecular and epidemiological trends of human bocavirus and adenovirus in children with acute gastroenteritis in Bangladesh during 2015 to 2019. *Journal of Medical Virology*. 2020;92(12):3194-201.

53. Okitsu S, Khamrin P, Takanashi S, Thongprachum A, Hoque SA, Takeuchi H, et al. Molecular detection of enteric viruses in the stool samples of children without diarrhea in Bangladesh. *Infection, Genetics and Evolution*. 2020;77.

54. Lasure N, Gopalkrishna V. Molecular epidemiology and clinical severity of Human Bocavirus (HBoV) 1-4 in children with acute gastroenteritis from Pune, Western India. *J Med Virol*. 2017;89(1):17-23.

55. Soares LS, Lima ABF, Pantoja KC, Lobo PS, Cruz JF, Guerra SFS, et al. Molecular epidemiology of human bocavirus in children with acute gastroenteritis from North Region of Brazil. *J Med Microbiol*. 2019;68(8):1233-9.

56. Netshikweta R, Chidamba L, Nadan S, Taylor MB, Page NA. Molecular epidemiology of human bocavirus infection in hospitalized children with acute gastroenteritis in South Africa, 2009-2015. *Journal of Medical Virology*. 2020;92(8):1124-32.

57. Adam MA, Wang J, Enan KA, Shen HW, Wang H, El Hussein AR, et al. Molecular Survey of Viral and Bacterial Causes of Childhood Diarrhea in Khartoum State, Sudan. *Frontiers in Microbiology*. 2018;9.

58. Rikhotso MC, Khumela R, Kabue JP, Traore-Hoffman IAN, Potgieter N. Predominance of Human Bocavirus Genotype 1 and 3 in Outpatient Children with Diarrhea from Rural Communities in South Africa, 2017-2018. *Pathogens*. 2020;9(4).

59. Zhou T, Chen YH, Chen JY, Hu PW, Zheng TL, Xu X, et al. Prevalence and clinical profile of human bocavirus in children with acute gastroenteritis in Chengdu, West China, 2012-2013. *Journal of Medical Virology*. 2017;89(10):1743-8.

60. Zhao M, Zhu RA, Qian Y, Deng J, Wang F, Sun Y, et al. Prevalence and
Phylogenetic Analysis of Human Bocaviruses 1-4 in Pediatric Patients with Various
Infectious Diseases. Plos One. 2016;11(8).
61. Kim GR, Kim SH, Jeon GW, Shin JH. Prevalence of eleven infectious viruses
causing diarrhea in korea. Japanese Journal of Infectious Diseases. 2020;73(6):427-30.
62. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of
the quality of nonrandomized studies in meta-analyses. Eur J Epidemiol.
2010;25(9):603-5.
63. Guido M, Tumolo MR, Verri T, Romano A, Serio F, De Giorgi M, et al. Human
bocavirus: Current knowledge and future challenges. World J Gastroenterol.
2016;22(39):8684-97.
64. Schildgen O. Human bocavirus: lessons learned to date. Pathogens. 2013;2(1):1-
12.
65. Modrow S, Wenzel JJ, Schimanski S, Schwarzbeck J, Rothe U, Oldenburg J, et al.
Prevalence of nucleic acid sequences specific for human parvoviruses, hepatitis A and
hepatitis E viruses in coagulation factor concentrates. Vox Sang. 2011;100(4):351-8.
66. Fabbiani M, Terrosi C, Martorelli B, Valentini M, Bernini L, Cellesi C, et al.
Epidemiological and clinical study of viral respiratory tract infections in children from
Italy. J Med Virol. 2009;81(4):750-6.
67. Söderlund-Venermo M, Lahtinen A, Jartti T, Hedman L, Kempainen K, Lehtinen
P, et al. Clinical assessment and improved diagnosis of bocavirus-induced wheezing in
children, Finland. Emerg Infect Dis. 2009;15(9):1423-30.
68. Longtin J, Bastien M, Gilca R, Leblanc E, de Serres G, Bergeron MG, et al. Human
bocavirus infections in hospitalized children and adults. Emerg Infect Dis.
2008;14(2):217-21.
69. Albuquerque MC, Pena GP, Varella RB, Gallucci G, Erdman D, Santos N. Novel
respiratory virus infections in children, Brazil. Emerg Infect Dis. 2009;15(5):806-8.
70. Ghietto LM, Cámara A, Cámara J, Adamo MP. High frequency of human
bocavirus 1 DNA in infants and adults with lower acute respiratory infection. J Med
Microbiol. 2012;61(Pt 4):548-51.

71. Ghietto LM, Cámara A, Zhou Y, Pedranti M, Ferreyra S, Frey T, et al. High prevalence of human bocavirus 1 in infants with lower acute respiratory tract disease in Argentina, 2007-2009. *Braz J Infect Dis.* 2012;16(1):38-44.
72. Arden KE, Chang AB, Lambert SB, Nissen MD, Sloots TP, Mackay IM. Newly identified respiratory viruses in children with asthma exacerbation not requiring admission to hospital. *J Med Virol.* 2010;82(8):1458-61.
73. Arden KE, McErlean P, Nissen MD, Sloots TP, Mackay IM. Frequent detection of human rhinoviruses, paramyxoviruses, coronaviruses, and bocavirus during acute respiratory tract infections. *J Med Virol.* 2006;78(9):1232-40.
74. Khamrin P, Malasao R, Chaimongkol N, Ukarapol N, Kongsricharoern T, Okitsu S, et al. Circulating of human bocavirus 1, 2, 3, and 4 in pediatric patients with acute gastroenteritis in Thailand. *Infect Genet Evol.* 2012;12(3):565-9.
75. Pham NT, Trinh QD, Chan-It W, Khamrin P, Nishimura S, Sugita K, et al. Human bocavirus infection in children with acute gastroenteritis in Japan and Thailand. *J Med Virol.* 2011;83(2):286-90.
76. Carrol ED, Mankhambo LA, Guiver M, Banda DL, Denis B, Dove W, et al. PCR improves diagnostic yield from lung aspiration in Malawian children with radiologically confirmed pneumonia. *PLoS One.* 2011;6(6):e21042.
77. Smuts H, Workman L, Zar HJ. Role of human metapneumovirus, human coronavirus NL63 and human bocavirus in infants and young children with acute wheezing. *J Med Virol.* 2008;80(5):906-12.
78. Li H, He M, Zeng P, Gao Z, Bian G, Yang C, et al. The genomic and seroprevalence of human bocavirus in healthy Chinese plasma donors and plasma derivatives. *Transfusion.* 2015;55(1):154-63.
79. Wang K, Wang W, Yan H, Ren P, Zhang J, Shen J, et al. Correlation between bocavirus infection and humoral response, and co-infection with other respiratory viruses in children with acute respiratory infection. *J Clin Virol.* 2010;47(2):148-55.
80. Martin ET, Taylor J, Kuypers J, Magaret A, Wald A, Zerr D, et al. Detection of bocavirus in saliva of children with and without respiratory illness. *J Clin Microbiol.* 2009;47(12):4131-2.

646 81. Mitui MT, Tabib SM, Matsumoto T, Khanam W, Ahmed S, Mori D, et al.
647 Detection of human bocavirus in the cerebrospinal fluid of children with encephalitis.
648 Clin Infect Dis. 2012;54(7):964-7.

649 82. Iaconelli M, Divizia M, Della Libera S, Di Bonito P, La Rosa G. Frequent
650 Detection and Genetic Diversity of Human Bocavirus in Urban Sewage Samples. Food
651 Environ Virol. 2016;8(4):289-95.

652 83. La Rosa G, Purpari G, Guercio A, Di Bella S, Gucciardi F, Proroga YTR, et al.
653 Detection of Human Bocavirus Species 2 and 3 in Bivalve Shellfish in Italy. Appl
654 Environ Microbiol. 2018;84(7).

655 84. Netshikweta R, Chidamba L, Nadan S, Taylor MB, Page NA. Molecular
656 epidemiology of human bocavirus infection in hospitalized children with acute
657 gastroenteritis in South Africa, 2009-2015. J Med Virol. 2020;92(8):1124-32.

658 85. Mohammadi M, Armin S, Yazdanpour Z. Human bocavirus infections and co-
659 infections with respiratory syncytial virus and Rotavirus in children with acute
660 respiratory or gastrointestinal disease. Braz J Microbiol. 2020;51(1):45-51.

661 86. Farahmand M, Moghoofei M, Dorost A, Shoja Z, Ghorbani S, Kiani SJ, et al.
662 Global prevalence and genotype distribution of norovirus infection in children with
663 gastroenteritis: A meta-analysis on 6 years of research from 2015 to 2020. Rev Med
664 Virol. 2021:e2237.

665 87. Shioda K, Kambhampati A, Hall AJ, Lopman BA. Global age distribution of
666 pediatric norovirus cases. Vaccine. 2015;33(33):4065-8.

667 88. Nordgren J, Svensson L. Genetic Susceptibility to Human Norovirus Infection: An
668 Update. Viruses. 2019;11(3).

669 89. Hickman D, Jones MK, Zhu S, Kirkpatrick E, Ostrov DA, Wang X, et al. The effect
670 of malnutrition on norovirus infection. mBio. 2014;5(2):e01032-13.

671 90. Taneja V. Sex Hormones Determine Immune Response. Front Immunol.
672 2018;9:1931.

673 91. Tymentsev A, Tikunov A, Zhirakovskaia E, Kurilshchikov A, Babkin I, Klemesheva
674 V, et al. Human bocavirus in hospitalized children with acute gastroenteritis in Russia
675 from 2010 to 2012. Infect Genet Evol. 2016;37:143-9.

676 92. Ong DS, Schuurman R, Heikens E. Human bocavirus in stool: A true pathogen or
677 an innocent bystander? J Clin Virol. 2016;74:45-9.

678 93. Martin ET, Kuypers J, McRoberts JP, Englund JA, Zerr DM. Human Bocavirus 1
679 Primary Infection and Shedding in Infants. J Infect Dis. 2015;212(4):516-24.

680 94. Martin ET, Fairchok MP, Kuypers J, Magaret A, Zerr DM, Wald A, et al. Frequent
681 and prolonged shedding of bocavirus in young children attending daycare. J Infect Dis.
682 2010;201(11):1625-32.

683 95. Yu JM, Li DD, Xu ZQ, Cheng WX, Zhang Q, Li HY, et al. Human bocavirus
684 infection in children hospitalized with acute gastroenteritis in China. J Clin Virol.
685 2008;42(3):280-5.

686 96. Brieu N, Guyon G, Rodière M, Segondy M, Foulongne V. Human bocavirus
687 infection in children with respiratory tract disease. Pediatr Infect Dis J.
688 2008;27(11):969-73.

689