Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Article Summary Line: Evidence of increasing dengue endemicity in Florida: Vector surveillance during dengue and West Nile virus outbreaks revealed widespread presence of other dengue virus serotypes in the absence of local index cases.

Running Title: Dengue and West Nile Virus outbreak in Florida

Keywords: Dengue virus, West Nile Virus, Arboviruses, Disease Vectors, Disease Outbreaks, COVID-19

Title: A molecular surveillance-guided vector control response to concurrent dengue and West Nile virus outbreaks in a COVID-19 hotspot of Florida

Authors: Heather Coatsworth¹, Catherine A. Lippi, Chalmers Vasquez, Jasmine B. Ayers, Caroline J. Stephenson, Christy Waits, Mary Florez, André B. B. Wilke, Isik Unlu, Johana Medina, Maria L. Alcaide, Sadie J. Ryan, John A. Lednicky, John C. Beier, William Petrie, Rhoel R. Dinglasan²

¹Corresponding author for review purposes only

²Corresponding author

Corresponding Author: Rhoel R. Dinglasan, University of Florida, 2055 Mowry Rd, Gainesville, FL 32611, USA, rdinglasan@epi.ufl.edu, (+1) 352-294-8448

Author Affiliations:

University of Florida, Gainesville, Florida, USA (H. Coatsworth, C. Lippi, J. B. Ayers, C. J. Stephenson, C. Waits, M. Florez, S. J. Ryan, J. A. Lednicky, R. R. Dinglasan)

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Miami-Dade Mosquito Control District, Miami, Florida, USA (C. Vasquez, I. Unlu, J. Medina,

W. Petrie)

University of Miami, Miami, Florida, USA (A. B. B. Wilke, M. L. Alcaide, J. C. Beier)

Navy Entomology Center of Excellence, Jacksonville, FL USA (C. Waits)

Abstract

Simultaneous dengue virus (DENV) and West Nile virus (WNV) outbreaks in Florida, USA, in 2020 resulted in 71 dengue virus serotype 1 and 86 WNV human cases. Our outbreak response leveraged a molecular diagnostic screen of mosquito populations for DENV and WNV in Miami-Dade County to quickly employ targeted mosquito abatement efforts. We detected DENV serotypes 2 and 4 in mosquito pools, highlighting the silent circulation of diverse dengue serotypes in mosquitoes. Additionally, we found WNV-positive mosquito pools in areas with no historical reports of WNV transmission. These findings demonstrate the importance of proactive, strategic arbovirus surveillance in mosquito populations to prevent and control outbreaks, particularly when other illnesses (e.g., COVID-19), which present with similar symptoms are circulating concurrently. Growing evidence for substantial infection prevalence of dengue in competent mosquito vectors in the absence of local index cases suggests a higher level of dengue endemicity in Florida than previously thought.

Introduction

Although West Nile Virus (WNV) is endemic in the continental United States, only a handful of states, especially Florida, are at risk of autochthonous dengue virus (DENV) transmission (1,2). Throughout 2020, Miami-Dade County was the epicenter of COVID-19 in

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Florida, but unbeknownst to the general U.S. population, the state experienced in parallel, concurrent outbreaks of DENV and WNV from May - December 2020, resulting in 71 local cases of dengue fever (DENV serotype 1) and 86 cases of WNV. Herein, we report the outcome of vector surveillance and control effort in response to the arbovirus outbreaks in Miami-Dade County (August- November 2020).

Methods

Sample collection and processing

Adult mosquitoes were collected with BG-Sentinel traps (Biogents AG, Regensburg, Germany) baited with dry ice, for 24 hours. Miami-Dade Mosquito Control District (MDMCD) trapped, morphologically identified, and sorted adult female mosquitoes by location into pools of 2-25 mosquitoes of the same species. Samples were kept cold to avoid RNA degradation, shipped on dry ice, and stored at -80°C. In total, 743 pools were collected for testing [548 for DENV (pools of *Aedes aegypti* and *Aedes albopictus*, N= 5,079 mosquitoes) and 188 for WNV (pools of *Anopheles crucians*, *Culex coronator*, *Culex nigripalpus*, and *Culex quinquefasciatus*, N= 2,589 mosquitoes)].

RNA extraction

Chilled, sterile 1X phosphate-buffered saline (PBS) and sterile glass beads were added to each sample in a biosafety cabinet (BSC). Each 1 mL sample was homogenized in a Bullet Blender (speed 8, 5 minutes) with repeated cooling on ice/cold block in the BSC. Samples were centrifuged (3,750xg, 3 minutes) and 140 µL of the homogenate supernatant was added to AVL lysis buffer (560 µL) (Qiagen). RNA was extracted using the QIAmp Viral RNA extraction kit (Qiagen) following the manufacturer's protocols with 2x40 µL elution steps. One pool of 5

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

uninfected *Ae. aegypti* (Orlando) mosquitoes was processed in parallel, serving as a negative extraction control to rule out contamination or spurious amplification.

Real-Time RT-PCR Virus Detection

Sample RNA was either tested for i) WNV or ii) DENV based on the mosquito species for a given pool. Samples designated for DENV screening were first run through a pan-dengue serotype screen, DENVA11 (Appendix Table 1), while the remaining samples were screened via a WNV assay (Appendix Table 2). Each sample, prepared in a PCR hood with static air, was run as technical duplicates. Each plate included a mosquito extraction control, a no template control, and either i) a positive WNV control or ii) dengue virus serotype 1, 2, 3 and 4 (DENV-1 through DENV-4) positive controls. All positive RNA controls were obtained from BEI resources diluted 1:10 (Appendix Tables 1,2,5). Analyses used either QuantaBio UltraPlex 1-Step ToughMix (4X) Low-ROX master mix (Appendix Table 3) or SuperScript™ III Platinum™ One-Step qRT-PCR (Appendix Table 4) on a BioRad CFX96 Touch Real-Time PCR Detection System at 50°C for 30 minutes (for Superscript reactions) or 50°C for 10 minutes (for QuantaBio reactions), 95°C for 2 minutes, and 45 cycles of: 95°C for 15 seconds and 60°C for 45 seconds. Both assays yielded similar results, which helped overcome availability bottlenecks during the COVID-19 pandemic. Samples that were DENV-positive via the DENVA11 assay were analyzed for serotype specificity (Appendix Tables 5). Any putative positive or inconclusive (i.e., only one replicate amplified) samples were re-run in confirmatory reactions, adding extra technical replicates and performing additional confirmatory runs as needed to confirm virus positivity.

Positive and negative sample designation

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

A sample was considered 'positive' based on positive technical duplicate results in two independent runs using the WNV assay (Appendix Table 2) and the DENV serotype-specific assay (Appendix Table 5). Samples were considered 'negative' based on no detectable Ct value in either technical duplicate after 45 cycles.

Mapping mosquito samples

To understand the spatial distribution of mosquito pools collected for this effort, we developed maps using ArcMap (v 10.6) at the zip code level, which is the smallest level of resolution obtainable with our datasets. All datasets were either publicly available: Miami-Dade County Boundaries, population density and land use (2010 US Census, Miami-Dade County's Open Data Hub), median household income (2010 US Census, Michigan Population Studies), persons living with HIV/AIDS (PLWH) (2018, AIDsVu), or obtained with permission from personal communications (2018 – 2020 mosquito prevalence c/o Chalmers Vasquez, Miami-Dade Mosquito Control District; 2009 – 2019 Imported DENV cases c/o Andrea Morrison, Florida Department of Health). Land-use types were manually concatenated to 11 different primary categories: Agricultural, Cemeteries, Commercial, Educational, Industrial, Marine, Recreational, Residential, Paved, Water, and Vacant. These types were further summed as: 'urban/built' (Commercial, Educational, Industrial, and Paved), 'agricultural/recreational' (Agricultural, Recreational), 'residential' (Residential) and 'other' (Cemeteries, Marine, Water, and Vacant) (Appendix Table 6).

Spatial statistical analysis of vector distribution and spatial visualization of overlapping PLWH and arbovirus hotspots

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

To explore the spatial relationship between positivity and overall mosquito sampling, the directional distribution and mean center (e.g., central tendencies, dispersion) were computed for DENV- and WNV-positive mosquito pools, for all collected DENV vectors (*Ae. aegypti* and *Ae. albopictus*) and all WNV vectors (*An. crucians*, *Cx. coronator*, *Cx. nigripalpus*, and *Cx. quinquefasciatus*), both individually and together. Kernel density estimation (KDE) with optimal distance bandwidths (3) was used to visualize the continuous density of DENV- and WNV-positive mosquito pools, as well as traps containing un-infected DENV and WNV vectors. To determine whether arbovirus-positive pools occurred in areas where there were imported DENV cases or in areas with a higher number of PLWH, local Moran's I, a local indicator of spatial association (LISA), analyses with inverse distance weighting were performed to detect and identify PLWH hotspots (areas of elevated incidence or prevalence) that potentially overlap with locations of arbovirus positive pools, as well as imported DENV cases for 2019 only, and for the 2009-2019 period. All spatial statistical analyses were completed in ArcMap (v 10.6).

Results

Molecular detection of arbovirus-positive mosquito pools

We found that 2.96% of female mosquito pool samples (22/743 pools, 382/7,668 mosquitoes) were positive for an arbovirus; 4 were DENV-2 positive, 9 were DENV-4 positive (2.43% DENV positivity), and 9 were WNV positive (4.79% WNV positivity) (Table 1 and Appendix 7). Results were confirmed and verified independently by the Florida Department of Health (FLDOH), and the inclusion of a pool of lab reared mosquitoes in every extraction and PCR run ruled out contamination during lab handling or spurious amplification of mosquito material.

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

For DENV, we tested 45 *Ae. albopictus* and 510 *Ae. aegypti* mosquito pools, and only *Ae. aegypti* pools were found to be positive for DENV-2 and/or DENV-4 (2.43% positivity).

For WNV, we tested five *An. crucians*, five *Cx. coronator*, 47 *Cx. nigripalpus* and 131 *Cx. quinquefasciatus* pools. Only *Cx. nigripalpus* pools were WNV positive (19.1% positivity), although CDC findings from earlier in 2020 (June), found three positive *Cx. quinquefasciatus* pools.

Mosquito pool samples that were found to be DENV- or WNV-positive were reported to the MDMCD within 48-72 hours. Miami-Dade Mosquito Control Division initiated door-to-door source reduction in the areas where positive mosquito pools were found. Vectobac WGD (Valent Biosciences), a larvicide product containing *Bacillus thuringiensis israelensis*, was applied at positive pool sites for four weeks. Adulticide treatment using chlorpyrifos was also conducted (weather permitting) using Mosquitomist (Clarke).

A spatial-climatic distribution of arbovirus-positive mosquito pools identified exigent and emerging areas of concern

DENV-positive mosquito pools were distributed throughout the county, with most (11/13) positive pools located in downtown Miami (N=3), Wynwood (N=2), and South Miami (N=6) (Figure 1A) and in areas with mid-to-high human population density (2,000 – 7,000 people/km²) (Figure 1B). At the zip code level, DENV-positive pools were primarily found in built and residential environments (59.42%) over more rural, (i.e., agricultural and recreational) areas (14.62%) (Figure 1C). Conversely, the thirteen WNV-positive pools (9 identified herein) had three primary infection foci – in Wynwood (N=4), high population density (>3000 people/km², urban built environment), in Homestead (N=8), low population density (<470

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

people/km², mostly agricultural and recreational land), and in North Miami (N=1), mid population density (1,000 – 3,000 people/km² and mixed land use) (Figure 1D-F). WNV-positive pools were primarily from agricultural and recreational areas (69.32%) and were less common in urban built and residential areas (15.98%).

The directional distribution and mean center computations of dengue vector distributions showed that DENV-positive mosquito pools were within the directional distribution of all dengue vector traps (Figure 2A). *Ae. aegypti* prevalence dominated the traps in much of the county and had a much larger directional distribution than *Ae. albopictus*, whose presence was concentrated in southern Miami-Dade County (Figure 2B). Although the directional distribution of all WNV vectors overlapped with the directional distribution of the WNV-positive mosquito pools, the latter extended well beyond the southernmost tip of the WNV vector directional ellipse (Figure 2C). *Culex quinquefasciatus*, *An. crucians* and *Cx. nigripalpus* had distinct directional distributions in the Miami Beach area, North Miami, and South Miami, respectively (Figure 2D). The distribution of *Cx. coronator* encompassed all three of these zones, lacking any strict boundary (Figure 2D).

The KDE maps show a clear high density of vector trapping efforts across Miami-Dade County except for the southernmost area (Appendix Figure 1A). The KDE map for DENV-2 positive pools showed high values surrounding each positive pool but low values between pools (Appendix Figure 1B), while the DENV-4 positive KDE map showed connected high values across all positive pools (Appendix Figure 1C). The KDE map for the WNV vector trapping effort showed a high concentration of vectors in the downtown Miami/Miami Beach areas, with areas of mid values dotted throughout the county (Appendix Figure 1E). The WNV-positive

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

pools showed two clear high-value foci, one in Homestead, and one in Miami Beach (Appendix Figure 1F).

DENV-positive mosquito pools were in areas of higher average maximum temperatures, lower average minimum temperatures, and higher total precipitation (Appendix Figure 2 A-C). WNV-positive pools displayed the same trend with respect to temperature but were found in areas of lower total precipitation (Appendix Figure 2D-F).

LISA analyses on imported DENV case data showed similar clusters for both the 2009-2019 imported DENV data (Appendix Figure 3A), as well as the 2019 only imported DENV dataset (Appendix Figure 3B). The northwestern area of Miami-Dade was a high-high cluster, indicating this area is a hotspot for imported DENV cases. Conversely, the Miami Beach area was a low-low cluster, suggesting a consistently lower-than-average number of imported DENV cases (a cold spot). The low-high outliers for both datasets occurred just below the hotspot in northwest Miami-Dade, while the high-low outlier for the decade-wide dataset was in South Miami, and the 2019 only data had high-low outliers throughout North Miami.

We explored the possibility of increased arbovirus risk to special populations in Miami-Dade (i.e., PLWH) by examining in parallel the LISA results for PLWH and the spatial distribution of DENV or WNV positive pools and observed that DENV-positive mosquito pools were in the low-low PLWH cluster (Appendix Figure 3C), while one of the WNV-positive pool foci was in a PLWH hotspot (Appendix Figure 3D).

Discussion

Ecological distribution of WNV-positive mosquito pools

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Although WNV-positive mosquito pools are rarely reported in Miami-Dade County, we found nine positive mosquito pools collected between June – November 2020; three other WNV positive mosquito pools were found in 2020 (collected earlier in June and analyzed by the CDC). All four mosquito species we tested can transmit WNV (4,5); however, we only found WNV-positive *Cx. nigripalpus* pools, an expected result, as it is the primary enzootic and epidemic vector of mosquito-borne viruses encephalitides such as WNV, St. Louis encephalitis virus, and eastern equine encephalitis virus throughout southern Florida (6). This result is intriguing, considering the overwhelmingly dominant collection of *Cx. quinquefasciatus* throughout the county. Differences in WNV vector positivity may be due to several factors including host preference and vector competence. Previous reports show that *Cx. quinquefasciatus* has a wide range of WNV competency that appears to partition according to spatial and climatic scales and is influenced by virus genetic background (7–11).

There appeared to be a centralized distribution of traps collecting *Culex* and *Anopheles* mosquitoes, while positive pool traps were dispersed throughout the county. The spatial partitioning of the WNV-positive mosquito pools suggests that the positive pools had a distinct spatial trend within the larger distribution of traps collecting *Culex* and *Anopheles* mosquitoes in Miami-Dade County. This might indicate that current trap coverage is not adequate to reliably detect WNV-positive mosquito pools. As WNV infects a wealth of other non-human animals (i.e., horses, alligators, birds, etc.) (12), controlling the spread of WNV requires additional zoonotic measures including implementing equine WNV vaccines, as well as testing dead birds to understand prevalence and virus hotspots throughout the county.

WNV-positive mosquito pools were primarily located in areas with lower precipitation, higher maximum temperature, and lower minimum temperature (range of 21-29°C), optimal for

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

multiple *Culex* vectors (10). As there was no clear overlap between the spatial distribution of WNV vectors and climatic variables, these general climatic conditions are likely ideal for all the WNV vectors we analyzed. WNV-positive mosquito pools were found primarily in agricultural and recreational areas. These areas include protected bird sanctuaries, as well as habitats for resident and migrating shorebirds, which are known WNV avian hosts (12).

Ecological distribution of DENV-positive mosquito pools

Although other dengue vectors (i.e., *Aedes spp.*) are present in Miami-Dade (13), our data suggest that *Ae. aegypti* is clearly the primary vector of concern. Evidence suggests that Floridian *Ae. aegypti* (Monroe County) and *Ae. albopictus* (Indian River County) are similarly competent for DENV-1 (14). DENV vector competency varies greatly based on DENV serotype and FL *Ae. aegypti* geographic origin (15), so extrapolations of *Ae. aegypti* and *Ae. albopictus* vector competence across FL may not be appropriate. Our findings may be a by-product of *Ae. aegypti* prevalence, as considerably more *Ae. aegypti* were collected than *Ae. albopictus* in traps across the entirety of Miami-Dade County in 2020, and in former years (13).

DENV-positive pools were primarily located in areas with higher precipitation, higher maximum temperature, and lower minimum temperature. The directional distribution of DENV-positive pools suggests that the current trap spread is more than adequate to reliably detect DENV-positive pools. However, since trap density within the DENV-positive pool distribution area is not as concentrated as in the other *Aedes* positive trap locales, additional traps in the ellipse margins could prove useful.

Epidemiological relevance of WNV-positive mosquito pools

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

WNV-positive mosquito pools overlapped with the timing of 27 human symptomatic cases (June, July, August), 33 human asymptomatic blood donors (June, July August), as well as 15 WNV-positive birds (June, July), all in Miami-Dade County (16). One of the WNV-positive pool foci overlapped with a hotspot of PLWH (Appendix Figure 3D). This is especially concerning for PLWH, as these individuals are at a higher risk of neurotropism from WNV infection (17–21). WNV positive pools were primarily found in more rural areas with lower median household incomes, and low to medium population density (Figure 1 D-F). Low-income areas have previously been associated with higher WNV prevalence (22,23). These data underscore the need for tailored programs to protect and prevent infection in these at-risk populations.

Epidemiological relevance of DENV-positive mosquito pools

In the last five years, the number of local DENV cases in Florida has risen from zero (2017) to 71 (2020). A rise in cases can also be seen in the imported DENV case numbers (DENV-1 through DENV-4). In 2017, there were only 18 imported cases, 73 in 2018, 395 in 2019, 41 in 2020. This spike in dengue cases was mirrored in Miami-Dade County, where local case numbers moved from zero cases in 2017, to one in 2018, fourteen in 2019 and four in 2020, and travel-associated cases progressed from nine in 2017, to thirty-eight in 2018, 226 in 2019 and twenty in 2020. The large spike in imported cases in 2019 may have introduced other DENV serotypes into the resident Miami-Dade mosquito populations, acting as seeding events, which are known drivers of local DENV case incidence (24,25).

To examine whether this could be the case, we analyzed the available zip code-mapped historical (2009 – 2019) and 2019 imported DENV data using LISA analyses (Appendix Figure

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

3A and B), as well as the 2019 only imported DENV dataset (Appendix Figure 3B). The similarity in the decade-long and 2019 LISA analyses suggests that the high number of imported cases in 2019 may have driven the hotspot trends seen in our analyses. None of our DENV-positive mosquito pools overlapped with known imported DENV case hotspots. This could be due to a lack of reporting of zip code level imported DENV data. Alternatively, if mosquitoes bite individuals outside of their listed area of primary residence, such as where individuals spend time outdoors, where they work, or where they socialize, the direct overlap would not be observed.

Local human cases of DENV-1, DENV-2 and DENV-3 have been reported in FL (2,16,26–28). In 2020, only DENV-1 local cases were reported, despite our findings of DENV-2 and DENV-4 positive mosquito pools in the county. ‘Silent’ DENV circulation, defined as transmission between DENV asymptomatic individuals, as well as DENV maintenance in the vector population despite no reported human infection, likely represents the majority of transmission events (29). Finding additional DENV serotypes in the absence of a local human index case is not the first instance of silent DENV circulation in *Ae. aegypti* in the Americas. Previous reports have shown that DENV-4 was found in Manatee County, FL in 2018, and DENV-3 was detected in *Ae. aegypti* in Brazil despite no human index cases (30,31). This silent circulation is likely due to low but persistent vertical transmission in the mosquito population (31–33). This disparity could also be due to inherent differences in mosquito vector competence for DENV-1, -2 and -4, as mosquitoes collected in Miami-Dade had higher DENV-1 infection and transmission rates than mosquitoes from Miami-Dade infected with DENV-2 and -4 (15). Although the pathogenicity of the 2020 Miami-Dade human DENV-1 index strain is unknown,

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

the strain may be more infectious to or cause increased disease severity in humans than other circulating dengue strains.

Having multiple concurrent circulating serotypes puts individuals at an increased risk of DENV infection due to complications arising from immune enhancement (34,35). These risks include dengue fever and severe dengue (dengue hemorrhagic fever and dengue shock syndrome) and can be fatal (36). A woman in her 30's died from DENV-2 in Miami-Dade in 2019, and subsequent viral analysis suggested that infection occurred after DENV circulation had occurred in the Miami area (37).

There were more DENV-positive pools in areas with lower prevalence of PLWH, higher median household income and mid-high population density (Figure 1A-C, Appendix Figure 3C). Although there does not seem to be evidence for an association between income and DENV prevalence (38), DENV is a known urban-centric disease (39).

The perfect storm: arbovirus transmission in a global pandemic

Miami Dade County has been a hotspot of SARS-CoV-2 transmission throughout the pandemic, with the highest case rate and death toll in the state. Unfortunately, due to the non-standardized nature of COVID-19 case reporting throughout Florida (reports range from residence-based, testing locale-based and exposure-based), we could not directly map COVID-19 prevalence alongside mosquito pool positivity. The potential for immunocompromise resulting from acute SARS-CoV-2 infection to increase severity and/or transmissibility of other diseases (i.e. West Nile, dengue) in the Miami Dade population is concerning.

Due to the similar symptoms produced by COVID-19, West Nile, and dengue infection (fever, headache, muscle pain, nausea, vomiting, malaise), symptomatic WNV or DENV

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

infected individuals may have self-quarantined assuming they were positive for COVID-19 and never received a correct diagnosis. The high number of asymptomatic WNV positive individuals identified directly through blood donorship suggests that the proportion of asymptomatic WNV positive human carriers may be high, and previous reports show that the majority of DENV infected humans are also asymptomatic carriers (29). It should be noted that dengue is not currently part of routine blood donor screening in Miami-Dade.

The spread of the positive pools is intriguing, as prior arbovirus positive pools (for WNV and Zika virus) in Miami-Dade County were solely concentrated within the Miami Beach, Little River and Wynwood locales. Our data suggest a much wider arbovirus positive pool spread, ranging across the county.

Together, these data highlight a lack of vector containment and the importance of continued surveillance for arboviruses in their vector species. This is especially true when infection risk and continued transmission is increased due to: i) the presence of other illnesses (e.g., COVID-19) that present similarly in a clinic to arbovirus infection, ii) the high incidence of PLWH in areas of known WNV positivity resulting in potentially worse health outcomes following infection, and iii) climatic conditions favoring above-average *Ae. aegypti* and *Cx. nigripalpus* vector indices. The large proportion of asymptomatic WNV cases highlights the critical need for improved DENV diagnostics, as no dengue rapid diagnostic tests are currently cleared for use in the United States (37). Prevention and control measures are imperative to prevent future and expanded DENV and WNV outbreaks; targeted vector control efforts through continued mosquito pathogen screening and subsequent population specific insecticide spraying are necessary to prevent arbovirus spread throughout the county.

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Acknowledgments

This research was supported in part by U.S. Centers for Disease Control and Prevention (CDC) grant 1U01CK000510-03, Southeastern Regional Center of Excellence in Vector Borne Diseases Gateway Program. The CDC did not have a role in the design of the study, the collection, analysis, or interpretation of data, or writing the manuscript. Christy Waits is a contractor employee of the U.S. Government. This work was prepared as part of her official duties. Title 17, U.S.C., §105 provides that copyright protection under this title is not available for any work of the U.S. Government. Title 17, U.S.C., §101 defines a U.S. Government work as a work prepared by a military Service member or employee of the U.S. Government as part of that person's official duties. The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U. S. Government. The Department of the Navy, Department of Defense, or the U. S. Government did not have a role in the design of the study, the collection, analysis, or interpretation of data, or writing of the manuscript.

Author Bio

Dr. Coatsworth is a post-doctoral associate at the University of Florida. Her primary research interests include arbovirology, vector biology, and vector-borne disease transmission.

References

1. Franco C, Hynes NA, Bouri N, Henderson DA. The Dengue Threat to the United States. [cited 2021 Sep 10]; Available from: www.liebertpub.com
2. Rey JR. Dengue in Florida (USA). *Insects* 2014, Vol 5, Pages 991-1000 [Internet]. 2014 Dec 16 [cited 2021 Oct 7];5(4):991–1000. Available from: <https://www.mdpi.com/2075->

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

4450/5/4/991/htm

3. Fotheringham AS, Brunsdon C, Charlton M. Quantitative Geography: Perspectives on Spatial Data Analysis. 6th ed. London: Sage Publications; 2000.

4. Turell MJ, Sardelis MR, Dohm DJ, O'guinn ML. Potential North American Vectors of West Nile Virus.

5. Unlu I, Kramer WL, Roy AF, Foil LD. Detection of West Nile Virus RNA in Mosquitoes and Identification of Mosquito Blood Meals Collected at Alligator Farms in Louisiana. J Med Entomol [Internet]. 2010 [cited 2021 Sep 10];47(4):625–33. Available from: <https://academic.oup.com/jme/article/47/4/625/992551>

6. Day JF, Tabachnick WJ, Smartt CT. Factors That Influence the Transmission of West Nile Virus in Florida. J Med Entomol [Internet]. 2015;52(5):743–54. Available from: <https://academic.oup.com/jme/article/52/5/743/828059>

7. Anderson SL, Richards SL, Tabachnick WJ, Smartt, Chelsea T. Effects of West Nile Virus Dose and Extrinsic Incubation Temperature on Temporal Progression of Vector Competence in *Culex pipiens quinquefasciatus*. Source J Am Mosq Control Assoc [Internet]. 2010;26(1):103–7. Available from: <https://doi.org/10.2987/09-5926.1>

8. Richards SL, Lord CC, Pesko KN, Tabachnick WJ. Environmental and biological factors influencing *Culex pipiens quinquefasciatus* Say (Diptera: Culicidae) vector competence for West Nile virus. Am J Trop Med Hygiene [Internet]. 2010 [cited 2021 Sep 10];83(1):126–34. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2912589/pdf/tropmed-83-126.pdf>

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

- 367 9. Keyel AC, Timm OE, Backenson PB, Id CP, Quinones S, Mcdonough KA, et al. Seasonal
368 temperatures and hydrological conditions improve the prediction of West Nile virus
369 infection rates in Culex mosquitoes and human case counts in New York and Connecticut.
370 2019; Available from: <https://doi.org/10.1371/journal.pone.0217854>
- 371 10. Shocket MS, Verwillow AB, Numazu MG, Slamani H, Cohen JM, El Moustaid F, et al.
372 Transmission of west nile and five other temperate mosquito-borne viruses peaks at
373 temperatures between 23°C and 26°C. Elife. 2020 Sep 1;9:1–67.
- 374 11. Richards SL, Anderson SL, Lord CC. Vector competence of Culex pipiens
375 quinquefasciatus (Diptera: Culicidae) for West Nile virus isolates from Florida. 2014;
376 Available from: <http://cdo.ncdc>.
- 377 12. Marra P., Griffing S, Caffrey C, Kilpatrick MA, McLean R, Brand C, et al. West Nile
378 Virus and Wildlife . Bioscience [Internet]. 2004 [cited 2021 Sep 11];54(5):393–402.
379 Available from: <https://academic.oup.com/bioscience/article/54/5/393/416853?login=true>
- 380 13. Wilke ABB, Vasquez C, Medina J, Carvajal A, Petrie W, Beier JC. Community
381 Composition and Year-round Abundance of Vector species of Mosquitoes make Miami-
382 Dade County, Florida a Receptive Gateway for Arbovirus entry to the United states.
383 Available from: <https://doi.org/10.1038/s41598-019-45337-2>
- 384 14. Richards SL, Anderson SL, Alto BW. Vector Competence of Aedes aegypti and Aedes
385 albopictus (Diptera: Culicidae) for Dengue Virus in the Florida Keys. J Med Entomol
386 [Internet]. 2012;49(4):942–6. Available from: <http://dx.doi.org/10.1603/ME11293>
- 387 15. Stephenson CJ, Coatsworth H, Waits CM, Nazario-Maldonado NM, Mathias DK,

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

- Dinglasan RR, et al. Geographic partitioning of dengue virus transmission risk in Florida. prep. 2021;
16. Morrison A, Rowe D, McDermott C, Stanek D, Heberlein-Larson L, LaCrue A, et al. Florida Arbovirus Surveillance [Internet]. Available from: http://www.floridahealth.gov/diseases-and-conditions/mosquito-borne-diseases/_documents/2020-week-53-arbovirus-surveillance-report.pdf
17. Klein RS. A Moving Target: The Multiple Roles of CCR5 in Infectious Diseases. 2007; Available from: <https://academic.oup.com/jid/article/197/2/183/809887>
18. Pang J, Thein T-L, Lye DC, Leo Y-S. Differential Clinical Outcome of Dengue Infection among Patients with and without HIV Infection: A Matched Case-Control Study. Am J Trop Med Hyg. 2015;92(6):1156–62.
19. WG G, DH M, JK L, S L, SF Y, WA F, et al. CCR5 deficiency increases risk of symptomatic West Nile virus infection. J Exp Med [Internet]. 2006 Jan 23 [cited 2021 Sep 20];203(1):35–40. Available from: <https://pubmed.ncbi.nlm.nih.gov/16418398/>
20. JK L, CY L, C G, C J, B J, H J, et al. Genetic deficiency of chemokine receptor CCR5 is a strong risk factor for symptomatic West Nile virus infection: a meta-analysis of 4 cohorts in the US epidemic. J Infect Dis [Internet]. 2008 Jan 15 [cited 2021 Sep 20];197(2):262–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/18179388/>
21. JK L, DH M, A L, GA F, D K, D F, et al. CCR5 deficiency is a risk factor for early clinical manifestations of West Nile virus infection but not for viral transmission. J Infect Dis [Internet]. 2010 Jan 15 [cited 2021 Sep 20];201(2):178–85. Available from:

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

- 409 <https://pubmed.ncbi.nlm.nih.gov/20025530/>
- 410 22. Harrigan RJ, Thomassen HA, Buermann W, Cummings RF, Kahn ME. Economic
411 Conditions Predict Prevalence of West Nile Virus. PLoS One [Internet].
412 2010;5(11):15437. Available from: www.plosone.org
- 413 23. Rothman SE, Jones JA, LaDeau SL, Leisnham PT. Higher West Nile Virus Infection in
414 Aedes albopictus (Diptera: Culicidae) and Culex (Diptera: Culicidae) Mosquitoes From
415 Lower Income Neighborhoods in Urban Baltimore, MD. J Med Entomol [Internet]. 2021
416 May 15 [cited 2021 Sep 11];58(3):1424–8. Available from:
417 <https://academic.oup.com/jme/article/58/3/1424/6013060>
- 418 24. Guzzetta G, Marques-Toledo CA, Rosà R, Teixeira M, Merler S. Quantifying the spatial
419 spread of dengue in a non-endemic Brazilian metropolis via transmission chain
420 reconstruction. Available from: www.nature.com/naturecommunications
- 421 25. Li N, Feng Y, Vrancken B, Chen Y, Dong L, Yang Q, et al. Assessing the impact of
422 COVID-19 border restrictions on dengue transmission in Yunnan Province, China: an
423 observational epidemiological and phylogenetic analysis. Lancet Reg Heal Pacific
424 [Internet]. 2021 [cited 2021 Sep 11];14:100259. Available from:
425 <https://doi.org/10.1016/j.lanwpc.2021.100259>
- 426 26. Trout A, Baracco G, Rodriguez M, Barber J, Leal A, Radke E, et al. Locally Acquired
427 Dengue --- Key West, Florida, 2009--2010. Morb Mortal Wkly Rep [Internet]. 2010 [cited
428 2021 Oct 7];59(19):577–81. Available from:
429 <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5919a1.htm>

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

27. Radke EG, Gregory CJ, Kintziger KW, Sauber-Schatz EK, Hunsperger EA, Gallagher GR, et al. Dengue Outbreak in Key West, Florida, USA, 2009. *Emerg Infect Dis* [Internet]. 2012 Jan [cited 2021 Oct 7];18(1):135. Available from: [/pmc/articles/PMC3310087/](https://pubmed.ncbi.nlm.nih.gov/210087/)
28. FD T, MN R, KD S, AS G, SF M, S I. Origin of the dengue virus outbreak in Martin County, Florida, USA 2013. *Virol Reports* [Internet]. 2014 Mar 1 [cited 2021 Oct 7];1–2(C):2–8. Available from: <https://europepmc.org/articles/PMC4318122>
29. Bosch QA ten, Clapham HE, Lambrechts L, Duong V, Buchy P, Althouse BM, et al. Contributions from the silent majority dominate dengue virus transmission. *PLOS Pathog* [Internet]. 2018 May 1 [cited 2021 Sep 11];14(5):e1006965. Available from: <https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1006965>
30. Boyles SM, Mavian CN, Finol E, Ukhanova M, Stephenson CJ, Hamerlinck G, et al. Under-the-Radar Dengue Virus Infections in Natural Populations of *Aedes aegypti* Mosquitoes. *mSphere*. 2020 Apr 29;5(2).
31. Ferreira-De-Lima H, Dos P, Andrade S, Matsumiya Thomazelli L, Toledo Marrelli M, Urbinatti PR, et al. Silent circulation of dengue virus in *Aedes albopictus* (Diptera: culicidae) resulting from natural vertical transmission. Available from: <https://doi.org/10.1038/s41598-020-60870-1>
32. Stephenson CJ, Coatsworth H, Kang S, Lednicky JA, Dinglasan RR. Transmission Potential of Floridian *Aedes aegypti* Mosquitoes for Dengue Virus Serotype 4: Implications for Estimating Local Dengue Risk. *mSphere* [Internet]. 2021 Aug 25 [cited 2021 Sep 11];6(4). Available from: <https://doi.org/10.1128/mSphere>

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

33. Ayers JB, Xie X, Coatsworth H, Stephenson CJ, Waits M, Shi P-Y, et al. Ecology and transmission of a dengue virus serotype 4 identified in 1 wild *Aedes aegypti* in Florida 2. Available from: <https://doi.org/10.1101/2021.07.11.451177>
34. Cummings DAT, Schwartz IB, Billings L, Shaw LB, Burke DS. Dynamic effects of antibody-dependent enhancement on the fitness of viruses [Internet]. 2005. Available from: www.pnas.org/cgi/doi/10.1073/pnas.0507320102
35. Recker M, Blyuss KB, Simmons CP, Tinh Hien T, Wills B, Farrar J, et al. Immunological serotype interactions and their effect on the epidemiological pattern of dengue Hospital for Tropical Diseases, 190 Ben. Available from: <http://rspb.royalsocietypublishing.org>.
36. Gubler DJ. Dengue and Dengue Hemorrhagic Fever. *Clin Microbiol Rev*. 1998;11(3):480–96.
37. Sharp TM, Morris S, Morrison A, de Lima Corvino D, Santiago GA, Shieh W, et al. Fatal Dengue Acquired in Florida. *N Engl J Med*. 2021;(384):2257–9.
38. Mulligan K, Dixon J, Sinn C-LJ, Elliott SJ. Is dengue a disease of poverty? A systematic review.
39. Gubler DJ. Dengue, Urbanization and Globalization: The Unholy Trinity of the 21st Century. *Trop Med Health*. 2011;39(4):3–11.

Supplementary Information

Contamination troubleshooting

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

471 We noted issues with DENV-2 amplification in the no template and mosquito extraction negative
472 controls with the CDC DENV-2 serotype-specific RT-qPCR assay for a few of the mosquito
473 pools. We addressed these concerns using standard contamination troubleshooting protocols:
474 preparing all qPCR reactions in a PCR hood, deep cleaning qPCR preparation areas, creating
475 new primer and probe dilutions, and using different primer and probe stocks. We also tested each
476 assay reagent for inherent DENV-2 RNA/cDNA contamination but did not find any evidence of
477 contamination. Due to these issues, a new DENV-2 serotype specific assay was implemented
478 (see DENV2 Alm, Supplementary Table 5) targeting a different region of the DENV-2 E
479 glycoprotein. No amplification of negative controls was seen with the new primer-probe
480 combination.

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Table 1. Positive mosquito pool results from screening mosquitoes collected in Miami-Dade County (Florida, USA) in 2020. WNV = West Nile virus, DENV-2 = dengue virus serotype 2, DENV-4 = dengue virus serotype 4

Collection Date	Sample ID	Overall sample result	Species	Pool Size
06/09/20	N/A*	WNV positive	<i>Culex quinquefasciatus</i>	18
06/23/20		WNV positive	<i>Culex quinquefasciatus</i>	4
		WNV positive	<i>Culex quinquefasciatus</i>	24
06/30/20		WNV positive	<i>Culex quinquefasciatus</i>	5
07/23/20	0723-30	WNV positive	<i>Culex nigripalpus</i>	25
	0723-31	WNV positive	<i>Culex nigripalpus</i>	25
	0723-39	WNV positive	<i>Culex nigripalpus</i>	25
	0723-44	WNV positive	<i>Culex nigripalpus</i>	25
07/30/20	0730-37	DENV-2 positive	<i>Aedes aegypti</i>	25
	0730-41	DENV-2 positive	<i>Aedes aegypti</i>	25
08/20/20	0820-06	WNV positive	<i>Culex nigripalpus</i>	25
	0820-07	WNV positive	<i>Culex nigripalpus</i>	25
	0820-08	WNV positive	<i>Culex nigripalpus</i>	25
	0820-09	WNV positive	<i>Culex nigripalpus</i>	25
	0820-11	WNV positive	<i>Culex nigripalpus</i>	25
08/28/20	0828-2	DENV-2 positive	<i>Aedes aegypti</i>	23
	0828-10	DENV-2 positive	<i>Aedes aegypti</i>	25
09/17/20	0917-4	DENV-4 positive	<i>Aedes aegypti</i>	13
	0917-8	DENV-4 positive	<i>Aedes aegypti</i>	8
	0917-11	DENV-4 positive	<i>Aedes aegypti</i>	8
	0917-12	DENV-4 positive	<i>Aedes aegypti</i>	4
	0917-13	DENV-4 positive	<i>Aedes aegypti</i>	10
	0917-16	DENV-4 positive	<i>Aedes aegypti</i>	3
	0917-27	DENV-4 positive	<i>Aedes aegypti</i>	6
09/22/20	0922-2	DENV-4 positive	<i>Aedes aegypti</i>	5
09/29/20	0929-07	DENV-4 positive	<i>Aedes aegypti</i>	2

*Pathogen detection conducted by the CDC.

Figure 1. Spatial distribution of arbovirus positive mosquito pools (purple – DENV, orange – WNV). A and D: Positive pool spread throughout Miami-Dade County – DENV positive pools (A), or WNV positive pools (D). B and E: Positive pool spread overlayed on 2010 Miami-Dade population density (individuals/km²) – DENV positive pools (B), or WNV positive pools (E). C and F: Positive pool spread overlayed on 2010 Miami-Dade land use data – DENV positive pools (C) or WNV positive pools (F).

Figure 2. Spatial analysis of A: DENV positive pool distribution (purple) within traps containing DENV vectors (black), B: Distribution of traps containing *Aedes aegypti* (purple) or *Aedes albopictus* (pink), C: WNV positive pool distribution (orange) within traps containing WNV vectors (black), D: Distribution of traps containing *Culex quinquefasciatus* (dark orange), *Culex nigripalpus* (light orange), *Culex coronator* (yellow), or *Anopheles crucians* (brown).

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Appendix

For Appendix Tables 1, 2, and 5: Positive controls used were the following: DENV-1:NR-50530, DENV-2: NR-50531, DENV-3: NR-50532, DENV-4: NR-50533, and WNV: NR-50434. Limits of quantification (LOQs) were determined by standard curve analysis of diluted RNA positive controls

Appendix Table 1. Pan-dengue Real-Time RT-PCR assay (assay courtesy of an active collaboration with Dr. Remi Charrel (Aix Marseille University))

Description	Name	Sequence	Fluor/ Quencher	Position	Product size (bp)	Target	Limit of quantification (LOQ)
Dengue All 1 Forward	DenAll-1 For	AGGACYAGAGGTTAGAGGAGA		DENV1: 10590 - 10610 DENV2: 10579 - 10599 DENV3: 10564 - 10584 DENV4: 10506 - 10526			
Dengue All 1 Reverse	DenAll-1 Rev	CGYTCTGTGCCTGGAWTGAT		DENV1: 10697 - 10678 DENV2: 10685 - 10666 DENV3: 10479 - 10499 DENV4: 10611 - 10592	DENV1: 108 DENV2: 107 DENV3: 106 DENV4: 106	DENV sfRNA	DENV1: 30 copies/μL DENV2: 279 copies/μL DENV3: 66 copies/μL DENV4: 20.1 copies/μL
Dengue All 1 Probe	DenAll-1 Probe	ACAGCATATTGACGCTGGGARA GACC	FAM/BHQ- 1	DENV1: 10629 - 10654 DENV2: 10617 - 10642 DENV3: 10601 - 10626 DENV4: 10543 - 10568			
Dengue All 2 Forward	DenAll-2 For	GGACTAGAGGTTAGAGGAGACC CC		DENV1: 10591 - 10614 DENV2: 10580 - 10603 DENV3: 10565 -	DENV1: 81 DENV2: 80 DENV3: 79 DENV4: 79	DENV sfRNA	

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

				10588
				DENV4:
				10507 -
				10530
				DENV1:
				10671 -
				10651
				DENV2:
				10659 -
				10639
Dengue All	DenAll-2	GAGACAGCAGGATCTCTGGTC		DENV3:
2 Reverse	Rev			10643 -
				10623
				DENV4:
				10585 -
				10565
				DENV1:
				10633 -
				10650
				DENV2:
				10619 -
				10636
Dengue All	DenAll-2	AGCATATTGACGCTGGGA	FAM/BHQ-	DENV3:
2 Probe	Probe		1	10583 -
				10600
				DENV4:
				10545 -
				10562

Appendix Table 2. West Nile Virus Real-Time RT-PCR assay (courtesy of Dr. Thomas Unnasch, University of South Florida)

Description	Name	Sequence	Fluor/ Quencher	Position	Product size (bp)	Target	Limit of quantification (LOQ)
WNV A forward primer	10668F	CAGACCACGCTACGGC G		10668 - 10684	103	WNV 3'UTR	1.5 x 10 ⁻⁶ ng
WNV A reverse primer	10770R	CTAGGGCCGCGTGCGG		10770 - 10756			
WNV A probe	WNV A probe	CTGCGGAGAGTGCACT CTGCGAT	FAM/BHQ- 1	10691 - 10714			

Appendix Table 3. Reaction volumes for each assay using 1-step qRT-PCR using QuantaBio UltraPlex 1-Step ToughMix (4X) Low-ROX master mix. Volumes are representative of one 25µL reaction. Primer and probes were diluted to working solutions of 10µM for assay use. Final primer and probe concentrations/reaction are denoted by []

Assay	Reagent	Volume (in µL)
Dengue serotype specific – Dengue 1	Water	10.3
	4X Reaction mix	6.25
	Dengue 1 forward primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 1 reverse primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 1 probe [10µM]	0.45 [Final conc. 0.18µM]
	Template	3

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Dengue serotype specific – Dengue 2	Water	12.8
	4X Reaction mix	6.25
	Dengue 2 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 2 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 2 probe [10µM]	0.45 [Final conc. 0.18µM]
	Template	3
Dengue serotype specific – Dengue 3	Water	10.3
	4X Reaction mix	6.25
	Dengue 3 forward primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 3 reverse primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 3 probe [10µM]	0.45 [Final conc. 0.18µM]
	Template	3
Dengue serotype specific – Dengue 4	Water	12.8
	4X Reaction mix	6.25
	Dengue 4 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 4 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 4 probe [10µM]	0.45 [Final conc. 0.18µM]
	Template	3
West Nile Virus	Water	11.6
	4X Reaction mix	6.25
	WNV A forward primer [10µM]	0.9375 [Final conc. 0.375µM]
	WNV A reverse primer [10µM]	0.9375 [Final conc. 0.375µM]
	WNV A 4 probe [10µM]	0.3125 [Final conc. 0.125µM]
	Template	5
Dengue All	Water	7.75
	2X Reaction mix	6.25
	DENVAll-1 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-1 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-1 probe [10µM]	0.50 [Final conc. 0.2µM]
	DENVAll-2 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-2 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-2 probe [10µM]	0.50 [Final conc. 0.2µM]
	Template	5

512

513 **Appendix Table 4.** Reaction volumes for each assay using SuperScript™ III Platinum™ One-
514 Step qRT-PCR Kit. Volumes are representative of one 25µL reaction. Primer and probes were
515 diluted to working solutions of 10µM for assay use. Final primer and probe
516 concentrations/reaction are indicated []

Assay	Reagent	Volume (in µL)
Dengue serotype specific – Dengue 1	Water	3.55
	2X Reaction mix	12.5
	Dengue 1 forward primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 1 reverse primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 1 probe [10µM]	0.45 [Final conc. 0.18µM]
	RT/Platinum Taq	0.5
Dengue serotype specific – Dengue 2	Water	6.05
	2X Reaction mix	12.5
	Dengue 2 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 2 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 2 probe [10µM]	0.45 [Final conc. 0.18µM]
	RT/Platinum Taq	0.5
Dengue serotype specific –	Template	3
	Water	3.55

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Dengue 3	2X Reaction mix	12.5
	Dengue 3 forward primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 3 reverse primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 3 probe [10µM]	0.45 [Final conc. 0.18µM]
	RT/Platinum Taq	0.5
	Template	3
Dengue serotype specific – Dengue 4	Water	6.05
	2X Reaction mix	12.5
	Dengue 4 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 4 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	Dengue 4 probe [10µM]	0.45 [Final conc. 0.18µM]
	RT/Platinum Taq	0.5
	Template	3
West Nile Virus	Water	4.8125
	2X Reaction mix	12.5
	WNV A forward primer [10µM]	0.9375 [Final conc. 0.375µM]
	WNV A reverse primer [10µM]	0.9375 [Final conc. 0.375µM]
	WNV A 4 probe [10µM]	0.3125 [Final conc. 0.125µM]
	RT/Platinum Taq	0.5
	Template	5
Dengue All	Water	3.0
	2X Reaction mix	12.5
	DENVAll-1 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-1 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-1 probe [10µM]	0.50 [Final conc. 0.2µM]
	DENVAll-2 forward primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-2 reverse primer [10µM]	1.25 [Final conc. 0.5µM]
	DENVAll-2 probe [10µM]	0.50 [Final conc. 0.2µM]
	RT/Platinum Taq	0.5
	Template	3
Alm Dengue 2 serotype specific	Water	3.55
	2X Reaction mix	12.5
	Dengue 2 forward primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 2 reverse primer [10µM]	2.5 [Final conc. 1µM]
	Dengue 2 probe [10µM]	0.45 [Final conc. 0.18µM]
	RT/Platinum Taq	0.5
	Template	3

Appendix Table 5. Serotype-specific assays, including both CDC (Santiago et al., 2013, doi:10.1371/journal.pntd.0002311) and Alm dengue 2 serotype-specific assays (Alm et al., 2015, doi: 10.1186/s12879-015-1226-z). “Alm” is used to refer to the publication describing the assay primers and probe to differentiate it from the CDC DENV-2 target

Description	Name	Sequence	Fluor/ Quencher	Position	Product size (bp)	Target	Limit of quantification (LOQ)
Dengue 1 forward primer	D1-F_CDC	CAAAAGGAAGTCGYGCAATA		8936-8955			
Dengue 1 reverse primer	D1-R_CDC	CTGAGTGAATTCTCTCTGCTRAAC		9023-9047	112	DENV 1 NS5	3 copies/µL
Dengue 1 probe	D1-Pr_CDC	CATGTGGYTGAGGAGCRCGC	FAM/BH Q-1	8961-8979			
Dengue 2 forward primer	D2-F_CDC	CAGGCTATGGCACYGTCACGAT		1426-1447			
Dengue 2 reverse primer	D2-R_CDC	CCATYTGACAGCACCACATCTC		1482-1504	78	DENV 2 E	27.9 copies/µL

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Dengue 2 probe	D2-Pr_CDC	CTCYCCRAGAACGGGCCTC GACTTCAA	FAM/BH Q-1	1454- 1480			
Dengue 2 Alm forward primer	DENV2_Alm _F	GCAGAAACACAACATGGAAC RATAGT		1873- 1898			
Dengue 2 Alm reverse primer	DENV2_Alm _R	TGATGTAGCTGTCTCCRAAT GG		2050- 2071	199	DENV 2 E	27.9 copies/ μL
Dengue 2 Alm probe	DENV2_Alm _probe	TCAACATAGAAGCAGAACC	FAM/BH Q-1	2030- 2048			
Dengue 3 forward primer	D3-F_CDC	GGACTRGACACACGCACCCA		701-720	74	DENV 3 prM	66 copies/ μL
Dengue 3 reverse primer	D3-R_CDC	CATGTCTCTACCTTCTCGAC TTGYCT		749-775			
Dengue 3 probe	D3-Pr_CDC	ACCTGGATGTCGGCTGAAG GAGCTTG	FAM/BH Q-1	722-747			
Dengue 4 forward primer	D4-F_CDC	TTGTCCTAATGATGCTRGTC G		884-904			
Dengue 4 reverse primer	D4-R_CDC	TCCACCYGAGACTCCTTCCA		953-973	89	DENV 4 prM	20.1 copies/ μL
Dengue 4 probe	D4-Pr_CDC	TYCCTACYCCTACGCATCGC ATTCCG	FAM/BH Q-1	939-965			

522

523 Appendix Table 6. Summed categories of land use description types from the 2010 US Census

Land Use Description	Summed Category
Agriculture, Other (Exotic Birds, Monkeys, Research Facilities).	Agricultural
Fallow.	Agricultural
Farm Storage Areas (Storage Structures or Lots for Farm Implements).	Agricultural
Fish Farms (Includes Tropical Fish Aquariums, Fish and Alligator Farms).	Agricultural
Groves.	Agricultural
Horse Training and Stables.	Agricultural
Pasture (Grazing, Animal Farming, Dairy Farms and Animal Feed Lots), excluding Horse and Poultry.	Agricultural
Plant Nurseries (Includes Sod Farms and Ornamental Nurseries).	Agricultural
Poultry.	Agricultural
Row and Field Cropland.	Agricultural
Cemeteries.	Cemeteries
Marine commercial (includes private commercial [non-recreational] marinas and repair yards on public or private land).	Commercial
Office and/or Business and other services (ground level) / Residential (upper levels). Low-density < 15 dwellings per acre or 4 floors.	Commercial
Office Building.	Commercial
Office/Business/Hotel/Residential. Substantial components of each use present. Treated as any combination of the mentioned uses with a hotel as part of development.	Commercial
Sales and Services (Wholesale facilities, Spot commercial, strip commercial, neighborhood shopping centers/plazas). Excludes office facilities.	Commercial
Shopping Centers (Regional and Community).	Commercial
Colleges and Universities, Including Research Centers, Public and Private.	Educational
Private Schools, Including Playgrounds (K-12, Vocational Ed., Day Care and Child Nurseries).	Educational
Public Schools, Including Playgrounds (K-12, Vocational Ed., Day Care and Child Nurseries).	Educational
Antenna Arrays.	Industrial
Communications (Radio, TV, Cable, and Phone), excluding Antenna Arrays.	Industrial
Electric Power (Generator and Substation, and Service Yards).	Industrial

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Extraction, Excavation, Quarrying, Rock-Mining, excluding the resulting water body (see 917).	Industrial
Industrial Extensive.	Industrial
Industrial intensive, Commercial Condominium type of use.	Industrial
Industrial Intensive, heavy-light manufacturing, and warehousing-storage type of use.	Industrial
Industrial Intensive, Office type of use.	Industrial
Junk Yard.	Industrial
Major Approved Projects.	Industrial
Major Transmission Lines.	Industrial
Oil and Gas Storage (Tank Farms).	Industrial
Sewerage Treatment Plants.	Industrial
Solid Waste Disposal and Transfer (Includes Dumps, Solid Waste Land Fills, Resource Recovery Plants and Facilities, Trash Transfer Stations).	Industrial
Water Supply Plants.	Industrial
Wellfields.	Industrial
Canal right-of-way.	Marine
Coastal Water (Bay only) within Biscayne Bay National Park (Excluding Ocean Waters).	Marine
Coastal Water (Bay only) within the Biscayne Bay Urban Aquatic Preserve (Excluding Ocean Waters).	Marine
Coastal Water (Ocean only) within Biscayne National Park (Excluding Bay Waters).	Marine
Coastal Waters within Everglades National Park.	Marine
Inland water bodies (Lakes, Ponds, and Watercourses) associated with industrial areas, industrial parks, and new industrial development.	Marine
Inland water bodies (Lakes, Rock Pits) associated with extraction, excavation, quarrying and rock-mining activities.	Marine
Inland water bodies (Lakes, Watercourses) associated with residential developments.	Marine
Marina complexes (docks, piers, moorings, ramps, boat lifts and hoists, boat maintenance and repair, boat storage, fueling operations) for recreational craft located within Parks and Preserves and other small craft harbor complexes used primarily for recreation.	Marine
Other inland water bodies (Lakes, Ponds, Watercourses other than rivers and canals), including road borrow pits.	Marine
Remaining Bay Waters (Excluding Ocean).	Marine
Remaining Ocean Waters (Excluding Bay).	Marine
Rivers and Canals (Water).	Marine
Water Conservation Areas (Properties owned by SFWMD; Big Cypress national preserve, Everglades, and Francis Taylor WMA (also known as conservation areas 3A and 3B)).	Marine
Beaches.	Recreational
Biscayne National Park.	Recreational
County Operated Parks.	Recreational
Cultural (auditoriums, convention centers, exhibition centers, museums, art galleries, libraries).	Recreational
Everglades National Park.	Recreational
Golf courses, Public and Private.	Recreational
Municipal Operated Parks.	Recreational
Other Nature Preserves and Protected Areas (State Mangrove Preserves, Turkey Point Wilderness Area, Great Cypress Swamp Preserves, and acquired government owned EEL sites).	Recreational
Private Recreational Camps/Areas not associated with private Residential Developments (Boy Scout/Girl Scout Camps, Private Recreational Camps). Includes private tennis courts and pools that are part of the recreational complex.	Recreational
Private Recreational Facilities Associated with private Residential Developments, except marinas/yacht basins, includes landscape and open spaces associated to residential, commercial and office developments.	Recreational
Recreational Vehicle Parks/Camps.	Recreational
Sports Stadiums, Arenas, and Tracks.	Recreational
Mobile Home Parks and Permanent Mobile Homes.	Residential
Multi-Family, High Density (Over 25 DU/Gross Acre).	Residential
Multi-Family, Low-Density (Under 25 DU/Gross Acre).	Residential
Residential MF-- government-owned or government subsidized multi-family residential or elderly housing	Residential

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

Residential predominantly (condominium/ rental apartments with lower floors Office and/or Retail. High density > 15 dwelling units per ac, multi-story buildings (Generally more than 5 stories).	Residential
Residential SF--government-owned or government subsidized multi-family residential or elderly housing	Residential
Single-Family, High Density (Over 5 DU/Gross Acre, other than Townhouses, Duplexes and Mobile Homes).	Residential
Single-Family, Low-Density (Under 2 DU/Gross Acre).	Residential
Single-Family, Med.-Density (2-5 DU/Gross Acre).	Residential
Townhouses.	Residential
TRANSIENT-RESIDENTIAL (HOTEL-MOTEL)	Residential
Two-Family (Duplexes).	Residential
Governmental/Public Administration (Other than Military or Penal).	Commercial
Hospitals, clinics, medical offices and/or dental facilities	Commercial
Houses of Worship and Religious, and associated uses (parking, retreat houses, residencies, childcare, etc.).	Commercial
Military Facilities.	Industrial
Nursing homes, Assisted living facilities, and Adult congregate living quarters	Commercial
Penal and Correctional.	Commercial
Social Services, and Charitable institutions (Shrines, Elks, Moose, Lions Club).	Commercial
Airports (other than Military and Small Grass Airports).	Industrial
Bus/Truck/Freight Forwarding Terminals.	Industrial
Highways and Expressways right-of-way and associated open and landscaped areas excluding paved expressways and ramps.	Paved
Ocean Ship Terminals and Port Facilities, Bay and River Based.	Marine
Parking - Public and Private Garages and Lots.	Paved
Paved Highways, Expressways and Ramps.	Paved
Private Drives.	Paved
Railroads - Terminals, Trackage, and Yards.	Paved
Road Maintenance and Storage Yards, and Motor Pools.	Paved
Small Grass Airports (Includes Crop Dusting Activities).	Industrial
Street right-of-way and entrance features both public and private, and utility easements.	Paved
Streets and Roads, except Expressways and Private Drives.	Paved
TRANSPORTATION, COMMUNICATION, AND UTILITIES	Industrial
Vacant Government owned or controlled.	Vacant
Vacant, Non-Protected, Privately-Owned.	Vacant
Vacant, Protected, Government-Owned or controlled. EEL sites included	Vacant
Vacant, Protected, Privately-Owned. Proposed and designated EEL sites until acquired or protected under any other conservation or environmental mechanism.	Vacant

524

525 Appendix Table 7. WNV-positive and DENV-positive samples

Sample	Sample Designation	Replicate #	Ct value	Avg Ct	Assay Run	Positive Control (1:10)	Positive Control Avg Ct
0723-30	WNV positive	1	39.78		WNV Run 1	WNV	26.05
		2	41.77				26.01
		1	45.49		WNV Run 2	WNV	23.74
		2	43.48				23.77
0723-31	WNV positive	1	38.18		WNV Run 1	WNV	26.05
		2	36.56				26.01
		1	37.52		WNV Run 2	WNV	23.74
		2	42.73				23.77
0723-39	WNV positive	1	32.69		WNV Run 1	WNV	26.05

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

		2	33.11				26.01
		1	33.31		WNV Run 2	WNV	23.74
		2	32.99				23.77
0723-44	WNV positive	1	29.33		WNV Run 1	WNV	26.05
		2	29.5				26.01
		1	28.66		WNV Run 2	WNV	23.74
		2	28.6				23.77
0820-06	WNV positive	1	15.11		WNV Run 1	WNV	24.26
		2	15				24.42
		1	13.98		WNV Run 2	WNV	23.74
		2	14.08				23.77
0820-07	WNV positive	1	32.88		WNV Run 1	WNV	24.26
		2	32.75				24.42
		1	31.43		WNV Run 2	WNV	23.74
		2	31.22				23.77
0820-08	WNV positive	1	36.06		WNV Run 1	WNV	24.26
		2	38.35				24.42
		1	35.96		WNV Run 2	WNV	23.74
		2	36				23.77
0820-09	WNV positive	1	37.82		WNV Run 1	WNV	24.26
		2	37.02				24.42
		1	35.88		WNV Run 2	WNV	23.74
		2	34.74				23.77
0820-11	WNV positive	1	39.66		WNV Run 1	WNV	24.26
		2	41.72				24.42
		1	39.45		WNV Run 2	WNV	23.74
		2	38.42				23.77
		1	43.27		WNV Run 2	WNV	23.74
		2	Neg				23.77
0730-37	DENV-2 positive	1	Neg	N/A	DENV1 serotype specific Run 1	DENV-1	28.42
		2	Neg				28.47
		1	33.7	33.78	DENV2 serotype specific Run 1	DENV-2	25.06
		2	33.86				24.78
		1	Neg	N/A	DENV3 serotype specific Run 1	DENV-3	30.49
		2	Neg				30.38
		1	Neg	N/A	DENV4 serotype specific Run 1	DENV-4	27.88
		2	Neg				28.08
		1	37.47	36.62	DENV2 serotype specific Run 1	DENV-2	23.96
		2	35.77		specific Run 2		24.10
0730-41	DENV-2 positive	1	Neg	N/A	DENV1 serotype specific Run 1	DENV-1	28.42
		2	Neg				28.47
		1	27.21	27.205	DENV2 serotype specific Run 1	DENV-2	25.06
		2	27.20				24.78
		1	Neg	N/A	DENV3 serotype specific Run 1	DENV-3	30.49
		2	Neg				30.38
		1	Neg	N/A	DENV4 serotype specific Run 1	DENV-4	27.88
		2	Neg				28.08
		1	29.11	29.15	DENV2 serotype specific Run 1	DENV-2	23.96
		2	29.19		specific Run 2		24.10
0828-2	DENV-2 positive	1	38.97	38.97	DENVAll Run 1	DENV-1	28.90
							29.13
						DENV-2	24.75
							24.91
		2	Neg			DENV-3	32.37
							32.32
						DENV-4	28.83
							28.79
		1	Neg	N/A	DENV1 serotype specific Run 1	DENV-1	30.51
		2	Neg				30.82
		1	32.35	32.43	DENV2 serotype	DENV-2	27.63

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

		2	32.51		specific Run 1		27.47
		1	Neg	N/A	DENV3 serotype	DENV-3	32.47
		2	Neg		specific Run 1		32.61
		1	Neg	N/A	DENV4 serotype	DENV-4	29.74
		2	Neg		specific Run 1		29.74
		1	31.35	31.505	DENV2 serotype	DENV-2	27.12
		2	31.66		specific Run 2		26.78
0828-10	DENV-2 positive	1	Neg	37.87	DENVAll Run 1	DENV-1	28.90
							29.13
						DENV-2	24.75
							24.91
		2	37.87			DENV-3	32.37
							32.32
						DENV-4	28.83
							28.79
		1	Neg	N/A	DENV1 serotype	DENV-1	30.51
		2	Neg		specific Run 1		30.82
		1	31.45	31.455	DENV2 serotype	DENV-2	27.63
		2	31.46		specific Run 1		27.47
		1	Neg	N/A	DENV3 serotype	DENV-3	32.47
		2	Neg		specific Run 1		32.61
		1	Neg	N/A	DENV4 serotype	DENV-4	29.74
		2	Neg		specific Run 1		29.74
		1	30.43	30.485	DENV2 serotype	DENV-2	27.12
		2	30.54		specific Run 2		26.78
0917-4	DENV-4 positive	1	34.7	34.7	DENVAll Run 1	DENV-1	31.24
							31.02
						DENV-2	25.67
							26.01
		2	Neg			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22
		1	41.18	39.735	DENVAll Run 2	DENV-1	32.26
							32.37
						DENV-2	27.28
							27.57
		2	38.29			DENV-3	33.38
							34.10
						DENV-4	32.02
							32.28
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53
		1	37.46	37.745	DENV4 serotype	DENV-4	26.88
		2	38.03		specific Run 1		26.58
		1	Neg	N/A	DENV4 serotype	DENV-4	27.52
		2	Neg		specific Run 2		27.77
		1	35.03	36.34	DENV4 serotype	DENV-4	27.02
		2	37.65		specific Run 3		27.04
0917-8	DENV-4 positive	1	Neg	37.4	DENVAll Run 1	DENV-1	31.24
							31.02
						DENV-2	25.67
							26.01
		2	37.4			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

		1	42.04	42.32	DENVAll Run 2	DENV-1	32.26
							32.37
						DENV-2	27.28
							27.57
		2	42.6			DENV-3	33.38
							34.10
						DENV-4	32.02
							32.28
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53
		1	40.53	38.95	DENV4 serotype	DENV-4	26.88
		2	37.37		specific Run 1		26.58
		1	Neg	39.19	DENV4 serotype	DENV-4	27.52
		2	39.19		specific Run 2		27.77
		1	38.82	38.9	DENV4 serotype	DENV-4	27.02
		2	38.98		specific Run 3		27.04
0917-11	DENV-4 positive	1	33.14	33.12	DENVAll Run 1	DENV-1	31.24
							31.02
						DENV-2	25.67
							26.01
		2	33.09			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53
		1	38.09	37.96	DENV4 serotype	DENV-4	26.88
		2	37.82		specific Run 1		26.58
		1	Neg	40.38	DENV4 serotype	DENV-4	27.52
		2	40.38		specific Run 2		27.77
		1	37.16	36.98	DENV4 serotype	DENV-4	27.02
		2	36.80		specific Run 3		27.04
0917-12	DENV-4 positive	1	33.47	33.05	DENVAll Run 1	DENV-1	31.24
							31.02
						DENV-2	25.67
							26.01
		2	32.63			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53
		1	Neg	37.55	DENV4 serotype	DENV-4	26.88
		2	37.55		specific Run 1		26.58
		1	42.13	41.54	DENV4 serotype	DENV-4	27.52
		2	40.94		specific Run 2		27.77
		1	36.04	35.92	DENV4 serotype	DENV-4	27.02
		2	35.79		specific Run 3		27.04
0917-13	DENV-4 positive	1	32.84	33.42	DENVAll Run 1	DENV-1	31.24

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

						DENV-2	31.02
							25.67
		2	33.99			DENV-3	26.01
							30.19
						DENV-4	30.38
							29.49
							29.22
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53
		1	Neg	38.46	DENV4 serotype	DENV-4	26.88
		2	38.46		specific Run 1		26.58
		1	40.29	40.50	DENV4 serotype	DENV-4	27.52
		2	40.7		specific Run 2		27.77
		1	38.75	37.99	DENV4 serotype	DENV-4	27.02
		2	37.23		specific Run 3		27.04
0917-16	DENV-4 positive	1	35.95	37.09	DENVAll Run 1	DENV-1	31.24
							31.02
						DENV-2	25.67
							26.01
		2	38.23			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53
		1	36.57	36.57	DENV4 serotype	DENV-4	26.88
		2	Neg		specific Run 1		26.58
		1	39.41	40.94	DENV4 serotype	DENV-4	27.52
		2	42.46		specific Run 2		27.77
		1	35.05	35.50	DENV4 serotype	DENV-4	27.02
		2	35.94		specific Run 3		27.04
0917-27	DENV-4 positive	1	36.42	36.42	DENVAll Run 1	DENV-1	31.24
							31.02
						DENV-2	25.67
							26.01
		2	Neg			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22
		1	Neg	N/A	DENVAll Run 2	DENV-1	32.26
							32.37
						DENV-2	27.28
							27.57
		2	Neg			DENV-3	33.38
							34.10
						DENV-4	32.02
							32.28
		1	Neg	N/A	DENV1 serotype	DENV-1	31.84
		2	Neg		specific Run 1		31.97
		1	Neg	N/A	DENV2 serotype	DENV-2	28.28
		2	Neg		specific Run 1		26.43
		1	Neg	N/A	DENV3 serotype	DENV-3	30.75
		2	Neg		specific Run 1		30.53

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

		1	36	36.33	DENV4 serotype specific Run 1	DENV-4	26.88
		2	36.65				26.58
		1	Neg	38.29	DENV4 serotype specific Run 2	DENV-4	27.52
		2	38.29				27.77
		1	37.12	37.45	DENV4 serotype specific Run 3	DENV-4	27.37
		2	37.77				27.16
0922-2	DENV-4 positive	1	33.91	34.28	DENVAll Run 1	DENV-1	31.24
						DENV-2	31.02
							25.67
							26.01
		2	34.64			DENV-3	30.19
							30.38
						DENV-4	29.49
							29.22
		1	Neg	N/A	DENV1 serotype specific Run 1	DENV-1	31.84
		2	Neg				31.97
		1	Neg	N/A	DENV2 serotype specific Run 1	DENV-2	28.28
		2	Neg				26.43
		1	Neg	N/A	DENV3 serotype specific Run 1	DENV-3	30.75
		2	Neg				30.53
		1	36.11	35.72	DENV4 serotype specific Run 1	DENV-4	26.88
		2	35.33				26.58
		1	36.75	37.51	DENV4 serotype specific Run 2	DENV-4	27.52
		2	38.26				27.77
		1	35.08	35.17	DENV4 serotype specific Run 3	DENV-4	27.02
		2	35.25				27.04
0929-07	DENV-4 positive	1	Neg	38.87	DENVAll Run 1	DENV-1	31.11
						DENV-2	31.14
							26.75
							26.70
		2	38.87			DENV-3	32.47
							32.44
						DENV-4	30.46
							30.75
		1	Neg	40.23	DENVAll Run 2	DENV-1	32.26
							32.37
						DENV-2	27.28
							27.57
		2	40.23			DENV-3	33.38
							34.10
						DENV-4	32.02
							32.28
		1	Neg	N/A	DENV1 serotype specific Run 1	DENV-1	29.37
		2	Neg				29.47
		1	Neg	N/A	DENV2 serotype specific Run 1	DENV-2	28.28
		2	Neg				26.43
		1	Neg	N/A	DENV3 serotype specific Run 1	DENV-3	30.26
		2	Neg				30.06
		1	38.84	38.32	DENV4 serotype specific Run 1	DENV-4	27.37
		2	37.8				27.16
		1	39.59	38.27	DENV4 serotype specific Run 2	DENV-4	31.25
		2	36.94				31.34

526

527 **Appendix Figure 1.** Kernel density estimation (KDE) of uninfected vector populations and
528 arbovirus positive mosquito pools (purple – DENV, orange – WNV). A: KDE on traps that
529 collected DENV vectors, B: KDE on all DENV positive pools, C: KDE on all DENV-2 positive

Reserved space. Do not place any text in this section. Include the mandatory author checklist or your manuscript will be returned. Use *continuous* line numbering in your manuscript.

pools, D: KDE on all DENV-4 positive pools, E: KDE on traps that collected WNV vectors, F: KDE on WNV positive pools.

Appendix Figure 2. Yearly temperature and precipitation maps for Miami-Dade County in 2020 overlaid with arbovirus positive mosquito pools (purple – DENV, orange – WNV). A and D: Maximum average temperature for 2020 in Miami-Dade visualized with DENV positive pools (A) or WNV positive pools (D). B and E: Minimum average temperature for 2020 in Miami-Dade visualized with DENV positive pools (B) or WNV positive pools (E). C and F: Total precipitation in Miami-Dade in 2020 visualized with DENV positive pools (C) or WNV positive pools (F).

Appendix Figure 3. Local Moran's I, a local indicator of spatial association (LISA), analyses of imported DENV cases and PLWH overlaid with arbovirus positive mosquito pools (purple – DENV, orange – WNV). A: LISA from DENV imported cases (2009 – 2019) visualized with 2020 DENV positive pools, B: LISA from 2019 DENV imported cases visualized with 2020 DENV positive pools, C: LISA from PLWH visualized with 2020 DENV positive pools, D: LISA from PLWH visualized with 2020 WNV positive pools.

A

B

C

LEGEND

- DENV-2 Positive Pool
- DENV-4 Positive Pool
- WNV Positive Pool

Population Density

- ≤ 470
- ≤ 1245
- ≤ 2135
- ≤ 3682
- ≤ 7131

Land Use

- Agricultural
- Recreational
- Residential
- Commercial
- Educational
- Industrial
- Paved
- Cemeteries
- Vacant
- Marine
- Water

D

E

F

A **B** **C** **D**

A

B

C

D

E

F

A

B

C

D

E

F

A **B** **C** **D**