Multi-Omics Signatures of Alcohol Use Disorder in the Dorsal and Ventral Striatum

Lea Zillich¹ (MSc), Eric Poisel¹ (BSc), Josef Frank¹ (PhD), Jerome C. Foo¹ (PhD), Marion M. Friske² (MSc), Fabian Streit¹ (PhD), Lea Sirignano¹ (MSc), Stefanie Heilmann-Heimbach³ (PhD), André Heimbach³ (PhD), Per Hoffmann^{3,4} (PhD), Franziska Degenhardt⁵ (MD), Anita C. Hansson² (PhD), Georgy Bakalkin⁶ (PhD), Markus M. Nöthen³ (MD), Marcella Rietschel^{1#} (MD), Rainer Spanagel^{2#} (PhD), Stephanie H. Witt^{1,7#} (PhD)

¹ Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany

² Institute of Psychopharmacology, Central Institute of Mental Health, Medical Faculty Mannheim, University of Heidelberg, Mannheim, Germany

³ Institute of Human Genetics, University of Bonn, School of Medicine & University Hospital Bonn, Bonn, Germany

⁴ Department of Biomedicine, University of Basel, Basel, 4003, Switzerland

⁵ Department of Child and Adolescent Psychiatry, University Hospital Essen, University of Duisburg-Essen, Essen, Germany

⁶ Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden

⁷ Center for Innovative Psychiatric and Psychotherapeutic Research, Biobank, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany

#contributed equally

corresponding author:

Prof. Dr. Rainer Spanagel
Central Institute of Mental Health
Institute of Psychopharmacology
J5
68159 Mannheim
Germany
Phone: +49 621 1703 6251
Fax: +49 621 1703 6255
E-Mail: Rainer.Spanagel@zi-mannheim.de

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Alcohol Use Disorder (AUD) is a major contributor to global mortality and morbidity. Postmortem human brain tissue enables the investigation of molecular mechanisms of AUD in the neurocircuitry of addiction. We aimed to identify differentially expressed (DE) genes in the ventral and dorsal striatum between individuals with AUD and controls, and to integrate the results with findings from genome- and epigenome-wide association studies (GWAS/EWAS) to identify functionally relevant molecular mechanisms of AUD. DNA-methylation and gene expression (RNA-seq) data was generated from postmortem brain samples of 48 individuals with AUD and 51 controls from the ventral striatum (VS) and the dorsal striatal regions caudate nucleus (CN) and putamen (PUT). We identified DE genes using DESeq2, performed gene-set enrichment analysis (GSEA), and tested enrichment of DE genes in results of GWASs using MAGMA. Weighted correlation network analysis (WGCNA) was performed for DNA-methylation and gene expression data and gene overlap was tested. In the dorsal striatum, we discovered differential expression (FDR<0.05) for a total of 50 genes. In the VS, DE genes at FDR<0.25 were overrepresented in a recent GWAS of problematic alcohol use. The *ARHGEF15* gene was upregulated in all three brain regions. GSEA in CN and VS pointed towards cell-structure associated GO-terms and in PUT towards immune pathways. The WGCNA modules most strongly associated with AUD showed strong enrichment for immune response and inflammation pathways. Our integrated analysis of multi-omics data sets provides further evidence for the importance of immune-and inflammation-related processes in AUD.

22 Introduction

23 Alcohol Use Disorder (AUD) is a major contributor to the global disease burden, with a
 24 prevalence of ~17% among 12-month alcohol users in the US^{1, 2} and an estimated heritability
 25 of 49%³. Knowledge about the molecular mechanisms can foster understanding of causes and
 26 promote prevention. Recent genome-wide association studies (GWASs) have identified 29
 27 genetic loci associated with Problematic Alcohol Use (PAU), a proxy of AUD⁴. While GWASs
 28 identify increasing numbers of disease-associated loci, the functional interpretation of many
 29 of these findings remains inconclusive. Analyzing the transcriptome can extend the
 30 understanding of the molecular mechanisms underlying AUD, by identifying associated gene
 31 expression patterns. Findings can in turn be integrated with results from GWASs and
 32 epigenome-wide association studies (EWASs) to identify the pathomechanisms underlying
 33 disease.

34 Processes in the central nervous system are considered to play a major role in the etiology of
 35 addiction, and the transition from chronic alcohol consumption to AUD⁵. Therefore, it is of
 36 particular interest to examine molecular changes associated with addiction in brain tissue. So
 37 far, only few studies have been conducted in postmortem human brain tissue to identify
 38 transcriptional changes associated with AUD⁶⁻⁸. These studies mainly focused on the
 39 prefrontal cortex (PFC) one important part of the neurocircuitry of addiction^{9, 10}. The first
 40 transcriptome-wide study in the PFC found DE genes implicated in neuronal processes, such
 41 as myelination, neurogenesis, and neural diseases, as well as cellular processes, such as cell
 42 adhesion and apoptosis¹¹. In Brodmann Area 9 downregulation of calcium signaling pathways
 43 has been observed in individuals with AUD compared to controls¹². In the same study, a
 44 weighted gene co-expression analysis (WGCNA) pointed towards modules associated with
 45 AUD case/control status, which were enriched for nicotine and opioid signaling, as well as
 46 immune processes. Another study in the PFC (Brodmann Area 8) showed that co-expression
 47 networks associated with lifetime alcohol consumption were enriched for GWAS signals of
 48 alcohol dependence⁶.

Despite the importance of striatal regions in addiction processes, genome-wide human omics studies of these brain regions are still missing. The striatum is divided into the ventral striatum (VS), consisting of the nucleus accumbens and olfactory tubercle; and the dorsal striatum, comprising the caudate nucleus (CN) and putamen (PUT)¹³. The nucleus accumbens is involved in mediating motivational processes such as aversion and reward, which play a significant role in the development and maintenance of substance use disorders (SUD)¹³. In addition to regulating motor function, the CN and PUT are involved in cognitive processes relevant for addiction, such as executive functioning and cognitive control, reinforcement learning and habit formation¹⁴. Analyses of omics data from striatal regions could complement the knowledge on global molecular changes in the neurocircuitry of addiction in AUD.

In a recent EWAS of AUD in postmortem brain tissue, we identified differentially methylated CpG-sites and regions in the ventral and dorsal striatum¹⁵. Previous studies have shown the utility of integrating epigenetic and transcriptomic data in postmortem brain tissue of SUDs using weighted correlation network analysis (WGCNA)¹⁶. WGCNA clusters genes or CpG-sites into co-expressed or co-methylated modules based on correlation matrices. By relating modules to each other, WGCNA can be used for data integration, providing more insights than descriptive overlap. For example, whereas a descriptive comparison of histone H3 lysine 4 trimethylation (H3K4me3) and mRNA expression in individuals with AUD and cocaine use disorder revealed no consistent overlap between H3K4me3 trimethylation and gene expression¹⁷, a network analysis identified overlapping modules pointing towards co-expressed genes associated with H3K4me3 trimethylation⁶. Modules associated with AUD were enriched for CNS functions, such as synaptic transmission and regulation of neurogenesis⁶. WGCNA has also been used for integrating epigenetic and transcriptomic data and investigating their association with opioid use disorder (OUD) in postmortem human brain, identifying immune-related transcriptional regulation to be enriched in co-expressed and co-methylated modules¹⁸.

The aim of the present study was to investigate differential gene expression associated with AUD status in the ventral and dorsal striatum, relate these to GWAS findings, and to integrate

the findings with DNA-methylation data using a network approach (WGCNA) in order to identify functionally relevant molecular mechanisms in AUD.

Materials and Methods

Samples

Postmortem human brain tissue from CN, PUT and VS of a total of 48 individuals with AUD and 51 control individuals was obtained from the New South Wales Tissue Resource Centre at the University of Sydney. After quality control (QC), the sample sizes for each brain region ranged between 63 and 77. Phenotypic information was assessed by next-of-kin interviews. Inclusion criteria for this study were: age > 18 years, Western European Ancestry, no history of severe psychiatric or neurodevelopmental disorders, or SUDs other than AUD and nicotine use disorder. AUD was defined as meeting DSM-IV criteria for alcohol dependence and consuming 80g of alcohol a day or more (control group: <20g/day). Descriptive information can be found in Table 1 and Supplementary Table S1.

DNA extraction and Methylation Profiling

DNA extraction, methylation profiling, and QC was performed as described in Zillich, Frank¹⁵. In brief, DNA was extracted using the DNeasy extraction kit (Qiagen, Hilden, Germany); the Illumina HumanMethylation EPIC BeadChip and the Illumina HiScan array scanning system (Illumina, San Diego, CA) were used to determine DNA-methylation levels. We used an updated and customized version of the CPACOR pipeline to extract beta values from raw intensities¹⁹. Criteria for the removal of samples and probes can be found in Zillich, Frank¹⁵.

RNA extraction and -sequencing

RNA was extracted from frozen tissue according to the manufacturer's protocol using the Qiagen RNeasy microKit (Qiagen, Hilden, Germany). The RNA Integrity Number (RIN) of all samples was determined using a TapeStation 4200 (Agilent, Santa Clara, CA). RIN values of

273 samples were larger than 5.5, for which libraries were prepared using the TruSeq Stranded Total RNA Library Prep Kit (Illumina, San Diego, CA) and which were sequenced on the NovaSeq 6000 (Illumina) at the Life & Brain Center in Bonn, Germany with read lengths of 2x100bp and a sequencing depth of 62.5 M read pairs per sample on average. Technical replicates were sequenced for all but four samples.

Statistical Analyses

All analyses apart from QC and read mapping were performed using R version 3.6.1²⁰. An overview of the analysis workflow can be found in Figure 1.

Mapping and Quantification

Sequencing quality was determined using FastQC²¹ and 24 samples (11 cases and 13 controls) were excluded due to insufficient sequencing quality (e.g. strong overrepresentation of sequences, GC distribution). Raw reads were mapped to the human genome (hg38) using HISAT2 (v.2.1.0)²². Quantification was performed with the featureCounts function of the Rsubread package (v.2.0.1)²³, with hg38 annotation.

Differential Gene Expression Analysis

Differential gene expression was determined using DESeq2 (v.1.26.0)²⁴. Minimal pre-filtering was applied, removing genes with normalized counts <10 for more than two samples. Technical replicates were merged prior to differential expression analysis using the collapseReplicates function as implemented in DESeq2. For the differential gene expression analysis, we included age, sex, RIN, pH-value of the brain, and postmortem interval (PMI) as covariates, because of their known influence on gene expression²⁵⁻²⁷. Results were filtered for differentially expressed (DE) genes with an absolute log2 fold change larger than 0.02. P-values were adjusted for multiple testing using Benjamini-Hochberg correction (FDR<0.05)²⁸. All downstream analyses were performed for DE genes with FDR<0.25.

Pathway Analysis

Gene-set enrichment analysis was performed using the R package fgsea (v.1.12.0)²⁹, for which DE genes were ranked according to p-value. Enrichment analysis was performed for Gene-Ontology (GO) terms³⁰ and Hallmark genesets³¹ and the results were adjusted using FDR correction.

WGCNA

Weighted correlation network analyses (WGCNA, v.1.70-3)¹⁶ were performed to identify modules of co-expressed genes and co-methylated CpG-sites. We assessed the relationship of these modules with AUD case/control status and tested the overlap between associated modules. WGCNA clusters the input matrix according to a dynamic tree cutting algorithm, using a soft power threshold that approximates the criterion of scale-free topology ($R_{signed}^2 > 0.80$). Resulting soft power thresholds for expression networks were 6 for CN, 5 for PUT, and 14 for VS; for methylation networks, all power thresholds were 2.

To identify methylation networks associated with gene expression, beta values from normalized intensities of all samples from which gene expression data were available were filtered for promoter-associated CpG-sites based on the manufacturer's manifest (Illumina, San Diego, CA). The resulting 105 796 CpG-sites were used as input.

For the RNA-seq data, count matrices were normalized using the DESeq2 function normalizeCounts and variance stability transformation was applied.

Networks were constructed using following settings: minimum module size=30, mergeCutHeight=0.25, maxBlockSize=36 000. In WGCNA, modules are labeled using colors.

In the results section modules are labeled according to type of data, brain region, and color assigned in the analysis, e.g. "e-VS-pink" for module "pink" from the WGCNA analysis of gene expression data in the ventral striatum. For each module, its eigengene was calculated and correlated with AUD status. For modules associated with AUD status, we performed enrichment analysis using the GOenrichmentAnalysis function implemented in the WGCNA

package for expression data and the R package missMethyl (v.1.20.4)³² for methylation modules.

Expression and Methylation Data Integration

To identify genes both DE and differentially methylated, we analyzed the overlap of DE genes (FDR<0.25) with the results of an EWAS (p<0.001) in the same sample¹⁵. We prioritized CpG-sites based on their functional relevance in gene expression regulation. Thus, promoter-associated CpG-sites were used in the analysis.

At the module level, gene-set overlap tests were performed using the R package GeneOverlap (v.1.22.0)³³. Here, Fisher's exact test is used to identify significant overlap. For each brain region, the overlap of the AUD-associated co-expression and co-methylation modules was tested.

GWAS Enrichment Analysis

We analyzed enrichment of DE genes with an FDR<0.25, and genes in AUD-associated WGCNA modules in GWAS summary statistics using Multi-marker Analysis of GenoMic Annotation (MAGMA, v.1.08b)³⁴. We performed GWAS enrichment analysis for several SUDs, such as alcohol use disorder and problematic alcohol use⁴, cannabis use disorder³⁵, and a recent GWAS comparing individuals with opioid use disorder with unexposed controls³⁶. Results with p<0.05 were considered statistically significant.

Results

Differential Gene Expression

Gene expression analysis of postmortem brain tissue from AUD cases and controls revealed DE genes in all three investigated brain regions. DE genes were assessed at two different FDR thresholds (0.05 and 0.25). In the caudate nucleus, 1 326 DE genes were identified at FDR<0.25 (49 at FDR<0.05). Tubulin Tyrosine Ligase Like 4 (*TTLL4*, log2FC=0.11, p=2.3*10⁻

⁸) and GATA Binding Protein 2 (*GATA2*, log2FC=-0.27, p=8.6*10⁻⁷) were the most significantly upregulated and downregulated genes, respectively. In the putamen, 84 DE genes were detected at FDR<0.25 (1 at FDR<0.05). Top DE genes were found to be Transcription Elongation Factor A Like 2 (*TCEAL2*, log2FC=0.09, p=5.8*10⁻⁵) and Desmin (*DES*, log2FC=-0.86, p=2.6*10⁻⁶). One gene, cardiomyopathy associated 5 (*CMYA5*) showed an upregulation in both caudate nucleus and putamen. Nine genes were downregulated in both dorsal striatal regions, with *HLA-DOB* having the highest log2FC in both regions. In the ventral striatum, 5 genes were DE at FDR<0.25, but none at FDR<0.05. Strongest differential gene expression in the ventral striatum was observed for Ankyrin Repeat And Ubiquitin Domain Containing 1 (*ANKUB1*) which was upregulated in AUD cases (log2FC=1.35, p=5.8*10⁻⁵). The most significantly downregulated gene in the VS of AUD cases was Caseinolytic Mitochondrial Matrix Peptidase Chaperone Subunit B (*CLPB*, log2FC=-0.11, p=5.2*10⁻⁶). *ARHGEF15* (Rho Guanine Nucleotide Exchange Factor 15) was upregulated in all three brain regions at FDR<0.10. The Top 5 DE genes are listed in Table 2; DE genes significant at FDR<0.10 are listed in Supplementary Table S2 (CN), S3 (PUT), and S4 (VS). Overlap between DE genes in the different brain regions is shown in Figure 2A.

Pathway Analysis

Pathway analysis using a pre-ranked enrichment analysis revealed significant enrichment of dorsal striatum DE genes for several GO terms and Hallmark gene-sets. Genes in the CN were found to be related to cilia- and microtubule-associated GO-terms, while none of the Hallmark gene-sets was significantly enriched. GO-term and Hallmark gene-set analysis in PUT samples showed enrichment for immune processes, such as “acute inflammatory response to antigenic stimuli” (padj=0.006) and “adaptive immune response” (padj=0.006). In the VS the most significantly enriched GO-terms were also related to cilia and microtubules, as well as antigen processing. All GO-terms and Hallmark gene-set with FDR <0.10 are listed in Supplementary Tables S5 (CN), S6 (PUT), and S7 (VS).

201 WGCNA

202 Expression

203 In the CN, 21 modules with a median size of 352 genes (range: 64-7 259) were identified.
 204 Module “e-CN-magenta”, consisting of 328 genes, showed the strongest association with AUD
 205 status ($r=0.42$, $p=2.89 \times 10^{-4}$). In PUT module “e-PUT-black”, of the 25 modules (median size
 206 249 genes, range: 33-5 381) identified, was most strongly correlated with AUD ($r=0.41$,
 207 $p=2.31 \times 10^{-4}$). For expression data from the ventral striatum, 16 modules with a median size of
 208 429 genes (range: 35-9 708) were identified; module “e-VS-pink” had the strongest association
 209 with AUD ($r=0.41$, $p=0.009$). Interestingly, in a GO-term analysis the three AUD associated
 210 modules were all enriched for immune processes, such as “defense response” and
 211 “inflammation response”. There was also a wide overlap of the genes in the three modules:
 212 174 (22.54%) were partially shared between all three modules corresponding to the three brain
 213 regions, while another 21.76% were shared between at least two modules (Figure 2B).
 214 In the GWAS-enrichment analyses, the WGCNA modules “e-CN-magenta” and “e-PUT-black”
 215 were not enriched for signals from SUD-GWASs. “e-VS-pink” was enriched for genes
 216 associated with Cannabis Use Disorder ($p=0.043$).

217 Methylation

218 In the CN, WGCNA resulted in 36 modules with a median size of 346 CpG-sites (range: 66-
 219 41 423). Module “m-CN-red”, consisting of 2 117 CpG-sites, showed the strongest association
 220 with AUD case control status ($r=-0.27$, $p=0.021$). This module was most highly enriched for
 221 the biological processes “cell activation” ($p=1.52 \times 10^{-5}$) and “leukocyte activation” ($p=2.09 \times 10^{-5}$).
 222 In PUT 177 modules were identified (median size=57 CpG-sites, range: 30-42 248). Module
 223 “m-PUT-plum” consisted of 70 CpG-sites and was significantly associated with AUD
 224 case/control status ($r=-0.29$, $p=0.023$) and enriched for the biological processes “positive
 225 regulation of I- κ B kinase/NF- κ B signaling” ($p=0.002$) and “regulation of I- κ B kinase/NF- κ B
 226 signaling” ($p=0.005$). WGCNA in the VS methylation data resulted in 85 modules (median

size=178 CpG-sites, range: 35-30 370). The module with the strongest association with AUD was “m-VS-lavender” ($r=-0.29$, $p=0.023$), which consisted of 117 CpG-sites and was enriched for the molecular function “natural killer cell lectin-like receptor binding” ($p=3.43 \times 10^{-4}$) and the biological process “susceptibility to natural killer cell mediated cytotoxicity” ($p=3.65 \times 10^{-4}$). The top 10 enriched GO-terms for all AUD-associated modules can be found in Supplementary Tables S8-S10.

Expression and Methylation Data Integration

In the CN, 12 genes showed both differential methylation and differential gene expression (see Supplementary Table S11). No overlap was observed in the VS and PUT.

At the module-level, co-expression module “e-CN-magenta” showed significant overlap with the methylation modules “m-CN-red” ($p=0.003$) and “m-CN-midnightblue” ($p=0.014$) (Figure 2C), while expression module “e-CN-purple” did not show significant overlap with the methylation modules in CN. Of the 3 AUD-associated expression modules in the VS, only “e-VS-salmon” showed significant overlap with the methylation module “m-VS-turquoise” ($p=0.003$), but not “m-VS-lavender”. No overlap was observed for gene expression and DNA-methylation in PUT.

GWAS Enrichment Analysis

In the VS, but not in the dorsal striatum, we observed enrichment of DE genes in GWAS signal of PAU. In the dorsal striatum DE genes were enriched for GWAS signal from a study comparing individuals with OUD to unexposed controls. None of the DE genes in any of the brain regions showed enrichment for signals from a GWAS of Cannabis Use Disorder or AUD. Results from the respective analyses are depicted in Figure 3.

From the WGCNA modules showing the strongest association with AUD, only module e-VS-pink showed significant enrichment for GWAS signals of CUD. Results of this analysis can be found in Supplementary Table S12.

Discussion

In the present study, we identified DE genes, co-expression networks, and pathways associated with AUD in the dorsal and ventral striatum. The results were integrated with DNA-methylation data and results from GWASs of SUDs.

We discovered that one gene (*ARHGEF15*) was consistently upregulated in all investigated brain regions of AUD cases compared to controls. *ARHGEF15* encodes a specific guanine nucleotide exchange factor for the activation of Ras homolog family member A (RhoA), a GTPase, which has been linked to higher blood pressure and hypertension over the Rho/ROCK signaling cascade³⁷. It is postulated that the Rho Guanine Nucleotide Exchange Factor 15 negatively regulates excitatory synapse development by suppressing the synapse-promoting activity of EPHB2. EPHB2 deficiency has been linked to depression-like behaviors and memory impairments in animal studies³⁸. In line with this, genetic variation within *ARHGEF15* has been associated with hematocrit, red blood cell count, and hemoglobin concentration³⁹, but also with psychiatric traits, such as neuroticism and worries⁴⁰ as well as bipolar disorder⁴¹.

Among the genes that were downregulated in both dorsal striatal regions, *HLA-DOB* displayed the highest fold change. HLAs of the Major Histocompatibility Class II are an essential part of the acquired immune system presenting antigens to T-lymphocytes (for review: Howell, Carter⁴²). The most significantly downregulated gene in the VS is *CLPB*, a mitochondrial chaperone, which has been associated with progressive brain atrophy⁴³ and with the cellular response to alcohol-induced stress⁴⁴. In a recent GWAS, *CLPB* was associated with the amount of alcohol consumed on a typical day ($p=9.67 \times 10^{-5}$, $N=116\,163$)⁴⁵.

DE genes in the ventral striatum were enriched for GWAS signals of PAU, but not AUD. This could be a result of the larger sample size of the PAU GWAS, but also point towards differences in genetic variation as responsible for differential expression.

Our results from the pathway and network analyses further underline immune-related effects of chronic alcohol exposure; the pathway and network modules most strongly associated with

AUD case-control status were also enriched for immune system and inflammation processes. This was observed for all three brain regions, and both in expression and methylation data, providing further evidence for the important role of immune processes in AUD. These results strongly reflect the well-described effect of chronic alcohol exposure on different aspects of the innate and acquired immune systems⁴⁶. Chronic alcohol exposure accelerates the inflammatory response and reduces anti-inflammatory cytokines⁴⁶. An activated immune response in response to chronic alcohol exposure has been shown on the cell level⁴⁷, as well as on the transcription⁴⁷, and protein levels^{48, 49}. In a previous EWAS, we found strong enrichment of immune processes in differentially methylated CpG-sites associated with alcohol withdrawal⁵⁰. Neuroinflammation has been repeatedly associated with AUD and both the glutamate excitotoxicity and the production of acetaldehyde, key processes in AUD metabolism, have been suggested to produce an inflammatory response in the brain⁵¹. On a phenotypic level, there is also widespread overlap between symptoms of inflammation and of SUDs, such as anhedonia, depression, and decreased cognitive functioning⁵². In addition, in postmortem human brain studies in the PFC, hippocampus, and orbitofrontal cortex, increased mRNA levels of *HMGB1* encoding a proinflammatory cytokine and toll-like receptor genes have been associated with alcohol consumption in AUD cases, providing evidence for chronic neuroinflammation in response to alcohol⁵³⁻⁵⁵. Notably, there is an overlap of findings not only on the single-gene level but also on the level of pathways and networks/modules. This overlap underlines that alcohol consumption has common biological effects in different brain regions, i.e., most prominently, effects on immune and inflammation processes.

Several limitations apply to our study. First, we cannot distinguish between effects being a consequence of chronic alcohol consumption or addiction. Second, although we corrected for PMI, which can influence tissue quality as a confounding factor, it cannot be ruled out that other characteristics not easily accounted for, such as cause of death, or blood alcohol level for which the majority of individuals have missing data, influenced gene expression.

In summary, the present study provides further evidence from multi-omics data sets for the importance of immune-and inflammation-related processes in AUD. Notably, drugs that

309 reduce neuroinflammation to reduce drinking, such as phosphodiesterases, may be promising
310 approaches for novel treatment options for AUD. Recently published randomized controlled
311 trials suggest that a phosphodiesterase inhibitor reduces heavy drinking whereas an antibiotic
312 compound was not effective^{56, 57}. A deeper understanding of the underlying mechanisms will
313 enhance the discovery of drug targets and drive forward the development of precision
314 medicine within in this field.

Acknowledgments

The study was supported by the German Federal Ministry of Education and Research (BMBF), “A systems-medicine approach towards distinct and shared resilience and pathological mechanisms of substance use disorders” (01ZX01909), “Towards Targeted Oxytocin Treatment in Alcohol Addiction” (031L0190). ERA-NET: Psi-Alc (01EW1908), and the Deutsche Forschungsgemeinschaft (DFG) – Project-ID 402170461 – TRR 265⁵⁸.

We thank Elisabeth Röbel and Claudia Schäfer-Arnold for technical assistance.

Tissues were received from the New South Wales Brain Tissue Resource Centre at the University of Sydney which is supported by the University of Sydney. Research reported in this publication was supported by the National Institute of Alcohol Abuse and Alcoholism of the National Institutes of Health under Award Number R28AA012725. The content is solely the responsibility of the authors and does not represent the official views of the National Institutes of Health.

Conflict of Interest

The authors have nothing to disclose.

Supplementary Information

Supplementary information is available at MP’s website.

Data Availability

Raw data and summary statistics are available on request.

References

1. World Health Organization. *Global status report on alcohol and health 2018*. World Health Organization 2019.
2. Grant BF, Chou SP, Saha TD, Pickering RP, Kerridge BT, Ruan WJ *et al*. Prevalence of 12-Month Alcohol Use, High-Risk Drinking, and DSM-IV Alcohol Use Disorder in the United States, 2001-2002 to 2012-2013: Results From the National Epidemiologic Survey on Alcohol and Related Conditions. *JAMA Psychiatry* 2017; **74**(9): 911-923.
3. Verhulst B, Neale MC, Kendler KS. The heritability of alcohol use disorders: a meta-analysis of twin and adoption studies. *Psychological medicine* 2015; **45**(5): 1061.
4. Zhou H, Sealock JM, Sanchez-Roige S, Clarke T-K, Levey DF, Cheng Z *et al*. Genome-wide meta-analysis of problematic alcohol use in 435,563 individuals yields insights into biology and relationships with other traits. *Nature Neuroscience* 2020; **23**(7): 809-818.
5. Volkow ND, Koob GF, McLellan AT. Neurobiologic Advances from the Brain Disease Model of Addiction. *New England Journal of Medicine* 2016; **374**(4): 363-371.
6. Farris SP, Arasappan D, Hunnicke-Smith S, Harris RA, Mayfield RD. Transcriptome organization for chronic alcohol abuse in human brain. *Mol Psychiatry* 2015; **20**(11): 1438-1447.
7. Ponomarev I, Wang S, Zhang L, Harris RA, Mayfield RD. Gene coexpression networks in human brain identify epigenetic modifications in alcohol dependence. *J Neurosci* 2012; **32**(5): 1884-1897.
8. Liu J, Lewohl JM, Dodd PR, Randall PK, Harris RA, Mayfield RD. Gene expression profiling of individual cases reveals consistent transcriptional changes in alcoholic human brain. *J Neurochem* 2004; **90**(5): 1050-1058.
9. Koob GF, Volkow ND. Neurocircuitry of Addiction. *Neuropsychopharmacology* 2010; **35**(1): 217-238.
10. Noori HR, Spanagel R, Hansson AC. Neurocircuitry for modeling drug effects. *Addict Biol* 2012; **17**(5): 827-864.
11. Liu J, Lewohl JM, Harris RA, Iyer VR, Dodd PR, Randall PK *et al*. Patterns of gene expression in the frontal cortex discriminate alcoholic from nonalcoholic individuals. *Neuropsychopharmacology* 2006; **31**(7): 1574-1582.
12. Kapoor M, Wang JC, Farris SP, Liu Y, McClintick J, Gupta I *et al*. Analysis of whole genome-transcriptomic organization in brain to identify genes associated with alcoholism. *Transl Psychiatry* 2019; **9**(1): 89.
13. Volkow ND, Morales M. The Brain on Drugs: From Reward to Addiction. *Cell* 2015; **162**(4): 712-725.

14. Galandra C, Basso G, Cappa S, Canessa N. The alcoholic brain: neural bases of impaired reward-based decision-making in alcohol use disorders. *Neurol Sci* 2018; **39**(3): 423-435.
15. Zillich L, Frank J, Streit F, Friske MM, Foo JC, Sirignano L *et al*. Epigenome-wide Association Study of Alcohol Use Disorder in Five Brain Regions. *medRxiv* 2021: 2021.2008.2001.21261118.
16. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC bioinformatics* 2008; **9**(1): 559.
17. Zhou Z, Yuan Q, Mash DC, Goldman D. Substance-specific and shared transcription and epigenetic changes in the human hippocampus chronically exposed to cocaine and alcohol. *Proc Natl Acad Sci U S A* 2011; **108**(16): 6626-6631.
18. Liu A, Dai Y, Mendez EF, Hu R, Fries GR, Najera KE *et al*. Genome-Wide Correlation of DNA Methylation and Gene Expression in Postmortem Brain Tissues of Opioid Use Disorder Patients. *International Journal of Neuropsychopharmacology* 2021.
19. Lehne B, Drong AW, Loh M, Zhang W, Scott WR, Tan ST *et al*. A coherent approach for analysis of the Illumina HumanMethylation450 BeadChip improves data quality and performance in epigenome-wide association studies. *Genome Biol* 2015; **16**(1): 37.
20. Team RC. R: A language and environment for statistical computing. 2013.
21. Andrews S. FastQC: A quality control tool for high throughput sequencing data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>, 2010.
22. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* 2019; **37**(8): 907-915.
23. Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Research* 2019; **47**(8): e47-e47.
24. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* 2014; **15**(12): 550.
25. Durrenberger PF, Fernando S, Kashefi SN, Ferrer I, Hauw J-J, Seilhean D *et al*. Effects of Antemortem and Postmortem Variables on Human Brain mRNA Quality: A BrainNet Europe Study. *Journal of Neuropathology & Experimental Neurology* 2010; **69**(1): 70-81.
26. Trabzuni D, Ryten M, Walker R, Smith C, Imran S, Ramasamy A *et al*. Quality control parameters on a large dataset of regionally dissected human control brains for whole genome expression studies. *Journal of Neurochemistry* 2011; **119**(2): 275-282.
27. Birdsill AC, Walker DG, Lue L, Sue LI, Beach TG. Postmortem interval effect on RNA and gene expression in human brain tissue. *Cell and Tissue Banking* 2011; **12**(4): 311-318.

28. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: series B* 1995; **57**(1): 289-300.
29. Sergushichev AA. An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation. *bioRxiv* 2016: 060012.
30. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM *et al.* Gene Ontology: tool for the unification of biology. *Nature Genetics* 2000; **25**(1): 25-29.
31. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov Jill P, Tamayo P. The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Systems* 2015; **1**(6): 417-425.
32. Phipson B, Maksimovic J, Oshlack A. missMethyl: an R package for analyzing data from Illumina's HumanMethylation450 platform. *Bioinformatics* 2016; **32**(2): 286-288.
33. Shen L. GeneOverlap: Test and visualize gene overlaps. R package version 1.28.0, 2021.
34. de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. *PLOS Computational Biology* 2015; **11**(4): e1004219.
35. Johnson EC, Demontis D, Thorgeirsson TE, Walters RK, Polimanti R, Hatoum AS *et al.* A large-scale genome-wide association study meta-analysis of cannabis use disorder. *The Lancet Psychiatry* 2020; **7**(12): 1032-1045.
36. Polimanti R, Walters RK, Johnson EC, McClintick JN, Adkins AE, Adkins DE *et al.* Leveraging genome-wide data to investigate differences between opioid use vs. opioid dependence in 41,176 individuals from the Psychiatric Genomics Consortium. *Mol Psychiatry* 2020; **25**(8): 1673-1687.
37. Wirth A. Rho kinase and hypertension. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 2010; **1802**(12): 1276-1284.
38. Zhen L, Shao T, Luria V, Li G, Li Z, Xu Y *et al.* EphB2 Deficiency Induces Depression-Like Behaviors and Memory Impairment: Involvement of NMDA 2B Receptor Dependent Signaling. *Frontiers in Pharmacology* 2018; **9**(862).
39. Astle WJ, Elding H, Jiang T, Allen D, Ruklisa D, Mann AL *et al.* The Allelic Landscape of Human Blood Cell Trait Variation and Links to Common Complex Disease. *Cell* 2016; **167**(5): 1415-1429.e1419.
40. Nagel M, Jansen PR, Stringer S, Watanabe K, de Leeuw CA, Bryois J *et al.* Meta-analysis of genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic loci and pathways. *Nature Genetics* 2018; **50**(7): 920-927.
41. Ikeda M, Takahashi A, Kamatani Y, Okahisa Y, Kunugi H, Mori N *et al.* A genome-wide association study identifies two novel susceptibility loci and trans population polygenicity associated with bipolar disorder. *Mol Psychiatry* 2018; **23**(3): 639-647.
42. Howell WM, Carter V, Clark B. The HLA system: immunobiology, HLA typing, antibody screening and crossmatching techniques. *Journal of clinical pathology* 2010; **63**(5): 387-390.

43. Saunders C, Smith L, Wibrand F, Ravn K, Bross P, Thiffault I *et al.* CLPB Variants Associated with Autosomal-Recessive Mitochondrial Disorder with Cataract, Neutropenia, Epilepsy, and Methylglutaconic Aciduria. *The American Journal of Human Genetics* 2015; **96**(2): 258-265.
44. Tóth ME, Vigh L, Sántha M. Alcohol stress, membranes, and chaperones. *Cell Stress and Chaperones* 2014; **19**(3): 299-309.
45. Watanabe K, Stringer S, Frei O, Umičević Mirkov M, de Leeuw C, Polderman TJC *et al.* A global overview of pleiotropy and genetic architecture in complex traits. *Nat Genet* 2019; **51**(9): 1339-1348.
46. Szabo G, Saha B. Alcohol's Effect on Host Defense. *Alcohol research : current reviews* 2015; **37**(2): 159-170.
47. McClintick JN, Tischfield JA, Deng L, Kapoor M, Xuei X, Edenberg HJ. Ethanol activates immune response in lymphoblastoid cells. *Alcohol* 2019; **79**: 81-91.
48. Donnadieu-Rigole H, Mura T, Portales P, Duroux-Richard I, Bouthier M, Eliaou JF *et al.* Effects of alcohol withdrawal on monocyte subset defects in chronic alcohol users. *Journal of leukocyte biology* 2016; **100**(5): 1191-1199.
49. Yen CH, Ho PS, Yeh YW, Liang CS, Kuo SC, Huang CC *et al.* Differential cytokine levels between early withdrawal and remission states in patients with alcohol dependence. *Psychoneuroendocrinology* 2017; **76**: 183-191.
50. Witt SH, Frank J, Frischknecht U, Treutlein J, Streit F, Foo JC *et al.* Acute alcohol withdrawal and recovery in men lead to profound changes in DNA methylation profiles: a longitudinal clinical study. *Addiction (Abingdon, England)* 2020; **115**(11): 2034-2044.
51. de Timary P, Stärkel P, Delzenne NM, Leclercq S. A role for the peripheral immune system in the development of alcohol use disorders? *Neuropharmacology* 2017; **122**: 148-160.
52. Crews FT, Lawrimore CJ, Walter TJ, Coleman LG. The role of neuroimmune signaling in alcoholism. *Neuropharmacology* 2017; **122**: 56-73.
53. Crews FT, Qin L, Sheedy D, Vetreno RP, Zou J. High mobility group box 1/Toll-like receptor danger signaling increases brain neuroimmune activation in alcohol dependence. *Biol Psychiatry* 2013; **73**(7): 602-612.
54. Coleman LG, Jr., Zou J, Crews FT. Microglial-derived miRNA let-7 and HMGB1 contribute to ethanol-induced neurotoxicity via TLR7. *J Neuroinflammation* 2017; **14**(1): 22.
55. Vetreno RP, Qin L, Coleman Jr LG, Crews FT. Increased Toll-like Receptor-MyD88-NFkB-Proinflammatory neuroimmune signaling in the orbitofrontal cortex of human alcohol use disorder. *Alcoholism: Clinical and Experimental Research* 2021; **00**: 1-15.
56. Grodin EN, Bujarski S, Towns B, Burnette E, Nieto S, Lim A *et al.* Ibudilast, a neuroimmune modulator, reduces heavy drinking and alcohol cue-elicited neural activation: a randomized trial. *Translational Psychiatry* 2021; **11**(1): 355.

57. Petrakis IL, Ralevski E, Gueorguieva R, Sloan ME, Devine L, Yoon G *et al.* Targeting neuroinflammation with minocycline in heavy drinkers. *Psychopharmacology (Berl)* 2019; **236**(10): 3013-3021.
58. Heinz A, Kiefer F, Smolka MN, Endrass T, Beste C, Beck A *et al.* Addiction Research Consortium: Losing and regaining control over drug intake (ReCoDe)—From trajectories to mechanisms and interventions. *Addict Biol* 2020; **25**(2): e12866.

Figure Legends

Figure 1. Analysis workflow of the present study.

Figure 2. Venn Diagrams of gene overlap of A) DE genes at FDR < .25 in caudate nucleus (CN), putamen (PUT), and ventral striatum (VS), B) genes forming WGCNA expression-modules showing the strongest association with AUD status for CN, PUT, and VS, C) genes forming WGCNA expression-module “e-CN-magenta” and those forming the methylation-modules “m-CN-red” and “m-CN-midnightblue”.

Figure 3. Bar plots depicting the $-\log_{10}$ transformed p values of GWAS enrichment analysis for DE genes and the WGCNA expression and methylation modules showing the strongest association with AUD in all three brain regions. Color represents the GWAS: dark blue = alcohol use disorder (AUD), light blue = problematic alcohol use (PAU), grey = Cannabis use disorder (CUD) and red = opioid use disorder (OUD). Dotted line represents nominal significance.

Tables

Table 1.

Descriptive statistics of demographic data.

Characteristic	Cases	Controls	<i>p</i>
N	48	51	
Age, years	55.58 (10.62)	57 (10.64)	0.51
Sex (M/F)	31/17	37/14	
pH-value	6.53 (0.26)	6.65 (0.25)	0.026*
PMI (hours)	37.07 (15.79)	30.7 (15.57)	0.047*
Blood Alcohol level (N)	7	0	
Blood Alcohol Level (g/100ml)	0.21 (0.21)		
Samples per Brain Region			
Caudate Nucleus	36	37	
Putamen	35	42	
Ventral Striatum	31	32	

Data are presented as count (n/n; n (%)) or mean ($\pm$ SD), PMI: post-mortem interval, pH: pH-value of the brain, p: p-value of t-Test comparing cases and controls.

**significant difference between cases and controls*

Table 2.

Entrez Gene ID	Gene Name	baseMean	log2(FC)	lfcSE	Stat	P-Value	FDR
Caudate Nucleus							
9654	<i>TTLL4</i>	1125.89	0.11	0.02	5.59	2.33*10 ⁻⁸	0.0005
2624	<i>GATA2</i>	51.17	-0.27	0.05	-4.92	8.58*10 ⁻⁷	0.0091
25904	<i>CNOT10</i>	695.68	0.06	0.01	4.84	1.27*10 ⁻⁶	0.0091
222256	<i>CDHR3</i>	1483.68	0.19	0.04	4.75	1.99*10 ⁻⁶	0.0106
375611	<i>SLC26A5</i>	63.80	0.28	0.06	4.62	3.81*10 ⁻⁶	0.0163
Putamen							
1674	<i>DES</i>	22.06	-0.86	0.18	-4.70	2.64*10 ⁻⁶	0.0486
2050	<i>EPHB4</i>	118.04	-0.19	0.05	-4.19	2.78*10 ⁻⁵	0.0939
9144	<i>SYNGR2</i>	499.84	-0.20	0.05	-4.19	2.76*10 ⁻⁵	0.0939
55741	<i>EDEM2</i>	348.56	-0.07	0.02	-4.23	2.30*10 ⁻⁵	0.0939
84245	<i>MRI1</i>	662.19	-0.13	0.03	-4.13	3.57*10 ⁻⁵	0.0939
Ventral Striatum							
81570	<i>CLPB</i>	1188.85	-0.11	0.02	-4.56	5.16*10 ⁻⁶	0.0653
22899	<i>ARHGEF15</i>	54.23	-0.26	0.06	-4.49	7.04*10 ⁻⁶	0.0653
55584	<i>CHRNA9</i>	5.58	-1.17	0.26	-4.44	9.14*10 ⁻⁶	0.0653
100463488	<i>MTRNR2L10</i>	4.91	-2.07	0.50	-4.16	3.23*10 ⁻⁵	0.1730
389161	<i>ANKUB1</i>	30.26	1.35	0.34	4.02	5.79*10 ⁻⁵	0.2480

Top 5 differentially expressed genes in caudate nucleus, putamen and ventral striatum.

Figures