Simulation and predictions of a new COVID-19 pandemic wave in Ukraine with the use of generalized SIR model

Igor Nesteruk

Institute of Hydromechanics, National Academy of Sciences of Ukraine, Kyiv, Ukraine

Igor Sikorsky Kyiv Polytechnic Institute, Kyiv, Ukraine.
inesteruk@yahoo.com

ABSTRACT

A new wave of the COVID-19 pandemic in Ukraine, which began in the summer of 2021, was characterized by almost exponential growth of smoothed daily numbers of new cases. This is a matter of great concern and the need to immediately predict the dynamics of the epidemic in order to assess the possible maximum values of new cases, the risk of infection and the number of deaths. The generalized SIR-model and corresponding parameter identification procedure was used to simulate and predict the dynamics of this epidemic wave. The new COVID-19 epidemic wave in Ukraine will begin to subside in mid-October 2021, but its duration will be quite long. Unfortunately, new cases may appear by the summer of 2022.

Keywords: COVID-19 pandemic, epidemic dynamics in Ukraine, mathematical modeling of infection diseases, SIR model, parameter identification, statistical methods.

Introduction

The COVID-19 pandemic dynamics in Ukraine was discussed in [1-14]. To predict the first wave of the pandemic, the classical SIR model [15-17] and the statistics-based method of its parameter identification [18] were used. To simulate new epidemic waves, a numerical method of their detection [3, 19], a generalized SIR-model [20] and a corresponding parameter identification procedure [21] were developed. In particular, eleven epidemic waves in Ukraine were calculated [4, 7-10].

The calculations of the 11-th pandemic wave (based on the accumulated numbers cases reported by Ukrainian national statistics [22, 23] in the period May 23 – June 5, 2021) predicted the end of this wave on August 25, 2021 with the number of cases 2,226,797 (see [10]). As of August 25, 2021 the real number of cases accumulated in Ukraine was 2,278,171 (see Table 1). It means that the predicted saturation level was exceeded only 2.26% (after 81 days of observation). The obtained high accuracy of the method allows us to hope for a fairly accurate forecast for the next 12th epidemic wave in Ukraine,

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Data

We will use the data set regarding the accumulated numbers of laboratory-confirmed COVID-19 cases in Ukraine from national sources [22, 23]. The corresponding numbers V_j and moments of time t_j (measured in days) are shown in Table 1. The values for the period T_{c11} : May 23 – June 5, 2021 have been used in [10] for SIR simulations of the tenth epidemic wave in Ukraine. Here we use the dataset, corresponding to the period T_{c12} : September 29 – October 12, 2021 to simulate the 12-th wave. Other V_j and t_j values will be used to control the accuracy of predictions.

Table 1. Cumulative numbers of laboratory-confirmed Covid-19 cases in Ukraine V_j in the summer and autumn of 2021 according to the national statistics, [22, 23].

Day in corresponding month of 2021	Number of cases in May, V_j	Number of cases in June, V_j	Number of cases in July, V_j	Number of cases in August, V_j	Number of cases in September, V_j	Number of cases in October, V_j
1	2083180	2206836	2236497	2253534	2290848	2447222
2	2085938	2209417	2237202	2254361	2293541	2455189
3	2088410	2211683	2237579	2255345	2296155	2460010
4	2090986	2213580	2237823	2256397	2297534	2469856
5	2097024	2214517	2238364	2257478	2298307	2482518
6	2105428	2215052	2238974	2258532	2300504	2497643
7	2114138	2216654	2239591	2259151	2303276	2514005
8	2119510	2218039	2240246	2259451	2306939	2529913
9	2122327	2219824	2240753	2260232	2310554	2541257
10	2124535	2221427	2241043	2261354	2314423	2550089
11	2129073	2222701	2241217	2262601	2316619	2562085
12	2135886	2223558	2241698	2263864	2317824	2578394
13	2143448	2223978	2242245	2265217	2321156	-
14	2150244	2224992	2242868	2265912	2325796	-
15	2153864	2226037	2243605	2266329	2331540	-
16	2156000	2227225	2244196	2267219	2338164	-
17	2160095	2228192	2244495	2268666	2344398	-
18	2165233	2229044	2244677	2270226	2348381	-
19	2170398	2229523	2245275	2271826	2350646	-
20	2175382	2229846	2245930	2273558	2355805	-
21	2179988	2230142	2246656	2274561	2362559	-
22	2182521	2230977	2247419	2275171	2370425	-
23	2183855	2231914	2248164	2275863	2379483	-
24	2186463	2232790	2248450	2276590	2387750	-
25	2189858	2233546	2248663	2278171	2392397	-
26	2193367	2233996	2249344	2280203	2395404	-
27	2196673	2234281	2250061	2282285	2401956	-
28	2199769	2234463	2250907	2284191	2411622	-
29	2201472	2235096	2251869	2284940	2423379	-
30	2202494	2235801	2252785	2286296	2435413	-
31	2204631	-	2253269	2288371	-	-

To estimate the mortality rate (ratio of accumulated number of cases V_j to the accumulated number of deaths D_j), let us take figures D_j corresponding different days: 52,286 (June 26, 2021); 52,665 (July 13, 2021); 52,981 (August 2, 2021); 53,789 (August 30, 2021); 54,550 (September 15, 2021); 57,526 (October 5, 2021); 59,052 (October 11, 2021), [22, 23]. Taking corresponding V_j values from Table 1, we can calculate the mortality rates $m_i = D_j * 1000 / V_j$ (per thousand of cases) for the listed days: 23.40; 23.49; 23.50; 23.52; 23.40; 23.17; 23.05. Thus the mortality rate is rather stable (its variation during June to October 2021 is only 0.47). We will use the average value $m = 23.36$ to predict the number of deaths in Ukraine during the new 12-th pandemic wave.

Generalized SIR model and data smoothing procedure

The generalized SIR-model relates the number of susceptible S , infectious I and removed persons R for a particular epidemic wave i [8, 20]. The exact solution of the set of non-linear differential equations uses the function

$$V(t) = I(t) + R(t), \quad (1)$$

corresponding to the number of victims or the cumulative confirmed number of cases [8, 20]. Its derivative:

$$\frac{dV}{dt} = \alpha_i SI \quad (2)$$

yields the estimation of the average daily number of new cases. When the registered number of victims V_j is a random realization of its theoretical dependence (1), the exact solution presented in [8, 20] depends on five parameters (α_i is one of them). The details of the optimization procedure for their identification can be found in [21].

Since daily numbers of new cases are random and characterized by some weekly periodicity, we will use the smoothed daily number of accumulated cases:

$$\bar{V}_i = \frac{1}{7} \sum_{j=i-3}^{j=i+3} V_j, \quad (3)$$

and its numerical derivative:

$$\left. \frac{d\bar{V}}{dt} \right|_{t=t_i} \approx \frac{1}{2} (\bar{V}_{i+1} - \bar{V}_{i-1}) \quad (4)$$

to estimate the smoothed number of new daily cases [3, 4, 9, 19].

Results and discussion

The optimal values of the general SIR model and other characteristics of the 12-th pandemic wave in Ukraine are calculated and listed in Table 2 (middle column). The same characteristics of the 11-th wave calculated in [10] are presented in the last column of Table 2 for comparison. The corresponding SIR curves are shown in Figure by blue and black lines. It can be seen that the optimal values of SIR parameters are very different for 11-th and 12-th pandemic waves. Close values were obtained only for the average times of spreading the infection $1/\rho_{11}=4.1$ days (in comparison with $1/\rho_{12}=3.3$ days for the twelfth wave). The assessment of the 12-th epidemic wave duration corresponding the moment when the number of infectious persons becomes less that unit) is very pessimistic (June 16, 2022). A similar long epidemic wave was also predicted for India [24].

Table 2. Optimal values of parameters and other characteristics of the eleventh and twelfth COVID-19 pandemic wave in Ukraine.

Characteristics	12 th epidemic wave, $i=12$	11 th epidemic wave, $i=11$, [10]
Time period taken for calculations T_{ci}	September 29 – October 12, 2021	May 23 – June 5, 2021
I_i	25,261.4089164122	10,190.8327995721
R_i	2,399,050.73394073	2,180,167.16720043
N_i	3,790,400	2,258,464
ν_i	1,137,541.61656928	60,891.3982283695
α_i	2,63670321714285e-07	3.98540619731056e-06
ρ_i	0.299935964004210	0.242676955862249
$1/\rho_i$	3.33404499630449	4.12070440082342
r_i	0.997536473683354	0.996838194153353
$S_{i\infty}$	845,264	31,667
$V_{i\infty}$	2,945,136	2,226,797
Final day of the epidemic wave	June 16, 2022	August 25, 2021

The saturation level (final size) of the 12-th wave can approach 3 millions. As of the end of 2021 the accumulated numbers of laboratory-confirmed cases and deaths can exceed 2,943,700 and 68,764, respectively (see blue and magenta solid lines in Figure). The numbers of infectious persons and average daily new cases (see blue dashed and dotted lines) will stop to increase around 17 and 14 October 2021, respectively with the maximal values of 45,545 and 13,930, respectively (see blue dashed and dotted lines). As of the end of 2021 these values are expected to be around 505 and 113, respectively (see blue dashed and dotted lines).

To illustrate the accuracy of simulations the accumulated number of cases (red “stars”) and averaged daily numbers of new cases (eq. (4), red “crosses”) are also presented in Figure. Comparisons with corresponding solid and dotted lines show that the theoretical curves are in good agreement with the observations of the pandemic dynamics (markers). To estimate the accumulated number of deaths during the 12-th wave, we have multiplied the theoretical $V(t)=I(t)+R(t)$ values by the coefficient $m=23.36$ calculated before. The magenta line represents expected accumulated numbers of deaths multiplied by 10.

Figure. 11-th and 12-th waves of the COVID-19 pandemic in Ukraine in the summer and autumn of 2021.

The results of SIR simulations of the eleventh [10] and twelfth waves are shown by black and blue lines respectively. Numbers of victims $V(t)=I(t)+R(t)$ – solid lines; numbers of infected and spreading $I(t)$ multiplied by 5 – dashed; derivatives dV/dt (eq. (2)) multiplied by 100 – dotted. The magenta line represents the estimation of the accumulated number of deaths during the 12-th epidemic wave multiplied by 10. Black and blue “circles” correspond to the accumulated numbers of cases registered during the periods of time taken for SIR simulations (May 23 – June 5, 2021) and September 29 – October 12, 2021, respectively. “Stars” corresponds to V_j values beyond these time periods. The red “crosses” show the first derivative (4) multiplied by 100.

Conclusions

The new COVID-19 epidemic wave in Ukraine will begin to subside in mid-October 2021, but its duration will be quite long. Unfortunately, new cases may appear by the summer of 2022.

Acknowledgements

The author is grateful to Oleksii Rodionov for his help in collecting and processing data.

References

1. Nesteruk I., Kydybyn I., Demelmair G. GLOBAL STABILIZATION TRENDS OF COVID-19 PANDEMIC. KPI Science News. 2020. No. 2. pp. 55-62. DOI: 10.20535/kpi-sn.2020.2.205124
2. Nesteruk I. Simulations and predictions of COVID-19 pandemic with the use of SIR model. Innov Biosyst Bioeng. 2020. vol. 4. no. 2. 110–121. doi: 10.20535/ibb.2020.4.2.204274. <http://ibb.kpi.ua/article/view/204274>
3. Nesteruk I. Coronasummer in Ukraine and Austria. [Preprint.] ResearchGate. 2020 June. DOI: 10.13140/RG.2.2.32738.56002
4. Nesteruk I. *COVID-19 pandemic dynamics*. Springer Nature. 2021. DOI: 10.1007/978-981-33-6416-5
5. <https://link.springer.com/book/10.1007/978-981-33-6416-5>
5. Pardhan S., Drydak N. Associating the Change in New COVID-19 Cases to GDP per Capita in 38 European Countries in the First Wave of the Pandemic. Front Public Health. 2021 Jan 20;8:582140. doi: 10.3389/fpubh.2020.582140.
6. Chintala S., Dutta R., Tadmor D. COVID-19 spatiotemporal research with workflow-based data analysis. Infect Genet Evol. 2021 Mar;88:104701. doi: 10.1016/j.meegid.2020.104701.
7. Nesteruk I., Benlagha N. PREDICTIONS OF COVID-19 PANDEMIC DYNAMICS IN UKRAINE AND QATAR BASED ON GENERALIZED SIR MODEL. Innov Biosyst Bioeng. 2021. vol. 5. no. 1. pp. 37–46. DOI: 10.20535/ibb.2021.5.2.230487 <http://ibb.kpi.ua/article/view/230487>
8. Nesteruk I. Visible and real sizes of new COVID-19 pandemic waves in Ukraine Innov Biosyst Bioeng. 2021. vol. 5. no. 2. pp. 85–96. DOI: 10.20535/ibb.2021.5.2.230487 <http://ibb.kpi.ua/article/view/230487>
9. Nesteruk I. Detections and SIR simulations of the COVID-19 pandemic waves in Ukraine. Comput. Math. Biophys. 2021;9:46–65. <https://doi.org/10.1515/cmb-2020-0117>
10. Nesteruk I. Will a natural collective immunity of Ukrainians restrain new COVID-19 waves? MedRxiv, July 2021. DOI: [10.1101/2021.07.20.21260840](https://doi.org/10.1101/2021.07.20.21260840)
11. Nesteruk I, Rodionov O. The impact of demographic factors on the accumulated number of COVID-19 cases per capita in Europe and the regions of Ukraine in the summer of 2021. medRxiv, July 2021. DOI: [10.1101/2021.07.04.21259980](https://doi.org/10.1101/2021.07.04.21259980)
12. Nesteruk I, Rodionov O, Nikitin AV. The impact of seasonal factors on the COVID-19 pandemic waves. medRxiv, August 2021. DOI: [10.1101/2021.08.06.21261665](https://doi.org/10.1101/2021.08.06.21261665)

13. Nesteruk I. Comparison of the COVID-19 pandemics dynamics in Ukraine and Israel in the summer of 2021. Research Gate, September 2021.
https://www.researchgate.net/publication/354497030_Comparison_of_the_COVID-19_pandemics_dynamics_in_Ukraine_and_Israel_in_the_summer_of_2021
14. Nesteruk I. Effects of testing and vaccination levels on the dynamics of the COVID-19 pandemic and the prospects for its termination. MedRxiv, September 2021. DOI:
<https://doi.org/10.1101/2021.09.20.21263823>
15. Kermack WO, McKendrick AG. A Contribution to the mathematical theory of epidemics. J Royal Stat Soc Ser A. 1927;115:700-21.
16. Murray JD. Mathematical Biology I/II. New York: Springer; 2002.
17. Langemann D, Nesteruk I, Prestin J. Comparison of mathematical models for the dynamics of the Chernivtsi children disease. Mathematics in Computers and Simulation. 2016;123:68-79. DOI: 10.1016/j.matcom.2016.01.003
18. Nesteruk I. Statistics based models for the dynamics of Chernivtsi children disease. Naukovi Visti NTUU KPI. 2017;5:26-34. DOI: 10.20535/1810-0546.2017.5.108577
19. Nesteruk I. Identification of the New Waves of the COVID-19 Pandemic. In book: COVID-19 Pandemic Dynamics, Springer Nature, 2021. DOI: 10.1007/978-981-33-6416-5_8.
https://link.springer.com/chapter/10.1007/978-981-33-6416-5_8
20. Nesteruk I. General SIR Model and Its Exact Solution. In book: COVID-19 Pandemic Dynamics, Springer Nature, 2021. DOI: 10.1007/978-981-33-6416-5_9.
https://link.springer.com/content/pdf/10.1007%2F978-981-33-6416-5_9.pdf
21. Nesteruk I. Procedures of Parameter Identification for the Waves of Epidemics. In book: COVID-19 Pandemic Dynamics, Springer Nature, 2021. DOI: 10.1007/978-981-33-6416-5_10.
https://link.springer.com/chapter/10.1007%2F978-981-33-6416-5_10
22. Coronavirus in Ukraine - Statistics - Map of infections, graphs [Internet]. Index.minfin.com.ua. 2021. <https://index.minfin.com.ua/ua/reference/coronavirus/ukraine/>
23. Cabinet of Ministers of Ukraine – Home [Internet]. <https://www.kmu.gov.ua/>
24. Nesteruk I. The COVID-19 pandemic storm in India. [Preprint] medRxiv Posted May 08, 2021. doi: <https://doi.org/10.1101/2021.05.06.21256523>