INFERRING THE COVID-19 INFECTION FATALITY RATE IN THE COMMUNITY-DWELLING POPULATION: A SIMPLE BAYESIAN EVIDENCE SYNTHESIS OF SEROPREVALENCE STUDY DATA AND IMPRECISE MORTALITY DATA

Harlan Campbell

Department of Statistics, University of British Columbia
Vancouver, British Columbia, Canada
harlan.campbell@stat.ubc.ca
ORCID ID: 0000-0002-0959-1594

Paul Gustafson

Department of Statistics, University of British Columbia
Vancouver, British Columbia, Canada
ORCID ID: 0000-0002-2375-5006

October 20, 2021

Acknowledgments - *This work was supported by the European Union's Horizon 2020 research and innovation programme under ReCoDID grant agreement No 825746 ( ) and by the Canadian Institutes of Health Research, Institute of Genetics (CIHR-IG) under Grant Agreement No 01886-000. We wish to thank Andrew Levin, Thomas Debray, Valentijn de Jong, and Thomas Jaenisch for their valuable feedback. We also wish to thank Ariel Karlinsky, Nana Owusu-Boaitey, Lauren Maxwell, and Sayali Arvind Chavan for help with data collection. Finally, thank you to the many fellow researchers who helped by providing information about various data including, amongst others, Maurizio Napolitano, Shiwani Mahajan, Miguel Quartin, Stefano Lombardo, and Timon Gaertner.*

ABSTRACT - Estimating the COVID-19 infection fatality rate (IFR) has proven to be particularly challenging –and rather controversial– due to the fact that both the data on deaths and the data on the number of individuals infected are subject to many different biases. We consider a Bayesian evidence synthesis approach which, while simple enough for researchers to understand and use, accounts for many important sources of uncertainty inherent in both the seroprevalence and mortality data. With the understanding that the results of one’s evidence synthesis analysis may be largely driven by which studies are included and which are excluded, we conduct two separate parallel analyses based on two lists of eligible studies obtained from two different research teams. The results from both analyses are rather similar. With the first analysis, we estimate the COVID-19 IFR to be 0.31% (95% credible interval of (0.16%, 0.53%)) for a typical community-dwelling population where 9% of the population is aged over 65 years and where the gross-domestic product at purchasing-power parity (GDP at PPP) per capita is \$17.8k (the approximate worldwide average). With the second analysis, we obtain 0.32% (95% credible interval of (0.19%, 0.47%)). Our results suggest that, as one might expect, lower IFRs are associated with younger populations (and may also be associated with wealthier populations). For a typical community-dwelling population with the age and wealth of the United States we obtain IFR estimates of 0.43% and 0.41%; and with the age and wealth of the European Union, we obtain IFR estimates of 0.67% and 0.51%. **Keywords:** COVID-19; evidence synthesis; Bayesian inference; infection fatality rate.

Above all, what’s needed is humility in the face of an intricately evolving body of evidence. The pandemic could well drift or shift into something that defies our best efforts to model and characterize it.

Siddhartha Mukherjee, *The New Yorker*
February 22, 2021

1 Introduction

The infection fatality ratio (IFR), defined as the proportion of individuals infected who will go on to die as a result of their infection, is a crucial statistic for understanding SARS-CoV-2 and the ongoing COVID-19 pandemic. Estimating the COVID-19 IFR has proven to be particularly challenging –and rather controversial– due to the fact that both the data on deaths and the data on the number of individuals infected are subject to many different biases.

SARS-CoV-2 seroprevalence studies can help provide a better understanding of the true number of infections in a given population and for this reason several researchers have sought to leverage seroprevalence study data to infer the COVID-19 IFR [30]. In particular, Ioannidis [52], Levin et al.

[66], Brazeau et al. [21], and O’Driscoll et al. [91] have all undertaken analyses, of varying degrees of complexity, in which they combine data from multiple seroprevalence studies with available mortality statistics to derive IFR estimates.

The analyses of both Brazeau et al. [21] and O’Driscoll et al. [91] are done using rather complex Bayesian models which rely on numerous detailed assumptions. For instance, Brazeau et al. [21] use a Bayesian “statistical age-based model that incorporates delays from onset of infection to seroconversion and onset of infection to death, differences in IFR and infection rates by age, and the uncertainty in the serosample collection time and the sensitivity and specificity of serological tests.” O’Driscoll et al. [91] employ a Bayesian ensemble model which assumes “a gamma-distributed delay between onset [of infection] and death” and assumes different risks of infection for “individuals aged 65 years and older, relative to those under 65.” While these analyses go to great lengths to account for the various sources of uncertainty in the data, the complexity of the models will no doubt make it challenging for other researchers to fit these models to different data in a constantly evolving pandemic.

In contrast, the analyses of Ioannidis [52] and Levin et al. [66] are decidedly more simple. For each seroprevalence study under consideration, Ioannidis [52] counts the cumulative number of deaths (from the beginning of the pandemic) until 7 days after the study mid-point (or until the date the study authors suggest), and divides this number of deaths by the estimated number of (previous or current) infections to obtain a study-specific IFR estimate. A “location specific” IFR estimate is then obtained by taking a weighted (by the study’s sample size) average of the study-specific IFR estimates for a given location (i.e., for a given country or state). Ioannidis [52] then calculates the median of all the location specific IFR estimates. No uncertainty interval for this estimate is provided. As such, it is impossible to determine what level of confidence one should place in Ioannidis [52]’s estimates.

The analysis of Levin et al. [66] is based on a standard frequentist random-effects meta-analysis model. For each age-group and seroprevalence study under consideration, Levin et al. [66] calculate a 95% confidence interval (CI) for a study-specific IFR by counting the cumulative number of deaths up until 4 weeks after the study mid-point and dividing this number of deaths by the estimated upper and lower bounds of the number of infected individuals. The meta-analysis model then combines each of these study-specific IFRs. While this analysis provides standard confidence intervals and is relatively straightforward, it does not take into account certain important sources of uncertainty (to be discussed in Section 2).

The analysis method we propose is simple enough for researchers to easily understand and use, while accounting for important sources of uncertainty inherent in both the seroprevalence data and the mortality data. Similar Bayesian models have been used previously for evidence synthesis of seroprevalence data for other infectious diseases (e.g., Brody-Moore [22]). We will apply the method

in an analysis with the objective of estimating the average COVID-19 IFR in a community-dwelling population with a certain approximate age-composition and wealth.

A major part in any evidence synthesis is determining which studies to consider within the analysis. Determining appropriate inclusion and exclusion criteria for seroprevalence studies is a rather complicated and delicate issue when it comes to estimating the COVID-19 IFR [12, 53]. Reviewing and evaluating the merits of the hundreds of available seroprevalence studies also involves a tremendous amount of review work and time. Fortunately, both Chen et al. [29] and Arora et al. [6] have done comprehensive and thorough reviews to ascertain study quality (i.e., risk of bias). We will work from these two lists to conduct two separate parallel analyses. This approach –conducting two analyses based on two distinct and independent literature reviews– will allow us to better understand the impact of different inclusion and exclusion criteria [39]. We will review the data and how it was obtained in Section 3, following a review of the methods in Section 2. In Section 4, we summarize the results of our analysis and conclude in Section 5.

2 Methods

2.1 Bayesian model for evidence synthesis

Suppose we have data from K different seroprevalence studies. Then, for $k = 1, \dots, K$, let:

- T_k be the total number of individuals tested in the k -th study;
- CC_k be the total number of confirmed cases (of past or current infection) resulting from those tested in the k -th study;
- P_k be the number of individuals at risk of infection in the population of interest for the k -th study; and
- D_k be the total number of observed deaths (cumulative since pandemic onset) in the population of interest that are attributed to infection.

We do not observe the following latent (i.e., unknown) variables; for $k = 1, \dots, K$, let:

- C_k be the total number of infected people (cases) in the k -th population;
- IR_k be the true infection rate (proportion of the k -th population which has been infected), which is the expected value of C_k/P_k ; and
- IFR_k be the true underlying infection fatality rate, which is the expected value of D_k/C_k (given C_k).

We will make a series of simple binomial assumptions such that, for $k = 1, \dots, K$:

$$CC_k \sim \text{Binom}(T_k, C_k/P_k), \quad (1)$$

$$C_k \sim \text{Binom}(P_k, IR_k), \quad \text{and} \quad (2)$$

$$D_k|C_k \sim \text{Binom}(C_k, IFR_k). \quad (3)$$

We wish to emphasize the importance of the third “ $D|C$ ” binomial distribution above. Failing to account for the conditional distribution of the deaths given the cases may lead to inappropriately precise estimates of the IFR. For example, Streeck et al. [132] (in their original preprint (*medRxiv*, May 8, 2020)) calculate an uncertainty interval for the IFR by dividing the number of deaths ($D = 7$) by the upper and lower bounds of the 95% CI for the number of infections (95% CI for $C = [1,551, 2,389]$). Doing so, they obtain a relatively narrow 95% CI for the IFR: [0.29%, 0.45%] ($= [7/1,551, 7/2,389]$). In the published version of their article (*Nature Communications*, November 17, 2020), an alternative interval “accounting for uncertainty in the number of recorded deaths” is provided. This alternative interval, which essentially takes into account the $D|C$ binomial distribution, is substantially wider: [0.17%; 0.77%]. In a very similar way, Levin et al. [66] also fail to take into account the $D|C$ binomial distribution when estimating study-specific IFRs resulting in spuriously precise study-specific IFR estimates.

Having established simple binomial distributions for the study-specific IRs and IFRs, we define a simple random-effects model such that, for $k = 1, \dots, K$:

$$g(IFR_k) \sim \mathcal{N}(\theta_0 + \theta_1 Z_{1k} + \theta_2 Z_{2k}, \tau^2), \quad \text{and} \quad (4)$$

$$g(IR_k) \sim \mathcal{N}(\beta, \sigma^2), \quad (5)$$

where θ_0 represents the mean $g(\text{infection fatality rate})$, τ^2 represents between group infection fatality rate heterogeneity, β represents the mean $g(\text{infection rate})$, σ^2 describes the variability in infection rates across the K groups, Z_{1k} and Z_{2k} are covariates of interest that may be related to the infection fatality rate by means of the θ_1 and θ_2 parameters, and $g()$ is a given link function. In our analysis, we define $g()$ as the complimentary log-log link function (cloglog), though there are other sensible choices including the logit and probit functions. As for the two covariates, Z_{1k} and Z_{2k} , we will define these as the centered and scaled logarithm of the proportion of the population aged over 65 years ($65yo_k$), and of the GDP (at purchasing power parity (PPP)) per capita (GDP_k), respectively.

The model is considered within a Bayesian framework requiring the specification of priors for the unknown parameters. Our strategy for priors is to assume weakly informative priors. Beta, Normal, and half-Normal priors (following the recommendations of Gelman et al. [40] and Kümmerer et al. [63]) are set accordingly: $g^{-1}(\theta) \sim \text{Beta}(0.3, 3)$; $g^{-1}(\beta) \sim \text{Beta}(1, 30)$; $\theta_1 \sim \mathcal{N}(0, 10)$; $\theta_2 \sim \mathcal{N}(0, 10)$; $\sigma \sim \text{half-}\mathcal{N}(0, 10)$; and $\tau \sim \text{half-}\mathcal{N}(0, 10)$. Note that the performance of any

Bayesian estimator will depend on the choice of priors and that this choice can substantially influence the posterior when few data are available [15, 64]. In the Supplementary Material 6.6, we show results obtained with an alternative set of priors as a sensitivity analysis.

2.2 Uncertainty in infection rates

While some seroprevalence studies report the exact number of individuals tested and the exact number of confirmed cases amongst those tested, to obtain estimates for the infection rate there are typically numerous adjustments made (e.g., adjusting for imperfect diagnostic test accuracy, adjusting for clustering of individuals within a household). For this reason, the sample size of a given study might not be a reliable indicator of its precision and weighting a study's contribution in an evidence synthesis based solely on its sample size (as in e.g., Ioannidis [52]) may not be appropriate.

Rather than work with the raw testing numbers published in the seroprevalence studies, we calculate effective data values for T_k and CC_k based on a binomial distribution that corresponds to the reported 95% CI for the IR. By “inverting uncertainty intervals” in this way, we are able to properly use the adjusted numbers provided. (This is a similar approach to the strategy employed by Kümmerer et al. [63].) Tables 1 and 2 list the 95% uncertainty intervals obtained from each of the seroprevalence studies in our two parallel analyses and Tables 3 and 4 list the corresponding values for T_k and CC_k .

It must be noted that, as Ioannidis [52] cautions, it is possible that under our “inverting uncertainty intervals” approach, poorly conducted seroprevalence studies which fail to make proper adjustments (and thereby have spuriously narrower uncertainty intervals) receive more weight in our analysis, while high-quality studies, which make proper adjustments, are unfairly penalized. Ioannidis [52] notes that the strategy of “weighting the study-specific infection fatality rates by the sample size of each study” avoids giving more weight to studies “with seemingly narrower confidence intervals because of poor or no adjustments, while still giving more weight to larger studies.” Since we are restricting our analysis to only those supposedly high quality seroprevalence studies, we hope to largely avoid this issue. Weighting studies based on their true precision is obviously the goal in any evidence synthesis, and we recognize that this is particularly difficult when so many studies may misrepresent the precision of their estimates [19, 23].

2.3 Uncertainty in mortality

Matching prevalence estimates with a relevant number of fatalities is a difficult task. Prevalence estimates obtained from a seroprevalence study do not typically correspond to a specific date. Instead, these estimates will correspond to a window of time during which testing occurred. This period may be only a few days for some studies (e.g., 4 days for Petersen et al. [98]), but can also

be several weeks or months for others (e.g., 135 days for Ward et al. [145]). Tables 1 and 2 list the sampling window start and end dates for each of the studies in our two parallel analyses.

Evidently, a longer sampling window will lead to greater uncertainty when it comes to establishing the relevant number of deaths. It can be difficult to account for this uncertainty and analyses will often simply select a specific date at which to count deaths based on some simple rule of thumb. For example, Ioannidis [52] considers the number of deaths at 7 days after the mid-point of the sampling window (or as the relevant number of deaths discussed by the seroprevalence study's authors). As another example, Meyerowitz-Katz and Merone [80] take the number of deaths as recorded at 10 days after the end of the sampling window. While these two particular analytical choices are not all that different, each may lead to a substantially different number of deaths for a given study if the study was conducted during a period of time during which the number of deaths was rapidly accelerating. Levin et al. [66], who consider the number of deaths up until 4 weeks after the sampling window mid-point, acknowledge this limitation noting that: "matching prevalence estimates with subsequent fatalities is not feasible if a seroprevalence study was conducted in the midst of an accelerating outbreak."

In order to account for the uncertainty in selecting the relevant number of deaths for a given seroprevalence study, we propose considering the number of deaths as interval censored data. Tables 3 and 4 list numbers for an interval corresponding to the number of deaths recorded 14 days after the start of the sampling window and 14 days after the end of sampling window for each seroprevalence study. While we might not know exactly what number of deaths is most appropriate, we can be fairly confident that the appropriate number lies somewhere within this interval. (Note that some intervals in Tables 3 and 4 have also been widened to account for other sources of uncertainty in the number of deaths; see details in the Supplementary Material Section 6.3). The 14 day offset allows for the known delay between the onset of infection and death, taking into consideration the delay between the onset of infection and the development of detectable antibodies [70, 149].

3 The data

3.1 Seroprevalence data

As the COVID-19 pandemic has progressed, a rapidly increasing number of SARS-CoV-2 seroprevalence studies have been conducted worldwide [6]. However, many of these studies have produced biased estimates or are otherwise unreliable due to a variety of different issues with study design, and/or with data collection, and/or with inappropriate statistical analysis [19]. We seek to restrict our analysis to high quality studies which used probability-based sampling methods. Such studies are less likely to suffer from substantial biases [123]. Based on the reviews of Chen et al. [29] and of Arora et al. [6], we compiled two separate sets of studies for analysis (these are listed in Tables 1

and 2, respectively). With the understanding that the results of an evidence synthesis may be largely driven by which studies are included/excluded, we will use these two separate sets to conduct two separate analyses. Note that the data collected for both analyses are relevant to the time before the widespread availability of COVID-19 vaccinations.

Figure 1: Flowchart of seroprevalence studies considered for Chen et al. -based analysis (based on the review of Chen et al. [29]).

Chen et al. [29] reviewed the literature for articles published between December 1, 2019, and December 22, 2020, and identified more than 400 unique seroprevalence studies. For each of these, study quality was established using a scoring system developed on the basis of a seroepidemiological protocol from the Consortium for the Standardization of Influenza Seroepidemiology [49]. In total, Chen et al. [29] identified 38 articles which considered a sample based on a “general population” and which obtained a study quality grade of A or B (see full list in Table S8 of Chen et al. [29]). We consider these 38 articles as a starting point for inclusion for our analysis. After excluding those studies which are duplicates ($n=2$), those that used a “convenience” or “non-probability” based sampling method (according to the classification of Arora et al. [6]) ($n=8$), a study no longer considered accurate based on new information about the accuracy of the antibody test used ($n=1$), a study that has a very narrowly defined target population ($n=1$), studies for which relevant death data could not be found ($n=5$), and studies which did not provide a 95% uncertainty interval ($n=2$), we were left with a set of $K = 19$ studies for analysis; see Figure 1 and details in the Supplementary Material.

Arora et al. [6] conducted the Serotracker “living systematic review” of COVID-19 seroprevalence studies whereby the results of the review are continuously updated on serotracker.com/data. For each study reviewed, the risk of bias was evaluated based on an assessment using the Joanna Briggs Institute Critical Appraisal Guidelines for Prevalence studies [19, 84]. For analysis, we consider the 45 studies listed on serotracker.com/data (as of June 5, 2021), that are categorized as having a “low risk of bias” and are categorized as targeting “household and community samples.” After excluding those studies which are duplicates ($n=3$), one study that used a “convenience” or “non-probability” based sampling method (according to the classification of Arora et al. [6]) ($n=1$), those studies no longer considered accurate based on new information about the accuracy of the antibody test used ($n=2$), those that have very narrowly defined target populations ($n=2$), those for which relevant death data could not be found ($n=8$), and those which did not provide a 95% uncertainty interval for the estimated prevalence ($n=1$), we are left with a set of $K = 28$ studies for analysis; see Figure 2 and details in the Supplementary Material.

Figure 2: Flowchart of seroprevalence studies considered for Serotracker-based analysis (based on the Serotracker review Arora et al. [6]).

For each of the seroprevalence studies included in each of the two analysis sets, we recorded the 95% uncertainty interval for the infection rate as reported in the study article. If an article reported on multiple phases of a study (e.g., a longitudinal series of different surveys), or reported different results for different areas instead of an overall estimate (e.g., a series of different estimates for different regions), we selected only the first set of estimates. Furthermore, if a study reported more than one 95% uncertainty interval (e.g., different intervals corresponding to different adjustments

and assumptions), we selected the lowest value amongst the different lower bounds and the highest value amongst the different upper bounds. These numbers are recorded in Tables 1 and 2 under *IR interval*. Based on these numbers, we calculated effective data values for the number of tests (T_k) and the number of confirmed cases (CC_k) which are listed in Tables 3 and 4 alongside population numbers (P_k) and numbers corresponding to the proportion of the population over 65 years old ($65yo_k$) and the GDP (PPP) per capita (GDP_k); see Supplemental Material for details and data sources.

3.2 Mortality data

Mortality data was obtained from various sources (e.g., academic, government, health authority); see details in Supplementary Material (Section 6.3). If a seroprevalence study referenced a specific source for mortality data, we used the referenced source for our numbers whenever possible. If no source was referenced or suggested, we considered publicly available data sources.

For many populations, there were concerns that cause of death information may be very inaccurate and lead to biased COVID-19 mortality statistics. To overcome this issue, many have suggested looking to “excess deaths” by comparing aggregate data for all-cause deaths from the time during the pandemic to the years prior [65]. For populations with a large discrepancy between the number of deaths attributed to COVID-19 and the number of excess deaths –as suggested, when possible, by a large undercount ratio (UCR) derived by Karlinsky and Kobak [57]– we used the number of deaths attributed to COVID-19 for the lower bound of the D_k interval and used numbers based on excess deaths for the upper bound of the D_k interval.

India, Pakistan, Palestine, Ethiopia, and China are the only countries represented in the studies that we assessed for data availability that were not included in Karlinsky and Kobak [57]’s analysis. There was evidence of substantial under-reporting of COVID-19 deaths in India [10, 104] while little could be gathered about the reliability of official mortality data for Pakistan, Palestine¹, Ethiopia², and China (but do see [72] and [137]). As such, we excluded the Qutob et al. [107] (“Palestinian population residing in the West Bank”) and He et al. [47] (“Wuhan, China”) studies from the Serotracker-based analysis and excluded the Alemu et al. [1] (“Addis Ababa, Ethiopia”) and the Nisar et al. [90] (“Two neighborhoods of Karachi, Pakistan”) studies from the Chen et al.-based analysis. For India, Mukherjee et al. [83] and Purkayastha et al. [105] estimate UCRs for the entire country as well as for each individual Indian state and union territory. We used these UCRs to adjust the death numbers for seroprevalence studies in India (see Supplementary Material for details).

¹Official regional death numbers for Palestine are available from the Palestinian government dashboard (see <https://corona.ps/details>; accessed July 28, 2021).

²Official regional death numbers for Ethiopia have been made available previously (e.g., http://web.archive.org/web/2020*/https://www.covid19.et/covid-19/ and the Twitter account: https://twitter.com/Harun_Asefa/status/125906983287793280; accessed August 4, 2021).

There are two countries represented within our data that were identified by Karlinsky and Kobak [57] as having large discrepancies between the official number of deaths attributed to COVID-19 and the number of excess deaths Iran (with UCR=2.4) and Russia (with UCR=4.5). As such, for the Barchuk et al. [11] (“Saint Petersburg, Russia”) and for the Khalagi et al. [58] (“Iran”), we used numbers based on excess deaths (see Supplementary Material for details).

Finally, our target of inference is the IFR for the community-dwelling population and does not apply to people living in long-term care (LTC) facilities (also known as “nursing homes” or, in France as “Établissement d’hébergement pour personnes âgées dépendantes” (EHPAD)). The spread of COVID-19 is substantially different in LTC facilities than in the general population, and residents of LTC facilities are particularly vulnerable to severe illness and death from infection; see Danis et al. [31]. With this in mind, we made adjustments (when appropriate/possible) to the mortality numbers used in our analysis in order to exclude deaths of LTC residents; see Supplementary Material for details. Modeling the spread and mortality of COVID-19 within LTC facilities will require unique approaches and should be considered in a separate analysis; see the recommendations of Pillemer et al. [100].

Authors	Location	Sampling (mm/dd)	IR interval (%)	In both analyses
Barchuk et al. [11]	Saint Petersburg, Russia	05/27 - 06/26	[5.60, 12.90]	yes
Biggs et al. [18]	DeKalb and Fulton, GA, USA (A)	04/28 - 05/03	[1.40, 4.50]	yes*
Bruckner et al. [24]	Orange County, CA, USA	07/10 - 08/16	[8.10, 15.50]	yes
Carrat et al. [27]	Ile-de-France, France	05/04 - 06/14	[8.90, 11.30]	
Mahajan et al. [74]	Connecticut, USA	06/10 - 07/29	[1.70, 6.30]	
Murhekar et al. (A) [86]	India	05/11 - 06/04	[0.34, 1.13]	yes
Office of National Stat [92]	England, UK (A)	04/26 - 09/08	[5.40, 7.10]	
Petersen et al. [98]	Faroe Islands, Denmark	04/27 - 05/01	[0.10, 1.20]	yes
Pollan et al. [102]	Spain	04/27 - 05/11	[3.30, 6.60]	yes
Samore et al. [116]	Four counties in UT, USA	05/04 - 06/30	[0.10, 1.60]	
Santos-Hovener et al. [117]	Kupferzell, Germany	05/20 - 06/09	[10.40, 14.00]	yes
Sharma et al. [120]	Delhi, India	08/01 - 08/07	[27.65, 29.14]	yes
Snoeck et al. [126]	Luxembourg	04/15 - 05/05	[1.23, 2.77]	
Sood et al. [127]	Los Angeles County, CA, USA	04/10 - 04/14	[2.52, 7.07]	
Statistics Jersey (A) [129]	Jersey, UK (A)	04/29 - 05/05	[1.80, 4.40]	yes*
Streeck et al. [132]	Gangelt, Germany	03/31 - 04/06	[12.31, 24.40]	yes
Stringhini et al. (A) [133]	Geneva, Switzerland (A)	04/06 - 05/09	[8.15, 13.95]	yes*
Vos et al. [141]	Netherlands	03/31 - 05/11	[2.10, 3.70]	yes
Ward et al. [145]	England, UK (B)	06/20 - 07/13	[5.78, 6.14]	yes

Table 1: Seroprevalence studies selected for the analysis based on the list compiled by Chen et al. [29] (listed in alphabetical order of authors), with geographic location of sampling, sampling dates, and 95% uncertainty interval for the infection rate (IR interval). Also noted, under “In both analyses”, is whether or not each study is included in the Serotracker-based analysis (i.e., is also in Table 2). Studies with yes* are alternate versions of studies that are included in the Serotracker-based analysis. Note that sampling for all studies took place during mid-2020, before the widespread availability of COVID-19 vaccinations.

Authors	Location	Sampling (mm/dd)	IR interval (%)	In both analyses
Álvarez-Antonio et al. [2]	Iquitos, Peru	07/13 - 07/18	[67.00, 73.00]	
Bajema et al. [9]	DeKalb and Fulton, GA, USA (B)	04/28 - 05/03	[1.49, 6.67]	yes*
Barchuk et al. [11]	Saint Petersburg, Russia	05/27 - 06/26	[5.60, 12.90]	yes
Bruckner et al. [24]	Orange County, CA, USA	07/10 - 08/16	[8.10, 15.50]	yes
Chan et al. [28]	Rhode Island, USA	05/05 - 05/22	[1.00, 6.20]	
Gov. of Andorra [43]	Andorra	05/04 - 05/28	[10.50, 11.50]	
Kar et al. [56]	Puducherry District, India	08/11 - 08/16	[3.50, 6.40]	
Khalagi et al. [58]	Iran	08/03 - 10/31	[13.30, 15.20]	
Malani et al. [76]	Tamil Nadu, India	10/19 - 11/30	[30.40, 32.80]	
Melotti et al. [79]	Gardena Valley, Italy	05/26 - 06/08	[25.20, 28.60]	
MoHoI [81]	Israel	06/28 - 09/17	[5.30, 5.60]	
Murhekar et al. (A) [86]	India	05/11 - 06/04	[0.34, 1.13]	yes
Nawa et al. [89]	Utsunomiya City, Japan	06/14 - 07/05	[0.08, 2.28]	
Pagani et al. [93]	Castiglione d'Adda, Italy	05/18 - 06/07	[17.20, 29.10]	
Petersen et al. [98]	Faroe Islands, Denmark	04/27 - 05/01	[0.10, 1.20]	yes
Pollan et al. [102]	Spain	04/27 - 05/11	[3.30, 6.60]	yes
Radon et al. [108]	Munich, Germany	11/02 - 01/31	[2.90, 4.30]	
Reyes-Vega et al. [111]	Lima, Peru	06/28 - 07/09	[22.50, 28.20]	
Richard et al. [112]	Geneva, Switzerland (B)	04/06 - 06/30	[6.80, 8.90]	yes*
Santos-Hovener et al. [117]	Kupferzell, Germany	05/20 - 06/09	[10.40, 14.00]	yes
Selvaraju et al. [118]	Chennai, Tamil Nadu, India	07/01 - 07/31	[14.80, 22.60]	
Sharma et al. [120]	Delhi, India	08/01 - 08/07	[27.65, 29.14]	yes
Statistics Jersey (B) [130]	Jersey, UK (B)	06/21 - 06/27	[2.80, 5.20]	yes*
Streeck et al. [132]	Gangelt, Germany	03/31 - 04/06	[12.31, 24.40]	yes
Vos et al. [141]	Netherlands	03/31 - 05/11	[2.10, 3.70]	yes
Ward et al. [145]	England, UK (B)	06/20 - 07/13	[5.78, 6.14]	yes
Warszawski et al. [146]	Metropolitan France	05/02 - 06/02	[3.90, 5.00]	
Yoshiyama et al. [150]	Tokyo, Japan	06/01 - 06/07	[0.01, 0.37]	

Table 2: Seroprevalence studies selected for the analysis based on the list compiled by Serotracker (listed in alphabetical order of authors), with geographic location of sampling, sampling dates, and 95% uncertainty interval for the infection rate (IR interval). Also noted, under “In both analyses”, is whether or not each study is included in the Chen et al.-based analysis (i.e., is also in Table 1). Studies with yes* are alternate versions of studies that are included in the Chen et al.-based analysis. Note that sampling for all studies took place during mid-to-late 2020, before the widespread availability of COVID-19 vaccinations.

4 Results

The model as described in Section 2, was fit to the two datasets as described in Section 3. We fit the model using JAGS (just another Gibbs sampler) [62], with 5 independent chains, each with 2 million draws (20% burn-in, thinning of 100); see Supplementary Material (Section 6.4) for details and JAGS code.

We report posterior median estimates and 95% highest probability density (HPD) credible intervals (CrI). Figure 3 (for the Chen et al.-based analysis) and Figure 4 (for the Serotracker-based analysis) plot the point estimates and credible intervals obtained for IFR_k , for k in $1, \dots, K$, respectively; see Figures 5 and 6 for IR_k are in the Supplementary Material. In these figures the seroprevalence studies are listed in order of their “fitted” IFR values (the posterior median of $g^{-1}(\theta_0 + \theta_1 Z_{1k} + \theta_2 Z_{2k})$),

Location	P_k	D_k		T_k	CC_k	$65yo_k$ (%)	GDP_k \$
		lower	upper				
Saint Petersburg, Russia	5351935	1596	5128	233	21	18	30144
DeKalb and Fulton, GA, USA (A)	1823234	122	136	419	11	12	58933
Orange County, CA, USA	3010232	512	839	285	33	15	79287
Ile-de-France, France	12213447	6816	7427	2414	243	14	82574
Connecticut, USA	2837877	1136	1179	251	9	22	80729
India	1366417750	4172	45766	1632	11	6	4735
England, UK (A)	56550000	25975	30709	3100	193	18	43310
Faroe Islands, Denmark	52154	0	0	592	3	17	60421
Spain	47351567	17710	17767	643	31	20	42362
Four counties in UT, USA	2194298	40	97	393	2	10	60050
Kupferzell, Germany	6247	1	3	1263	153	16	63885
Delhi, India	19800000	4188	26901	13966	3965	4	12817
Luxembourg	632275	47	58	1214	23	14	122166
Los Angeles County, CA, USA	10039107	529	606	316	14	14	79287
Jersey, UK (A)	107800	17	18	648	19	17	48365
Gangelt, Germany	12597	8	8	153	27	18	53751
Geneva, Switzerland (A)	504128	122	155	442	48	16	68964
Netherlands	17344874	2621	4186	1652	47	20	59685
England, UK (B)	56550000	29526	30038	10635	634	18	43310

Table 3: The Chen et al. based dataset required for the Bayesian evidence synthesis model.

for k in $1, \dots, K$, marked on the plot by the $\times$ symbols). Results obtained for the other model parameters are listed in Table 5.

In general, the Chen et al.-based analysis and the Serotracker-based analysis provide mostly similar results. Notably, the Serotracker-based analysis considers a much more geographically diverse set of seroprevalence studies and several studies that appear to be prominent outliers (e.g., “Tamil Nadu”, “Castiglione d’Adda, Italy”, and “Utsunomiya City, Japan”), see Figure 4. These outliers could be due to infection rates in these populations being markedly different for the elderly relative to the general population.³ With regards to heterogeneity, fitting the model without any covariates, one obtains $\hat{\tau} = 0.62$ (Chen et al.-based analysis) and $\hat{\tau} = 0.90$ (Serotracker-based analysis). This suggests that the two covariates, $65yo_k$ and GDP_k , account for approximately 45% ($= (0.62^2 - 0.46^2)/(0.62^2)$; Chen et al.-based analysis) and 13% ($= (0.90^2 - 0.84^2)/(0.90^2)$; Serotracker-based analysis) of the heterogeneity in the IFR.⁴

³Malani et al. [76] (“Tamil Nadu”) note that: “Seroprevalence among the elderly (70+: 25.8%) is significantly lower than among the working age populations (age 40-49: 31.6%; $p < 0.001$) or the young (18-29: 30.7%; $p < 0.001$).” Pagani et al. [93] (“Castiglione d’Adda, Italy”) also report that the elderly are more likely to have been infected (“strong association observed between IgG seroprevalence and age”). On the other hand, note that among the 181 participants in the Nawa et al. [89] study (“Utsunomiya City, Japan”) who were aged 65 years or older, none were positive. This suggests that infection in the elderly may be lower than in the general population. However, since the Nawa et al. [89] found only 3 positive cases out of a total of 742 individuals tested, inference on this is limited.

⁴For reference, Levin et al. [66] conclude that 87% of the heterogeneity in the IFR (of advanced economies) can be explained by variations in age composition and age-specific prevalence of COVID-19. However, note that the linear regression analysis used to obtain this 87% result is done without an intercept term (see Levin et al. [66] - Figure 6). A linear regression with intercept results in a value of 43%. The intraclass correlation coefficient (ICC) [140] between Levin et al. [66]’s predicted IFRs and the observed IFRs is 0.65 (after removing one outlier, “Portugal”), suggesting that the extent of agreement is reasonably high but nowhere near perfect.

Location	P_k	D_k lower	D_k , upper	T_k	CC_k	$65yo_k$ (%)	GDP_k \$
Iquitos, Peru	467000	1581	1942	895	626	4	5896
DeKalb and Fulton, GA, USA (B)	1823234	122	136	196	7	12	58933
Saint Petersburg, Russia	5351935	1596	5128	233	21	18	30144
Orange County, CA, USA	3010232	512	839	285	33	15	79287
Rhode Island, USA	1059361	147	189	163	5	18	58416
Andorra	77543	35	35	11236	1235	17	49900
Puducherry District, India	1250000	145	393	840	41	6	9152
Iran	82913906	19804	98071	5197	740	6	12937
Tamil Nadu, India	83697770	11183	29773	5765	1821	7	7191
Gardena Valley, Italy	10700	15	15	2612	702	20	71853
Israel	9216900	187	995	10578	577	11	40747
India	1366417750	4172	45766	1632	11	6	4735
Utsunomiya City, Japan	517527	0	0	228	2	26	42931
Castiglione d'Adda, Italy	4605	39	40	190	43	26	59291
Faroe Islands, Denmark	52154	0	0	592	3	17	60421
Spain	47351567	17710	17767	643	31	20	42362
Munich, Germany	1563090	219	780	2692	96	17	65345
Lima, Peru	10804609	21109	28846	893	226	9	19313
Geneva, Switzerland (B)	504128	122	156	2509	196	16	68964
Kupferzell, Germany	6247	1	3	1263	153	16	63885
Chennai, Tamil Nadu, India	10900000	1295	5960	380	70	7	7191
Delhi, India	19800000	4188	26901	13966	3965	4	12817
Jersey, UK (B)	107800	20	20	998	39	17	48365
Gangelt, Germany	12597	8	8	153	27	18	53751
Netherlands	17344874	2621	4186	1652	47	20	59685
England, UK (B)	56550000	29526	30038	10635	634	18	43310
Metropolitan France	64897954	17396	19038	5377	238	21	49551
Tokyo, Japan	13960236	270	275	1314	2	20	45796

Table 4: The Serotracker based dataset required for the Bayesian evidence synthesis model.

Our estimates of $\hat{\theta}_1 = 0.61$ (Chen et al.-based analysis) and $\hat{\theta}_1 = 0.42$ (Serotracker-based analysis) suggest that older populations are more likely to have higher IFRs. This is as expected since age is known to be a very important risk factor [148, 152]. Our estimate of $\hat{\theta}_2 = -0.28$ (Chen et al.-based analysis) and $\hat{\theta}_2 = -0.11$ (Serotracker-based analysis) suggest that wealthier populations may be more likely to have lower IFRs. However, the wide credible intervals obtained for the θ_2 parameter (in both analyses) suggest a much less definitive conclusion. There are several reasons which might explain this result. As with any observational data analysis, the estimate of θ_2 may suffer from bias due to unobserved confounding and statistical power may be compromised by the presence of outliers and insufficient heterogeneity in the GDP per capita metric across the different populations included in our analyses.

We can infer (by determining the posterior median of $g^{-1}(\theta_0 + \theta_1 z_{1*} + \theta_2 z_{2*})$, for selected values of z_{1*} and z_{2*}) the *typical* IFR amongst populations (be they included in our study or not) having a given proportion of the populace aged over 65 and a given GDP per capita. Thus we calculate posterior point and interval estimates corresponding to age and wealth values that match the population of the entire world (World), the United States (USA), and the European Union (EU) (as listed by the

Parameter	Chen et al. analysis	Serotracker analysis
θ_0	-5.50, with 95% CrI of (-5.78, -5.22)	-5.56, with 95% CrI of (-5.93, -5.19)
θ_1	0.61, with 95% CrI of (0.21, 1.00)	0.42, with 95% CrI of (-0.40, 1.16)
θ_2	-0.28 with 95% CrI of (-0.67, 0.14)	-0.11, with 95% CrI of (-0.86, 0.69)
τ	0.46 with 95% CrI of (0.25, 0.75)	0.84, with 95% CrI of (0.55, 1.23)
σ	1.17 with 95% CrI of (0.80, 1.68)	1.53, with 95% CrI of (1.13, 2.05)

Table 5: Parameter estimates (posterior medians and 95% HPD credible intervals) obtained from the Chen et al.-based analysis and the Serotracker-based analysis.

World Bank’s World Development Indicators (WDI)); see “World”, “USA”, and “EU” rows in Figures 3 and 4. For 65yo = 9% and GDP = \$17,811, the approximate worldwide values, we obtain, from the Chen et al.-based analysis, an across-population average IFR estimate of 0.31%, with a 95% CrI of (0.16%, 0.53%). With the Serotracker-based analysis, we obtain a similar estimate of 0.32%, with a 95% CrI of (0.19%, 0.47%). For 65yo = 16% and GDP = \$65,298, the USA values, we obtain across-population average IFR estimates of 0.43%, with a 95% CrI of (0.31%, 0.56%) (Chen et al.-based analysis) and of 0.41%, with a 95% CrI of (0.22%, 0.67%) (Serotracker-based analysis). Finally, for 65yo = 20% and GDP = \$47,828, the EU values, we obtain across-population average IFR estimates 0.67%, with a 95% CrI of (0.41%, 0.96%) (Chen et al.-based analysis) and of 0.51%, with a 95% CrI of (0.27%, 0.79%) (Serotracker-based analysis). Note that for the “World” predictions, the Serotracker-based analysis has the more precise estimates, while the Chen et al.-based estimates are more precise for the “USA” predictions.

While the infection-rate estimates obtained from the seroprevalence studies should be relatively reliable (due to having satisfied the risk of bias assessments of either Chen et al. [29] or Arora et al. [6]), the mortality data we collected may be less reliable depending on the target population and source. The data which were not obtained from official and reliable sources may be particularly suspect. With this in mind, as a sensitivity analysis, we repeated both analyses with these data excluded; see results in Figures 7 and 8 in the Supplementary Material Section 6.6. Without the excluded studies, we are unable to provide a reasonable “World” estimate (see the extremely wide credible intervals). However, the “USA” and “EU” estimates are relatively similar. We also repeated the two analyses using a different set of priors to verify that our results were not overly sensitive to our particular choice of priors. The results of this alternative analysis are very similar to the results of our original analyses; see Figures 9 and 10 in the Supplementary Material Section 6.6.

Our estimates are somewhat similar to those obtained in other analyses. Brazeau et al. [21], using data from 10 representative seroprevalence studies (identified after screening 175 studies), infer “the overall IFR in a typical low-income country, with a population structure skewed towards younger individuals, to be 0.23% (0.14%-0.42% 95% prediction interval range).” For a “typical high income country, with a greater concentration of elderly individuals,” Brazeau et al. [21] obtain an estimate

of 1.15% (95% prediction interval of 0.78%-1.79%). Ioannidis [52], using data from seroprevalence studies with sample sizes greater than 500 (and including deaths of LTC residents), obtains a “median infection fatality rate across all 51 locations” of 0.27% and (and of 0.23% following an ad-hoc correction to take into account “that only one or two types of antibodies” may have been tested in some seroprevalence studies). Levin et al. [66], who restricted their analysis to populations in “advanced economies” (and included deaths of LTC residents) provide age-group specific estimates and country-specific estimates. For instance, for the 45–54 year old age group, Levin et al. [66] estimate the IFR to be 0.23% (95% CI of 0.20%–0.26%), and for the 55–64 year old age group, 0.75% (95% CI of 0.66%–0.87%). For Spain, Levin et al. [66] estimate an IFR of 1.90%. For comparison, we estimate the IFR for the community-dwelling population (i.e., excluding deaths of LTC residents) of Spain to be 0.77% (in both analyses). This is similar to the 0.83% estimate obtained by Pastor-Barriuso et al. [94] and the 0.75% estimate obtained by Brazeau et al. [21] (both of these excluding deaths of LTC residents).

Specifically with regards to the United States, Sullivan et al. [135] estimate the IFR for adults to be 0.85% (95% CrI of 0.76%-0.97%) based on a US nationwide seroprevalence survey conducted between August and December, 2020.⁵ Pei et al. [95] using a rather complex Bayesian “metapopulation” model conclude that, for the United States during 2020, the IFR likely “decreased from around 1% in March to about 0.25% in December.” For comparison, our “USA” predictions of 0.43% and of 0.41% are based on data obtained mostly between April, 2020 and August, 2020 (see dates in Tables 1 and 2).

5 Conclusion

Estimation of the IFR can be incredibly challenging due to the fact that it is a ratio of numbers where both the numerator and the denominator are subject to a wide range of biases. Our proposed method seeks to address some of these biases in a straightforward manner. In our analysis, proper handling of the various sources of uncertainty was a primary focus [151].

With regards to the numerator, we considered the number of deaths as interval censored data so as to account for the uncertainty in selecting the most relevant number of deaths. While we consider this an improvement over other methods that use a single fixed number, we acknowledge that the specific choice of a 14 day offset is somewhat arbitrary and that the data for deaths also suffer from other sources of bias. Ioannidis [53] notes that the time between infection and death may vary substantially “and may be shorter in developing countries where fewer people are long-sustained by medical support.” In addition to official numbers, we used mortality data based on “excess deaths” statistics for Russia and Iran, since official mortality statistics appeared to be potentially

⁵While Sullivan et al. [135] used a nationwide representative sampling frame, the results may be subject to sizable selection bias given the low response rate of 12.6%.

Figure 3: Results from the Chen et al.-based analysis: posterior median estimates (black circles) for the IFR_k variables (for $k = 1, \dots, 19$) with 95% HPD CrIs. Studies are listed from top to bottom in order of increasing fitted values (these values are indicated by $\times$). Also plotted, under the labels “World (65 yo=9%, GDP=17.8k)”, “USA (65 yo=16%, GDP=65.3k)”, “EU (65 yo=20%, GDP=47.8k)”, are the posterior median estimate and 95% HPD CrIs for the typical IFR corresponding to values for the proportion of the population aged 65 years and older of 9% and for GDP per capita of \$17,811 (the worldwide values), of 16% and of \$65,298 (the USA values), and of 20% and of \$47,828 (the EU values).

Figure 4: Results from the Serotracker-based analysis: posterior median estimates (black circles) for the IFR_k variables (for $k = 1, \dots, 28$) with 95% HPD CrIs. Studies are listed from the top to the bottom in order of increasing fitted values (these values are indicated by $\times$). Also plotted, under the labels “World (65 yo=9%, GDP=17.8k)”, “USA (65 yo=16%, GDP=65.3k)”, “EU (65 yo=20%, GDP=47.8k)”, are the posterior median estimate and 95% HPD CrIs for the typical IFR corresponding to values for the proportion of the population aged 65 years and older of 9% and for GDP per capita of \$17,811 (the worldwide values), of 16% and of \$65,298 (the USA values), and of 20% and of \$47,828 (the EU values).

highly inaccurate. We also used adjusted mortality numbers for India based on the best available information. These adjustments are certainly not perfect and we note that “excess deaths” statistics may also suffer from substantial inaccuracies [82].

With regards to the denominator, we looked to data from “high-quality” seroprevalence studies in an effort to avoid biased estimates. However, these data are also not perfect. Seroprevalence studies are severely limited by the representativeness of the individuals they test. Certain groups of individuals who may have very high infection rates are unlikely to be tested in a seroprevalence study (e.g., homeless people). On the other hand, those individuals who have reason to believe they may have been infected, may be more likely to volunteer to participate in a seroprevalence study [123]. It is also likely that seroreversion (loss of detectable antibodies over time) may lead to a seroprevalence study underestimating the true number of infections if the time between the main outbreak and the subsequent antibody testing is substantial [96]. Notably, Axfors and Ioannidis [7] employ a “X-fold”-based correction factor to adjust seroprevalence estimates for this type of bias.

The need to improve the quality and reporting of seroprevalence studies cannot be overemphasized.⁶ A major limitation of evidence synthesis is often summarized by the expression “garbage in, garbage out” [34], meaning that if one includes biased studies in one’s analysis, the analysis results will themselves be biased [122]. In our two analyses, we only included data from 19 and 28 out of potentially hundreds of seroprevalence studies due primarily to the fact that so few studies were considered reliable and at low risk of bias. Excluding low-quality/biased studies from our analysis was necessary, at least to a certain degree, in order to obtain valid estimates. However, as a consequence of our strict exclusion criteria, much of the world’s population is severely under-represented in our data. In related work, Levin et al. [67] review the available literature and “informally assess studies for risk of bias” in an attempt to estimate the COVID-19 IFR specifically for developing countries. If the quality of studies were to be correlated with unmeasured factors that impact the IFR, excluding studies based on their perceived quality could lead to unmeasured confounding at a meta-analytic level [54]. Novel methods which allow evidence syntheses to appropriately incorporate biased data are urgently needed. Recently, Campbell et al. [25] proposed a partially identified model to combine seroprevalence study data with biased data from official statistics.

Outside of biased data, perhaps the foremost challenge in evidence synthesis using observational data is that necessarily one is forced to make an array of design choices around inclusion/exclusion criteria, statistical modeling approaches, and prior specifications [39]. With the two separate

⁶Bobrovitz et al. [19] conclude that a majority of COVID-19 seroprevalence studies are “at high risk of bias [...], often for not statistically correcting for demographics or for test sensitivity and specificity, using non-probability sampling methods, and using non-representative sample frames.” Shook-Sa et al. [123] note that “it is very difficult to estimate the level of bias introduced by convenience sampling.”

analyses and the various additional sensitivity analyses, we were quite encouraged by the stability of our results to perturbations of these inputs.

Reducing the uncertainty around the severity of COVID-19 was of great importance to policy makers and the public during the early stages of the pandemic [35, 51, 71] and immense efforts have been made in the collection and analysis of data (e.g., Williamson et al. [148]). And yet, even after more than a year, there is still a large amount of uncertainty and unexplained heterogeneity surrounding the COVID-19 IFR, particularly with respect to populations in less affluent countries. While a certain amount of heterogeneity is to be expected [48], identifying factors associated with higher IFRs is the ultimate goal and investigating potential variables that can account for the observed heterogeneity may lead to important insights [16, 54].

We prioritized simplicity in our modeling so as to promote transparency in our findings, and to facilitate adaptations to similar, but not identical, data structures. While “simple” is a relative term, note that the entire dataset used for our analyses fits on a single page (in Tables 3 and 4) and that the entire JAGS MCMC code fits on less than a single page (see Supplementary Material). One model extension that could be pursued would involve age stratification of IFR.

Including age-stratification in the model could represent a substantial improvement given that infection in some populations is far from homogeneous (e.g., about 95% of Singapore’s COVID-19 infections were among young migrant workers (as of September, 2020), which explains the incredibly low case fatality rate [37]). If a factor, such as age, impacts both the risk of infection and the risk of death given infection, then estimating the IFR as we have done in our analysis could be subject to confounding [147]. Age-group specific seroprevalence/mortality data is available for certain geographic areas [113] (although not always consistently reported) and such data could inform an extended version of our model, thereby offering an alternative to the approach described by Levin et al. [66] for estimating age-group specific IFRs.

Finally, we must emphasize that the IFR is a moving target. As the pandemic changes, so to does the IFR. Our estimates are based on data from 2020, most of which were obtained more than a year ago. It is likely that, with continual viral mutation of SARS-CoV-2, advances in treatment, and the availability of vaccines, the current IFR in many places is now markedly different than it was earlier in 2020 [95, 99, 143].

Key Messages

- The COVID-19 IFR is estimated to be about 0.32% for a typical community-dwelling population where 9% of individuals are over 65 years old and where the GDP per capita is \$17.8k (the approximate worldwide averages). For a typical community-dwelling population with the age and wealth of the United States we estimate the IFR to be approximately 0.42%.
- Any estimation of the COVID-19 IFR should take into account the various uncertainties and potential biases in both the mortality data and the seroprevalence data.
- Bayesian methods with interval censoring are well suited for complex evidence synthesis tasks such as estimating the COVID-19 IFR.

Funding - This work was supported by the European Union's Horizon 2020 research and innovation programme under ReCoDID grant agreement No 825746 (🇪🇺) and by the Canadian Institutes of Health Research, Institute of Genetics (CIHR-IG) under Grant Agreement No 01886-000.

Conflicts of interest/Competing interests - None declared.

Availability of data and material - Data used for the analysis is available in the Supplementary Material and at <https://github.com/harlanhappydog/BayesianSeroMetaAnalysis>.

Code availability - Code used for the analysis is available in the Supplementary Material and at <https://github.com/harlanhappydog/BayesianSeroMetaAnalysis>.

References

- [1] Berhanu Nega Alemu, Adamu Addissie, Gemechis Mamo, Negussie Deyessa, Tamrat Abebe, Abdulnasir Abagero, Wondimu Ayele, Workeabeba Abebe, Tewodros Haile, Rachel Argaw, Wondwossen Amogne, Ayele Belachew, Zelalem Desalegn, Brhanu Teka, Eva Kantelhardt, Mesfin Wossen, Saro Abdella, Getachew Tollera, and Lia Tadesse. Seroprevalence of anti-SARS-CoV-2 antibodies in Addis Ababa, Ethiopia. *bioRxiv*, 2020. doi:10.1101/2020.10.13.337287.
- [2] Carlos Álvarez-Antonio, Graciela Meza-Sánchez, Carlos Calampa, Wilma Casanova, Cristian Carey, Freddy Alava, Hugo Rodríguez-Ferrucci, and Antonio M Quispe. Seroprevalence of anti-sars-cov-2 antibodies in iquitos, peru in july and august, 2020: a population-based study. *The Lancet Global Health*, 9(7):e925–e931, 2021.

- [3] Carlos Álvarez-Antonio, Graciela Meza-Sánchez, Carlos Calampa, Wilma Casanova, Cristian Carey, Freddy Alava, Hugo Rodríguez-Ferrucci, and Antonio Marty Quispe. Seroprevalence of anti-SARS-CoV-2 antibodies in Iquitos, Loreto, Peru. *medrxiv*, 2021.
- [4] Abhishek Anand, Justin Sandefur, Arvind Subramanian, et al. Three new estimates of India’s all-cause excess mortality during the COVID-19 pandemic. *Center for Global Development - Working Paper*, 2021.
- [5] Ayesha Appa, Saki Takahashi, Isabel Rodriguez-Barraquer, Gabriel Chamie, Aenor Sawyer, Elias Duarte, Jill Hakim, Keirstinne Turcios, Joanna Vinden, Owen Janson, et al. Universal PCR and antibody testing demonstrate little to no transmission of SARS-CoV-2 in a rural community. In *Open Forum Infectious Diseases*, 2020.
- [6] Rahul K Arora, Abel Joseph, Jordan Van Wyk, Simona Rocco, Austin Atmaja, Ewan May, Tingting Yan, Niklas Bobrovitz, Jonathan Chevrier, Matthew P Cheng, et al. Serotracker: a global SARS-CoV-2 seroprevalence dashboard. *The Lancet Infectious Diseases*, 21(4): e75–e76, 2021.
- [7] Cathrine Axfors and John PA Ioannidis. Infection fatality rate of COVID-19 in community-dwelling populations with emphasis on the elderly: An overview. *medRxiv*, 2021.
- [8] Insa Backhaus, Nico Dragano, Friedrich Boege, Klaus Göbels, Derik Hermesen, Nadine Lübke, and Jörg Timm. First results of the SERODUS study (“erste ergebnisse der SERODUS-studie”). <https://www.uniklinik-duesseldorf.de/patienten-besucher/klinikeninstitutezentren/institut-fuer-medizinische-soziologie/das-institut/forschung/serodus>, 2021.
- [9] Kristina L Bajema, F Scott Dahlgren, Travis W Lim, Nicolette Bestul, Holly M Biggs, Jacqueline E Tate, Claudio Owusu, Christine M Szablewski, Cherie Drenzek, Jan Drobeniuc, et al. Comparison of estimated severe acute respiratory syndrome coronavirus 2 seroprevalence through commercial laboratory residual sera testing and a community survey. *Clinical Infectious Diseases*, 2020.
- [10] Murad Banaji. Estimating COVID-19 infection fatality rate in Mumbai during 2020. *medRxiv*, 2021.
- [11] Anton Barchuk, Dmitriy Skougarevskiy, Kirill Titaev, Daniil Shirokov, Yulia Raskina, Anastasia Novkunkskaya, Petr Talantov, Artur Isaev, Ekaterina Pomerantseva, Svetlana Zhikrivetskaya, et al. Seroprevalence of SARS-CoV-2 antibodies in Saint Petersburg, Russia: a population-based study. *medRxiv*, 2020.
- [12] Hilda Bastian. Peering through the smoke at a duel over COVID’s infection fatality rate. <https://absolutelymaybe.plos.org/2021/05/31/>

peering-through-the-smoke-at-a-duel-over-covids-infection-fatality-rate/, 2021.

- [13] Adeline Beaumont, Cécile Durand, Martine Ledrans, Valérie Schwoebel, Harold Noel, Yann Le Strat, Donatien Diulus, Léa Colombain, Marie Médus, Philippe Guedet, et al. High seroprevalence of anti-sARS-CoV-2 antibodies after the first wave of the COVID-19 pandemic in a vulnerable population in Perpignan, France. *medRxiv*, 2021.
- [14] Eran Bendavid, Bianca Mulaney, Neeraj Sood, Soleil Shah, Emilia Ling, Rebecca Bromley-Dulfano, Cara Lai, Zoe Weissberg, Rodrigo Saavedra, James Tedrow, et al. COVID-19 antibody seroprevalence in Santa Clara County, California. *medRxiv*, 2020.
- [15] James O Berger. *Statistical decision theory and Bayesian analysis*. Springer Science & Business Media, 2013.
- [16] Josse A Berlin. Invited commentary: benefits of heterogeneity in meta-analysis of data from epidemiologic studies. *American Journal of Epidemiology*, 142(4):383–387, 1995.
- [17] Rupam Bhattacharyya, Ritoban Kundu, Ritwik Bhaduri, Debashree Ray, Lauren J Beesley, Maxwell Salvatore, and Bhramar Mukherjee. Incorporating false negative tests in epidemiological models for SARS-CoV-2 transmission and reconciling with seroprevalence estimates. *Scientific Reports*, 11(1):1–14, 2021.
- [18] Holly M Biggs, Jennifer B Harris, Lucy Breakwell, F Scott Dahlgren, Glen R Abedi, Christine M Szablewski, Jan Drobeniuc, Nirma D Bustamante, Olivia Almendares, Amy H Schnell, et al. Estimated community seroprevalence of SARS-CoV-2 antibodies—two Georgia counties, April 28–May 3, 2020. *Morbidity and Mortality Weekly Report*, 69(29):965, 2020.
- [19] Niklas Bobrovitz, Rahul Krishan Arora, Christian Cao, Emily Boucher, Michael Liu, Claire Donnici, Mercedes Yanes-Lane, Mairead Whelan, Sara Perlman-Arrow, Judy Chen, et al. Global seroprevalence of SARS-CoV-2 antibodies: a systematic review and meta-analysis. *PloS one*, 16(6):e0252617, 2021.
- [20] Lysandro Pinto Borges, Aline Fagundes Martins, Mônica Santos de Melo, Makson Gleydson Brito de Oliveira, José Melquiades de Rezende Neto, Marcos Barbosa Dósea, Bruna Cecília Maia Cabral, Rafael Fontes Menezes, Aryanne Araujo Santos, Igor Leonardo Santos Matos, et al. Seroprevalence of SARS-CoV-2 IgM and IgG antibodies in an asymptomatic population in Sergipe, Brazil. *Revista Panamericana de Salud Pública*, 44, 2020.
- [21] N Brazeau, Robert Verity, Sara Jenks, Han Fu, Charles Whittaker, Peter Winskill, Ilaria Dorigatti, Patrick Walker, Steven Riley, Ricardo P Schnekenberg, et al. Report 34: COVID-19 infection fatality ratio: estimates from seroprevalence. <https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/covid-19/report-34-ifr/>, 2020.

- [22] Peter Brody-Moore. Bayesian hierarchical meta-analysis of asymptomatic Ebola seroprevalence. CMC Senior Theses. 2228. https://scholarship.claremont.edu/cmc_theses/2228, 2019.
- [23] Naomi C Brownstein and Yian Ann Chen. Predictive values, uncertainty, and interpretation of serology tests for the novel coronavirus. *Scientific Reports*, 11(1):1–12, 2021.
- [24] Tim A Bruckner, Daniel M Parker, Scott M Bartell, Veronica M Vieira, Saahir Khan, Andrew Noymer, Emily Drum, Bruce Albala, Matthew Zahn, and Bernadette Boden-Albala. Estimated seroprevalence of SARS-CoV-2 antibodies among adults in Orange County, California. *Scientific Reports*, 11(1):1–9, 2021.
- [25] Harlan Campbell, Perry de Valpine, Lauren Maxwell, Valentijn MT de Jong, Thomas Debray, Thomas Jänisch, and Paul Gustafson. Bayesian adjustment for preferential testing in estimating the COVID-19 infection fatality rate: Theory and methods. *Annals of Applied Statistics*, 2020.
- [26] Mireia G Carrasco, Cristina Royo-Cebrecos, Irene Ambatlle, Kyla Serres, Joel López, David Vilanova, Vanesa Arroyo, Marc Pons, Odile Sarroca, Eva Heras, et al. Assessment of the andorran response to sars-cov-2 pandemic.
- [27] Fabrice Carrat, Xavier de Lamballerie, Delphine Rahib, Helene Blanche, Nathanael Lapidus, Fanny Artaud, Sofiane Kab, Adeline Renuy, Fabien Szabo de Edelenyi, Laurence Meyer, et al. Seroprevalence of SARS-CoV-2 among adults in three regions of france following the lockdown and associated risk factors: a multicohort study. *medRxiv*, 2020.
- [28] Philip A Chan, Ewa King, Yizhen Xu, William Goedel, Leanne Lasher, Matt Vargas, Ken Brindamour, Richard Huard, Ailis Clyne, James McDonald, et al. Seroprevalence of SARS-CoV-2 antibodies in Rhode Island from a statewide random sample. *American Journal of Public Health*, 111(4):700–703, 2021.
- [29] Xinhua Chen, Zhiyuan Chen, Andrew S Azman, Xiaowei Deng, Ruijia Sun, Zeyao Zhao, Nan Zheng, Xinghui Chen, Wanying Lu, Tingyu Zhuang, et al. Serological evidence of human infection with SARS-CoV-2: a systematic review and meta-analysis. *The Lancet Global Health*, 9(5):E598–E609, 2021.
- [30] Hannah Clapham, James Hay, Isobel Routledge, Saki Takahashi, Marc Choisy, Derek Cummings, Bryan Grenfell, C Jessica E Metcalf, Michael Mina, Isabel Rodriguez Barraquer, et al. Seroepidemiologic study designs for determining SARS-CoV-2 transmission and immunity. *Emerging Infectious Diseases*, 26(9):1978, 2020.
- [31] Kostas Danis, Laure Fonteneau, Scarlett Georges, Côme Daniau, Sibylle Bernard-Stoecklin, Lisa Domegan, Joan O’Donnell, Siri Helene Hauge, Sara Dequeker, Eline Vandael, et al.

- High impact of COVID-19 in long-term care facilities, suggestion for monitoring in the EU/EEA, May 2020. *Eurosurveillance*, 25(22):2000956, 2020.
- [32] Owen Dyer. COVID-19: Peru’s official death toll triples to become world’s highest. *The BMJ*, 373(n1442), 2021.
- [33] Margarita Estévez-Abe and Hiroo Ide. COVID-19 and Japan’s small death toll in long-term care facilities. *Journal of Aging and Social Policy*. https://programs.wcfia.harvard.edu/files/us-japan/files/margarita_estevez-abe_covid19_and_japanese_ltcfs.pdf, 2021.
- [34] Hans J Eysenck. An exercise in mega-silliness. *American Psychologist*, 33(5):517, 1978.
- [35] Jeremy Samuel Faust. Comparing COVID-19 deaths to flu deaths is like comparing apples to oranges. *Scientific American*, <https://tinyurl.com/ydxx8e18>, 2020.
- [36] Barbara Fraser. COVID-19 strains remote regions of peru. *The Lancet*, 395(10238):1684, 2020.
- [37] John Geddie and Aradhana Aravindan. Why is Singapore’s COVID-19 death rate the world’s lowest. Reuters - <https://www.reuters.com/article/health-coronavirus-singapore-explainer-idUSKBN2680TF>, 2020.
- [38] Anne Gégout-Petit, Hélène Jeulin, Karine Legrand, Agathe Bochnakian, Pierre Vallois, Evelyne Schvoerer, and Francis Guillemin. Seroprevalence of SARS-CoV-2, symptom profiles and seroneutralization during the first COVID-19 wave in a suburban area, France. *medRxiv*, 2021.
- [39] Andrew Gelman and Eric Loken. The garden of forking paths: Why multiple comparisons can be a problem, even when there is no “fishing expedition” or “p-hacking” and the research hypothesis was posited ahead of time. *Department of Statistics, Columbia University*, 348, 2013.
- [40] Andrew Gelman et al. Prior distributions for variance parameters in hierarchical models (comment on article by Browne and Draper). *Bayesian Analysis*, 1(3):515–534, 2006.
- [41] Shabnam Ghasemyani, Mehdi Jafari, Ahmad Ahmadi Teymurlouy, and Reza Fadayevatan. Components of elderly long-term care system in Iran and selected countries: A comparative study. *Health Scope*, 10(3), 2021.
- [42] Sabine Gleich, Verena Turlik, Susann Schmidt, and Doris Wohlrab. COVID-19 deaths among residents of inpatient nursing homes in munich-causes and places of death. *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz*, 2021.

- [43] Govern d'Andorra. Els tests d'anticossos permeten diagnosticar 78 positius de la COVID-19, que podrien haver contagiats unes 360 persones. <https://tinyurl.com/dybe27uk>, 2020.
- [44] Daniel F Gudbjartsson, Gudmundur L Norddahl, Pall Melsted, Kristbjorg Gunnarsdottir, Hilma Holm, Elias Eythorsson, Asgeir O Arnthorsson, Dadi Helgason, Kristbjorg Bjarnadottir, Ragnar F Ingvarsson, et al. Humoral immune response to SARS-CoV-2 in Iceland. *New England Journal of Medicine*, 383(18):1724–1734, 2020.
- [45] Pedro C Hallal, Fernando P Hartwig, Bernardo L Horta, Mariângela F Silveira, Claudio J Struchiner, Luís P Vidaletti, Nelson A Neumann, Lucia C Pellanda, Odir A Dellagostin, Marcelo N Burattini, et al. SARS-CoV-2 antibody prevalence in Brazil: results from two successive nationwide serological household surveys. *The Lancet Global Health*, 8(11): e1390–e1398, 2020.
- [46] Yannick Hansen and Emery P. Dalesio. Covid deaths continue among elderly in care homes. Luxembourg Times, at: <https://www.luxtimes.lu/en/luxembourg/covid-deaths-continue-among-elderly-in-care-homes-6058bf8cde135b923604daba>, 2021.
- [47] Zhenyu He, Lili Ren, Juntao Yang, Li Guo, Luzhao Feng, Chao Ma, Xia Wang, Zhiwei Leng, Xunliang Tong, Wang Zhou, et al. Seroprevalence and humoral immune durability of anti-SARS-CoV-2 antibodies in Wuhan, China: a longitudinal, population-level, cross-sectional study. *The Lancet*, 397(10279):1075–1084, 2021.
- [48] Julian PT Higgins. Commentary: Heterogeneity in meta-analysis should be expected and appropriately quantified. *International Journal of Epidemiology*, 37(5):1158–1160, 2008.
- [49] Peter W Horby, Karen L Laurie, Benjamin J Cowling, Othmar G Engelhardt, Katharine Sturm-Ramirez, Jose L Sanchez, Jacqueline M Katz, Timothy M Uyeki, John Wood, Maria D Van Kerkhove, et al. CONSISE statement on the reporting of seroepidemiologic studies for influenza (ROSES-I statement): an extension of the STROBE statement. *Influenza and other respiratory viruses*, 11(1):2–14, 2017.
- [50] Charles Huamaní, Lucio Velasquez, Sonia Montes, Ana Mayanga-Herrera, and Antonio Bernabe-Ortiz. Population-based seroprevalence of SARS-CoV-2 antibodies in a high-altitude setting in Peru. *Available at SSRN 3760458*, 2021.
- [51] John P.A. Ioannidis. First Opinion: A fiasco in the making? as the coronavirus pandemic takes hold, we are making decisions without reliable data. STAT, <https://tinyurl.com/uj539o4>, 2020.
- [52] John PA Ioannidis. Infection fatality rate of COVID-19 inferred from seroprevalence data. *Bulletin of the World Health Organization*, 99(1):19, 2021.

- [53] John PA Ioannidis. Reconciling estimates of global spread and infection fatality rates of COVID-19: an overview of systematic evaluations. *European Journal of Clinical Investigation*, page e13554, 2021.
- [54] John PA Ioannidis and Joseph Lau. Can quality of clinical trials and meta-analyses be quantified? *The Lancet*, 352(9128):590, 1998.
- [55] Faheem Arshad Premkumar Raja Jayeeta Rajagopalan, Saadiya Hurzuk and Suvarna Alladi. The COVID-19 Long-Term Care situation in India. <https://ltccovid.org/wp-content/uploads/2020/05/LTC-COVID-situation-in-India-30th-May-1.pdf>; accessed Oct. 15, 2021, 2020.
- [56] Sitanshu Sekhar Kar, Sonali Sarkar, Sharan Murali, Rahul Dhodapkar, Noyal Mariya Joseph, and Rakesh Aggarwal. Prevalence and time trend of SARS-CoV-2 infection in Puducherry, India, August–October 2020. *Emerging Infectious Diseases*, 27(2):666, 2021.
- [57] Ariel Karlinsky and Dmitry Kobak. Tracking excess mortality across countries during the COVID-19 pandemic with the world mortality dataset. *eLife*, 10:e69336, jun 2021. ISSN 2050-084X. doi:10.7554/eLife.69336. URL <https://doi.org/10.7554/eLife.69336>.
- [58] Kazem Khalagi, Safoora Gharibzadeh, Davood Khalili, Mohammad Ali Mansournia, Siamak Mirab Samiee, Saeide Aghamohamadi, Maryam Mir-Mohammad-Ali Roodaki, Seyed Mahmoud Hashemi, Katayoun Tayeri, Hengameh Namdari Tabar, et al. Prevalence of COVID-19 in iran: Results of the first survey of the iranian COVID-19 serological surveillance program. *Clinical Microbiology and Infection*, 2021.
- [59] Jahidur Rahman Khan, Nabil Awan, Md Islam, Olav Muurlink, et al. Healthcare capacity, health expenditure, and civil society as predictors of COVID-19 case fatalities: A global analysis. *Frontiers in Public Health*, 8:347, 2020.
- [60] S Khan, Mariya Amin Qurieshi, Inaamul Haq, Sabhiya Majid, Javid Ahmad, Taha Ayub, Anjum Bashir Fazili, Ashfaq Ahmad Bhat, Abdul Majeed Ganai, Yasmeen Jan, et al. Results of a population-based survey to estimate the seroprevalence of SARS-CoV-2 specific IgG antibodies in Kashmir, India, seven months after the appearance of the first COVID-19 case. 2021.
- [61] Dmitry Kobak. Excess mortality reveals Covid’s true toll in Russia. *Significance*, 18(1):16, 2021.
- [62] John Kruschke. *Doing Bayesian data analysis: A tutorial with R, JAGS, and Stan*. Academic Press, 2014.

- [63] Matthias Kümmerer, Philipp Berens, and Jakob Macke. A simple Bayesian analysis of the infection fatality rate in Gangel, and an uncertainty aware extrapolation to infection-counts in Germany. <https://matthias-k.github.io/BayesianHeinsberg.html>, 2020.
- [64] Paul C Lambert, Alex J Sutton, Paul R Burton, Keith R Abrams, and David R Jones. How vague is vague? A simulation study of the impact of the use of vague prior distributions in mcmc using winbugs. *Statistics in Medicine*, 24(15):2401–2428, 2005.
- [65] David A Leon, Vladimir M Shkolnikov, Liam Smeeth, Per Magnus, Markéta Pechholdová, and Christopher I Jarvis. COVID-19: a need for real-time monitoring of weekly excess deaths. *The Lancet*, 395(10234):E81, 2020.
- [66] Andrew T Levin, William P Hanage, Nana Owusu-Boaitey, Kensington B Cochran, Seamus P Walsh, and Gideon Meyerowitz-Katz. Assessing the age specificity of infection fatality rates for COVID-19: systematic review, meta-analysis, and public policy implications. *European Journal of Epidemiology*, pages 1–16, 2020.
- [67] Andrew T Levin, Nana Owusu-Boaitey, Sierra Pugh, Bailey K Fosdick, Anthony B Zwi, Anup Malani, Satej Soman, Lonni Besançon, Ilya Kashnitsky, Sachin Ganesh, et al. Assessing the burden of covid-19 in developing countries: Systematic review, meta-analysis, and public policy implications. *medRxiv*, 2021.
- [68] Zhongjie Li, Xuhua Guan, Naiying Mao, Huiming Luo, Ying Qin, Na He, Zhen Zhu, Jianxing Yu, Yu Li, Jianhua Liu, et al. Antibody seroprevalence in the epicenter wuhan, hubei, and six selected provinces after containment of the first epidemic wave of COVID-19 in china. *The Lancet Regional Health-Western Pacific*, 8:100094, 2021.
- [69] Ruijie Ling, Yihan Yu, Jiayu He, Jixian Zhang, Sha Xu, Renrong Sun, Tao Li, Hong-Long Ji, and Huan-Qiang Wang. Seroprevalence and epidemiological characteristics of immunoglobulin M and G antibodies against SARS-CoV-2 in asymptomatic people in Wuhan, China. *China (6/15/2020)*, 2020.
- [70] Natalie M Linton, Tetsuro Kobayashi, Yichi Yang, Katsuma Hayashi, Andrei R Akhmetzhanov, Sung-mok Jung, Baoyin Yuan, Ryo Kinoshita, and Hiroshi Nishiura. Incubation period and other epidemiological characteristics of 2019 novel coronavirus infections with right truncation: a statistical analysis of publicly available case data. *Journal of Clinical Medicine*, 9(2):538, 2020.
- [71] Marc Lipsitch. First Opinion: We know enough now to act decisively against COVID-19. social distancing is a good place to start. STAT, <https://tinyurl.com/yx4gf9mr>, 2020.
- [72] Jiangmei Liu, Lan Zhang, Yaqiong Yan, Yuchang Zhou, Peng Yin, Jinlei Qi, Lijun Wang, Jingju Pan, Jinling You, Jing Yang, et al. Excess mortality in Wuhan city and other parts of

- china during the three months of the COVID-19 outbreak: findings from nationwide mortality registries. *The BMJ*, 372, 2021.
- [73] Mahajan et al. SARS-CoV-2 infection hospitalization rate and infection fatality rate among the non-congregate population in Connecticut. *The American Journal of Medicine*, 2021.
- [74] Mahajan et al. Seroprevalence of SARS-CoV-2-specific IgG antibodies among adults living in Connecticut: Post-infection prevalence (pip) study. *The American Journal of Medicine*, 134(4):526–534, 2021.
- [75] Hussaini Majiya, Mohammed Aliyu-Paiko, Vincent Tochukwu Balogu, Dickson Achimugu Musa, Ibrahim Maikudi Salihu, Abdullahi Abubakar Kawu, Ishaq Yakubu Bashir, Aishat Rabiou Sani, John Baba, Amina Tako Muhammad, et al. Seroprevalence of COVID-19 in Niger State. *medRxiv*, 2020.
- [76] Anup Malani, Daksha Shah, Gagandeep Kang, Gayatri Nair Lobo, Jayanthi Shastri, Manoj Mohanan, Rajesh Jain, Sachee Agrawal, Sandeep Juneja, Sofia Imad, et al. Seroprevalence of SARS-CoV-2 in slums versus non-slums in Mumbai, India. *The Lancet Global Health*, 9(2):e110–e111, 2021.
- [77] Valerio Marra and Miguel Quartin. A Bayesian estimate of the COVID-19 infection fatality rate in Brazil based on a random seroprevalence survey. *International Journal of Infectious Diseases*, 2021. doi:<https://doi.org/10.1016/j.ijid.2021.08.016>.
- [78] Colleen C McLaughlin, Margaret K Doll, Kathryn T Morrison, William L McLaughlin, Terry O'Connor, Anton M Sholukh, Emily L Bossard, Khamsone Phasouk, Emily S Ford, Kurt Diem, et al. High community SARS-CoV-2 antibody seroprevalence in a ski resort community, Blaine County, Idaho, US. preliminary results. *medRxiv*, 2020.
- [79] Roberto Melotti, Federica Scaggiante, Michela Falciani, Christian X Weichenberger, Luisa Foco, Stefano Lombardo, Alessandro De Grandi, Dorothee von Laer, Angelika Mahlknecht, Peter P Pramstaller, et al. Prevalence and determinants of serum antibodies to SARS-CoV-2 in the general population of the Gardena Valley. *medRxiv*, 2021.
- [80] Gideon Meyerowitz-Katz and Lea Merone. A systematic review and meta-analysis of published research data on COVID-19 infection-fatality rates. *International Journal of Infectious Diseases*, 101:138–148, 2020. ISSN 1201-9712. doi:<https://doi.org/10.1016/j.ijid.2020.09.1464>.
- [81] MoHoI Ministry of Health of Israel. National sero-epidemiological coverage for 19-c. https://www.gov.il/BlobFolder/reports/de-covid19-28062020-17092020/he/files_publications_corona_DE-covid19.pdf, 2020.

- [82] Peter Morfeld, Barbara Timmermann, J Valérie Groß, Philip Lewis, Pierluigi Cocco, and Thomas C Erren. COVID-19: Heterogeneous excess mortality and “burden of disease” in Germany and Italy and their states and regions, January–June 2020. *Frontiers in Public Health*, 9:477, 2021.
- [83] Bhramar Mukherjee, Soumik Purkayashtha, Ritoban Kundu, and Ritwik Bhaduri. Estimating the infection fatality rate from SARS-CoV-2 in India. Available at SSRN 3798552, 2021.
- [84] Zachary Munn, Sandeep Moola, Karolina Lisy, Dagmara Riitano, and Catalin Tufanaru. Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data. *International Journal of Evidence-Based Healthcare*, 13(3):147–153, 2015.
- [85] Manoj V Murhekar, Tarun Bhatnagar, Sriram Selvaraju, V Saravanakumar, Jeromie Wesley Vivian Thangaraj, Naman Shah, Muthusamy Santhosh Kumar, Kiran Rade, R Sabarinathan, Smita Asthana, et al. SARS-CoV-2 antibody seroprevalence in India, August–September, 2020: findings from the second nationwide household serosurvey. *The Lancet Global Health*, 9(3):e257–e266, 2021.
- [86] Murhekar et al. Prevalence of SARS-CoV-2 infection in India: Findings from the national serosurvey, May-June 2020. *Indian Journal of Medical Research*, 152(1):48, 2020.
- [87] Murhekar et al. SARS-CoV-2 antibody prevalence in India: Findings from the second nationwide household serosurvey, August-September 2020. *The Lancet Global Health*, 9(3): E257–E266, 2020.
- [88] Vivek Naranbhai, Christina C Chang, Wilfredo F Garcia Beltran, Tyler E Miller, Michael G Astudillo, Julian A Villalba, Diane Yang, Jeffrey Gelfand, Bradley E Bernstein, Jared Feldman, et al. High seroprevalence of anti-SARS-CoV-2 antibodies in Chelsea, Massachusetts. *The Journal of Infectious Diseases*, 222(12):1955–1959, 2020.
- [89] Nobutoshi Nawa, Jin Kuramochi, Shiro Sonoda, Yui Yamaoka, Yoko Nukui, Yasunari Miyazaki, and Takeo Fujiwara. Seroprevalence of sARS-CoV-2 IgG antibodies in Utsunomiya City, Greater Tokyo, after first pandemic in 2020 (u-corona): a household-and population-based study. *medRxiv*, 2020.
- [90] Muhammad Imran Nisar, MBBS Nadia Ansari, and Farah Khalid. Serial population-based sero-surveys for COVID-19 in low and high transmission. *medRxiv*, 2020.
- [91] Megan O’Driscoll, Gabriel Ribeiro Dos Santos, Lin Wang, Derek AT Cummings, Andrew S Azman, Juliette Paireau, Arnaud Fontanet, Simon Cauchemez, and Henrik Salje. Age-specific mortality and immunity patterns of SARS-CoV-2. *Nature*, pages 1–6, 2020.

- [92] Office of National Statistics. Coronavirus (COVID-19) infection survey: England. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/datasets/coronaviruscovid19infectionsurveydata>, 2020.
- [93] Gabriele Pagani, Andrea Giacomelli, Federico Conti, Dario Bernacchia, Rossana Rondanin, Andrea Prina, Vittore Scolari, Arianna Rizzo, Martina Beltrami, Camilla Caimi, et al. Prevalence of SARS-CoV-2 in an area of unrestricted viral circulation: Mass seroepidemiological screening in Castiglione d’Adda, Italy. *PloS one*, 16(2):e0246513, 2021.
- [94] Roberto Pastor-Barriuso, Beatriz Perez-Gomez, Miguel A Hernan, Mayte Perez-Olmeda, Raquel Yotti, Jesus Oteo, Jose Luis Sanmartin, Inmaculada Leon-Gomez, Aurora Fernandez-Garcia, Pablo Fernandez-Navarro, et al. SARS-CoV-2 infection fatality risk in a nationwide seroepidemiological study. *medRxiv*, 2020.
- [95] Sen Pei, Teresa K Yamana, Sasikiran Kandula, Marta Galanti, and Jeffrey Shaman. Overall burden and characteristics of COVID-19 in the United States during 2020. *medRxiv*, 2021.
- [96] Michael J Peluso, Saki Takahashi, Jill Hakim, J Daniel Kelly, Leonel Torres, Nikita S Iyer, Keirstinne Turcios, Owen Janson, Sadie E Munter, Cassandra Thanh, et al. SARS-CoV-2 antibody magnitude and detectability are driven by disease severity, timing, and assay. *medRxiv*, 2021.
- [97] Mayte Perez-Olmeda, Jose Maria Saugar, Aurora Fernández-García, Beatriz Perez-Gomez, Marina Pollan-Santamaria, Ana Avellon, Roberto Pastor-Barriuso, Nerea Fernández-de Larrea, Mariano Martín, Israel Cruz, et al. Evolution of antibodies against SARS-CoV-2 over seven months: experience of the nationwide seroprevalence ENE-COVID study in Spain. *medRxiv*, 2021.
- [98] Maria Skaalum Petersen, Marin Strøm, Debes Hammershaimb Christiansen, Jógvan Páll Fjallabak, Eina Hansen Eliassen, Malan Johansen, Anna Sofía Veyhe, Marnar Frídhheim Kristiansen, Shahin Gaini, Lars Fodgaard Møller, et al. Seroprevalence of SARS-CoV-2-specific antibodies, Faroe Islands. *Emerging Infectious Diseases*, 26(11):2760, 2020.
- [99] Patrick Pietzonka, Erik Brorson, William Bankes, Michael E Cates, Robert L Jack, and Ronjoy Adhikari. Bayesian inference across multiple models suggests a strong increase in lethality of COVID-19 in late 2020 in the UK. *medRxiv*, 2021.
- [100] Karl Pillemer, Lakshminarayanan Subramanian, and Nathaniel Hupert. The importance of long-term care populations in models of COVID-19. *JAMA*, 324(1):25–26, 2020.
- [101] Martyn Plummer. JAGS Version 4.0.0 user manual, 2015.

- [102] Marina Pollán, Beatriz Pérez-Gómez, Roberto Pastor-Barriuso, Jesús Oteo, Miguel A Hernán, Mayte Pérez-Olmeda, Jose L Sanmartín, Aurora Fernández-García, Israel Cruz, Nerea Fernández de Larrea, et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *The Lancet*, 396(10250):535–544, 2020.
- [103] Hossein Poustchi, Maryam Darvishian, Zahra Mohammadi, Amaneh Shayanrad, Alireza Delavari, Ayad Bahadorimonfared, Saeid Eslami, Shaghayegh Haghjooy Javanmard, Ebrahim Shakiba, Mohammad Hossein Somi, et al. SARS-CoV-2 antibody seroprevalence in the general population and high-risk occupational groups across 18 cities in iran: a population-based cross-sectional study. *The Lancet Infectious Diseases*, 21(4):473–481, 2021.
- [104] Priyanka Pulla. What counts as a COVID-19 death? *The BMJ*, 370, 2020.
- [105] Soumik Purkayastha, Ritoban Kundu, Ritwik Bhaduri, Daniel Barker, Michael Kleinsasser, Debashree Ray, and Bhramar Mukherjee. Estimating the wave 1 and wave 2 infection fatality rates from SARS-CoV-2 in India. *medRxiv*, 2021.
- [106] Xinyue Qi, Shouhao Zhou, and Martyn Plummer. A note on Bayesian modeling specification of censored data in JAGS. *arXiv preprint arXiv:2012.02074*, 2020.
- [107] Nouar Qutob, Faisal Awartani, Zaidoun Salah, Mohammad Asia, Imad Abu Khader, Khaled Herzallah, Nadeen Balqis, and Husam Sallam. Seroprevalence of SARS-CoV-2 in the West Bank region of Palestine: a cross-sectional seroepidemiological study. *BMJ open*, 11(2): e044552, 2021.
- [108] Katja Radon, Elmar Saathoff, Michael Pritsch, Jessica Michelle Guggenbuehl Noller, Inge Kroidl, Laura Olbrich, Verena Thiel, Maximilian Diefenbach, Friedrich Riess, Felix Forster, et al. Protocol of a population-based prospective COVID-19 cohort study Munich, Germany (koco19). *medRxiv*, 2020.
- [109] Sudarshan Ramaswamy, Aditya Athotra, PK Kasar, Rajesh Tiwari, Shashi Prabha Tomar, Ratnesh Kuraria, Arvind Rai, Meera Dhuria, Arti Bahl, Anil Digambar Patil, et al. Cross-sectional study on sero-prevalence of SARS-CoV-2 infection in Jabalpur, Madhya Pradesh, India. *Journal of Communicable Diseases (E-ISSN: 2581-351X & P-ISSN: 0019-5138)*, 53 (1):82–88, 2021.
- [110] Shay Reicher, Ronit Ratzon, Shay Ben-Sahar, Sharon Hermoni-Alon, David Mossinson, Yotam Shenhar, Michael Friger, Yaniv Lustig, Sharon Alroy-Preis, Emilia Anis, et al. Nationwide seroprevalence of antibodies against SARS-CoV-2 in Israel. *European Journal of Epidemiology*, pages 1–8, 2021.

- [111] Mary F Reyes-Vega, M Gabriela Soto-Cabezas, Fany Cárdenas, Kevin S Martel, Andree Valle, Juan Valverde, Margot Vidal-Anzardo, María Elena Falcón, César V Munayco, Peru COVID-19 Working Group, et al. SARS-CoV-2 prevalence associated to low socioeconomic status and overcrowding in an lmic megacity: A population-based seroepidemiological survey in Lima, Peru. *EClinicalMedicine*, 34:100801, 2021.
- [112] Aude Richard, Ania Wisniak, Javier Perez-Saez, Henri Garrison-Desany, Dusan Petrovic, Giovanni Piumatti, Helene Baysson, Attilio Picazio, Francesco Pennacchio, David De Ridder, et al. Seroprevalence of anti-SARS-CoV-2 IgG antibodies, risk factors for infection and associated symptoms in Geneva, Switzerland: a population-based study. *medRxiv*, 2020.
- [113] Tim Riffe, A Acosta, and Virginia Zarulli. COVerAGE-DB: A global demographic database of COVID-19 cases and deaths. *International Journal of Epidemiology*, 2021.
- [114] Eli S Rosenberg, James M Tesoriero, Elizabeth M Rosenthal, Rakkoo Chung, Meredith A Barranco, Linda M Styer, Monica M Parker, Shu-Yin John Leung, Johanne E Morne, Danielle Greene, et al. Cumulative incidence and diagnosis of SARS-CoV-2 infection in New York. *Annals of Epidemiology*, 48:23–29, 2020.
- [115] Cristina Royo-Cebrecos, David Vilanova, Joel López, Vanesa Arroyo, Marc Pons, Guillem Francisco, Mireia G Carrasco, Josep M Piqué, Sergi Sanz, Carlota Dobaño, et al. Mass SARS-CoV-2 serological screening, a population-based study in the Principality of Andorra. *The Lancet Regional Health-Europe*, 5:100119, 2021.
- [116] Matthew Samore, Adam Looney, Brian Orleans, Tom Greene, Nathan Seegert, Julio C Delgado, Angela Presson, Chong Zhang, Jian Ying, Yue Zhang, et al. SARS-CoV-2 seroprevalence and detection fraction in Utah urban populations from a probability-based sample. *medRxiv*, 2020.
- [117] Claudia Santos-Hövenner, Hannelore K Neuhauser, Angelika Schaffrath Rosario, Markus Busch, Martin Schlaud, Robert Hoffmann, Antje Gößwald, Carmen Koschollek, Jens Hoebel, Jennifer Allen, et al. Serology-and PCR-based cumulative incidence of SARS-CoV-2 infection in adults in a successfully contained early hotspot (comolo study), Germany, May to June 2020. *Eurosurveillance*, 25(47):2001752, 2020.
- [118] Sriram Selvaraju, Muthusamy Santhosh Kumar, Jeromie Wesley Vivian Thangaraj, Tarun Bhatnagar, Velusamy Saravanakumar, Chethrapilly Purushothaman Girish Kumar, Krithikaa Sekar, Ezhilarasan Ilayaperumal, Ramasamy Sabarinathan, Murugesan Jagadeesan, et al. Population-based serosurvey for severe acute respiratory syndrome coronavirus 2 transmission, chennai, india. *Emerging Infectious Diseases*, 27(2):586, 2021.

- [119] Maryam Shakiba, Seyed Saeed Hashemi Nazari, Fardin Mehrabian, Seyed Mahmoud Rezvani, Zahra Ghasempour, and Abtin Heidarzadeh. Seroprevalence of COVID-19 virus infection in Guilan province, Iran. *medRxiv*, 2020.
- [120] Nandini Sharma, Pragya Sharma, Saurav Basu, Sonal Saxena, Rohit Chawla, Kumar Dushyant, Nutan Mundeja, Z SK Marak, Sanjay Singh, Gautam Kumar Singh, et al. The seroprevalence and trends of SARS-CoV-2 in Delhi, India: A repeated population-based seroepidemiological study. *medRxiv*, 2020.
- [121] Gal Horowitz Shuli Brammli-Greenberg Sharona Tsadok-Rosenbluth, Boaz Hovav. Update on COVID-related mortality in care homes in Israel, 12th October 2020. <https://ltccovid.org/2020/10/14/update-on-covid-related-mortality-in-care-homes-in-israel-12th-october-2020/>, 2020.
- [122] Donald Sharpe. Of apples and oranges, file drawers and garbage: Why validity issues in meta-analysis will not go away. *Clinical Psychology Review*, 17(8):881–901, 1997.
- [123] Bonnie E Shook-Sa, Ross M Boyce, and Allison E Aiello. Estimation without representation: early severe acute respiratory syndrome coronavirus 2 seroprevalence studies and the path forward. *The Journal of Infectious Diseases*, 222(7):1086–1089, 2020.
- [124] Mariângela F Silveira, Aluísio JD Barros, Bernardo L Horta, Lúcia C Pellanda, Gabriel D Victora, Odir A Dellagostin, Claudio J Struchiner, Marcelo N Burattini, Andréia RM Valim, Evelise M Berlezi, et al. Population-based surveys of antibodies against SARS-CoV-2 in Southern Brazil. *Nature Medicine*, 26(8):1196–1199, 2020.
- [125] Mariangela Freitas Silveira, Marilia Mesenburg, Odir Antonio Dellagostin, Natasha R Oliveira, Mara Andrade Colares Maia, Francisco Denis Souza Santos, Andre M Vale, Aluisio Barros, Luis Paulo Vidaletti, Fernando C Barros, et al. Time-dependent decay of detectable antibodies against SARS-CoV-2: A comparison of elisa with two batches of a lateral-flow test. *The Brazilian Journal of Infectious Diseases*, 101601, 2021.
- [126] Chantal J Snoeck, Michel Vaillant, Tamir Abdelrahman, Venkata P Satagopam, Jonathan D Turner, Katy Beaumont, Clarissa PC Gomes, Joelle Veronique Fritz, Valerie E Schröder, Anne Kaysen, et al. Prevalence of SARS-CoV-2 infection in the Luxembourgish population: the CON-VINCE study. *medRxiv*, 2020.
- [127] Neeraj Sood, Paul Simon, Peggy Ebner, Daniel Eichner, Jeffrey Reynolds, Eran Bendavid, and Jay Bhattacharya. Seroprevalence of SARS-CoV-2-specific antibodies among adults in Los Angeles County, California, on april 10-11, 2020. *JAMA*, 323(23):2425–2427, 2020.

- [128] Nicola A Spiers. Rapid response: Comparing REACT with the ONS Infection Survey. *The BMJ*, 2020. URL <https://www.bmj.com/content/371/bmj.m3850/rr>.
- [129] Statistics Jersey. SARS-CoV-2: prevalence of antibodies in Jersey (community survey round 2). <https://tinyurl.com/6htjn3af>, 2020.
- [130] Statistics Jersey. SARS-CoV-2: prevalence of antibodies in Jersey (community survey round 3). <https://tinyurl.com/hxd8sadb>, 2020.
- [131] Paola Stefanelli, Antonino Bella, Giorgio Fedele, Serena Pancheri, Pasqualina Leone, Paola Vacca, Arianna Neri, Anna Carannante, Cecilia Fazio, Eleonora Benedetti, et al. Prevalence of SARS-CoV-2 IgG antibodies in an area of northeastern Italy with a high incidence of COVID-19 cases: a population-based study. *Clinical Microbiology and Infection*, 27(4): 633–e1, 2021.
- [132] Hendrik Streeck, Bianca Schulte, Beate M. Kümmerer, Enrico Richter, Tobias Höller, Christine Fuhrmann, Eva Bartok, Ramona Dolscheid-Pommerich, Moritz Berger, Lukas Wessendorf, Monika Eschbach-Bludau, Angelika Kellings, Astrid Schwaiger, Martin Coenen, Per Hoffmann, Birgit Stoffel-Wagner, Markus M. Nöthen, Anna M. Eis-Hübinger, Martin Exner, Ricarda Maria Schmithausen, Matthias Schmid, and Gunther Hartmann. Infection fatality rate of SARS-CoV-2 infection in a German community with a super-spreading event. *Nature Communications*, 11(1):5829, 2020. doi:10.1038/s41467-020-19509-y.
- [133] Silvia Stringhini, Ania Wisniak, Giovanni Piumatti, Andrew S Azman, Stephen A Lauer, Hélène Baysson, David De Ridder, Dusan Petrovic, Stephanie Schrempft, Kailing Marcus, et al. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (serocov-pop): a population-based study. *The Lancet*, 396(10247):313–319, 2020.
- [134] Silvia Stringhini, María-Eugenia Zaballa, Javier Perez-Saez, Nick Pullen, Carlos de Mestral, Attilio Picazio, Francesco Pennacchio, Ania Wisniak, Aude Richard, Helene Baysson, et al. Seroprevalence of anti-SARS-CoV-2 antibodies after the second pandemic peak. *The Lancet. Infectious Diseases*, 21(5):600–601, 2021.
- [135] Patrick Sullivan, Aaron Siegler, Kayoko Shioda, Eric W Hall, Heather Bradley, Travis H Sanchez, Nicole Luisi, Mariah Valentine-Graves, Kristin N Nelson, Mansour Fahimi, et al. SARS-CoV-2 cumulative incidence, United States, August-December 2020. *Clinical Infectious Diseases*, page ciab626, 2021.
- [136] Beatriz H Tess, Celso FH Granato, Maria CGP Alves, Maria C Pintao, Edgar Rizzatti, Marcia C Nunes, and Fernando C Reinach. SARS-CoV-2 seroprevalence in the municipality of São Paulo, Brazil, ten weeks after the first reported case. *medRxiv*, 2020.

- [137] The Economist. COVID-19 deaths in wuhan seem far higher than the official count. <https://www.economist.com/graphic-detail/2021/05/30/covid-19-deaths-in-wuhan-seem-far-higher-than-the-official-count>, 2021.
- [138] F Truc and Gianpiero Gervino. The effects of physical distancing and lockdown to restrain SARS-CoV-2 outbreak in the Italian municipality of Cogne. *Available at SSRN 3848514*, 2021.
- [139] Agne Ulyte, Thomas Radtke, Irene A Abela, Sarah R Haile, Christoph Berger, Michael Huber, Merle Schanz, Magdalena Schwarzmüller, Alexandra Trkola, Jan Fehr, et al. Clustering and longitudinal change in SARS-CoV-2 seroprevalence in school children in the canton of Zurich, Switzerland: prospective cohort study of 55 schools. *The BMJ*, 372, 2021.
- [140] Thomas R Vetter and Patrick Schober. Agreement analysis: what he said, she said versus you said. *Anesthesia & Analgesia*, 126(6):2123–2128, 2018.
- [141] Eric R A Vos, Gerco den Hartog, Rutger M Schepp, Patricia Kaaijk, Jeffrey van Vliet, Kina Helm, Gaby Smits, Alienke Wijmenga-Monsuur, Janneke D M Verberk, Michiel van Boven, Rob S van Binnendijk, Hester E de Melker, Liesbeth Mollema, and Fiona R M van der Klis. Nationwide seroprevalence of SARS-CoV-2 and identification of risk factors in the general population of the netherlands during the first epidemic wave. *Journal of Epidemiology & Community Health*, 75(6):489–495, 2021. ISSN 0143-005X. doi:10.1136/jech-2020-215678.
- [142] Patrick Alexander Wachholz, Alessandro Ferrari Jacinto, Ruth Caldeira de Melo, José Luis Dinamarca-Montecinos, and Paulo José Fortes Villas Boas. COVID-19: challenges in long-term care facilities for older adults in hispanic american countries. *Geriatrics, Gerontology and Aging*, 14(4):259–266, 2020.
- [143] Rochelle P Walensky, Henry T Walke, and Anthony S Fauci. SARS-CoV-2 variants of concern in the United States—Challenges and opportunities. *JAMA*, 325(11):1037–1038, 2021.
- [144] Xiaoli Wang, Wenjing Gao, Shujuan Cui, Yi Zhang, Ke Zheng, Ji Ke, Jun Lv, Canqing Yu, Dianjianyi Sun, Quanyi Wang, et al. A population-based seroprevalence survey of severe acute respiratory syndrome coronavirus 2 infection in Beijing, China. *medRxiv*, 2020.
- [145] Helen Ward, Graham Cooke, Christina J Atchison, Matthew Whitaker, Joshua Elliott, Maya Moshe, Jonathan C Brown, Barney Flower, Anna Daunt, Kylie EC Ainslie, et al. Declining prevalence of antibody positivity to SARS-CoV-2: a community study of 365,000 adults. *medRxiv*, 2020.

- [146] Josiane Warszawski, Nathalie Bajos, Laurence Meyer, et al. In May 2020, 4.5% of the population of metropolitan France had developed antibodies against SARS-CoV-2. <https://drees.solidarites-sante.gouv.fr/sites/default/files/2021-01/er1167-en.pdf>, 2020.
- [147] Daniel Westreich, Jessie K Edwards, Peter WG Tennant, Eleanor J Murray, and Maarten van Smeden. Choice of outcome in COVID-19 studies and implications for policy: Mortality and fatality. *American Journal of Epidemiology*, 2021.
- [148] Elizabeth J Williamson, Alex J Walker, Krishnan Bhaskaran, Seb Bacon, Chris Bates, Caroline E Morton, Helen J Curtis, Amir Mehrkar, David Evans, Peter Inglesby, et al. Factors associated with COVID-19-related death using opensafely. *Nature*, 584(7821): 430–436, 2020.
- [149] Joseph T Wu, Kathy Leung, Mary Bushman, Nishant Kishore, Rene Niehus, Pablo M de Salazar, Benjamin J Cowling, Marc Lipsitch, and Gabriel M Leung. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. *Nature Medicine*, 26(4):506–510, 2020.
- [150] Takashi Yoshiyama, Yasuki Saito, Kunitsugu Masuda, Yoshiko Nakanishi, Yasutoshi Kido, Kazuhiro Uchimura, Satoshi Mitarai, Tadaki Suzuki, Yu Nakagama, Hiroshi Kubota, et al. Prevalence of sARS-CoV-2-specific antibodies, Japan, June 2020. *Emerging Infectious Diseases*, 27(2):628, 2021.
- [151] Jon Zelner, Julien Riou, Ruth Etzioni, and Andrew Gelman. Accounting for uncertainty during a pandemic. *Patterns*, 2(8):100310, 2021.
- [152] Petra Zimmermann and Nigel Curtis. Why is COVID-19 less severe in children? a review of the proposed mechanisms underlying the age-related difference in severity of SARS-CoV-2 infections. *Archives of Disease in Childhood*, 106(5):429–439, 2021.

6 Supplementary Material

6.1 Excluded studies - Chen et al.-based analysis

Figure 1 shows a flowchart of the literature search and Table 6 lists all the excluded studies. Among the 38 articles obtained from the Chen et al. review, four studies represented results from different phases of the same study. For each of these we considered only the data from the earliest phase of the study. Stringhini et al. [133] and Richard et al. [112] are two publications that report the earlier and later phases, respectively, of the same study of Geneva, Switzerland. We considered only data from the earlier phase as reported in Stringhini et al. [133]. Murhekar et al. [86] and Murhekar et al.

[87] are two publications that report the earlier and later phases, respectively, of the same study in India. We considered only data from the first phase as reported in [86].

Note that, while similar in many ways, Ward et al. [145] and Office of National Statistics [92] are two different large-sample studies. The Ward et al. [145] study is based on the “REACT-2” survey which is led by Imperial College London, while the Office of National Statistics [92] study is based on the “Coronavirus (COVID-19) Infection Survey” which is conducted by a partnership between the University of Oxford, University of Manchester, Public Health England and Wellcome Trust. Spiers [128] discusses the differences between the two surveys.

We excluded eight studies that used “non-probability” or “convenience” sampling methods (e.g., studies in which participants were recruited using social media, or recruited in shopping centers). Specifically, we excluded: (1) McLaughlin et al. [78] who sampled individuals from a list of volunteers (Arora et al. [6] classifies the sampling method for this study as “self-referral”); (2) Rosenberg et al. [114] who sampled individuals at grocery stores (“convenience” [6]); (3) Appa et al. [5] who recruited volunteers with support from Bolinas community leaders (“self-referral” [6]); (4) Bendavid et al. [14] who recruited participants by placing targeted advertisements on Facebook (“stratified non-probability” [6]); (5) Gudbjartsson et al. [44] (“convenience” [6]); (6) Borges et al. [20] (“convenience” [6]); (7) Ling et al. [69] (“self-referral” [6]); and (8) Naranbhai et al. [88] (“convenience” [6]). Due to new information about the sensitivity of the Wondfo antibody tests (Silveira et al. [125]: “Our findings cast serious doubts about the use of this brand of rapid tests for epidemiological studies.”), we excluded the Hallal et al. [45] Brazil study (but do see: Marra and Quartin [77]). One study (Malani et al. [76]) was excluded due to a narrowly defined target population.

Two additional studies were not included because the articles failed to report 95% uncertainty intervals for the estimated infection rate in the target population [75, 144] and five additional studies were not included due to unavailable reliable mortality data for the specific target populations [1, 90, 103, 136, 119].

Author	Location	Reason for exclusion
Alemu et al. [1]	Addis Ababa, Ethiopia	Not included in Karlinsky and Kobak [57] analysis.
Appa et al. [5]	Bolinas, CA, USA	Convenience sampling (recruited volunteers with support from Bolinas community leaders (“self-referral” [6])).

Bendavid et al. [14]	Santa Clara county, CA, USA	Recruited participants by placing targeted advertisements on Facebook (“stratified non-probability”[6]).
Borges et al. [20]	Sergipe (10 cities), Brazil	Convenience sampling (“convenience” [6]) (but also: the 95% uncertainty interval not provided for the prevalence estimate).
Gudbjartsson et al. [44]	Iceland	Convenience sampling (“convenience” [6]).
Hallal et al. [45]	Brazil (83 cities)	Excluded due to new information about the sensitivity of the Wondfo antibody test [125].
Ling et al. [69]	Wuhan, China	Convenience sampling (“Self-referral” [6]) (but also: death data may be unreliable; see [72] and [137]).
Majiya et al. [75]	Nigeria	95% uncertainty interval not provided for the prevalence estimate.

Malani et al. [76]	Mumbai, India	The seroprevalence study provides two estimates: (1) for those living in the slums of the Matunga, Chembur West, and Dahisar Each wards; and (2) for those living in the non-slums of the Matunga, Chembur West, and Dahisar Each wards. These two target populations are too narrowly defined for available mortality data. Note, however, that some official mortality data appears at the ward level is available; see for example: “Ward-wise breakdown of positive cases” information in https://web.archive.org/web/20200625110946/http://stopcoronavirus.mcgm.gov.in/assets/docs/Dashboard.pdf (accessed August 4, 2021).
McLaughlin et al. [78]	Blaine, ID, USA	Convenience sampling (sampled individuals from a list of volunteers (“self-referral” Arora et al. [6])).
Murhekar et al. (B) [85]	India	Duplicate ([86] report the results from the same study).
Naranbhai et al. [88]	Chelsea, MA, USA	Convenience sampling (“convenience” [6]) (but also: 95% uncertainty interval not provided for the prevalence estimate).
Nisar et al. [90]	Two neighborhoods of Karachi, Pakistan	Not included in Karlinsky and Kobak [57] analysis.
Poustchi et al. [103]	Iran (18 cities)	Death data not found for the target population.
Richard et al. [112]	Geneva, Switzerland	Duplicate ([133] report results from the same study).

Rosenberg et al. [114]	New York state, USA	Convenience sampling (sampled individuals at grocery stores (“convenience” Arora et al. [6])).
Shakiba et al. [119]	Guilan Province, Iran	Death data not found.
Tess et al. [136]	Six districts of São Paulo, Brazil	Death data not found.
Wang et al. [144]	Beijing, China	95% uncertainty interval not provided for the prevalence estimate.

Table 6: List of excluded studies and reason for exclusion for the Chen et al. - based analysis.

6.2 Excluded studies - Serotracker-based analysis

Figure 2 shows a flowchart of the literature search and Table 7 lists all the excluded studies. Among the 45 articles obtained from the Serotracker review, six studies represented results from different phases of the same study. For each of these we considered only the data from the earliest phase of the study. For instance, Murhekar et al. [86] and Murhekar et al. [87] report the results from an earlier and a later phase of the same study in India. We only include Murhekar et al. [86] for our analysis. We excluded one additional study that used “convenience” sampling methods [131] and two studies that used the Wondfo antibody test [68, 124] (without adequate adjustment given the recent information about the sensitivity of the Wondfo test Silveira et al. [125]). We excluded two additional studies that had a narrowly defined target population[8, 139].

One additional study was not included because the article failed to report a 95% uncertainty interval for the estimated infection rate in the target population (Truc and Gervino [138] focused on estimating the longitudinal changes in antibodies levels rather than the prevalence rate in Cogne, Italy.). Note that the “Andorra” study [43] also failed to report a 95% uncertainty interval for the estimated infection rate in the target population. However, we chose to exceptionally include the study due to the fact that it represents one of the largest seroprevalence studies conducted: a total of 70,494 inhabitants (90.9% of the population of Andorra) participated in the study. In their published article Royo-Cebrecos et al. [115] explain that 95% confidence intervals are not provided because such intervals “would be extremely narrow and potentially misleading given that they do not account for the potential bias that non-participating individuals could cause on our central seroprevalence estimate.” We therefore defined, in order to be very cautious, a very wide (yet entirely arbitrary) 95% confidence interval of [10.5%, 11.5%] around the 11.0% point estimate. Eight additional studies were not included due to unavailable reliable mortality data for the specific target populations.

Author	Location	Reason for exclusion
Backhaus et al. [8]	Dusseldorf, Germany	Sample not representative of overall population: only "young people between the ages of 18-30" and individuals in the "fire brigade and rescue services".
Beaumont et al. [13]	Three neighbourhoods in the city of Perpignan, France	Death data not found for the target population (three specific neighbourhoods (Saint-Jacques, Haut-Vernet, and Nouveau Logis) in the city of Perpignan, France).
Gégout-Petit et al. [38]	Grand Nancy metropolitan area, France	Death data not found; however, note that coronadatascraper.com does have data for the larger region of Meurthe-et-Moselle, Grand Est, France.
He et al. [47]	Wuhan, China	Not included in Karlinsky and Kobak [57]'s analysis and see [72] and [137] who discuss issues as to why this is particularly challenging and controversial data to validate.
Huamaní et al. [50]	Cusco, Peru	Death data not found for the target population ("three settings in Cusco: (1) Cusco city, (2) the periphery of Cusco, and (3) Quillabamba city").

Khan et al. [60]	Vale Kashmir, India	Death data not found (official death numbers can be found on "Official Twitter handle of Department of Information and Public Relations, Govt of Jammu & Kashmir ("@diprjk"). While not specifically referenced, the numbers reported by this source are consistent with those used in Khan et al. [59].) Mukherjee et al. [83] does not provide an estimated underreporting factor for Kashmir noting that this is "[o]wing to lack of sufficient data".
Li et al. [68]	Wuhan City, Hubei-ex-Wuhan, and six provinces, China	Excluded due to new information about the sensitivity of the Wondfo antibody test Silveira et al. [125].
Murhekar et al. (B) [87]	India	Duplicate ([86] report the results from the same study).
Pérez-Olmeda et al. [97]	Spain	Duplicate ([102] report the results from the same study).
Poustchi et al. [103]	18 cities of Iran	Death data not found for the target population.
Qutob et al. [107]	Palestinian population residing in the West Bank	Not included in Karlinsky and Kobak [57]'s analysis.
Ramaswamy et al. [109]	Jabalpur Municipal Corporation ("City of Jabalpur")	While official death numbers are available for the District of Jabalpur (from the "Official Account of the Directorate of Health Services, Government of Madhya Pradesh. (e.g., https://twitter.com/healthminmp/status/1346107670869155843/photo/1 ; accessed July 27, 2021)), we could not find data for the City of Jabalpur.

Silveira et al. [124]	Regions of Brazil	Excluded due to new information about the sensitivity of the Wondfo antibody test [125].
Stefanelli et al. [131]	Five municipalities of the Autonomous Province of Trento, Italy	Convenience sampling.
Stringhini et al. (B) [134]	Geneva, Switzerland	Duplicate ([112] report the results from the same study).
Truc et al. [138]	Municipality of Cogne, Italy	95% uncertainty interval not provided for the prevalence estimate.
Ulyte et al. [139]	Zurich, Switzerland	Sample not representative of overall population: only school children.

Table 7: List of excluded studies and reason for exclusion for the Serotracker-based analysis.

6.3 Details on mortality and covariate data

- For Álvarez-Antonio et al. [3] (“Iquitos, Peru”) (see also Álvarez-Antonio et al. [2]), we used official (post-audit) numbers from the Peru Ministry of Health (MINSA) (see <https://www.datosabiertos.gob.pe/dataset/fallecidos-por-covid-19-ministerio-de-salud-minsa/resource/4b7636f3-5f0c-4404-8526>) for mortality numbers. We included deaths recorded for 4 districts (“SAN JUAN BAUTISTA”, “BELEN”, “PUNCHANA”, and “IQUITOS”) in the Loreto department. Note that recording the accurate number of deaths is perhaps particularly challenging in Iquitos; see Fraser [36]. In order to acknowledge that there is substantial uncertainty in the post-audit mortality numbers, we widened the interval by lowering the lower bound by 10% and increasing the upper bound by 10%. Limited information is available about mortality amongst LTC residents [142]. As such, no LTC-adjustments were made to the mortality numbers. Data for the proportion aged over 65 is from a 2004 World Health Organization report (see https://www.who.int/ageing/projects/intra/phase_two/alc_intra2_cp_peru.pdf; accessed August 2, 2021). GDP data is from the “USD per head, current prices, current PPP” value listed by the Organisation for Economic Co-operation and Development’s Gross Domestic Product, Large regions TL2 database (“OECD database”; see <https://stats.oecd.org/>, accessed August 10, 2021) for the Loreto region (PE16). Population data is from the published article [2].

- For Warszawski et al. [146] (“Metropolitan, France”), we obtained mortality data from the French government COVID-19 dashboard (see <https://dashboard.covid19.data.gouv.fr/vue-d-ensemble?location=FRA>; accessed on August 4, 2021), summing up the cumulative deaths for the 13 regions of Metropolitan France (i.e., European France). (235+394+1616+407+471+6710+3256+959+371+1562+469+836+56 and 257+427+1804+461+528+7358+3518+1032+411+1721+508+927+59). Note that this death total does not include deaths that occurred in LTC facilities. These are listed separately on the French government COVID-19 dashboard under “EHPAD et EMS”. Data for the proportion aged over 65 is for France from 2019 as listed by World Bank’s World Development Indicators (WDI) data (<https://data.worldbank.org/indicator/SP.P0P.65UP.TO.ZS?locations=FR>; accessed August 4, 2021). GDP data is from the OECD database. Population data is from INSEE in report titled “Population totale par sexe et âge au 1er janvier 2020, France métropolitaine” (see <https://www.insee.fr/fr/statistiques/1892088?sommaire=1912926>; accessed August 4, 2021).
- For Bajema et al. [9] (“DeKalb and Fulton, GA, USA (B)”), we obtained the number of deaths for DeKalb, and Fulton counties from the county-level COVID-19 dataset curated by the New York Times available at: github.com/nytimes/covid-19-data (accessed on April 28, 2021). According to the New York Times analysis, (see: <https://web.archive.org/web/20200627101803/https://www.nytimes.com/interactive/2020/us/coronavirus-nursing-homes.html>) about 45% of COVID-19 deaths in the state of Georgia were among nursing home residents (as of June 27, 2020). We therefore adjusted the numbers by 45% and obtained 122 ($=221-0.45*221$) and 136 ($=247-0.45*247$). Data for the proportion aged over 65 is from https://github.com/JieYingWu/COVID-19_US_County-level_Summaries/blob/master/data/README.md (accessed on April 29, 2021). GDP value is from value listed in the OECD database for the state of Georgia, USA (US13). Population is from the United States census (<https://www.census.gov/quickfacts/fultoncountygeorgia> and <https://www.census.gov/quickfacts/dekalbcountygeorgia> ; accessed July 28, 2021) (i.e., 759297 + 1063937). Note that both Bajema et al. [9] and Biggs et al. [18] report the results from the same seroprevalence study.
- For Barchuk et al. [11] (“Saint Petersburg, Russia”), we used excess death numbers as reported by Kobak [61] for the upper bound and official government numbers for the lower bound. Russian official government statistics appear to underestimate the true number of fatalities by a substantial factor [57]. No information was found for mortality amongst LTC residents in Russia; however see: <https://www.hrw.org/news/2020/06/02/russia-publish-data-about-covid-19-institutional-care>. Data for the pro-

portion aged over 65 is from Barchuk et al. [11] (see Table A3 “KOUZh-2018”). GDP value is from OECD database for the “Federal City of Saint Petersburg” (RU29) region. The population value is from https://en.wikipedia.org/wiki/Saint_Petersburg (accessed June 21, 2021)).

- For Biggs et al. [18] (“DeKalb and Fulton, GA, USA (A)”), the number of deaths for DeKalb, and Fulton counties was obtained from the county-level COVID-19 dataset curated by the New York Times available at: github.com/nytimes/covid-19-data (accessed on April 28, 2021). Data for the proportion aged over 65 is from https://github.com/JieYingWu/COVID-19_US_County-level_Summaries/blob/master/data/README.md (accessed on April 29, 2021). According to the New York Times analysis, (see: <https://web.archive.org/web/20200627101803/https://www.nytimes.com/interactive/2020/us/coronavirus-nursing-homes.html>; accessed Oct. 12, 2021) about 45% of COVID-19 deaths in the state of Georgia were among nursing home residents (as of June 27, 2020). We adjusted the numbers by 45% and obtained 122 ($=221-0.45*221$) and 136 ($=247-0.45*247$). Data for the proportion aged over 65 is from https://github.com/JieYingWu/COVID-19_US_County-level_Summaries/blob/master/data/README.md (accessed on April 29, 2021). GDP value is from value listed in the OECD database for the state of Georgia, USA (US13). Population is from the United States census (<https://www.census.gov/quickfacts/fultoncountygeorgia> and <https://www.census.gov/quickfacts/dekalbcountygeorgia>; accessed July 28, 2021) (i.e., 759297 + 1063937). Note that both Bajema et al. [9] and Biggs et al. [18] report the results from the same seroprevalence study.
- For Bruckner et al. [24] (“Orange County, CA, USA”), we obtained the number of cumulative deaths for Orange County from the Orange County Public Works (as referenced by Bruckner et al. [24]) at: data-ocpw.opendata.arcgis.com/datasets/2ec9342ffc814cf58161b1cca57365fd_0 (accessed on April 28, 2021). Orange County Public Works reports deaths of skilled nursing facility (SNF) residents separately. For 2020-07-24, there were a total of 759 deaths including 247 deaths of SNF residents. For 2020-08-30, there were a total of 1209 deaths including 370 deaths of SNF residents. We recorded mortality numbers accordingly. Data for the proportion aged over 65 is from https://github.com/JieYingWu/COVID-19_US_County-level_Summaries/blob/master/data/README.md (accessed on April 29, 2021). GDP value is from value listed for the state of California, USA (US06) from the OECD database. Population value obtained from the United States census (<https://www.census.gov/quickfacts/orangecountycalifornia>; accessed June 27, 2021).

- For Carrat et al. [27] (“Île-de-France, France”), we only consider Île-de-France phase of the study (see Supplementary Table 1 in Carrat et al. [27] for sampling dates). Data for the number of deaths for Île-de-France was obtained from the French government COVID-19 dashboard (see <https://dashboard.covid19.data.gouv.fr/vue-d-ensemble?location=FRA>; accessed on August 4, 2021). Note that this death total does not include deaths that occurred in long term care facilities (“EHPAD et EMS”). Data for the proportion aged over 65 is from <https://contrevues.paris/ile-de-france-comment-le-vieillissement-de-la-population-impacte-lhabitat/> (accessed on April 29, 2021). GDP value is from value listed for Île-de-France region (FR1) from OECD database. Population data from the Institut national de la statistique et des études économiques (INSEE) (see <https://www.insee.fr/fr/statistiques/5002478>; accessed July 29, 2021).
- For Chan et al. [28] (“Rhode Island, USA”), we obtained mortality data from the Rhode Island Department of Health available at <https://ri-department-of-health-covid-19-data-rihealth.hub.arcgis.com/> (accessed July 28, 2021). Note that Chan et al. [28] state that: “As of May 31, Rhode Island had reported 827 cumulative lab-confirmed SARS-CoV-2-involved deaths; 78.5% of these deaths were associated with congregate care facilities. We therefore reduced the mortality numbers by 78.5% ($= 684 \times (1 - 0.785)$ and $= 877 \times (1 - 0.785)$). Data for the proportion aged over 65 is from the U.S. Department of Commerce <https://www.census.gov/quickfacts/RI> (accessed on June 21, 2021). GDP value is from value listed for the state of Rhode Island, USA (US44) in the OECD database. The population value is obtained from the United States census (<https://www.census.gov/quickfacts/RI>; accessed July 28, 2021).
- For Govern d’Andorra [43] (“Andorra”) (see also Royo-Cebrecos et al. [115]), we obtained mortality numbers from the Government of Andorra (<https://www.govern.ad/coronavirus>; accessed on June 28, 2021) as tabulated by Wikipedia (https://en.wikipedia.org/wiki/COVID-19_pandemic_in_Andorra; accessed on June 28, 2021). We subtracted 31.45% of deaths ($35 = 51 - 0.3145 \times 51$ for both lower and upper bounds) following the reporting of Carrasco et al. [26] which found that 39 out of 124 COVID-19 deaths in Andorra (for the period between March, 2020 and April, 2021) were amongst residents in nursing homes. Carrasco et al. [26] note that: “218/354 (61.6%) residents in nursing homes were infected and the fatality rate was 39/218 (17.8%).” Data for the proportion aged over 65 is from https://www.indexmundi.com/andorra/demographics_profile.html (accessed on June 28, 2021) and GDP data is from The World Factbook (<https://statisticstimes.com/economy/countries-by-gdp-capita-ppp.php>; accessed on

June 28, 2021)). The population number is as stated in the Royo-Cebrecos et al. [115] which is the published article reporting on the study.

- For Kar et al. [56] (“Puducherry District, India”), data for the number of deaths for the Puducherry District was obtained from <https://covid19dashboard.py.gov.in/Reporting/District> (accessed on August 5, 2021). We multiplied the number recorded for 14 days after the end of the sampling window by a factor of 2.1 (based on the upper bound of Mukherjee et al. [83]’s estimated underreporting factor for Puducherry) in order to account for potential underreporting. As such, our interval is relatively wide and reflects the substantial uncertainty in the true number of deaths. There is insufficient information about deaths of LTC residents in India to make any LTC adjustments; see Jayeeta Rajagopalan and Alladi [55]. Data for the proportion aged over 65 is from <http://statisticstimes.com/demographics/india/puducherry-population.php> (accessed on June 22, 2021). GDP value is from value listed for Puducherry (IN34) from OECD database. Population for Puducherry District from Kar et al. [56] who note that: “Puducherry district, population $\approx$ 1.25 million, is located in southern India.”
- For Khalagi et al. [58] (“Iran”), Karlinsky and Kobak [57] suggest that official mortality data for Iran is highly inaccurate and suggest that there have been approx. 58,092 excess deaths in Iran for the period until Sept. 21, 2020. This coincides with the estimate provided by ghafari2020excess of 58,900 (95% CI 46,900 - 69,500) for the period from December 22, 2019 to September 21, 2020. Excess mortality data is only available in quarterly format from Kobak [61]. Therefore it is not possible to obtain mortality data for the specific dates we require: 2020-08-17 and 2020-11-14. Official statistics for these dates are 19,804 and 41,034 (see https://en.wikipedia.org/wiki/COVID-19_pandemic_in_Iran; accessed on June 28, 2021, which cites the Iranian Ministry of Health and Medical Education). We used 19,804 for the lower bound and multiplied 41,034 by Kobak [61]’s estimated undercount ratio of 2.4 to obtain an upper bound of 98071. (see <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7852240/pdf/nihpp-2021.01.27.21250604v3.pdf>; accessed on June 28, 2021). In Iran, LTC is provided mostly informally by family caregivers [41]. As such, no adjustments were made to exclude deaths of LTC residents. Data for the population and the proportion of the population aged over 65 is from the World Bank World Development Indicators (WDI) data (<https://data.worldbank.org/indicator/SP.POP.65UP.T0.ZS?locations=IR>; accessed on June 28, 2021). GDP value is also from the WDI data (see: <https://data.worldbank.org/indicator/NY.GDP.PCAP.PP.CD> (last updated 2021-04-26)).
- For Mahajan et al. [74] (“Connecticut, USA”), mortality data obtained from Wikipedia (https://en.wikipedia.org/wiki/COVID-19_pandemic_in_Connecticut; ac-

cessed on July 22, 2021), who reference (<https://portal.ct.gov/Coronavirus>). Mortality numbers (4,287 and 4,450) are multiplied by 26.5% (i.e. by 1079/4071) due to the fact that approx. 26.5% of COVID-19 deaths in Connecticut occurred in the community-dwelling population (i.e., “non-congregate” population) according to Mahajan et al. [73]. Data for the proportion aged over 65 is derived from Table 1 of Mahajan et al. [73]. GDP value is from value listed for the state of Connecticut, USA (US09) from the OECD database. Population value is obtained from [73].

- For Malani et al. [76] (“Tamil Nadu, India”), the population of Tamil-Nadu, India, was obtained from <https://www.indiacensus.net/states/tamil-nadu> (accessed July 27, 2021). Mortality data for the number of deaths for Tamil Nadu was obtained from the Tamil Nadu Government (see <https://stopcorona.tn.gov.in/archive/>; accessed June 24, 2021). We multiplied the number recorded for 14 days after the end of the sampling window of 11,909 by a factor of 2.5 (based on the upper bound of Mukherjee et al. [83]’s estimated underreporting factor for Tamil Nadu). There is insufficient information about deaths of LTC residents in India to make any LTC adjustments; see Jayeeta Rajagopalan and Alladi [55]. GDP value is from value listed for Tamil-Nadu (IN33) from the OECD database. Data for the proportion aged over 65 is from <https://statisticstimes.com/demographics/india/tamil-nadu-population.php> (accessed July 27, 2021).
- For Melotti et al. [79] (“Gardena Valley, Italy”), we obtained information from the report “Study on the prevalence of Covid-19 in the Val Gardena - June 2020” (see: https://astat.provincia.bz.it/it/news-pubblicazioni-info.asp?news_action=300&news_image_id=1074603; accessed July 27, 2021). In this report, it is noted that “the number of deaths from this disease in 17 people...”. Stefano Lombardo (in a personal communication) notes that the source of this number is Astat, Bolzano (“statistics on deaths by cause”) and has issued a correction (via personal communication) to indicate that the number should be 20 (“In spring 2020 the covid19-deaths in Gardena were estimated at 17; at the end of the year the official statistics closed at 20 (march+april).”) In a further personal communication (“R: R: INFERRING THE COVID-19 IFR”; Oct. 18, 2021), Lombardo clarified that out of these 20 deaths 3 occurred at home, 12 at the hospital, and 5 at a nursing home. The study looks specifically at three municipalities (Ortisei, Santa Cristina, and Selva) and we used age data from https://www.citypopulation.de/en/italy/trentinoaltoadige/bolzano/021089_selva_di_val_gardena/ and <http://italia.indettaglio.it/eng/trentinoaltoadige/ortisei.html> (accessed August 4, 2021). GDP value is from OECD database for the Province of Bolzano-Bozen (ITH1).

- For Ministry of Health of Israel [81] (“Israel”) (see also Reicher et al. [110]), mortality data obtained from https://en.wikipedia.org/wiki/COVID-19_pandemic_in_Israel (accessed July 27, 2021) which references Israel’s Ministry of Health’ official coronavirus updates, it’s Telegram channel, and Israel’s Corona National Information and Knowledge Center. According to Sharona Tsadok-Rosenbluth [121], a large proportion of COVID-19 deaths in Israel occurred in long-term care facilities (LTCFs). Specifically, by the end of July, 2020 (2020/07/28), the percentage of deaths that occurred in LTCFs was 51%, and by the end of September, 2020 (2020/09/29), the percentage of deaths that occurred in LTCFs was 40%. Based on these figures we adjusted the death numbers 382 and 1659 to: $187 (=382-0.51*382)$ and $995 (=1659-0.4*1659)$. Data for the proportion aged over 65 is from https://brookdale.jdc.org.il/wp-content/uploads/2018/02/MJB-Facts_and_Figures_Elderly-65_in_Israel-2018_English.pdf (accessed July 27, 2021) and GDP value is from OECD. Population for Israel is from World Bank’s World Development Indicators (WDI) data. Note that while [6] categorizes the sampling method of the Ministry of Health of Israel [81] study as “stratified probability,” Reicher et al. [110] note that the study used blood samples from “insured individuals who arrived at the HMOs [Health Maintenance Organizations] to undergo a blood test for any reason” and caution that, as a consequence, the “study might not reliably represent the entire population.”
- For Murhekar et al. [86] (“India”), we obtained the number of cumulative deaths for India, from the Our World in Data COVID-19 dataset available at: ourworldindata.org/coronavirus/country/india (accessed on April 28, 2021). We multiplied the number recorded for 14 days after the end of the sampling window of 12,573 by a factor of 3.64 (based on the upper bound of Purkayastha et al. [105]’s estimated underreporting factor for India (first wave)) in order to account for potential underreporting. As such, our interval is relatively wide and reflects the uncertainty in the true number of deaths: [4172, 45766]. (Purkayastha et al. [105]: “Estimates from epidemiological models: For wave 1 our estimate ... [is] an underreporting factor for cases estimated at 11.11 (95% CrI 10.71 – 11.47) and for deaths at 3.56 (95% CrI 3.48 – 3.64).”) There is insufficient information about deaths of LTC residents in India to make any LTC adjustments; see Jayeeta Rajagopalan and Alladi [55]. Data for the proportion aged over 65 is from World Bank’s World Development Indicators (WDI) data (see <https://data.worldbank.org/indicator/SP.POP.65UP.T0.ZS> (last updated 2021-03-19)). GDP value is from OECD database. Population value is from the World Bank (see <https://data.worldbank.org/indicator/SP.POP.TOTL?locations=IN>; accessed on April 28, 2021).
- For Nawa et al. [89], we concluded that 0 deaths had occurred based on official records for the Tochigi prefecture (see <https://www.nippon.com/en/japan-data/h00657/>; ac-

cessed July 12th, 2021). For the infection interval we use the lowest bound of the confidence interval given for the “unimputed dataset” and the upper bound of the confidence interval given for the “imputed dataset” from Table 1 of Nawa et al. [89]. Note that among the 181 participants in the Nawa et al. [89] study who were aged 65 years or older none were positive (0/181). This suggests that the prevalence amongst those at greatest risk of death (i.e., amongst the elderly) may be lower than in the greater population. However, since the Nawa et al. [89] found only 3 positive cases out of 742 individuals tested, inference on this is limited. Also note that, as of July 13, 2020, amongst the 36 individuals who tested positive by pcr-testing in Utsunomiya City, 4 were in their 70s and none were older than 80 (see <https://web.archive.org/web/20200714014725/https://www.city.utsunomiya.tochigi.jp/kurashi/kenko/kansensho/etc/1023506.html>). Data for proportion aged over 65 is from <https://www8.cao.go.jp/kourei/english/annualreport/2017/pdf/c1-1.pdf> (accessed July 12th, 2021). GDP value is from OECD database for Northern-Kanto, Koshin (JPC) region. Population value is from <https://www.city.utsunomiya.tochigi.jp/shisei/gaiyo/1007461.html> (accessed July 12th).

- For Office of National Statistics [92] (“England, UK (A)”), we used numbers obtained from the Office of National Statistics, dataset (“deaths registered weekly in England and Wales”; see publishedweek532020.xlsx, sheet “UK - Covid-19 - Weekly occurrences”) retrieved from <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/weeklyprovisionalfiguresondeathsregisteredinenglandandwales> (accessed Oct. 12, 2021). We subtracted those deaths attributed to Wales, and subtracted deaths attributed to nursing home residents as recorded in the “Deaths involving COVID-19 in the care sector, England and Wales” dataset (available at <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/deathsinvolvingcovid19inthecaresectorenglandandwales>; (accessed Oct. 12, 2021)). For 2020-05-15 (the closest available date to 2020-07-14), there were 41135 total deaths for England (= 43168 (England and Wales) - 2033 (Wales)) from which we subtracted 15160 deaths of care home residents (“England, ONS Data”) to obtain 25975. For 2020-09-25 (the closest available date to 2020-09-26), there were 50589 total deaths for England from which we subtracted 19880 deaths of care home residents (“England, ONS Data”) to obtain 30709. Data for the proportion aged over 65 is for England from 2019 as listed by LG Inform (see <https://tinyurl.com/4vhrb2uu>; accessed on April 29, 2021). GDP value is obtained by taking the average of values listed in the OECD database for the regions of “South West England”, “North East England”, “East of England”, “North West England”, and “South East England”. Population

value is from the Office of National Statistics report: “Population estimates for the UK, England and Wales, Scotland and Northern Ireland: mid-2020” (see Table 2) (<https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/bulletins/annualmidyearpopulationestimates/mid2020>; accessed July 30, 2021).

- For Pagani et al. [93] (“Castiglione d’Adda, Italy”), data was taken directly from the paper. Pagani et al. [93] does not cite a specific source for deaths but notes: “From the 1st of January to the 31st of March 2020, 76 deaths (1.65% of the population) have been recorded in CdA, of which 47 were officially attributed to COVID-19.” We have not been able to verify these numbers using a publicly available dataset. With regards to LTC deaths, Costantino Pesatori, the mayor of Castiglione D’Adda, is quoted as saying: “In our retirement home alone, we have had at least seven casualties.” (“Sono morti tanti anziani, è vero. Solo nella nostra casa di riposo abbiamo avuto almeno sette perdite.” See La Stampa: <https://www.lastampa.it/topnews/primo-piano/2020/03/16/news/la-strage-silenziosa-di-castiglione-d-adda-43-morti-in-tre-settimane-1.38599650>; accessed Oct. 12, 2021). With this in mind, we subtracted 7 and 8 deaths from the total of 47 cited in the paper. Note that it appears that 100% of residents in the nursing home were eventually infected (see https://www.ilcittadino.it/stories/Cronaca/covid-casa-di-riposo-di-castiglione-tutti-gli-anziani-contagiati_61445_96/). For population, Pagani et al. [93] states that “Castiglione d’Adda (CdA) is a town of 4605 inhabitants (according to data from the local registry office at the time of our study) [...]”. Data for the proportion aged over 65 is from https://www.citypopulation.de/en/italy/localities/lombardia/lodi/09801410001__castiglione_dadda/ (accessed July 27, 2021). GDP value is from OECD database for Lombardy region (ITC4).
- For Petersen et al. [98] (“Faroe Islands, Denmark”), information on deaths for the Faroe Islands was obtained from corona.fo/hagto1, the government information website concerning COVID19 in the Faroe Islands, which indicated zero deaths. GDP value is from value listed for Denmark from OECD database. Data for the proportion aged over 65 is from Index mundi (see <https://tinyurl.com/bb2pwx6>; accessed on April 29, 2021) which cites the CIA World Factbook as a source. Population value is from Petersen et al. [98] (“In the Faroe Islands, a geographic isolate of 52,154 inhabitants.”).
- For Pollán et al. [102] (“Spain”), data for the number of deaths was obtained from Wikipedia (en.wikipedia.org/wiki/COVID-19_pandemic_in_Spain; accessed on April 28, 2021) which sourced the information from the Centro Nacional de Epidemiología (cnecovid.isciii.es/covid19/). Note that, the number of deaths of for 2020-05-11 of 26,920 (14 days after the start of the sampling window), is actually higher than

the number of deaths for 2020-05-25 of 26,834 (14 days after the end of the sampling window). This may be due to a reporting issue which is noted by Wikipedia: “Figures for 2020-05-24 to 2020-06-17 include corrections in the validation of past data from several autonomous communities as a result of the transition to a new surveillance methodology implemented from 2020-05-11.” We subtracted 34% from the mortality numbers (26834, and 26920) to account for those deaths that occurred in long-term care facilities. This follows the practice done by Pastor-Barriuso et al. [94] who subtracted 9,909 deaths from a the total of 29,137; see discussion within Pastor-Barriuso et al. [94] for details. GDP value is from value listed for Spain from OECD database. Data for the proportion aged over 65 is from World Bank’s World Development Indicators (WDI) data (see <https://data.worldbank.org/indicator/SP.POP.65UP.TO.ZS> (last updated 2021-03-19)). Population value is from the World Bank for 2020 (<https://data.worldbank.org/indicator/SP.POP.TOTL?locations=ES>; accessed July 30, 2021).

- For Radon et al. [108] (“Munich, Germany”), the number of deaths was obtained from the official database of the city of Munich (see <https://www.muenchen.de/rathaus/Stadtfinfos/Coronavirus-Fallzahlen.html>; accessed June 22, 2021). Gleich et al. [42] conclude that: “One quarter of all COVID-19 deaths in Munich occurred in the context of nosocomial outbreaks in elderly, chronically ill residents of nursing facilities.” Based on this finding, we subtract 25% from the lower bound (292) and the upper bound (1040) to obtain the numbers: 219 and 780. Population data is from <https://web.archive.org/web/20201218195500/https://www.muenchen.de/rathaus/Stadtfinfos/Statistik/Bev-lkerung.html> (accessed June 22, 2021) and data for those aged over 65 from https://www.muenchen.de/rathaus/dam/jcr:459cce24-3894-4d4d-b144-4185b750a310/LHM.Stat_Faltkarte_2019_englisch.pdf (accessed June 22, 2021). Finally, GDP value is from value listed for Bavaria (GE2) from the OECD database.
- For Reyes-Vega et al. [111] (“Lima, Peru”), note that Peru audited death records and their data was updated on May 31, 2021; see Dyer [32] for details. We used official (post-audit) numbers from the Peru Ministry of Health (MINSA) (see <https://www.datosabiertos.gob.pe/dataset/fallecidos-por-covid-19-ministerio-de-salud-minsa/resource/4b7636f3-5f0c-4404-8526>; accessed June 22, 2021) for mortality numbers. Note that the target population is Lima Metropolitana, an area that includes both area that includes Peruvian provinces of Lima and Callao. As such, we sum deaths recorded for both Lima and Callao provinces. In order to acknowledge that there is substantial uncertainty in the post-audit mortality numbers, we widened the interval by lowering the lower bound

by 10% and increasing the upper bound by 10%. Limited information is available about mortality amongst LTC residents [142]. As such, no LTC-adjustments were made to the mortality numbers. GDP value is from value listed for Lima (PE15) from the OECD database. Data for the population aged over 65 from https://www.citypopulation.de/en/peru/admin/15__lima/ (accessed July 27, 2021). Population value obtained from https://www.minsa.gob.pe/reunis/data/poblacion_estimada.asp (accessed July 27, 2021). Note that Reyes-Vega et al. [111] state that: “The study area has an estimated 10.7 million inhabitants.”

- For Richard et al. [112] (“Geneva, Switzerland (B)”), note that this study is excluded from the Chen et al.-based analysis set but included in the Serotracker-based analysis set. For Richard et al. [112], mortality data for the canton of Geneva were obtained from an excel file made publicly available by a Swiss government website at: [ge.ch/document/covid-19-donnees-completes-debut-pandemie](https://www.ge.ch/document/covid-19-donnees-completes-debut-pandemie) (accessed on April 28, 2021). According to reporting from mid-Jun 2020 by the canton of Geneva (Département de la sécurité, de l’emploi et de la santé (DSES) - Direction générale de la santé - Service du médecin cantonal), about 45% of deaths were among nursing home residents (“résidents des établissements médico-sociaux (EMS)"); see document at <https://www.ge.ch/document/19696/annexe/61>. We therefore adjusted the numbers by 45%: 122 ($=222-0.45*222$) and 156 ($=283-0.45*283$). Data for the population and the proportion aged over 65 is from <https://www.bfs.admin.ch/bfs/en/home/statistics/regional-statistics/regional-portraits-key-figures/cantons/geneva.html> (accessed on April 28, 2021). GDP value is from value listed for Lake Geneva Region (CH01) from the OECD database.
- For Samore et al. [116] (“Four counties in UT, USA”), data for the number of deaths for the counties of Utah county, Salt Lake county, Davis county, and Summit county, was obtained from the county-level COVID-19 dataset curated by the New York Times available at: github.com/nytimes/covid-19-data (accessed on April 28, 2021). The Salt Lake Tribune reported on July 14, 2020 that 42% of COVID-19 deaths in Utah had been attributed to nursing homes (“According to the state health department, 95 of Utah’s 226 COVID-19 deaths have been attributed to the facilities.”; see <https://www.sltrib.com/news/2020/07/14/utah-nursing-homes-keep/>; accessed Oct. 12, 2021). This agrees with analysis by the New York Times (see: <https://web.archive.org/web/20200627101803/https://www.nytimes.com/interactive/2020/us/coronavirus-nursing-homes.html>; accessed Oct. 12, 2021). We therefore adjusted the mortality number by 42% and ob-

- tained 40 ($=69-69*0.42$) and 97 ($=168-168*0.42$). Data for the proportion aged over 65 is from https://github.com/JieYingWu/COVID-19_US_County-level_Summaries/blob/master/data/README.md (accessed on April 29, 2021). GDP value is from value listed for the state of Utah, USA (US49) in the OECD database. Population value is from US county-level census data for 2019 for the four counties (636235+1160437+355481+42145) (see <https://www.census.gov/data/datasets/time-series/demo/popest/2010s-counties-total.html>; accessed July 30, 2021).
- For Santos-Hövenner et al. [117] (“Kupferzell, Germany”), data for the number of deaths for Kupferzell, Germany was obtained directly from Santos-Hövenner et al. [117] which cites the Robert Koch Institute. Despite efforts, no publicly available dataset was found which could confirm these numbers specific these numbers. Santos-Hövenner et al. [117] report that 3 deaths occurred in the town for the period before July, 2020, and do not specify if these were nursing home residents. Santos-Hövenner et al. [117] do however specify the ages of those who died: “aged 59, 81 and 91 years.” Data for the proportion aged over 65 is from: <https://ugeo.urbistat.com/AdminStat/en/de/demografia/eta/kupferzell/20172564/4> (accessed on April 29, 2021). GDP value is from value listed for Baden-Württemberg (DE1) from the OECD database. Population value is from Wikipedia (<https://als.wikipedia.org/wiki/Kupferzell>; accessed July 30, 2021) which cites the Statistisches Landesamt Baden-Württemberg – Bevölkerung nach Nationalität und Geschlecht am 31. Dezember 2019 (CSV-Datei).
 - For Selvaraju et al. [118] (“Chennai, Tamil Nadu, India”), mortality data for the number of deaths for Chennai were obtained from the Greater Chennai Corporation (see <http://covid19.chennaicorporation.gov.in/>; specifically, numbers published for July 15, 2020 and August 14, 2020 were listed at: <https://twitter.com/chennaicorp/status/1283280173132128259> and <https://twitter.com/chennaicorp/status/1294140456868052993>; accessed June 22, 2021). We multiplied the number recorded for 14 days after the end of the sampling window of 2,384 by a factor of 2.5 (based the upper bound of Mukherjee et al. [83]’s estimated underreporting factor for Tamil Nadu) in order to account for potential underreporting. Note that Ariel Karlinsky has assembled excess mortality data for Chennai (see: https://github.com/akarlinsky/world_mortality/tree/main/local_mortality; accessed August 4, 2021). These data, taking into account the weekly numbers and five-year pre-pandemic trend, suggest an interval of [1344, 2454]. However, we prefer not to use this much narrower interval since the excess mortality data have not been adjusted to account for the potentially substantial underreporting of all-cause deaths (see discussion in Anand et al. [4]). There is insufficient information about deaths of LTC residents in India to make any LTC adjustments; see Jayeeta Rajagopalan and Alladi

- [55]. Data for the proportion aged over 65 is from: <https://statisticstimes.com/demographics/india/tamil-nadu-population.php> (accessed June 22, 2021). GDP value is from value listed for Tamil Nadu (IN33) from the OECD database. Population value is from <https://www.populationu.com/cities/chennai-population> (accessed June 22, 2021).
- For Sharma et al. [120] (“Delhi, India”), infection rate estimates are based on survey data from round 1 of the study (August 1-7). Data for the number of deaths for Delhi was obtained from Wikipedia (en.wikipedia.org/wiki/COVID-19_pandemic_in_Delhi; accessed on April 28, 2021) which sourced the information from the Delhi State Health Bulletin (<https://delhifightscorona.in/>). We multiplied the number recorded for 14 days after the end of the sampling window of 4,270 by a factor of 6.3 (based on Mukherjee et al. [83]’s estimated underreporting factor) in order to account for potential underreporting. As such, our interval is relatively wide and should reflect the uncertainty in the true number of deaths: [4,188, 26,901]. There is insufficient information about deaths of LTC residents in India to make any LTC adjustments; see Jayeeta Rajagopalan and Alladi [55]. Data for the proportion aged over 65 is from Statistics Times for 2011 (<http://statisticstimes.com/demographics/india/delhi-population.php>; accessed on April 29, 2021). GDP value is from value listed for the National Capital Territory of Delhi (IN03) from the OECD database. Note that we use 18.9 million as the estimated population of Delhi, the number cited by both Sharma et al. [120] and Bhattacharyya et al. [17], while Chen et al. [29] use a much different number: 30,290,936 (see Table S14 in Supplementary Appendix 2 of Chen et al. [29]).
 - Snoeck et al. [126] (“Luxembourg”) “recruited a representative sample of the Luxembourgish population” between April 16th and May 5th, and obtained a 95% CI of [1.23%, 2.77%]. Two different 95% CIs, obtained with and without adjustment for age, gender and canton are provided in the paper: [1.23%; 2.67%] and [1.34%; 2.77%]. As such, we record [1.23%; 2.77%] for our IR interval. Data for the number of deaths was obtained from the Our World in Data COVID-19 dataset available at: ourworldindata.org/coronavirus/country/luxembourg (accessed on April 28, 2021). Hansen and Dalesio [46] report that: “Of the 507 fatalities from Covid-19 in 2020, 241 had lived in housing offering care for the elderly, the Health Ministry reported to Parliament in January.” As such we adjusted the numbers by subtracting 47% to obtain 47 ($=89-0.47*89$) and 58 ($=109-0.47*109$). GDP value is from value listed for Luxembourg (LU00) from the OECD database. Data for the proportion aged over 65 is from World Bank’s World Development Indicators (WDI) data (see <https://data.worldbank.org/indicator/SP.POP.65UP.T0.ZS> (last updated 2021-03-19)). Population value from World Bank for Luxembourg, 2020 (see <https://data.worldbank.org/>).

- org/indicator/SP.POP.TOTL?locations=LU; accessed July 30, 2021). We note that, while Arora et al. [6] categorizes the sampling method for the Snoeck et al. [126] study as “stratified probability,” Snoeck et al. [126] explains that “the sample of participants was enrolled through the use of a non-probabilistic web panel” and caution that “[t]here are pros and cons of using web panels for surveys,” specifically that “online surveys may be biased samples because the respondents are self-selected.” Levin et al. [66] specifically identify Snoeck et al. [126] as an example of a “active recruitment” study.
- For Sood et al. [127] (“Los Angeles County, CA, USA”), data for the number of deaths for Los Angeles County, CA was obtained from the County of Los Angeles Public Health for 2020-04-24 (913 total deaths, and 384 deaths in institutional settings “as of 8pm 4/26”; see <https://web.archive.org/web/20200426230946/http://publichealth.lacounty.gov/media/coronavirus/locations.htm>; accessed Oct. 12, 2021) and for 2020-04-28 (1111 total deaths, and 505 in institutional settings “As of 8:00pm 4/29”; see <https://web.archive.org/web/20200501065125/http://publichealth.lacounty.gov/media/coronavirus/locations.htm>; accessed Oct. 12, 2021). To be clear, the County of Los Angeles Public Health reports deaths in institutional settings separately. Sood et al. [127] notes that “Residents of Los Angeles County, California, within a 15-mile (24 km) radius of the testing site were eligible for participation.” Data for the proportion aged over 65 is from https://github.com/JieYingWu/COVID-19_US_County-level_Summaries/blob/master/data/README.md (accessed on April 29, 2021). GDP value is from value listed for the state of California, USA, (US06) in the OECD database. Population value is from US county-level census data for 2019 (see <https://www.census.gov/data/datasets/time-series/demo/popest/2010s-counties-total.html>; accessed July 30, 2021).
 - For Statistics Jersey [129] (“Jersey, UK (A)”), data for the number of deaths was obtained from the Government of Jersey website (<https://www.gov.je/datasets/listopendata?listname=COVID19DeathsClassification>; accessed on April 28, 2021) summing both “probable COVID-19” deaths and “laboratory proven” COVID-19 deaths. According to reporting by the BBC on Apr. 27 2020 (see <https://t.co/2mj8YJbMsL?amp=1>; accessed Oct. 12, 2021), 7 out of 19 deaths occurred in care homes. With this in mind, we adjust our mortality numbers (27, and 28) by 37% to obtain 17 and 18. Data for the proportion aged over 65 is from https://www.indexmundi.com/jersey/demographics_profile.html (accessed on April 29, 2021). GDP value is from value listed for United Kingdom (GBR) in the OECD database. Population value is from Statistics Jersey [129].

- For Statistics Jersey [130] (“Jersey, UK (B)”) data for the number of deaths was obtained from the Government of Jersey website (<https://www.gov.je/datasets/listopendata?listname=COVID19DeathsClassification>; accessed on April 28, 2021) summing both “probable COVID-19” deaths and “laboratory proven” COVID-19 deaths. According to reporting by the BBC on Apr. 27 2020 (see <https://t.co/2mj8YJbMsL?amp=1>; accessed Oct. 12, 2021), 7 out of 19 deaths occurred in care homes. With this in mind, we adjust our mortality numbers (31, and 31) by 37% to obtain 20 and 20. Data for the proportion aged over 65 is from https://www.indexmundi.com/jersey/demographics_profile.html (accessed on April 29, 2021). GDP value is from value listed for United Kingdom (GBR) in the OECD database. Population value is from Statistics Jersey [130].
- For Streeck et al. [132], the number of deaths for Gangelt, Kreis Heinsberg, Germany, is directly noted in the Streeck et al. [132] article (n=8 deaths). No publicly available dataset was found, however we did find information on the Gangelt municipal bulletin (see www.gangelt.de/news/226-erster-corona-fall-in-nrw; accessed on April 28, 2021) which suggests that 8 is a reasonable number to consider. According to reporting, Gangelt did not suffer substantial (if any) fatalities in LTC facilities during the study period (MedWatch reports that during the study period: “there were no deaths of patients with Covid-19 in the Katharina-Kasper-Heim [nursing home] - nor in a nursing home with 80 residents, which is located in the Breberen district.” see: <https://medwatch.de/2020/05/08/wie-manche-virologen-vertrauen-verspielen/>; accessed on Oct. 18, 2021). Therefore, we did not subtract any deaths to account for nursing home residents. Note that the seroprevalence estimates for this study are based on the results of antibody tests as well as the results of PCR tests. As such the 14 day offset may not be as appropriate for this study as compared to other studies that consider only antibody tests. Data for the proportion aged over 65 is from Figure S1 of Streeck et al. [132]. GDP value is from value listed for the North Rhine-Westphalia region (DEA) from the OECD database. Population value is from Streeck et al. [132] (“In the German community of Gangelt (12,597 inhabitants, January 1, 2020)”).
- For Stringhini et al. [133] (“Geneva, Switzerland (A)”), mortality data for the canton of Geneva were obtained from an excel file made publicly available by a Swiss government website at: ge.ch/document/covid-19-donnees-completes-debut-pandemie (accessed on April 28, 2021). According to reporting from mid-June 2020 by the canton of Geneva (Département de la sécurité, de l’emploi et de la santé (DSES) - Direction générale de la santé - Service du médecin cantonal), about 45% of deaths were among nursing home residents (“résidents des établissements médico-sociaux

- (EMS)"); see document at <https://www.ge.ch/document/19696/annexe/61>. We adjusted the numbers by 45%: 122 ($=222-0.45*222$) and 155 ($=281-0.45*281$). Data for the proportion aged over 65 is from <https://www.bfs.admin.ch/bfs/en/home/statistics/regional-statistics/regional-portraits-key-figures/cantons/geneva.html>. GDP value is from value listed for Lake Geneva Region (CH01) from the OECD database. Population value is from the Swiss Federal Statistical Office (<https://www.bfs.admin.ch/bfs/en/home/statistics/regional-statistics/regional-portraits-key-figures/cantons/geneva.html>; accessed July 30, 2021).
- For Vos et al. [141], mortality data for the Netherlands was obtained from the National Institute for Public Health and the Environment <https://www.rivm.nl/coronavirus-covid-19/grafieken>; accessed Oct. 12, 2021. The numbers used are those listed under “Aantal overledenen naar datum van overlijden” (“Number of deceased by date of death”). Based on available information (see <https://www.verenso.nl/nieuws/archief/2020/update-registratie-verpleeghuizen-12-mei-2020> (accessed Oct. 12, 2021)) there were 1696 nursing home residents who died as of May 12, 2020. This represents about 30% of all reported COVID-19 deaths in the country (1696/5738) up until May 12, 2020. As such we will adjust our numbers by 30% : 2621 ($=3744-3744*0.30$) and 4186 ($=5980-5980*0.30$). The GDP value is from data listed for Netherlands (NLD) from the OECD database. Data for the proportion aged over 65 is also from World Bank’s World Development Indicators (WDI) data (see <https://data.worldbank.org/indicator/SP.POP.65UP.TO.ZS> (last updated 2021-03-19)). Population value is from the World Bank (<https://data.worldbank.org/indicator/SP.POP.TOTL?locations=NL>; accessed July 30, 2021).
 - For Ward et al. [145] (“England, UK (B)”), infection rate estimates are based on survey data from the first survey (20 June - 13 July). We used numbers obtained from the Office of National Statistics, dataset (“deaths registered weekly in England and Wales”; see publishedweek532020.xlsx, sheet “UK - Covid-19 - Weekly occurrences”) retrieved from <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/weeklyprovisionalfiguresondeathsregisteredinenglandandwales> (accessed Oct. 12, 2021). We subtracted those deaths attributed to Wales, and subtracted deaths attributed to nursing home residents as recorded in the “Deaths involving COVID-19 in the care sector, England and Wales” dataset (available at <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/deathsinvolvingcovid19inthecaresectorenglandandwales>; accessed Oct. 12, 2021)). For 2020-07-03 (the closest available date to 2020-07-04), there were 48718

total deaths for England (= 51207 (England and Wales) - 2489 (Wales)) from which we subtracted 19192 deaths of care home residents (“England, ONS Data”) to obtain 29526. For 2020-07-31 (the closest available date to 2020-07-27), there were 49593 total deaths for England from which we subtracted 19555 deaths of care home residents (“England, ONS Data”) to obtain 30038. Note that the numbers of cumulative deaths for England, from the UK coronavirus dashboard (<https://coronavirus.data.gov.uk/details/deaths?areaType=nation&areaName=England>; accessed on April 29, 2021) are somewhat different. For example, for 2020-07-04, the UK coronavirus dashboard reports 47,608 deaths for England “with COVID-19 in the death certificate” and 36,233 deaths for England that occurred “within 28 days of a positive test.” Data for the proportion aged over 65 is for England from 2019 as listed by LG Inform (see <https://tinyurl.com/4vhrb2uu>; accessed on April 29, 2021). GDP value is obtained by taking the average of values listed in the OECD database for the regions of “South West England”, “North East England” “East of England”, “North West England” and “South East England”. Population value is from the Office of National Statistics report: “Population estimates for the UK, England and Wales, Scotland and Northern Ireland: mid-2020” (see Table 2 at: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/bulletins/annualmidyearpopulationestimates/mid2020>; accessed July 30, 2021).

- For Yoshiyama et al. [150] (“Tokyo, Japan”), for the number of deaths for “2020-06-15” and “2020-06-21”, we used official numbers from the Japanese Ministry of Health, Labour, and Welfare (see: <https://www.mhlw.go.jp/content/10906000/000640012.pdf> and <https://www.mhlw.go.jp/content/10906000/000641965.pdf>; accessed August 6, 2021). Estévez-Abe and Ide [33] report: “As of May 8, only 14% of COVID-19-related deaths in Japan had occurred in LTCFs (Kyodo News Service, 2020).” We adjusted the numbers accordingly: 270 (=314-0.14*314) and 275 (=320-0.14*320). Data for the population and the proportion of the population aged over 65 is from the Tokyo metropolitan government (<https://www.metro.tokyo.lg.jp/tosei/hodohappyo/press/2021/01/28/01.html> and <https://www.metro.tokyo.lg.jp/ENGLISH/ABOUT/HISTORY/history03.htm>; accessed August 6, 2021) and GDP value is obtained by taking the average of values listed in the OECD database for the regions of “Northern-Kanto, Koshin” (JPC) and “Southern-Kanto” (JPD).

6.4 MCMC details and R code

Note that, in order to improve the MCMC mixing, we replace the binomial distribution for CC_k as described in (2), with

$$CC_k \sim \text{Binom}(T_k, IR_k), \quad (6)$$

for $k = 1, \dots, K$. For any sufficiently large P_k , this simplification will make little to no difference. Then, since the distributions of C_k and $D_k|C_k$ are both binomials (see (2) and (3)), we have that unconditionally:

$$D_k \sim \text{Binom}(P_k, IFR_k \times IR_k). \quad (7)$$

Note that Z_{1k} is set equal to the centred and scaled logarithm of $65yo_k$, and Z_{2k} is set equal to the centred and scaled logarithm of GDP_k .

The following JAGS-code is used in R for the analysis. See the JAGS user manual [101] for details about writing JAGS code (and also Qi et al. [106] for details about the specification of censored data in JAGS and the alternative modeling strategy necessary for correctly calculating the DIC).

```
library("rjags")
metaIFR <- "model {
# Priors:
icloglog_theta0 ~ dbeta(0.3, 30);
icloglog_beta ~ dbeta(1, 3);
theta0 <- log(-log(1-icloglog_theta0));
beta <- log(-log(1-icloglog_beta));
inv.var_sig <- (1/sigma)^2 ;
inv.var_tau <- (1/tau)^2 ;
sigma ~ dnorm(0, 1/10) T(0,);
tau ~ dnorm(0, 1/10) T(0,);
theta1 ~ dnorm(0, 1/10);
theta2 ~ dnorm(0, 1/10);

# Likelihood:
for(k in 1:K){
cc[k] ~ dbin(ir[k], tests[k]);
  censor.index[k] ~ dinterval(deaths[k], c(deaths_lower[k], deaths_upper[k]))
deaths[k] ~ dbin(ifr[k]*ir[k], pop[k]);
cloglog(ir[k]) <- cloglog_ir[k];
cloglog(ifr[k]) <- cloglog_ifr[k];
cloglog_ir[k] ~ dnorm(beta, inv.var_sig);
cloglog_ifr[k] ~ dnorm(theta0 + theta1*Z1[k] + theta2*Z2[k], inv.var_tau);
}}
```

6.5 Additional Figures

Figure 5: Analysis results from the Chen et al.-based analysis: posterior median estimates for the IR_k variables (for $k = 1, \dots, 19$) with 95% HPD CrIs. Studies are listed from top to bottom in the same order as in Figure 3.

Figure 6: Analysis results from the Serotracker-based analysis: posterior median estimates for the IR_k variables (for $k = 1, \dots, 28$) with 95% HPD CrIs. Studies are listed from top to bottom in the same order as in Figure 4.

6.6 Sensitivity analyses

As a sensitivity analysis, we repeated both analyses with an alternative set of priors. For these alternative analyses, we used: $g^{-1}(\theta_0) \sim \text{Uniform}(0, 1)$; $g^{-1}(\beta) \sim \text{Uniform}(0, 1)$; $\theta_1 \sim \mathcal{N}(0, 100)$; $\theta_2 \sim \mathcal{N}(0, 100)$; $\sigma \sim \text{half-}\mathcal{N}(0, 100)$ and $\tau \sim \text{half-}\mathcal{N}(0, 100)$. The results are plotted in Figures 9 and 10.

We also conducted alternative analyses excluding studies for which the mortality data were not obtained directly from official and reliable sources. See results in Figures 7 and 8. Without the excluded studies, we are unable to provide a reasonable “World” estimate (see the extremely wide credible intervals). However, the “USA” and “EU” estimates are relatively similar.

Figure 7: Chen et al.-based analysis results with only those studies for which mortality data was obtained from official sources known to be reliable: posterior median estimates (black circles) for the IFR_k variables (for $k = 1, \dots, 15$) with 95% HPD CrIs. Studies are listed from top to bottom in order of increasing fitted values (these values are indicated by $\times$). Also plotted, under the labels “World (65 yo=9%, GDP=17.8k)”, “USA (65 yo=16%, GDP=65.3k)”, “EU (65 yo=20%, GDP=47.8k)”, are the posterior median estimate and 95% HPD CrIs for the typical IFR corresponding to values for the proportion of the population aged 65 years and older of 9% and for GDP per capita of \$17,811 (the worldwide values), of 16% and of \$65,298 (the USA values), and of 20% and of \$47,828 (the EU values).

Figure 8: Serotracker-based analysis results with only those studies for which mortality data was obtained from official sources known to be reliable: posterior median estimates (black circles) for the IFR_k variables (for $k = 1, \dots, 16$) with 95% HPD CrIs. Studies are listed from top to bottom in order of increasing fitted values (these values are indicated by $\times$). Also plotted, under the labels “World (65 yo=9%, GDP=17.8k)”, “USA (65 yo=16%, GDP=65.3k)”, “EU (65 yo=20%, GDP=47.8k)”, are the posterior median estimate and 95% HPD CrIs for the typical IFR corresponding to values for the proportion of the population aged 65 years and older of 9% and for GDP per capita of \$17,811 (the worldwide values), of 16% and of \$65,298 (the USA values), and of 20% and of \$47,828 (the EU values).

Figure 9: Analysis results from the Chen et al.-based analysis with alternative priors: posterior median estimates (black circles) for the IFR_k variables (for $k = 1, \dots, 19$) with 95% HPD CrIs. Studies are listed from top to bottom according to increasing fitted values (these values are indicated by $\times$). Also plotted, under the labels “World (65 yo=9%, GDP=17.8K)”, “USA (65 yo=16%, GDP=65.3K)”, “EU (65 yo=20%, GDP=47.8K)”, are the posterior median estimate and 95% HPD CrIs for the typical IFR corresponding to values for the proportion of the population aged 65 years and older of 9% and for GDP per capita of \$17,811 (the worldwide values), of 16% and of \$65,297 (the USA values), and of 20% and of \$47,828 (the EU values).

Figure 10: Analysis results from the Serotracker-based analysis with alternative priors: posterior median estimates (black circles) for the IFR_k variables (for $k = 1, \dots, 28$) with 95% HPD CrIs. Studies are listed from top to bottom according to increasing fitted values (these values are indicated by $\times$). Also plotted, under the labels “World (65 yo=9%, GDP=17.8K)”, “USA (65 yo=16%, GDP=65.3K)”, “EU (65 yo=20%, GDP=47.8K)”, are the posterior median estimate and 95% HPD CrIs for the typical IFR corresponding to values for the proportion of the population aged 65 years and older of 9% and for GDP per capita of \$17,811 (the worldwide values), of 16% and of \$65,297 (the USA values), and of 20% and of \$47,828 (the EU values).