

Supplementary Figure 1. Donor age.
Age of the donors analyzed in the first cohort of individuals vaccinated with ChAdOx1, mRNA-1273, and BNT162b2. Mean age for each timepoint group is represented. Statistic comparisons were carried out for matched time PD between vaccines: +, ChAd vs MO; #, ChAd vs BNT; *, MO vs BNT.



Supplementary Figure 2. Time post-dose collection of samples related to the effect of the age of the donor in the humoral response.

(A) Time post-dose collection of samples for the groups represented in figure 4.

(B) IgG1 and IgA titers of seropositive samples collected after the first (PD1) or the second (PD2) dose of BNT vaccine. Only 2 to 3 weeks PD samples were analyzed in the same way as for figure 4.



Supplementary Figure 3. Relation between IgM and IgG1 or IgA titers.
(A) IgG1 ratio for IgM seronegative (395) and IgM seropositive (24) samples from all BNT donors, naïve or infected pre-vaccination.
(B) Linear regression adjustment of IgM and IgG1 or IgA ratios for all IgM seropositive BNT donors (24), naïve or infected pre-vaccination.



Supplementary Figure 4. Effect the donor's gender in the humoral response.
IgG1 and IgA titers of seropositive samples collected after the first (PD1) or the second (PD2) dose of BNT vaccine grouped by sex of the donor.