Supplementary Digital Content -

Table S1. Prevalence of psychosocial risks (PSR) among SARS-CoV-2 infected and non-infected healthcare workers and association with work-related psychological distress

	COVID-19 HCWs		Non-COVID-19 HCWs						
	Prevalence	High work-related		Prevalence High wo		h work-re	ork-related		
	of PSR	psych	nological	distress	of PSR	psychological c		distress	
	Total	Total	Men	Women	Total	Total	Men	Women	
N	4068	4068	838	3230	4152	4152	537	3615	
Psychosocial risks	Column %	Line %	Line %	Line %	Column %	Line %	Line %	Line %	
High psychological demands	37.5	56.5	53.4	57.2	39.8	67.1	61.2	68.0	
Low or moderate psychological demands	62.5	24.0	17.1	25.9	60.2	32.8	25.6	33.9	
Low or moderate decision authority	54.8	41.3	34.1	42.9	49.5	54.6	44.4	56.0	
High decision authority	45.2	30.0	25.7	31.3	50.6	38.5	34.9	39.0	
Low reward	32.5	53.7	48.2	55.0	35.0	63.7	53.1	65.1	
Moderate or high reward	67.5	27.7	22.0	29.4	65.0	37.2	32.9	37.8	
Low or moderate co-worker support	16.9	51.1	51.2	51.1	17.4	63.6	56.8	64.6	
High co-worker support	83.1	33.1	26.2	35.0	82.6	42.8	35.8	43.9	
Low or moderate supervisor support	22.6	55.5	50.3	56.6	22.5	66.4	60.3	67.4	
High supervisor support	77.4	30.5	24.9	32.0	77.6	40.7	33.1	41.8	
Difficult work-life balance	26.3	63.2	53.0	65.8	35.3	71.0	64.0	72.0	
Easy work-life balance	73.8	26.5	21.8	27.8	64.7	33.0	28.0	33.8	
Not the means to do quality work	73.9	43.9	36.5	45.8	81.0	52.1	44.7	53.1	
Having the means to do quality work	26.1	14.2	11.6	14.9	19.0	22.3	22.4	22.3	
Work against professional conscience	50.3	53.1	46.9	54.6	54.1	60.5	54.4	61.3	
Not working against professional	49.7	19.0	15.1	20.2	45.9	29.9	24.5	30.8	
conscience									
	Prevalence of PSR	Very h psvch	nigh work Nological (-related distress	Prevalence Very high wo of PSR psychologica		nigh work Nological (rk-related	
	Total	Total	Men	Women	Total	Total	Men	Women	
Psychosocial risks	Column %	Line %	Line %	Line %	Column %	Line %	Line %	Line %	
High psychological demands	37.5	17.1	16.0	17.4	39.8	24.8	15.8	26.1	
Low or moderate psychological demands	62.5	4.7	2.8	5.2	60.2	6.4	4.3	6.7	
Low or moderate decision authority	54.8	12.0	9.4	12.6	49.5	19.3	11.2	20.5	
High decision authority	45.2	6.1	5.5	6.3	50.6	8.2	6.5	8.4	
Low reward	32.5	18.3	17.1	18.6	35.0	24.8	15.4	26.1	
Moderate or high reward	67.5	5.0	3.2	5.5	65.0	7.7	5.5	8.0	
Low or moderate co-worker support	16.9	19.7	17.9	20.0	17.4	27.5	20.0	28.7	
High co-worker support	83.1	7.3	5.6	7.7	82.6	10.8	6.3	11.4	
Low or moderate supervisor support	22.6	19.9	14.7	21.0	22.5	27.0	20.6	28.0	

High supervisor support	77.4	6.3	5.6	6.4	77.6	9.8	5.1	10.5
Difficult work-life balance	26.3	22.4	20.0	23.0	35.3	26.6	18.6	27.7
Easy work-life balance	73.8	4.7	3.1	5.1	64.7	6.6	4.1	7.0
Not the means to do quality work	73.9	11.3	9.1	11.9	81.0	15.4	10.0	16.1
Having the means to do quality work	26.1	3.8	2.7	4.1	19.0	6.5	4.8	6.8
Work against professional conscience	50.3	14.6	12.1	15.2	54.1	19.6	12.5	20.6
Not working against professional conscience	49.7	4.0	3.3	4.2	45.9	6.7	4.9	7.0

Abbreviations: PSR=Psychosocial risk

Note 1: The psychosocial risks identified refer to Karasek's job demand-control-support model and Siegrist's effort-reward imbalance model, as well as to conflicts of values and difficulties in reconciling work and family.

Note 2: Chi-square test of bivariate association between each psychosocial risk and work-related psychological distress <0.05 for all the proportions in the table; chi-square test of bivariate association between each psychosocial risk and COVID-19 status <0.05 for all proportions except co-worker and supervisor support.

Table S2. Prevalence of psychosocial risks and items asked to measure them

	Men	Women	Total
Psychosocial risks and items to measure them	n = 1375	n = 6845	n = 8220
	%	%	%
1. Level of psychological demands			
Low	33.4	31.2	31.6
Moderate	30.0	29.8	29.8
High	36.6	39.1	38.6
Items to measure the level of psychological demands			
1.1 Constant pressure due to a heavy workload	37.0	40.9	40.2
1.2 Conflicting demands	32.7	32.1	32.2
1.3 Work that requires working very quickly	56.5	56.0	56.1
1.4 Not enough time to do the job	32.6	38.5	37.5
1.5 Work that requires hard work (mental of physical demands)	74.6	76.0	75.7
2. Level of decision authority (or job control)			
Low or moderate	49.2	52.7	52.1
High	50.8	47.3	47.9
Items to measure the level of decision authority			
2.1 Lack of choice over the way the job is done	67.6	65.5	65.8
2.2 Lack of influence in decisions about his/her work	64.3	60.7	61.3
3. Job strain			
Yes	21.8	24.8	24.3
No	78.2	75.2	75.7
4. Level of reward			
Low	31.0	34.3	33.8
Moderate	37.7	39.7	39.4
High	31.4	26.0	26.9
Items to measure the level of reward			
4.1 Job insecurity	17.8	18.0	18.0
4.2 Poor job promotion prospects	45.1	49.9	49.1
4.3 Unsatisfactory salary/income	50.4	55.8	54.9

	Men	Women	Total	
Psychosocial risks and items to measure them	n = 1375	n = 6845	n = 8220	
	%	%	%	
4.4 Do not receive the respect and prestige deserved at work	25.2	25.9	25.7	
5. Level of co-worker support				
Low or moderate	15.9	17.4	17.2	
High	84.2	82.6	82.9	
Items to measure the level of coworkers support				
5.1 No or little help and support from coworkers	9.5	12.2	11.8	
5.2 Little or no feeling of being part of a team at work	10.9	10.5	10.6	
6. Level of supervisor support				
Low or moderate	21.0	22.9	22.5	
High	79.0	77.2	77.5	
Items to measure the level of supervisor support				
6.1 No or little help and support from the immediate superior	16.2	18.1	17.8	
6.2 Immediate superior is rarely or never willing to listen	17.0	18.1	17.9	
7. Work and family/personal life conflict (difficulty maintaining a				
balance between work obligations and personal and family				
responsibilities)				
Yes	28.2	31.3	30.8	
No	71.9	68.7	69.2	
 Do not have the means to do quality work (always, often or sometimes) 	74.6	78.1	77.5	
9. Forced to work in a way that offends their professional	48.0	53.1	52.2	
conscience (always, often or sometimes)		0011	52.2	

Note 1: The psychosocial risks identified refer to Karasek's job demand-control-support model and Siegrist's effort-reward imbalance model, as well as to conflicts of values and difficulties in balancing work and family.

Table S3. Prevalence of psychosocial risks among SARS-CoV-2 infected healthcare workers by sex and type of occupation and comparison with 2014/15 Québec's population health survey (QPHS)

	Total	Se	ex	Type of occupation				
	N = 8220	Men n = 1375	en Women Nursin 1375 n = 6845 n = 214		HCSW n = 1572	Physicians n = 381	Admin n = 955	
Psychosocial risks	%	%	%	%	%	%	%	
High psychological demands	38.6	36.6	39.1	52.2	41.2	36.2	31.4	
QPHS 2014/15 all active workers	32.2	30.1	34.5					
QPHS 2014/15 HCSA workers	40.3	33.9	41.9					
Low or moderate decision authority	52.1	49.2	52.7	57.5	61.6	34.9	37.2	
QPHS 2014/15 all active workers	30.6	26.8	34.8					
QPHS 2014/15 HCSA workers	36.0	31.9	37.0					
Low reward	33.8	31.0	34.3	37.6	42.8	4.7	27.9	
QPHS 2014/15 all active workers	20.9	18.9	23.1					
QPHS 2014/15 HCSA workers	24.1	21.5	24.7				-	

	Total	Sex			Type of occupation			
	N = 8220	Men n = 1375	Women n = 6845	Nursing n = 2140	HCSW n = 1572	Physicians n = 381	Admin n = 955	
Low or moderate co-worker support	17.2	15.9	17.4	13.7	23.0	8.4	18.5	
QPHS 2014/15 all active workers	18.9	18.4	19.4					
QPHS 2014/15 HCSA workers	18.1	17.8	18.1					
Low or moderate supervisor support	22.5	21.0	22.9	27.0	24.7	12.6	17.1	
QPHS 2014/15 all active workers	21.2	21.1	21.2					
QPHS 2014/15 HCSA workers	23.9	22.9	24.1					
Difficult work-life balance	30.8	28.2	31.3	40.7	23.2	47.5	24.8	
Not the means to do quality work	77.5	74.6	78.1	87.2	69.2	79.5	73.9	
Work against their professional conscience	52.2	48.0	53.1	66.9	53.4	52.0	36.9	

Abbreviations: Adm= Administration and management staff; HCSW= healthcare support workers; HCSA workers= healthcare and social assistance workers; Nursing= nurses and nursing assistants; QPHS= Québec's population health survey

^a Source : Institut national de santé publique du Québec. Risques psychosociaux du travail : définition, indicateurs et résultats de l'Enquête québécoise sur la santé de la population 2014-2015. Institut national de santé publique du Québec. 2021; en préparation.

Table S4. Prevalence ratios of high (Kessler scale score \geq 7) work-related psychological distress according to the perception of the risk of acquiring COVID-19 at work and of some infection prevention and control measures

Picks factors avaluated	Prevalence a	(proportion who gree)	HIGH work-related psychological distress		
Risks factors evaluated	Covid-19 Non- HCWs 19 I		PR^{a}	95 % CI	
High or very high perceived risk of acquiring COVID-19 at work (reference= low or moderate)	29.2%	15.7%	1.3	1.2 - 1.4	
Personal protective equipment not available	4.7%	3.6%	0.9	0.8-1.1	
Insufficient infection prevention and control training	11.8%	13.5%	1.1	1.0 - 1.2	
Insufficient practical training about the use of personal protective equipment	26.6%	35.7%	1.0	1.0 - 1.1	
Lack of information about identification of COVID-19 confirmed or suspected patients	17.5%	18.4%	1.2	1.1 - 1.2	
Difficult access to COVID-19 screening test	9.0%	7.2%	1.1	1.0-1.2	
Physical environment not allowing physical distancing during breaks and meals	19.6%	22.4%	1.2	1.1 - 1.3	
Insufficient human resources to ensure patient care and HCW safety	36.4%	39.5%	1.5	1.4 - 1.6	
Persistent COVID-19 symptoms longer than 4 weeks	49.2%	NA	1.3	1.2 - 1.4	
COVID-19 status (reference= non-COVID-19)	51.8%	48.2	0.7	0.7 - 0.8	

Abbreviations: CI= confidence interval; HCW=Healthcare worker; PR= Prevalence ratio;

^a A robust Poisson model including all risk factors and adjusted for sex, age (18-44, 45-59, ≥60 years), race/ethnicity (white, black, other) and type of occupation (nursing, healthcare support worker, physician, administration and management staff, other). An interaction strategy was used to account for structural missing values for persistent COVID-19 symptoms among non-COVID-19 healthcare workers.