

1 Supplementary Material

2

3 Metagenomic analysis of antimicrobial resistance genes in wastewaters in Benin and Burkina Faso indicates a
4 serious health risk from untreated hospital wastewaters in low-income countries

5

6 Melina A. Markkanen^{1*}, Kaisa Haukka^{1,6}, Katariina M. M. Pärnänen¹, Victorien Tamegnon Dougnon², Isidore
7 Juste O. Bonkougou³, Zakaria Garba⁴, Halidou Tinto⁴, Anniina Sarekoski¹, Antti Karkman¹, Anu Kantele^{5,6†},
8 Marko P. J. Virta^{1*†}

9

10 ¹Department of Microbiology, University of Helsinki, Finland

11 ²Research Unit in Applied Microbiology and Pharmacology of Natural Substances, Polytechnic School of
12 Abomey-Calavi, University of Abomey-Calavi, Benin

13 ³Department of Biochemistry and Microbiology, University Joseph KI-ZERBO, Ouagadougou, Burkina Faso

14 ⁴Clinical Research Unit Nanoro, Institute for Research in Health Sciences, National Center for Scientific and
15 Technological Research, Burkina Faso

16 ⁵Meilahti Infectious Diseases and Vaccine Research Center, MeiVac, Department of Infectious Diseases,
17 University of Helsinki and Helsinki University Hospital, Helsinki, Finland

18 ⁶Human Microbiome Research Program, Faculty of Medicine, University of Helsinki, Helsinki, Finland

19

20 *Author information*

21 Corresponding authors:

22 Marko P. J. Virta

23 Melina A. Markkanen

24

25 † Equal contribution

26

27 Figure S1. Sample collection in Benin and Burkina Faso.

28 Figure S2. Examples of sampling sites.

29 Figure S3. PCoA plots describing the ordinations based on resistomes, taxonomical compositions, and mobilomes.

30 Table S1. Significance levels for ordination plots.

31 Figure S4. Unique and shared ARGs between hospital wastewaters in Benin, Burkina Faso, and Finland.

32 Figure S5. Comparative analysis of contigs that harbor plasmid-mediated *mcr-5.1* colistin resistance gene in four
33 different metagenomic samples

34 Table S2: Relative abundances of plasmid-mediated colistin ARGs in Benin and Burkina Faso.

35

36 Pages: 7

37 Tables: 2

38 Figures: 5

39

40



41

42 **Figure S1: Sample collection in Benin and Burkina Faso.** Sample collection sites are denoted with red dots on
 43 the map. The climate in the coastal region of Benin is hot and humid. The average daytime temperature in Cotonou
 44 varies between 28–32 °C and humidity between 77–84 %, with an annual rainfall of 1,308 mm. Burkina Faso is
 45 characterized by a dry tropical climate. The average daytime temperature in Ouagadougou varies between 31–
 46 39 °C, and the annual rainfall is 745 mm. During the sampling period in November/December, it typically rains
 47 only a little in Cotonou and not at all in Ouagadougou.

48

49

50

51



52

53

54

55

56 **Figure S2: Examples of sampling sites.** The samples collected in Benin included **a)** HWW from an open cesspit
 57 next to the wall at the hospital B (sample BH29); **b)** moist soil between two septic tanks (samples BH31 and
 58 BH34) at hospital B; **c)** the central septic tank collecting wastewater from different sections of hospital C (sample
 59 BH44); **d)** water pool surrounding the surgery room septic tank at hospital C (sample BH48); **e)** water from a tank
 60 used for washing hands at hospital C (sample BH52); **f)** tap water in a distant village in Benin (sample BSE93);
 61 and **g)** well water 100 m away from hospital A in Benin (sample BH11). **h)** In Burkina Faso, the HWW septic
 62 tank is shown for hospital F (sample BFH1). Photos by K. Haukka, A. Sarekoski and A. Butcher.

63

64


65

66

67

68

69



70
 71
 72

73 **Figure S3: PCoA plots describing the ordinations based on a) resistomes, b) taxonomical compositions, and**
 74 **c) mobilomes present in HWWs from Benin, Burkina Faso, and Finland.** Countries are coded as colors, and
 75 the specific hospitals are marked with a corresponding capitalized letter. Confidence ellipses represent 90 %
 76 confidence levels.

77


78 **Table S1: Significance levels for ordination plots.** The significance was measured using equal size sample
 79 groups (n=8/country) of randomly selected samples. The triplicated tests were mainly in line with each other,
 80 suggesting that sample selection did not significantly affect the results. The significance levels are denoted as
 81 follows: ‘****’ ≤ 0.001; ‘***’ = 0.001-0.01; ‘**’ = 0.01-0.05; ‘.’ = 0.05-0.1; ‘ ’ = 0.1-1.

ARGs, Horn									
	sample set #1			sample set #2			sample set #3		
Benin-Finland	0.29492	0.001	***	0.30982	0.002	**	0.33668	0.001	***
Benin-Burkina Faso	0.09321	0.121		0.17575	0.002	**	0.16236	0.007	**
Burkina Faso-Finland	0.24544	0.001	***	0.25117	0.001	***	0.25585	0.001	***

Taxonomical composition, Horn									
	sample set #1			sample set #2			sample set #3		
Benin-Finland	0.17282	0.002	**	0.20603	0.001	***	0.18496	0.002	**
Benin-Burkina Faso	0.08476	0.159		0.14289	0.007	**	0.08541	0.175	
Burkina Faso-Finland	0.16481	0.001	***	0.14788	0.006	**	0.16983	0.002	**

MGEs Horn									
	sample set #1			sample set #2			sample set #3		
Benin-Finland	0.31312	0.002	**	0.34187	0.001	***	0.34196	0.001	***
Benin-Burkina Faso	0.06114	0.497		0.10201	0.098	.	0.0994	0.094	.
Burkina Faso-Finland	0.20934	0.001	***	0.22952	0.001	***	0.20678	0.002	**


82
83
84




85
86
87
88
89
90
91

Figure S4: Unique and shared ARGs between hospital wastewaters in Benin, Burkina Faso, and Finland.
A total of 2,331 ARGs were detected among 67 hospital wastewater samples in the three countries. The number of unique ARGs was substantially higher (333) in Burkina Faso compared to Benin (59) or Finland (126)

92
93





94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

Figure S5: Comparative analysis of contigs that harbor plasmid-mediated *mcr-5.1* colistin resistance gene in four different metagenomic samples. Visualizations of the contigs were created using anvio (v7) metagenomics workflow for single profiles¹. Putative encoding genes are indicated by blue arrows. The gene names were determined based on BLAST search results. The recruited reads are shown as a dark grey histogram with an overlay line (pink) denoting the GC content. Single nucleotide variation is shown on top of the panel with colourful vertical lines. Inverse repeat sequences of Tn3-like transposase sequences (38 bp) are shown with dark pink (IRL: 5' GGGGTCGTCTCAGAAAACGGAAAAAATCGTACGCTAAG 3') and green triangles (IRR: 5'

112 CTTAGCGTACGATTTTTTCCGTTTTCTAGACGACCCC 3'²). The genetic environments are shown for
 113 samples a) BH02 b) BH48, c) BFH26 d) BFH41, and e) FH9.

114

115 **Table S2: Relative abundances of plasmid-mediated colistin ARGs in a) Benin and b) Burkina Faso.**

Gene variant	Relative abundance															
	Benin					Burkina Faso										
Gene variant	Feces samples		River water, drinking		Well water, drinking		Tap water		Street gutter, 100 m from hospital		Mold soil between HWW septic tanks		Water used for handwashing		Water middle surrounding the surgery room septic	
	BH20, Hosp. B	BH22, Hosp. B	BH24, Hosp. B	BH25, Hosp. B	BHE100	BSE74	BSE79	BH11, Hosp. A	BSE93	BH13 (water), Hosp. B	BH14 (sediment), Hosp. B	BH22, Hosp. B	BH52, Hosp. C	BH48, Hosp. C		
<i>mcr-1</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>mcr-3</i>	0	1.52 x 10 ³	0	4.74 x 10 ⁵	0	0	0	2.43 x 10 ⁵	0	0	0	6.79 x 10 ⁵	1.77 x 10 ³	1.96 x 10 ²	0	0
<i>mcr-3.17</i>	0	0	2.25 x 10 ⁵	0	3.22 x 10 ⁵	0	0	0	0	0	0	0	0	4.23 x 10 ⁴	0	0
<i>mcr-4</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>mcr-5</i>	0	0	2.22 x 10 ⁵	4.68 x 10 ⁵	0	1.10 x 10 ⁴	4.19 x 10 ⁵	0	0	5.48 x 10 ⁵	0	8.9 x 10 ⁵	7.76 x 10 ⁵	1.99 x 10 ³	0	0
<i>mcr-7</i>	0	0	0	0	0	3.73 x 10 ⁵	0	9.73 x 10 ⁵	2.25 x 10 ³	0	2.27 x 10 ⁵	3.94 x 10 ⁵	7.07 x 10 ⁵	0	0	0
<i>mcr-8</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>mcr-9</i>	0	0	0	0	3.23 x 10 ⁵	0	0	0	0	0	0	0	0	0	0	0
<i>mcr-10</i>	0	6.12 x 10 ³	0	0	0	0	4.25 x 10 ⁵	4.87 x 10 ³	0	0	0	0	3.94 x 10 ⁵	0	0	1.77 x 10 ³

Gene variant	Burkina Faso			
	Municipality channel, after biological treatment	Wetland receiving treated wastewater (from BFH26)	Connection point between HWW and WWTP channel	River receiving treated wastewater (from BFH41)
Gene variant	BFH26, Hosp. I	BFH27, Hosp. I	BFH41, Hosp. J	BFH42, Hosp. J
<i>mcr-1</i>	0	0	0	0
<i>mcr-3</i>	4.19 x 10 ⁵	8.70 x 10 ⁵	1.80 x 10 ⁴	2.65 x 10 ⁵
<i>mcr-3.17</i>	0	0	0	0
<i>mcr-4</i>	0	0	0	0
<i>mcr-5</i>	2.8 x 10 ²	4.30 x 10 ⁴	3.20 x 10 ³	2.36 x 10 ⁴
<i>mcr-7</i>	8.40 x 10 ⁵	0	7.59 x 10 ⁵	0
<i>mcr-8</i>	0	0	0	0
<i>mcr-9</i>	0	0	9.49 x 10 ⁶	0
<i>mcr-10</i>	0	1.74 x 10 ⁴	6.64 x 10 ⁵	7.97 x 10 ⁵

116

117 *References*

118

- 119 1. Murat Eren, A. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat. Microbiol.*
120 **6**, 3–6 (2021).
- 121 2. Borowiak, M. *et al.* Identification of a novel transposon-associated phosphoethanolamine transferase
122 gene, *mcr-5*, conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica*
123 serovar Paratyphi B. *J. Antimicrob. Chemother.* **72**, 3317–3324 (2017).
- 124