REWARD PROCESSING AND ANHEDONIA

Objective measures of reward sensitivity and motivation in people with high vs low anhedonia

Chloe Slaney¹, Adam M. Perkins², Robert Davis³, Ian Penton-Voak^{4,6}, Marcus R Munafò^{4,5,6}, Conor J. Houghton⁷, Emma S.J. Robinson^{1*}.

¹ School of Physiology, Pharmacology & Neuroscience, Faculty of Life Sciences, University of Bristol, Bristol, BS8 1TD, UK.

² Department of Psychological Medicine, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, SE5 8AF, UK.

³ Department of Psychology, Goldsmiths, University of London, London, SE14 6NW, UK.

⁴ School of Psychological Science, University of Bristol, 12a Priory Road, Bristol, BS8 1TU, UK.

⁵ MRC Integrative Epidemiology Unit at the University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK.

⁶ National Institute for Health Research Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol, Bristol, UK.

⁷ Department of Computer Science, Faculty of Engineering, University of Bristol, BS8 1UB, UK.

*** Author for correspondence:** Emma S.J. Robinson (E-mail: emma.s.j.robinson@bristol.ac.uk).

Address: School of Physiology, Pharmacology & Neuroscience, Biomedical Sciences Building, University Walk, University of Bristol, Bristol, UK,

Financial support. This research was funded in whole, or in part, by the Wellcome Trust [108899/B/15/Z]. For the purpose of Open Access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

Key words: anhedonia; reward; motivation; sensitivity; EEfRT; decision-making.

Word count: 4,526

Abstract

Background. Anhedonia – a diminished interest or pleasure in activities – is a core self-reported symptom of depression which is poorly understood and often resistant to conventional antidepressants. This symptom may occur due to dysfunction in one or more sub-components of reward processing: motivation, consummatory experience, and/or learning. However, the precise impairments remain elusive. Dissociating these components (ideally, using cross-species measures) and relating them to the subjective experience of anhedonia is critical as it may benefit fundamental biology research and novel drug development.

Methods. Using a battery of behavioural tasks based on rodent assays, we examined reward motivation (Joystick-Operated Runway Task, JORT; and Effort-Expenditure for Rewards Task, EEfRT) and reward sensitivity (Sweet Taste Test) in a non-clinical population who scored high ($N = 32$) or low ($N = 34$) on an anhedonia questionnaire (Snaith-Hamilton Pleasure Scale).

Results. Compared to the low anhedonia group, the high anhedonia group displayed marginal impairments in effort-based decision-making (EEfRT) and reduced reward sensitivity (Sweet Taste Test). However, we found no evidence of a difference between groups in physical effort exerted for reward (JORT). Interestingly, whilst the EEfRT and Sweet Taste Test correlated with anhedonia measures, they did not correlate with each other, lending support for the possibility of sub-groups within anhedonia.

Conclusions. Our findings suggest that reward motivation and reward sensitivity are dissociable when tested in the same group of participants, and that anhedonia is a heterogenous symptom associated with impairments in reward sensitivity and effort-based decision-making.

Objective measures of reward sensitivity and motivation in people with high vs low anhedonia

Anhedonia, a markedly diminished interest or pleasure in activities (DSM-5; American Psychiatric Association, 2013), is a core self-reported symptom of Major Depressive Disorder (MDD) that responds poorly to conventional treatments (Uher et al., 2012). Despite its importance, the behavioural and neurobiological basis of anhedonia remains poorly understood (Cooper, Arulpragasam, & Treadway, 2018). This could, in part, be due the over-reliance on questionnaires to measure this symptom. Objective measures, particularly those that can be applied in both clinical and preclinical animal research, may help to improve our mechanistic understanding of anhedonia and aid the development of targeted treatments (Der-Avakian & Markou, 2012; Pizzagalli, Jahn, & O'Shea, 2005; Thomsen, 2015; Treadway & Zald, 2011).

One way to objectively measure anhedonia is behaviourally. Anhedonia may occur due to dysfunction in one or more components of reward processing: motivation to obtain rewards ("wanting"), consumption of rewards ("liking"), and learning what predicts rewards ("learning") (Berridge & Robinson, 2003). Whilst the taxonomy of reward processing is debated, preclinical studies suggest that these sub-domains involve partially dissociable neural systems (Berridge & Robinson, 1998; Berridge et al., 2009). Despite advancements in preclinical research (Der-Avakian et al., 2013), few studies have tried to dissociate these different components in people and relate them to questionnaire measures of anhedonia (McCabe, 2018). Translating behavioural tasks from animals to humans and dissociating the precise components related to anhedonia is critical as it may provide an objective biomarker that could be used in the development and testing of new treatments (Der-Avakian, Barnes, Markou, & Pizzagalli, 2016). As most preclinical tasks focus on measuring reward motivation and consumption, we also focus on these two components here.

Reward motivation is the “incentive or desire to act or accomplish goals” (Der-Avakian et al., 2016). Recently, preclinical reward motivation tasks, such as effort-related choice tasks (Salamone, Correa, Yang, Rotolo, & Presby, 2018) and the Progressive Ratio Task (PRT; Hodos, 1961), have been adapted for human studies (Hershenberg et al., 2016; Treadway, Buckholtz, Schwartzman, Lambert, & Zald, 2009). For example, the Effort-Expenditure for Rewards Task (EEfRT; Treadway et al., 2009), requires participants to choose whether to exert more effort for a high reward or less effort for a low reward over many trials. Interestingly, people with MDD and people who have higher levels of anhedonia in a non-clinical population choose the high effort/high reward option less often than healthy controls suggesting impairments in effort-based decision-making (Treadway, Bossaller, Shelton, & Zald, 2012; Treadway et al., 2009). In the PRT, incrementally more effortful responses are required to obtain the same amount of reward, and motivation is operationalised as the point at which the individual gives up (i.e., their “breakpoint”) (Strauss et al., 2016). In this task, people with MDD give up earlier than healthy controls (Hershenberg et al., 2016), suggesting a reduced willingness to work for reward. Whilst there is accumulating evidence showing reduced reward motivation in MDD, few studies focus on physical effort-expenditure for reward (Halahakoon et al., 2020) (although see Cléry-Melin et al., 2011). To address this gap, we developed a novel reward motivation task (Joystick-Operated Runway Task; JORT; Perkins et al., 2009), designed to measure a person’s physical effort exerted for reward.

Reward sensitivity is the consummatory experience of reward. The most commonly used reward sensitivity task in rodents is the sucrose preference test (SPT; Willner et al., 1987), which measures an animal’s ability to detect, and show a preference for, a weak sucrose solution over water. Researchers have attempted to translate this task into humans using a Sweet Taste Test (Amsterdam, Settle, Doty, Abelman, & Winokur, 1987; Berlin, Givry-Steiner, Lecrubier, & Puech, 1998; Dichter, Smoski, Kampov-Polevoy, Gallop, & Garbutt, 2010). Although different variations have been used, they typically require participants to report the

intensity and pleasantness of different sucrose concentrations. Previous studies have reported similar pleasantness ratings of sucrose in people with MDD compared to healthy controls (Amsterdam et al., 1987; Berlin et al., 1998; Dichter et al., 2010). However, there is discrepancy across studies when measuring sucrose intensity. Some studies have reported poorer detection of sucrose in people with MDD compared to healthy controls (Amsterdam et al., 1987; Berlin et al., 1998), whilst others have failed to find any differences (Dichter et al., 2010). One potential explanation is that point scales, often used to measure ability to detect sucrose, may not be sensitive enough to reliably detect impairments (McCabe, 2018).

Despite progress in the development of translational measures (Thomsen, 2015), there are gaps in the literature. First, most studies examine only one sub-component of reward (e.g., motivation) using one assay. However, a battery of tasks designed to probe different reward components (e.g., motivation, sensitivity) can demonstrate whether different components can be dissociated and whether anhedonia is related to a general reward deficit or specific to certain sub-components (Husain & Roiser, 2018; Nielson et al., 2020). Second, most studies compare people with MDD to healthy controls, with few directly measuring anhedonia using anhedonia questionnaires (McCabe, 2018). This is surprising, as only ~ 37% people with MDD exhibit significant levels of anhedonia (Pelizza & Ferrari, 2009). In line with the Research Domain Criteria (RDoC), a symptom-based approach which examines behavioural tasks in relation to symptom-based questionnaires may provide more rigorous findings (Cuthbert, 2014; Insel et al., 2010).

Here, we compared individuals with high versus low levels of anhedonia in a non-clinical population (Snaith et al., 1995) on a battery of behavioural tasks designed to measure reward motivation (EEfRT and JORT) and reward sensitivity (Sweet Taste Test). A novel measure of reward motivation - the JORT - was designed to assess reward motivation in the absence of explicit decision-making. These tasks were chosen based on their potential to dissociate reward components and their potential for cross-species research. Based on previous research, we

predicted that compared to people with low levels of anhedonia, people with high levels of anhedonia will show reduced reward motivation and reward sensitivity.

Method

This study was pre-registered on the Open Science Framework (<https://osf.io/p4zt6>) and was approved by the Faculty of Biomedical Sciences Research Ethics Committee at the University of Bristol (Ref: 74082).

Screening

Individuals who scored high (score ≥ 25) or low (score < 18) on the Snaith-Hamilton Pleasure Scale (SHAPS; see Figure 1) in a non-clinical population were invited to take part in this study. These individuals scored in the top and bottom 25th percentile on the SHAPS, which was administered online to 380 volunteers. In the pre-registration, the original cut-off was a low score of < 17 , however, due to initial difficulties with recruitment, this was increased to < 18 during the study.

Participants

Whilst 101 participants (44 low anhedonia, 57 high anhedonia) met the online SHAPS criteria and took part in the study, only 66 (34 low, 32 high) still met their SHAPS criteria when re-tested at the study visit and were included in the primary analysis. For sample size calculations, see supplementary materials.

Eligibility criteria were aged ≥ 18 years, fluent in English, not suffering from a mental health condition or neurological illness (self-report), no current physical injuries, no allergy or intolerance to sugar, no disorder of taste or smell, not diabetic, not previously participated in a study using the JORT.

Participants were recruited using advertisements within the University of Bristol and volunteer databases. Participants were informed that they would receive £20 for their time (~ 2 hours) and an additional performance-based pay (up to £5) based on one of the two reward motivation tasks (chosen randomly).

Behavioural Measures

Joystick-Operated Runway Task (JORT). The JORT was originally developed as a translational measure of fear and anxiety, here it has been adapted as a measure of reward motivation. This is a computerised task which uses a force-sensing joystick to measure a person's physical effort to obtain a reward (Perkins et al., 2009; PH-JS1, Psyal, London). Each trial begins with a green dot (representing the participant) positioned along an onscreen runway. A cue in the top left-hand corner of the screen indicates the number of points on offer (0, 10, 100 or 1,000 points; see Figure 2). Following this, a target (black dot) appears and immediately accelerates away from the participant. Participants are informed that to win the points on offer, they must push the joystick to chase and catch the onscreen target. The speed of the participants cursor is linked to the force applied on the joystick (Perkins et al., 2009). Trials vary in the number of points on offer (0, 10, 100 and 1,000 points; visible to the participant) and the minimum effort required to win (50, 80, 100 or 120% of their maximum calibrated force; not visible). If the participant catches the target, they receive feedback that they have won the number of points on offer. Trials end either upon catching the target or after 7 seconds. The inter-trial interval varies pseudo-randomly between 3-7 seconds. After each trial, the next trial automatically begins. Force applied to the joystick is recorded every 15 ms.

Participants complete two blocks of 48 trials (96-total), separated by a short break (~ 5 minutes). Each block contains an equal number of reward-effort combinations ($N = 3$), except for one combination where two trials were presented due to experimenter error (0 points-80%

effort in second block). All participants complete the same pre-randomised order of trials. For a proof-of-concept study of the JORT, see supplementary materials.

Prior to the task, participants are given standardised verbal instructions before completing a calibration phase and four practice trials. In the calibration phase, participants push the joystick as “hard as they can” five times whenever “GO” is presented on the screen. Peak force reached during the calibration or practice trials (whichever is highest) is recorded as a person’s max calibrated force. To standardise the amount of effort required across participants on the main task, effort levels reflect the percentage of a participants max calibrated force.

Effort-Expenditure for Rewards Task. The EEfRT (Treadway et al., 2009) is a computerised effort-based decision-making task. On each trial, participants choose between an “easy” and “hard” option. The easy option requires 30 button presses within 7 seconds (using the index finger on their dominant hand) for 50p. The hard option requires 98 button presses (using the little finger on their non-dominant hand) within 21 seconds for 58p - £2. Participants have 5 seconds to choose, or they are randomly assigned to either option for that trial. Each trial also contains a visible probability cue (12%, 50%, 88%) which indicates the probability of winning money if they successfully complete the trial (i.e., win trials). The probability cue applies to both the easy and hard option, with an equal proportion across the experiment. Participants are informed that two “win trials” may be paid to them at the end of the experiment (performance-based pay). Participants complete the same pre-randomised order of trials for 20 minutes. For further details, see supplementary materials.

Sweet Taste Test. We employed a modified version of the Sweet Taste Test (Berlin et al., 1998), which measures detection threshold of sweet taste. On each trial, participants were given one sucrose concentration. They were asked to sip one mouthful of the solution, swish it around their mouth for 5 seconds and then spit it out (Berlin et al., 1998). Participants were asked if they could detect the presence of sugar in the solution (Berlin et al., 1998). In between

trials, participants sipped one mouthful of water, swished it around their mouth for 5 seconds and then spat this out. Prior to the task, participants completed two practice trials containing water to ensure that they were familiar with the procedure and to clean their palette. Seven concentrations of sucrose were used (0, 0.5, 1, 1.5, 2, 2.5, 5% w/v), each delivered in 15 mL. A staircase method with five reversals was employed. The first concentration administered was always the strongest concentration (5% sucrose) to ensure participants were aware of what to detect. A staircase method was used as it provides a fast determination of threshold. The detection threshold was calculated as the mean of the five boundaries (i.e., points at which they changed response from either detect to fail to detect, or vice versa).

Self-Report Measures

The primary measure of anhedonia was the SHAPS (Snaith et al., 1995). This is a 14-item questionnaire which asks participants to report their ability to experience pleasure in the last few days. Additional measures of anhedonia, apathy and depression were collected: Chapman Physical Anhedonia Scale (CPAS; Chapman et al., 1976), Temporal Experience of Pleasure Scale (TEPS; Gard et al., 2006), Beck Depression Inventory-II (BDI-II; Beck et al., 1996) and Apathy Evaluation Scale (AES; Marin et al., 1991). Higher scores indicate higher symptoms on all questionnaires except for the TEPS. For further details, see supplementary materials.

Procedure

Participants completed the behavioural tasks in the following order: EEfRT, JORT and the Sweet Taste Test, which took approximately one hour. Participants then completed the questionnaires: demographic form, SHAPS, CPAS, TEPS, BDI-II, AES.

Data Analysis

Primary Analysis. Data were analysed using SPSS 24 (IBM). Primary analyses were run on those who met the SHAPS cut-offs at the study visit ($N = 66$). Based on previous research, *a priori* potential co-variables were examined and their main effects retained in the model (Engqvist, 2005).

For the JORT, 2 x 4 (group x reward) mixed ANOVAs examined differences between groups in relative average force, maximum force, and reaction time. The primary outcome was relative average force. For the EEfRT, 2 x 3 (group x probability) mixed ANCOVAs examined differences between groups in mean proportion of hard-task choices and reaction time. Sex was included as a covariate (Treadway et al., 2009). The primary outcome was proportion of hard-task choices. For the Sweet Taste Test, an ANCOVA examined differences between groups in detection threshold. Current smoking status (3 levels: daily, less than daily, not at all) was included as a covariate (Sato, Endo, & Tomita, 2002). All secondary outcomes and data checks are reported in the supplementary materials.

Exploratory Analysis. Spearman correlations examined associations between tasks and self-report measures, and correlations between different tasks. As these were exploratory, no corrections for multiple comparisons were used.

Results

Data are available at the University of Bristol data repository, [data.bris](https://data.bris.ac.uk/), at <https://doi.org/10.5523/bris.1wlrhv4jzqs7q2egf8i7ruzta1>.

Participant Characteristics

For participant demographics, see Table 1. For distribution of scores on self-report measures and correlations between measures, see Supplementary Materials.

Joystick-Operated Runway Task

Based on *a priori* criteria, participants who succeeded in over 75% trials ($N = 4$) were excluded from the analysis. This is because these participants must have achieved at least one trial designed to be impossible (120% effort trials) and were therefore considered to have not successfully achieved their maximum force during the calibration. In total, 62 participants (33 low anhedonia, 29 high anhedonia) were included in the analysis.

Average force. There was strong evidence of a main effect of reward, participants exerted more force for higher reward magnitudes, $F_{(1.45, 87.14)} = 39.41$, $p < .001$, $\eta_p^2 = .40$. Bonferroni-corrected pairwise comparisons revealed evidence of a difference between all reward magnitudes ($ps \leq .003$). There was no evidence of a main effect of group ($F_{(1, 60)} = .004$, $p = .95$, $\eta_p^2 < .001$; see Figure 3) or reward x group interaction ($F_{(1.45, 87.14)} = .56$, $p = .52$, $\eta_p^2 = .009$).

Effort-Expenditure for Rewards Task

Consistent with Treadway (2009), an upper bound of the first 50 trials was used in the analysis. Trials where participants did not make a choice were not included in the analysis ($M = 49.20$, $SD = 1.90$, range = 39 to 50).

Proportion of hard-task choices. There was strong evidence of a main effect of probability, participants chose the hard-task choice more often when the probability of winning was higher, $F_{(2, 126)} = 209.12$, $p < .001$, $\eta_p^2 = .77$. Bonferroni-corrected comparisons revealed evidence of a difference between all probabilities ($ps < .001$). There was evidence of a main effect of group, $F_{(1, 63)} = 4.45$, $p = .039$, $\eta_p^2 = .07$: the high anhedonia group ($M = 28.88$, $SE = 2.42$) chose the hard-task choice less often than the low anhedonia group ($M = 36.41$, $SE = 2.58$; see Figure 3). There was no evidence of other main effects or interactions ($ps \geq .57$). However, removal of one outlier weakened the evidence of the main effect of group, $F_{(1, 62)} =$

2.64, $p = .109$, $n_p^2 = .041$). *Post-hoc* Mann-Whitney U tests indicated evidence of a difference between groups on low probability trials ($p = .045$), but this did not survive Bonferroni-corrections for multiple comparisons. There was no evidence of a difference between groups on medium or high probability trials ($ps \geq .28$).

Button Pressing Rate. There was no evidence of a difference between groups in mean button pressing rate (button-pressing speed) for the easy or hard-task choices ($ps \geq .11$).

Sweet Taste Test

Based on *a priori* criteria, participants who reported detection of sucrose in 0% sucrose concentrations were excluded. This excluded 5 participants; 61 participants (31 low anhedonia, 30 high anhedonia) were included in the analysis.

The analysis was conducted on log transformed data, which met ANOVA assumptions. There was evidence of a main effect of group, the high anhedonia group required a higher concentration of sucrose to detect its presence, $F_{(1,57)} = 4.56$, $p = .037$, $n_p^2 = .07$, see Figure 3. There was no evidence of a main effect of smoking status, $F_{(2,57)} = .41$, $p = .66$, $n_p^2 = .01$. Removing one outlier strengthened the evidence of a main effect of Group ($p = .015$, $n_p^2 = .10$). This finding did not substantially differ when using non-transformed data ($p = .020$) or using a non-parametric Mann-Whitney U test with the covariate excluded ($p = .016$).

Exploratory Analyses

Correlations between tasks and self-report measures. In line with the original EEfRT study (Treadway et al., 2009), we found evidence of a negative correlation between proportion of hard-task choices and trait anhedonia (CPAS; $r = -.29$, $p = .003$). In the Sweet Taste Test, we found evidence of positive a correlation between detection threshold and symptoms of depression (BDI-II), anhedonia (SHAPS and CPAS) and apathy (AES; $ps \leq .033$). In the JORT, to facilitate correlational analyses, individual slopes (force exerted across the four reward

magnitudes) were extracted for each participant. There was no clear evidence of correlations between self-report measures and the JORT, see Table 2.

Correlations between tasks. There was evidence of a positive correlation between proportion of hard-task choices (EEfRT) and average force slopes (JORT), $r_{(101)} = .21$, $p = .039$, participants who were more willing to choose the hard-task option exerted more force for higher compared to lower reward magnitudes on the JORT. There was no evidence of a correlation between the EEfRT and Sweet Taste Test ($r_{(92)} = -.02$, $p = .84$), nor the JORT and Sweet Taste Test ($r_{(92)} = -.07$, $p = .49$).

Discussion

We found evidence of a difference in participants performance across paradigms, supporting the hypothesis that reward motivation and reward sensitivity are dissociable components involving different underlying mechanisms. In relation to anhedonia (SHAPS), we found no evidence of a difference between the high and low anhedonia group in physical effort exerted for reward (JORT). However, compared to the low anhedonia group, the high anhedonia group displayed altered effort-based decision-making (EEfRT) and reduced reward sensitivity (Sweet Taste Test). Although the JORT did not find any effect between groups, there was a correlation between the EEfRT and JORT, suggesting that they may measure similar aspects of reward motivation. In this study, the EEfRT was more sensitive to changes in anhedonia scores, possibly because it involves both physical effort and cognitive decision-making.

Reward motivation and anhedonia

In the JORT, there was no evidence of a difference between the high and low anhedonia group, both groups exerted more physical force for higher reward trials. This suggests that

people with higher levels of anhedonia, at least in a non-clinical population, do not display deficits in their exertion of physical effort for reward. Using a human PRT, Hershenberg and colleagues (2016) reported that people with MDD had a lower breakpoint compared to healthy controls (Hershenberg et al., 2016). However, this task required cognitive effort and as the rodent PRT has failed to find a similar effect in some models of depression (chronic mild stress and maternal separation) (Barr & Phillips, 1998; Shalev & Kafkafi, 2002), its translational validity has been questioned. Nevertheless, Cléry-Melin et al. (2011) found that compared to healthy volunteers, people with depression exerted less physical force to maximize monetary rewards on a handgrip apparatus (Cléry-Melin et al., 2011). Whilst these studies may seem to conflict the findings presented here, it is important to note that our study included a non-clinical population, measuring anhedonia as a varying trait, as opposed to clinical depression.

In the EEfRT, the high anhedonia group displayed a reduced proportion of hard-task choices, compared to the low anhedonia group. However, evidence for this difference was weak (especially after removing one outlier) and we did not find evidence of a difference between groups on 50% probability trials, as predicted in our pre-registration. Nevertheless, our exploratory analysis did replicate the original Treadway (2009) finding that higher trait anhedonia (measured using the CPAS) was associated with reduced hard-task choices on the EEfRT (Treadway et al., 2009). Interestingly, both our study and the Treadway study found an association between the EEfRT and CPAS, but failed to find a robust association between the EEfRT and SHAPS (Treadway et al., 2009; Yang et al., 2014). One possible reason for this is that the SHAPS focuses more on consummation of reward (“I would be able to enjoy my favourite meal”) whereas the CPAS also includes questions on interest/desire for future rewards (“I have had very little desire to try new kinds of foods”) (Leventhal, Chasson, Tapia, Miller, & Pettit, 2006; Rizvi, Pizzagalli, Sproule, & Kennedy, 2016), with the latter being more relevant to the EEfRT (although see Tran et al., 2020). Overall, this finding highlights the value of employing multiple anhedonia questionnaires.

There was a correlation between the JORT and EEfRT suggesting that they may measure a similar construct of motivation. However, only the EEfRT was related to anhedonia. The EEfRT examines explicit decision-making (Treadway et al., 2009) (similar to rodent choice-based tasks) where a participant must weigh up the value of two options, whereas the JORT (more similar to the PRT) measures amount of physical effort exerted for reward. Speculatively, anhedonia may not be related to reduced exertion of physical effort for reward when engaged in an activity, but rather in the choice to engage in an effortful activity for reward (decision-making; Halahakoon et al., 2020). In relation to the rodent literature, these findings suggest that effort-based-choice tasks may provide a more sensitive measure of motivational impairments relevant to anhedonia than physical effort-for-reward tasks (e.g., PRT). Nevertheless, it is important to consider alternative explanations including methodological differences between tasks (e.g., presence of probabilistic cues, reward sizes) and the cognitive demands of the EEfRT (Cooper et al., 2019; Whitton, Merchant, & Lewandowski, 2020).

Reward sensitivity and anhedonia

In the Sweet Taste Test, the high anhedonia group displayed reduced sensitivity to sucrose compared to the low anhedonia group. Most previous studies using this task have compared people with clinical disorders such as MDD to healthy controls, and found conflicting results (Amsterdam et al., 1987; Berlin et al., 1998; Dichter et al., 2010). In relation to anhedonia, Bedwell (2019) reported that sucrose sensitivity did not relate to anhedonia measures in the typical population (Bedwell, Spencer, Chirino, & O'donnell, 2019). One potential explanation for these conflicting results is how sucrose sensitivity is measured. Most studies have measured sweet taste intensity on a point scale (Bedwell et al., 2019; Dichter et al., 2010), as opposed to detection threshold (i.e., point at which sucrose can be detected) (Berlin et al., 1998). It has been suggested that point scales may not be sensitive enough to reliably detect subtle impairments (McCabe, 2018). This study suggests that anhedonia, at least

in a non-clinical population, is related to reduced reward sensitivity. Future studies should examine whether this effect is specific to pleasant stimuli (e.g., sucrose) or whether it can also be observed using negative stimuli (e.g., bitter tastes), indicating a general sensitivity impairment.

Collectively, using a battery of translational tasks designed to probe different reward components (motivation, sensitivity), we support and extend preclinical findings by revealing that these components may also be dissociable in people. This is demonstrated by a lack of correlations between tasks: individuals who were less willing to choose the hard-task choice on the EEfRT did not display reduced sensitivity on the Sweet Taste Test. We also found that anhedonia was related to reduced reward motivation and reward sensitivity. However, this does not seem to reflect a general reward deficit, indexed by the lack of correlation between tasks. Instead, our findings suggest that anhedonia may be a heterogeneous symptom (i.e., there may be sub-groups of individuals who have dysfunction in different components of reward) (Thomsen, 2015; Treadway & Zald, 2013). Whilst this hypothesis has been proposed before, to our knowledge it has never been directly tested. This is because most studies employ a single reward processing task alongside an anhedonia questionnaire. Further studies employing a battery of behavioural tasks (motivation, sensitivity, and learning) and self-report measures in a clinical population are needed to directly address this question.

Limitations

This study examined anhedonia in a non-clinical, predominantly student, population. There are benefits of examining non-clinical populations including the reduced likelihood of cognitive deficits, medication use and comorbidity. However, anhedonia levels in the clinical population are higher and may be qualitatively different. Therefore, future studies employing a battery of translational behavioural tasks in a clinical population are needed. Second, there was a higher proportion of males in the high anhedonia group (53%) compared to the low anhedonia

group (21%), which may have influenced the differences between groups. However, including sex as a covariate in all analyses did not change the overall findings. Interestingly, few studies have investigated sex differences in anhedonia, and thus it is unknown whether the higher proportion of males in the high anhedonia group reflects a real sex difference in anhedonia (Chan et al., 2012; Xinhua Yang, Wang, Liu, Liu, & Harrison, 2020) or whether this is a result of sampling. Third, the data did not always meet the ANOVA assumptions, which could reduce the confidence in our findings. Nevertheless, ANOVA is relatively robust to minor deviations from normality if extreme outliers are not the cause of skew (Field, 2005).

To summarize, this study demonstrates a dissociation between different reward components (effort-for-reward, effort-based decision-making, and reward sensitivity) in humans using a battery of behavioural tasks. Examining these different components in relation to anhedonia, we found that symptoms of anhedonia in a non-clinical population were related to changes in reward sensitivity and effort-based decision-making. Interestingly, as performance on the effort-based decision-making (EEfRT) and reward sensitivity (Sweet Taste Test) tasks did not correlate, this could suggest heterogeneity within anhedonia (Treadway & Zald, 2013). Specifically, some people with anhedonia may display reduced sensitivity to reward, whilst others may display altered decision-making to engage in effortful activities for reward. If this finding is also found in clinical populations, it may have important implications for assessment and treatment of anhedonia.

Acknowledgements. The authors wish to thank Michael T. Treadway for providing us with the EEfRT.

Conflicts of Interest. R.D. is the owner of Psyal (<http://www.psyal.co.uk/>), a company that designs and sells products to the psychology community. This study includes one product purchased from Psyal: the JORT.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders (5th ed.)*. American Psychiatric Association.
- Amsterdam, J. D., Settle, R. G., Doty, R. L., Abelman, E., & Winokur, A. (1987). Taste and smell perception in depression. *Biological Psychiatry*, 22, 1481–1485.
- Barr, A. M., & Phillips, A. G. (1998). Chronic mild stress has no effect on responding by rats for sucrose under a progressive ratio schedule. *Physiology & Behavior*, 64, 591–597.
- Beck, A., Steer, R., & Brown, G. (1996). Beck Depression Inventory-II. *San Antonio*, 12–15.
- Bedwell, J. S., Spencer, C. C., Chirino, C. A., & O'donnell, J. P. (2019). The Sweet Taste Test: Relationships with Anhedonia Subtypes, Personality Traits, and Menstrual Cycle Phases. *Journal of Psychopathology and Behavioral Assessment*, 41, 235–248.
- Berlin, I., Givry-Steiner, L., Lecrubier, Y., & Puech, A. J. (1998). Measures of anhedonia and hedonic responses to sucrose in depressive and schizophrenic patients in comparison with healthy subjects. *European Psychiatry*, 13, 303–309.
- Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience? *Brain Research Reviews*, 28, 309–369.
- Berridge, K. C., & Robinson, T. E. (2003). Parsing reward. *Trends in Neurosciences*, 26, 507–513.
- Berridge, K. C., Robinson, T. E., & Aldridge, J. W. (2009). Dissecting components of reward: “liking”, “wanting”, and learning. *Current Opinion in Pharmacology*, 9, 65–73.
- Chan, R. C. K., Wang, Y., Yan, C., Zhao, Q., McGrath, J., Hsi, X., & Stone, W. S. (2012). A Study of Trait Anhedonia in Non-Clinical Chinese Samples: Evidence from the Chapman Scales for Physical and Social Anhedonia. *PLoS One*, 7.
- Chapman, L. J., Chapman, J. P., & Raulin, M. L. (1976). Scales for physical and social anhedonia. *Journal of Abnormal Psychology*, 85, 374–382.
- Cléry-Melin, M., Schmidt, L., Lafargue, G., Baup, N., Fossati, P., & Pessiglione, M. (2011). Why

Don't You Try Harder? An Investigation of Effort Production in Major Depression. *PLoS ONE*, 6.

Cooper, J. A., Arulpragasam, A. R., & Treadway, M. T. (2018). Anhedonia in depression: biological mechanisms and computational models. *Current Opinion in Behavioral Sciences*, 22, 128–135.

Cooper, J. A., Barch, D. M., Reddy, L. F., Horan, W. P., Green, M. F., & Treadway, M. T. (2019). Effortful Goal-Directed Behavior in Schizophrenia: Computational Subtypes and Associations With Cognition. *Journal of Abnormal Psychology*.

Cuthbert, B. N. (2014). The RDoC framework: facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry*, 13, 28–35.

Daghighi, A., Daghighi, V., Niazi, M., & Nadorff, M. R. (2019). The Association Between Anhedonia, Suicide Ideation, and Suicide Attempts: A Replication in a Persian Student Sample. *Suicide and Life-Threatening Behavior*, 49, 678–683.

Der-Avakian, A., Barnes, S. A., Markou, A., & Pizzagalli, D. A. (2016). Translational assessment of reward and motivational deficits in psychiatric disorders. *Current Topics in Behavioral Neurosciences*, 28, 231–262.

Der-Avakian, A., & Markou, A. (2012). The neurobiology of anhedonia and other reward-related deficits. *Trends in Neurosciences*, 35, 68–77.

Der-Avakian, A., Souza, M. D., Pizzagalli, D., & Markou, A. (2013). Assessment of reward responsiveness in the response bias probabilistic reward task in rats: implications for cross-species translational research. *Translational Psychiatry*, 3.

Dichter, G. S., Smoski, M. J., Kampov-Polevoy, A. B., Gallop, R., & Garbutt, J. C. (2010). Unipolar depression does not moderate responses to the sweet taste test. *Depression and Anxiety*, 27, 859–863.

Engqvist, L. (2005). The mistreatment of covariate interaction terms in linear model analyses of

- behavioural and evolutionary ecology studies. *Animal Behaviour*, 70, 967–971.
- Field, A. P. (2005). *Discovering statistics using IBM SPSS statistics : and sex and drugs and rock “n” roll*. SAGE.
- Gard, D. E., Gard, M. G., Kring, A. M., & John, O. P. (2006). Anticipatory and consummatory components of the experience of pleasure: A scale development study. *Journal of Research in Personality*, 40, 1086–1102.
- Halahakoon, D. C., Kieslich, K., O’Driscoll, C., Nair, A., Lewis, G., & Roiser, J. P. (2020). Reward-Processing Behavior in Depressed Participants Relative to Healthy Volunteers: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. American Medical Association.
- Hershenberg, R., Satterthwaite, T. D., Daldal, A., Katchmar, N., Moore, T. M., Kable, J. W., & Wolf, D. H. (2016). Diminished effort on a progressive ratio task in both unipolar and bipolar depression. *Journal of Affective Disorders*, 196, 97–100.
- Hodos, W. (1961). Progressive ratio as a measure of reward strength. *Science*, 134, 943–944.
- Husain, M., & Roiser, J. P. (2018). Neuroscience of apathy and anhedonia: A transdiagnostic approach. *Nature Reviews Neuroscience*, 19, 470–484.
- Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., ... Wang, P. (2010). Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. *American Journal of Psychiatry*, 167, 748–751.
- Leventhal, A. M., Chasson, G. S., Tapia, E., Miller, E. K., & Pettit, J. W. (2006). Measuring Hedonic Capacity in Depression: A Psychometric Analysis of Three Anhedonia Scales. *Journal of Clinical Psychology*, 62, 1545–1558.
- Marin, R. S., Biedrzycki, R. C., & Firinciogullari, S. (1991). Reliability and validity of the Apathy Evaluation Scale. *Psychiatry Research*, 38, 143–162.
- McCabe, C. (2018). Linking anhedonia symptoms with behavioural and neural reward responses in adolescent depression. *Current Opinion in Behavioral Sciences*, 22, 143–151.
- Nielson, D. M., Keren, H., O’Callaghan, G., Jackson, S. M., Douka, I., Vidal-Ribas, P., ...

- Stringaris, A. (2020). Great Expectations: A Critical Review of and Suggestions for the Study of Reward Processing as a Cause and Predictor of Depression. *Biological Psychiatry*.
- Pelizza, L., & Ferrari, A. (2009). Anhedonia in schizophrenia and major depression: state or trait? *Annals of General Psychiatry*, 8, 22.
- Perkins, A. M., Ettinger, U., Davis, R., Foster, R., Williams, S. C. R., & Corr, P. J. (2009). Effects of Lorazepam and Citalopram on Human Defensive Reactions: Ethopharmacological Differentiation of Fear and Anxiety. *Journal of Neuroscience*, 29, 12617–12624.
- Pizzagalli, D. A., Jahn, A. L., & O'Shea, J. P. (2005). Toward an objective characterization of an anhedonic phenotype: A signal-detection approach. *Biological Psychiatry*, 57, 319–327.
- Rizvi, S. J., Pizzagalli, D. A., Sproule, B. A., & Kennedy, S. H. (2016). Assessing anhedonia in depression: Potentials and pitfalls. *Neuroscience & Biobehavioral Reviews*, 65, 21–35.
- Salamone, J. D., Correa, M., Yang, J.-H., Rotolo, R., & Presby, R. (2018). Dopamine, Effort-Based Choice, and Behavioral Economics: Basic and Translational Research. *Frontiers in Behavioral Neuroscience*, 12, 52.
- Sato, K., Endo, S., & Tomita, H. (2002). Sensitivity of Three Loci on the Tongue and Soft Palate to Four Basic Tastes in Smokers and Non-smokers. *Acta Oto-Laryngologica*, 122, 74–82.
- Shalev, U., & Kafkafi, N. (2002). Repeated maternal separation does not alter sucrose-reinforced and open-field behaviors. *Pharmacology, Biochemistry, and Behavior*, 73, 115–122.
- Snaith, R. P., Hamilton, M., Morley, S., Humayan, A., Hargreaves, D., & Trigwell, P. (1995). A scale for the assessment of hedonic tone the Snaith-Hamilton Pleasure Scale. *The British Journal of Psychiatry*, 167, 99–103.
- Strauss, G. P., Whearty, K. M., Morra, L. F., Sullivan, S. K., Ossenfort, K. L., & Frost, K. H. (2016). Avolition in schizophrenia is associated with reduced willingness to expend effort for reward on a Progressive Ratio task. *Schizophrenia Research*, 170, 198–204.

- Thomsen, K. R. (2015). Measuring anhedonia: impaired ability to pursue, experience, and learn about reward. *Frontiers in Psychology*.
- Tran, T., Hagen, A. E. F., Hollenstein, T., & Bowie, C. R. (2020). Physical- and Cognitive-Effort-Based Decision-Making in Depression: Relationships to Symptoms and Functioning. *Clinical Psychological Science*.
- Treadway, M. T., Bossaller, N. A., Shelton, R. C., & Zald, D. H. (2012). Effort-Based Decision-Making in Major Depressive Disorder: A Translational Model of Motivational Anhedonia. *Journal of Abnormal Psychology*, 121, 553–558.
- Treadway, M. T., Buckholtz, J. W., Schwartzman, A. N., Lambert, W. E., & Zald, D. H. (2009). Worth the “EEfRT”? The Effort Expenditure for Rewards Task as an Objective Measure of Motivation and Anhedonia. *PLoS ONE*, 4.
- Treadway, M. T., & Zald, D. H. (2011). Reconsidering anhedonia in depression: Lessons from translational neuroscience. *Neuroscience and Biobehavioral Reviews*, 35, 537–555.
- Treadway, M. T., & Zald, D. H. (2013). Parsing Anhedonia: Translational Models of Reward-Processing Deficits in Psychopathology. *Current Directions in Psychological Science*, 22, 244–249.
- Uher, R., Perlis, R. H., Henigsberg, N., Zobel, A., Rietschel, M., Mors, O., ... McGuffin, P. (2012). Depression symptom dimensions as predictors of antidepressant treatment outcome: replicable evidence for interest-activity symptoms. *Psychological Medicine*, 42, 967–980.
- Whitton, A. E., Merchant, J. T., & Lewandowski, K. E. (2020). Dissociable mechanisms underpinning effort-cost decision-making across the psychosis spectrum. *Schizophrenia Research*, 224, 133–140.
- Willner, P., Towell, A., Sampson, D., Sophokleous, S., & Muscat, R. (1987). Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. *Psychopharmacology*, 93, 358–364.

- Winer, E. S., Drapeau, C. W., Veilleux, J. C., & Nadorff, M. R. (2016). The Association between Anhedonia, Suicidal Ideation, and Suicide Attempts in a Large Student Sample. *Archives of Suicide Research*, 20, 265–272.
- Yang, X. hua, Huang, J., Zhu, C. ying, Wang, Y. fei, Cheung, E. F. C., Chan, R. C. K., & Xie, G. rong. (2014). Motivational deficits in effort-based decision making in individuals with subsyndromal depression, first-episode and remitted depression patients. *Psychiatry Research*, 220, 874–882.
- Yang, X., Wang, D., Liu, S., Liu, G., & Harrison, P. (2020). Trajectories of state anhedonia and recent changes in anhedonia in college students: Associations with other psychiatric syndromes. *Journal of Affective Disorders*, 262, 337–343.

Table 1

Demographic data and questionnaire scores for participants in the high and low anhedonia group.

	Low Group		High Group		Group Difference		
	Mean	SD / %	Mean	SD / %	<i>F</i>	<i>p</i> -value	Post-hoc
Age	24.0	5.1	22.8	8.6	.41	.52	n/a
Males (<i>N</i>)	7	21%	17	53%	$X^2 = 7.54$	.006	H > L
Smoke Status	2.7	0.7	2.7	0.6	$X^2 = .23$	.89	n/a
Income	2.3	0.6	2.2	0.6	$X^2 = .64$	.73	n/a
Money Concern	12	36%	16	50%	$X^2 = 1.23$	.27	n/a
SHAPS	15.4	1.0	27.6	2.2	838.34	<.001	H > L
CPAS	9.6	5.7	18.0	7.1	27.67	<.001	H > L
TEPS – A	50.5	4.6	40.3	5.2	71.10	<.001	L > H
TEPS – C	40.6	5.3	30.3	6.2	52.80	<.001	L > H
BDI-II	6.4	5.5	11.4	7.6	9.41	.003	H > L
Apathy Scale	24.6	4.4	32.4	6.2	33.74	<.001	H > L

Note. Current smoke status (1=daily, 2=less than daily, 3=not at all); income (1=high, 2=medium, 3=low); money concern, participants who reported “yes” to finance concerns; SHAPS, Snaith-Hamilton Pleasure Scale; TEPS, Temporal Experience of Pleasure Scale (A, anticipatory; C, consummatory); BDI-II, Beck Depression Inventory-II; n/a, not applicable. Welch’s ANOVA and Chi Square to compare groups.

Table 2

Spearman's correlations between behavioural tasks and self-report measures.

Variable		JORT	EEfRT	Sweet Taste Test
SHAPS	<i>R</i>	-.09	-.14	.27**
	<i>p-value</i>	.38	.16	.008
CPAS	<i>R</i>	-.03	-.29**	.22*
	<i>p-value</i>	.79	.003	.033
TEPS-A	<i>R</i>	.001	.16	-.15
	<i>p-value</i>	.99	.12	.17
TEPS-C	<i>R</i>	-.07	.09	-.19
	<i>p-value</i>	.50	.39	.076
BDI-II	<i>R</i>	-.04	-.06	.27*
	<i>p-value</i>	.70	.55	.010
Apathy Scale	<i>R</i>	.01	-.04	.23*
	<i>p-value</i>	.91	.66	.026

Note. Sweet Taste Test - mean detection threshold; EEfRT, Effort-Expenditure for Rewards

Task – overall proportion of hard-task choices; JORT, Joystick-Operated Runway Task – slope

of average force across four reward magnitudes; SHAPS, Snaith-Hamilton Pleasure Scale;

CPAS, Chapman Physical Anhedonia Scale; TEPS, Temporal Experience of Pleasure Scale (A,

Anticipatory; C, Consummatory); BDI-II, Beck Depression Inventory-II; Apathy Scale, Apathy

Evaluation Scale. *N* = 91–101.

Figure 1. CONSORT diagram of study selection process.

Figure 2. Example trial on the JORT. The green dot (representing the participant) is initially displayed along with a cue indicating the number of points on offer. This is followed by the presentation of the target (a black dot). The participant then receives feedback if they catch the target (e.g., “You win 10 points”) or no feedback if they fail to catch the target.

REWARD PROCESSING AND ANHEDONIA

Figure 3. Differences between the high and low anhedonia group in the (A) Joystick-Operated Runway Task: relative average force exerted for each reward magnitude ($N = 33$ low, 29 high) (B) Effort-Expenditure for Rewards Task: mean proportion of hard-task choices across different levels of probability ($N = 34$ low, 32 high) and (C) Sweet Taste Test ($N = 31$ low, 30 high). Points represent participants. Error bars represent *SEM*. Data presented with statistical outliers included.