The Relationship Between Food Security and Dietary Patterns Status with COVID-19 in Northeastern Iran: Protocol for a Case-Control Study

Running Title: Food Security, Dietary Status, and COVID-19

Sepideh Badri-Fariman¹, Bahram Pourghassem-Gargari^{2*}, Mahtab Badri-Fariman³, Mohammad Pourfridoni⁴, Milad Daneshi-Maskooni^{5*}

¹Department of Community Nutrition, School of Nutrition and Food Sciences, Tabriz University of Medical Sciences, Tabriz, Iran.

²**First Corresponding Author**, PhD, Professor, Department of Biochemistry and Diet Therapy, Nutrition Research Center, Faculty of Nutrition, Tabriz University of Medical Sciences, Tabriz, Iran. Email: bahrampg@yahoo.com, Tell: +989143165247

³MSPH, Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran.

⁴ Student Research Committee, Jiroft University of Medical Sciences, Jiroft, Iran.

⁵**Second Corresponding Author**, PhD, Assistant Professor, Department of Nutrition, School of Medicine, Jiroft University of Medical Sciences, Jiroft, Kerman, Iran. Email: miladdaneshi@gmail.com, Tell: +989190026295

Abstract

Background: Food insecurity is defined as the limited or uncertain availability of enough food for a consistently active and healthy life. COVID-19 is a highly transmissible viral infection with high mortality due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and, or the uncommon severe pneumonia. This study assesses the relationship between food security and dietary patterns status with COVID-19 in the North Khorasan province, Iran.

Methods: This case-control study will be conducted in the men and women aged 20-60 years improved from COVID-19 infection. The cases (n=124) and controls (n=124) were selected according to the eligibility criteria, including recently improved COVID-19 according to the positive COVID-19 PCR test. People referred to public and private laboratories or employees of public and private factories, offices, and departments of hospitals and universities (for the cases) and negative PCR tests without any clinical signs of COVID-19 infection (for the controls). The North Khorasan province was the target place. The groups are matched for age, sex, and body mass index (BMI). The assessments will include anthropometric measurements and general demographic, USDA 18-item food security (18item-FSSM), and 147-item food frequency (FFQ) questionnaires. Finally, the determination of the relationship between food security and dietary patterns status and associated socioeconomic factors with COVID-19 is done. P-value will be <0.05.

Discussion: This study would be the first assessment of the relationship between food security and dietary patterns status with COVID-19 disease. It may help planners and policymakers to manage food insecurity and unhealthy dietary patterns and later increasing the immune system and decreasing the incidence of COVID-19. Further studies are suggested.

Keywords: Case-control protocol study, COVID-19, Food security, Dietary pattern, Northeastern Iran

Background

A new epidemic of coronavirus infection began in late 2019 in Wuhan, China, and was originally called nCoV 2019, then renamed COVID-19 by the World Health Organization (WHO) on February 11, 2020. Previous CoV epidemics include acute respiratory syndrome (CoV-SARS), which began in China in 2002, and middle east respiratory syndrome (CoV-MERS), first reported in 2012. These epidemics began with animal-to-human infections, and the direct cause of death is generally due to unusually severe pneumonia. However, there are other abnormal causes such as acute respiratory distress syndrome, acute heart injury, and RNAemia that can lead to death (1-4). COVID-19 is a highly transmissible viral infection with total mortality due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (5). The most common symptoms at the onset of the disease are fever, cough, and fatigue, while other symptoms include sputum production, headache, diarrhea, shortness of breath, and lymphopenia (4). The relative incidence of severe cases is about 20%, and mortality is approximately 3% (6). The WHO estimates that seasonal flu alone worldwide causes 3-5 million cases of severe illness each year that requires hospitalization and 290,000-650,000 deaths, and overall, acute respiratory diseases estimated 2.38 million deaths worldwide in 2016 (7). Severe lower respiratory tract infections were the most common cause of death from sepsis from 1990 to 2017 worldwide (6). According to recent studies, COVID-19 is associated with increased production of pro-inflammatory cytokines, reactive protein C, increased risk of pneumonia, sepsis, acute respiratory distress syndrome, and heart failure (2). The COVID-19 epidemic is perhaps the most significant global health crisis of our time, with unprecedented human challenges. One of these challenges is food insecurity. More than 820 million people went to bed hungry even before the COVID-19 epidemic, of which 110 million lived in acute food insecurity (United Nations, 2020).

Food security is defined by the Food and Agriculture Organization as physical and economic access to sufficient safe and nutritious foods to meet [one's] dietary needs and food preferences for an active and healthy life. Particularly in high-income countries, households without sufficient financial resources may turn to cheap, low-nutrient foods to buy food (8). For this reason, food insecurity can be defined as limited or uncertain access nutritionally to adequate and safe food or limited or uncertain ability to gain acceptable foods in socially acceptable ways (9). The range of food insecurity varies from anxiety about access to food at the household level to a high degree of hunger among children who have no food to eat (10). Food insecurity can be chronic, seasonal, transient, or at the level of household, region or nation and is a complex and multidimensional phenomenon that in addition to quantity and quality, also includes social, cultural and psychological dimensions (11). It is not limited to people who do not have enough energy and nutrients, but also occurs when people do not have the right to choose food, are afraid of running out of food, or make significant changes in their food preferences (12). Food insecurity may trigger a complex cycle of poorer dietary intake and compensatory overconsumption of poor quality foods, predisposing an individual to increased metabolic risk. Furthermore, food insecurity rises household stress, forces trade-offs between food and medical care, increases the risk of medication non-adherence, and leads to poorer chronic disease self-management, all-important pathways toward worse metabolic and chronic disease outcomes (13). So what food insecurity at the household level and the individual is a social disease with obvious health consequences (14). It is known that in the general population, food insecurity and chronic conditions are associated with risk factors for chronic disease. Food insecurity, inadequate intake of essential nutrients, impaired development of mental, psychological, and behavioral dysfunction in children and adults and is associated with decreased resistance to disease. The inability to buy nutritious and adequate food and the resulting psychological and emotional stress can have adverse health effects or

exacerbate illnesses caused by other risk factors. Food insecurity is indirectly associated with obesity, high blood pressure, and impaired lipids, and blood sugar (15,16). Chronic food shortages lead to nutritional deficiencies, loss of food diversity, and low power modes that affect the ability of physical activity. In addition, the risk of high blood pressure increases due to the presence of high sodium and low potassium elements in highly processed, inexpensive foods. As well as lack of food may also lead to an inability to maintain hydration (17). In addition, food insecurity is a physiologically stressful and emotional condition. This stress leads to an increase in cortisol, which is associated with an increase in the accumulation of body fat (incredibly visceral fat), which is a risk factor for diabetes and cardiovascular disease (18). Asia has less than half the prevalence of food insecurity, according to the US Economic Search Service, and according to recent meta-analyses, the prevalence of food insecurity in Iran is 49% (19, 20). Income is a determining factor in food insecurity and hunger. Special events that affect the household budget such as job loss, not having a permanent job, increasing family members or loss of food aid, as well as factors that affect diet such as ethnicity, differences in eating habits in various regions, and age and education of the head of the household are effective in creating food insecurity (21). In separate studies in the world and Iran, the relationship between food insecurity and diseases such as obesity, reduced disease resistance, hyperlipidemia, diabetes, mental retardation, premenstrual syndrome, acne, depression, gastrointestinal cancer, upper extremities, osteoporosis, starvation, malnutrition, weakened immune system and reduced quality of life (9).

Also, in recent years, a lot of attention has been paid to dietary patterns. Because dietary patterns take into account the internal relationships between dietary compounds while maintaining the diversity of all food intake and by using this method, people's nutritional behavior, which is a broad picture of food and nutrient intake, can be better reflected

concerning disease, while dietary patterns are conceptually a helpful tool for disseminating dietary recommendations for public health. With these interpretations, dietary patterns can be considered a method that can show more aspects of the relationship between diet and infectious diseases. The dietary pattern of the populations is different due to geographical differences, socio-economic status, the culture of eating habits. Therefore it is necessary to do it separately in each place, while the passage of time creates the need for new study (22).

The role that nutrition plays in supporting the immune system has been well established. A wealth of clinical data suggests that the supply of calories, proteins particularly essential amino acids and vitamins, including vitamins A, B6, B12, C, D, E, and folate. Trace elements such as zinc, iron, selenium, magnesium, copper, and omega-3 fatty acids, eicosapentaenoic acid, and docosahexaenoic acid play an essential complementary role in supporting the immune system. Bad consumption of these nutrients is overall, leading to reduced resistance to infections and therefore an increased burden of infectious diseases. Inadequate consumption of these nutrients is widespread, leading to reduced resistance to infections and thus an increased burden of infectious diseases (7).

According to the recent pandemic of COVID-19 based on the statistics of the WHO and the relationship between food insecurity and the occurrence of malnutrition and its subsequent effects on weakening the immune system and increasing the risk of infectious diseases.

Aim and objectives

This study will aim to compare the food security and dietary patterns status with the possibility of developing COVID-19 symptoms in newly improved patients and non-infected individuals.

Primary objectives

- 1) To determine and compare food security status in groups of people improved from COVID-19 and those who non-infected.

- 2) To identify and compare the status of dietary patterns in groups of people improved from COVID-19 and non-patients.
- 3) To investigate the relationship between food security and COVID-19.
- 4) To investigate the relationship between dietary patterns and COVID-19.

Secondary objectives

We are determining and comparing independent quantitative and qualitative variables (household size, number of living children, employment status, household economic status, being covered by support organizations, receiving food aid, homeownership status, education status, race or ethnicity, and blood groups) in Groups of people improved from COVID-19 and non-patients.

Methods/Design

Study design and Setting: The design of this study will be a case-control study which will be performed on men and women aged 20-60 years living in North Khorasan province, Iran, in a 1:1 ratio between case and control groups. Participants will be selected according to the study criteria from among those who refer to public and private laboratories, employees of offices, factories, administrative departments of universities, and hospitals. To minimize bias, case and control groups will be matched for the age (every ten years), gender (male and female), and body mass index (BMI) (as defined by the WHO; lean ≤ 18.5 , normal (18.5-24.9), overweight (25-29.9), grade 1 and grade 2 obesity (30-34.9 and 35-40 respectively)).

Case group: Men and women aged 20-60 years who had a positive PCR test for COVID-19, and now improved.

Control group: Men and women aged 20-60 years who have a negative PCR test for COVID-19 and no clinical signs of COVID-19. These individuals will be selected from similar sites in the case group.

To prevent possible errors and to ensure the accuracy of the test result, in general questionnaires completed by individuals in the case and control groups, will be asked questions such as specific and non-specific symptoms of COVID-19 such as gastrointestinal symptoms, respiratory symptoms, fever, lack of sense of smell and Taste, dry cough, etc.

Inclusion criteria

- Pathologically confirmed COVID-19 (group of people improved by COVID-19)
- Being in the age range of 20 to 60 years
- The willingness of individuals to participate in the study and completion of the consent form
- Lack of mental illness and physical disabilities
- No COVID-19 infection so far (group of non-infected people)
- A maximum of 3 months has passed since the onset of COVID-19

Exclusion criteria

- Being in the age range of ≤ 20 or ≥ 60 years
- The unwillingness of individuals to participate in the study and refusal to complete the consent form
- Lack of cooperation until the end of the study
- Re-infection (group of people improved by COVID-19)
- Observe the symptoms of COVID-19 during the study
- People who have not answered more than half of the food frequency questionnaire
- People whose total daily energy intake is less than or more than three standard deviations from the average energy

Variables assessed in this study: Variables evaluated in this study, include anthropometric assessments, general demographic questionnaires, USDA 18-item food security questionnaire

(item-FSSM18), and a valid 147-item food frequency questionnaire (FFQ) based on face-to-face interviews by a researcher and a colleague of a senior nutrition expert.

Dietary assessments: The usual nutritional intake will be assessed using a valid and reliable 147-item semi-quantitative food frequency questionnaire (FFQ) (23-25). Trained dietitians, researchers, and a colleague of a senior nutrition expert will be asked the participants to report their intake frequency for each food item consumed during the past year in terms of the day, week, month, and year. The reported frequency according to the desired serving size of each food item or household measure will be converted into grams per day (26).

Information on food security during the last 12 months using a validated and reliable food insecurity questionnaire that includes 18 questions and its validity and reliability has been confirmed (27) by conducting a face-to-face interview by a researcher and a colleague of a senior nutrition expert collected. Rating of 18-item USDA household food security status questionnaire is that is given positive rate to answers “often true,” “sometimes true,” “almost every month,” “some months,” and “yes” and zero score to responses “not true,” “does not know or refused,” “only 1 or 2 months”, and “no.” Finally, scores 0-2 in the secure food group, 3-7 in the insecure food group without hunger, 8-12 in the insecure food group with moderate hunger, and 13 and higher in the insecure food group with severe hunger are situated (28).

Anthropometric measurements: Weight, height, and waist circumference are determined by using a digital scale, stadiometer, and none elastic tape, respectively. They are measured thus: weight without shoes, with minimal clothing, and with a 100-g accuracy; height without shoes, standing, heels against the wall, flat and forward head, and with 0.5-cm accuracy; and waist circumference with minimal clothing, midway between the last rib and the iliac crest.

Other assessments: Demographic and socio-economic information of this study using a demographic questionnaire including age, level of education, marital status, employment

status, family economic status, number of children, ethnicity, blood type, receiving food aid, covered by health insurance and support organizations Being, medical history and taking dietary supplements will be collected. Also, this questionnaire has two more questions for the case group than the control group. These two questions include the length of time since the disease and the state of recovery from the illness (recovery and quarantine at home or hospitalization and recovery in the hospital or hospitalization and recovery in the ICU).

Statistical analysis: Statistical data analysis will be performed using SPSS software (version 16.0; SPSS Inc., Chicago, IL, USA). Mean, and standard deviation will be used to describe quantitatively continuous and discrete variables. The Kolmogorov-Smirnov test will be used to test the normality of quantitative data. Independent t-test will be used to compare the mean of standard quantitative variables between case and control groups and Mann-Whitney test will be used to compare the mean of abnormal quantitative variables between case and control groups. Also, Chi-square and Fisher tests will be used to compare the distribution of qualitative variables among the groups. The significance p-value will be less than 0.05.

The sample size: First, a pilot study to get acquainted with the research environment, how patients respond to questionnaires, changes needed General questionnaire of demographic characteristics and socio-economic factors, determining the number of samples and the accuracy of the study was conducted. According to this pilot study performed on 13 patients recovering from COVID-19 whose positive PCR test was confirmed by Esfarayen Central Laboratory, the percentage of food insecurity was 69% ($P_2 = 0.69$) and based on the pilot study performed on 13 people who had never had COVID-19 and whose PCR test result was negative were 46% ($P_1 = 0.46$). Then, the sample size was calculated 124 cases and an equal number of controls with 5% drop, 95% power, and 95% confidence interval and using the following formula:

$$n = \frac{2(Z_{1-\alpha/2} + Z_{1-\beta})^2 [P(1-P)]}{(P_1 - P_2)^2}$$

$$RR \approx OR = \frac{\frac{P_2}{1-P_2}}{\frac{P_1}{1-P_1}}$$

$$P = \frac{P_1 + P_2}{2}$$

Ethics: All protocols used in this study were approved by the Ethics Committee of Tabriz University of Medical Sciences (Ethical Code: IR.TBZMED.REC.1399.1053).

Discussion:

This study is the first case-control interview-based study to examine the food security status and dietary patterns of individuals in the case group (improved by COVID-19) and the control group (individuals who have never had a COVID-19 infection). The practical purpose of this project is to provide information to planners to design the next steps in reducing food insecurity, modifying the diet, and using the appropriate diet, boosting the immunity system, reducing the incidence of COVID-19 infection, and perhaps other infectious and viral diseases.

The study's strengths are case-control design, protocol publication, and the assessment of dietary patterns and food security status.

The study's limitations are self-reporting of dietary intakes, and lack of cooperation of some patients in the end, which may lead to patients being replaced.

Conclusion:

This study would be the first assessment of the relationship between food insecurity and dietary patterns status with COVID-19. It can help planners and policymakers manage food insecurity and poor and inadequate nutritional intakes. It may boost the immune system and decrease the incidence of viral or bacterial infections, especially COVID-19. Further studies are suggested.

Study Status:

Recruitment of the participants will begin as soon as.

Abbreviations:

BMI: body mass index; **FI:** food insecurity; **FFQ:** food frequency questionnaire

Declaration:

-Ethical Approval and Consent to participate: This study is being conducted with the approval of the ethics committee of the Tabriz University of Medical Sciences (Ethical Code: IR.TBZMED.REC.1399.1053). A written informed consent form (in Persian) is being obtained from all the participants. Participation is free, and a patient can withdraw at whatever point the person feels they are unable to continue. There is no bar to receiving the other health care services of the center. The personal information of participants will be kept secret before, during, and after the study.

-Consent for Publication: Not applicable.

-Availability of Supporting Data: The datasets used and, or analyzed during the current study are available from the corresponding author on a reasonable request.

-Competing Interests: There is no potential conflict of interests for research, authorship, and publication.

-Funding: Not applicable.

- Authors' Contributions: SBF, BPG, and MDM conceived and developed the idea for the study and writing the original draft. MBF contributed to data collection. MP has contributed to the review and editing of the draft. All authors read and approved the final manuscript.

-Consent to Publish: Not applicable.

-Acknowledgements: This MSc thesis was supported by the Tabriz University of Medical Sciences. The cooperation of the Jiroft University of Medical Sciences is acknowledged.

References

1. Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA, Cummings DA, et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *New England Journal of Medicine*. 2013;369(5):407-16.
2. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, et al. Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients*. 2020;12(4):988.
3. Zhong N, Zheng B, Li Y, Poon L, Xie Z, Chan K, et al. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People's Republic of China, in February 2003. *The Lancet*. 2003;362(9393):1353-8.
4. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *Journal of autoimmunity*. 2020:102433.
5. Hastie CE, Mackay DF, Ho F, Celis-Morales CA, Katikireddi SV, Niedzwiedz CL, et al. Vitamin D concentrations and COVID-19 infection in UK Biobank. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020.
6. Bernheim A, Mei X, Huang M, Yang Y, Fayad ZA, Zhang N, et al. Chest CT findings in coronavirus disease-19 (COVID-19): relationship to duration of infection. *Radiology*. 2020:200463.
7. Calder PC, Carr AC, Gombart AF, Eggersdorfer M. Optimal nutritional status for a well-functioning immune system is an important factor to protect against viral infections. *Nutrients*. 2020;12(4):1181.
8. Vercammen KA, Moran AJ, McClain AC, Thorndike AN, Fulay AP, Rimm EB. Food security and 10-year cardiovascular disease risk among US adults. *American journal of preventive medicine*. 2019;56(5):689-97.
9. Dorosty AR. Food insecurity and chronic diseases: the editorial. *Nutrition and Food Sciences Research*. 2016;3(3):1-2.
10. Janzadeh H, Mozaffari-Khosravi H, Javadi M. The Association of Food Insecurity, Inflammation, and Several Socioeconomic Factors with Type 2 Diabetes: A Case-Control Study. *Journal of Nutrition and Food Security*. 2020;5(1):38-46.
11. Behzadifar M, Behzadifar M, Abdi S, Malekzadeh R, Salmani MA, Ghoreishinia G, et al. Prevalence of food insecurity in Iran: a systematic review and meta-analysis. *Archives of Iranian medicine*. 2016;19(4):288-94.
12. Saiz Jr AM, Aul AM, Malecki KM, Bersch AJ, Bergmans RS, LeCaire TJ, et al. Food insecurity and cardiovascular health: Findings from a statewide population health survey in Wisconsin. *Preventive medicine*. 2016;93:1-6.
13. Golovaty I, Tien PC, Price JC, Sheira L, Seligman H, Weiser SD. Food insecurity may be an independent risk factor associated with nonalcoholic fatty liver disease among low-income adults in the United States. *The Journal of Nutrition*. 2020;150(1):91-8.
14. Leung CW, Epel ES, Ritchie LD, Crawford PB, Laraia BA. Food insecurity is inversely associated with diet quality of lower-income adults. *Journal of the Academy of Nutrition and Dietetics*. 2014;114(12):1943-53. e2.
15. Weigel MM, Armijos RX, Hall YP, Ramirez Y, Orozco R. The household food insecurity and health outcomes of US-Mexico border migrant and seasonal farmworkers. *Journal of Immigrant and Minority Health*. 2007;9(3):157-69.
16. Cook JT, Frank DA, Levenson SM, Neault NB, Heeren TC, Black MM, et al. Child food insecurity increases risks posed by household food insecurity to young children's health. *The Journal of nutrition*. 2006;136(4):1073-6.
17. Androgué H, Madias M. Mechanisms of disease. Sodium and potassium in the pathogenesis of disease. *N Engl J Med*. 2007;356:1956-78.
18. Hamelin A-M, Beaudry M, Habicht J-P. Characterization of household food insecurity in Quebec: food and feelings. *Social science & medicine*. 2002;54(1):119-32.
19. Daneshi-Maskooni M, Sakineh S-B, Badri-Fariman M, Erfan A, Mohammadi Y, Jafarnejad S, et al. Questionnaire-based prevalence of food insecurity in Iran: a review article. *Iranian Journal of Public Health*. 2017;46(11):1454.
20. Kraege V, Vollenweider P, Waeber G, Sharp S, Vallejo M, Infante O, et al. Professor Masoud Mirzaei, MD PhD Short resume. Policy. 2019.

21. Dave JM, Evans AE, Saunders RP, Watkins KW, Pfeiffer KA. Associations among food insecurity, acculturation, demographic factors, and fruit and vegetable intake at home in Hispanic children. *Journal of the American Dietetic Association*. 2009;109(4):697-701.
22. Esmaeel Zadeh A, Kimiyagar M, Mehrabi Y, Azadbakht L. Dietary patterns and risk of chronic diseases. *Iran J Diabet Lipid Disord*. 2005;5(1):1-9.
23. Mirmiran P, Esfahani FH, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public health nutrition*. 2010;13(5):654-62.
24. Keshteli AH, Esmaillzadeh A, Rajaie S, Askari G, Feinle-Bisset C, Adibi P. A dish-based semi-quantitative food frequency questionnaire for assessment of dietary intakes in epidemiologic studies in Iran: design and development. *International journal of preventive medicine*. 2014;5(1):29.
25. Asghari G, Rezazadeh A, Hosseini-Esfahani F, Mehrabi Y, Mirmiran P, Azizi F. Reliability, comparative validity and stability of dietary patterns derived from an FFQ in the Tehran Lipid and Glucose Study. *British journal of nutrition*. 2012;108(6):1109-17.
26. Ghaffarpour M, Houshiar-Rad A, Kianfar H. The manual for household measures, cooking yields factors and edible portion of foods. Tehran: Nashre Olume Keshavarzy. 1999;7:213.
27. Rafiei M, Nord M, Sadeghizadeh A, Entezari MH. Assessing the internal validity of a household survey-based food security measure adapted for use in Iran. *Nutrition journal*. 2009;8(1):28.
28. Daneshi Maskooni M, Dorosty Motlagh AR, Hosseini SM, Zendehtdel K, Kashani A, Safarpour M, et al. Food insecurity and some associated socioeconomic factors among upper gastrointestinal cancer patients. *Int Res J Appl Basic Sci*. 2013;4(2):482-6.
29. Najibi N, Dorosty Motlagh AR, Sadrzadeh Yeganeh H, Eshraghian MR, Daneshi M, Azizi S. Food insecurity status and some associated socioeconomic factors among newly diagnosed patients with type 2 diabetes in Shiraz, 2012. *Journal of Arak University of Medical Sciences*. 2013 Apr 10;16(1):98-106.
30. Safarpour M, Dorosty Motlagh A, Hosseini SM, Ranjbar Noshari F, Safarpour M, Daneshi Maskooni M, Azizi S, Kashani A. Prevalence and outcomes of food insecurity and its relationship with some socioeconomic factors. *Knowledge Health*. 2014 Jan 1;8(4):193-8.
31. Daneshi-Maskooni M, Badri-Fariman M, Habibi N, Dorosty-Motlagh A, Yavari H, Kashani A, Hosseini M. The relationship between food insecurity and esophageal and gastric cancers: a case-control study. *Journal of research in health sciences*. 2017;17(2):381.
32. Azizi S, Sadrzadeh Yeganeh H, Hosseini SM, Ahmadi A, Daneshi Maskooni M, Safarpour M, Najibi N. Food insecurity and some associated socioeconomic factors among women with metabolic syndrome referred to clinics of shiraz university of medical sciences. *Armaghan Danesh*. 2013 May 10;18(2):95-106.
33. Najibi N, Firoozi R, Shahrezaee S, Eshraghian M, Daneshi-Maskooni M, Dorosty-Motlagh A. Food insecurity is an important risk factor for type 2 diabetes: a case-control study of new referrals to the University clinics, Shiraz, Southern Iran. *BMC public health*. 2019 Dec;19(1):1-8.
34. Parsavala S, Dorosty Motlagh A, Daneshi Maskooni M, Eshraghian M, Siassi F. Food insecurity and some associated socioeconomic factors among high school girls of Tehran. *Int Res J Appl Basic Sci*. 2013;7(8):508-11.
35. Motlagh AR, Safarpour P, Maskooni MD, Hosseini M, Noshari FR. Food Insecurity and Primary School Girl Students' Intelligence Quotients: A Case-Control Study. *Journal of research in health sciences*. 2018;18(2):411.
36. Daneshi-Maskooni M, Dorosty-Motlagh A, Hosseini SM, Zendehtdel K, Kashani A, Azizi S, Safarpour M. Prevalence of Food Insecurity and Some Associated Socioeconomic Factors among Patients with Upper Gastrointestinal Cancers. *medical journal of Mashhad University of medical sciences*. 2013;56(1):56-61.
37. Maskooni MD, Motlagh AD, Izadi A, Kashani A, Azizi S, Bahreini M, Najibi N. Is Food Security Associated to Nutritional Status among Hospitalized Cancer and non-Cancer Patients. *Int Res J Appl Basic Sci*. 2013;5(6):801-4.
38. Badri-Fariman M, Naeini AA, Mirzaei K, Moeini A, Hosseini M, Bagheri SE, Daneshi-Maskooni M. Association Between the Food Security Status and Dietary Patterns With Polycystic Ovary Syndrome (PCOS) in Overweight and Obese Iranian Women: A Case-Control Study.

39. Pasdar Y, Nachvak SM, Darbandi M, Morvaridzadeh M, Rezaeian S, Daneshi Maskooni M. A Population-based Cross-sectional Study of Food Insecurity and the Influential Factors in Households in Kermanshah, Iran. *Journal of Human Environment and Health Promotion*. 2019 Sep 10;5(3):116-20.
40. Azizi S, SadrzadehYeganeh H, Hosseini M, Daneshimaskooni M. RELATIONSHIP BETWEEN FOOD INSECURITY AND METABOLIC SYNDROME IN WOMEN. *ijdd*. 2014; 13 (2):144-152