Predictors of COVID testing among Australian youth: Insights from the Longitudinal Study of Australian Children

Md Irteja Islam ^{1,2,3}, Verity Chadwick ⁴, Alexandra Martiniuk ^{2,5,6}

¹ Maternal and Child Health Division, International Centre for Diarrheal Disease Research, Bangladesh (icddr,b), 68, Mohakhali, Dhaka 1212, Bangladesh.

² Sydney School of Public Health, Faculty of Medicine and Health, The University of Sydney, Edward Ford Building, A27 Fisher Road, NSW 2006, Australia.

³ Centre for Health Research and School of Business, The University of Southern Queensland, West Street, Darling Heights, QLD 4350, Australia.

⁴ Royal North Shore Hospital, Reserve Rd, St Leonard's NSW 2065, Australia.

⁵ Office of the Chief Scientist, The George Institute for Global Health, Level 5/1 King Street, Newtown NSW 2042, Australia.

⁶ Dalla Lana School of Public Health, The University of Toronto, 155 College St Room 500, Toronto, ON M5T 3M7, Canada.

Corresponding author:

Md Irteja Islam PhD

Assistant Scientist

Maternal and Child Health Division (Level 7)

International Centre for Diarrheal Disease Research, Bangladesh (icddr,b)

68, Mohakhali, Dhaka 1212, Bangladesh

Email: irteja.islam@icddr.org

Word count: 3358

ABSTRACT

Background Testing has played a crucial role in reducing the spread of COVID. Although COVID symptoms tend to be less severe in children and adolescents, a key concern is young people's role in the transmission of the virus given their highly social lifestyles. In this study, we aimed to identify the predictors associated with COVID testing in Australian youth using data from the Longitudinal Study of Australian Children (LSAC).

Methods We used the latest wave 9C1 of the LSAC, where data were collected from 16–21-year-old Australians via an online survey between October and December 2021. In total, 2291 Australian youths responded to the questions about COVID testing and COVID symptom severity. Data was stratified by living with/without parents, and bivariate and logistic regression analyses examined predictor variables (age, sex, country of birth, remoteness, education level, employment, relationship status, number of household members, living with parents, receiving the COVID financial supplement from government and index of relative socio-economic advantage and disadvantage) and their distributions over the outcome variable COVID testing.

Results Youths aged 16-17 were more likely to live at home than youths aged 20-21 years. The strongest predictor of COVID testing was living in major cities (regardless of living with or without parents). Changed household composition was significantly associated with COVID testing among the youths living in the parental home. While among the respondents living without their parents, living with multiple household members and low or no cohesion among household members was associated with higher rates of COVID testing.

Conclusion Our study revealed young people have been very good at getting tested for COVID. To further incentivise testing in this age group, we should consider providing this age group with continued financial and social support while awaiting the outcome of the test and during any isolation.

Keywords

COVID-19, SARS-COV-2, Coronavirus, Australia, adolescents, young adults, testing

Strengths and limitations of this study

- Large national cohort of young people strengthened the findings of the study and allowing us to examine the factors associated with COVID testing for the first time in Australia.
- A broad-based assessment of potential predictors of COVID testing, including sociodemographic and coronavirus specific factor.
- Cross-sectional observational design limits causal inference.
- Self-reported information about COVID testing can be subject to recall as well as social desirability bias.

Introduction

Public health interventions for COVID including testing and isolating, social distancing and bans on mass gatherings, wearing of masks, improving ventilation, closing schools, and working remotely, have been effective worldwide in reducing the spread of the SARS-CoV-2 virus.¹ With the introduction of COVID vaccines, governments, medical and public health experts have been devising exit strategies to resume more 'pre-COVID' ways of life. While the vaccines are effective in reducing hospitalisation and mortality rates,² there are some caveats. Evidence demonstrates that vaccinated people can still transmit the virus to others, there is decreased efficacy in the elderly and immunocompromised,³ immunity tends to wane over time,⁴ some people have not had access to a vaccine and some people may choose not to get vaccinated.⁵

Testing, tracing, and isolation/quarantine (TTIQ) play crucial roles in epidemic control, which requires fast and comprehensive testing and tracing, and then sufficient compliance with isolation/quarantine.⁶ At this point in time, regular testing is mandatory in some contexts, such as some workplaces or during travel.^{7, 8} However, in other situations we must rely on individuals to recognise the need to be tested and to promptly attend for a test.

It is well established that all people, but more so children and young people, can be asymptomatic carriers of SARS-CoV-2.⁹ Although COVID symptoms and outcomes tend to be less severe in children and adolescents, a key concern is that young people may be important community reservoirs for the transmission of the virus to household members, grandparents, and other community members. Further, adolescents and young adults tend to have a higher number of in-person connections on a daily basis,¹⁰ and often live in crowded houses¹¹ and work multiple jobs.¹² These factors increasing the likelihood of adolescents and young adults playing a crucial role in SARS-CoV-2 transmission as societies open up.¹³⁻¹⁵

Limiting young people from attending education institutions and places of employment cannot be a long-term solution. School and university closures adversely affect young people as they miss out on education (particularly those from disadvantaged backgrounds),¹⁶ opportunities to develop social skills¹⁷ and find role models from outside the home.¹⁸ Indeed, 30% of Australians aged 18 to 34 years experienced high or very high levels of psychosocial distress in June 2021, almost double the rate reported by this age group in 2018-19.¹⁹ Distress was greatest in Melbourne and Sydney where the tighter lockdowns occurred, while the prevalence of high or very high psychosocial distress in other age groups remained relatively unchanged when compared to the year prior to COVID.¹⁹

Throughout the whole pandemic period to date, just under half of Australians reported they would definitely get a COVID test if experiencing mild respiratory symptoms.^{20, 21} NSW statistics suggest that approximately one quarter of daily COVID tests have been for individuals aged 10-29 years of age.²² Although testing is crucial for all age groups, testing is arguably more vital in the highly social and mobile adolescent and young adult age group.¹⁰ However, there is little knowledge regarding predictors of adolescents and young adults seeking COVID testing. Here we examine what factors are associated with rates of COVID testing in 16–21-year-old Australians.

Methods

This paper is being reported according to STROBE guidelines for observational studies.²³

Study design and Sample

We used data from Growing Up in Australia: The Longitudinal Study of Australian Children (LSAC), a population-based cross-sequential cohort study carried out by the Australian Government Department of Social Services (DSS) in a partnership with the Australian Institute of Family Studies (AIFS) and the Australian Bureau of Statistics (ABS). The LSAC sampled participants from the Medicare enrolment database, Australia's universal health insurance scheme. A multi-stage cluster sampling design was used. First, representative postcodes were selected employing probability proportion to size method, stratified by state or territory and by capital city statistical division vs. rest of state to guarantee geographically proportionate samples across urban and rural areas. Second, children were randomly selected from a selection of 311 postcodes, around 40 and 20 children per postcode in the large and small states, respectively.²⁴

The LSAC collected data biennially since 2004 from two cohorts - the birth cohort (B cohort) aged 0–1 year at baseline (n=5107) and the kindergarten cohort (K cohort) aged 4–5 years at baseline (n=4983). Out of the total 10090 children recruited during Wave 1, 2958 responded in Wave 9C1 (i.e., the latest online survey data collected during COVID pandemic in between October-November 2020 among 16-17-year-olds B cohort and 20-21-year-olds K cohort). In the LSAC, data were collected primarily from the child's parents (typically from the mother or Parent 1) with additional information from other caregivers (such as Parent 2, usually the father) and the children themselves (when aged 10 years or more for some specific questions). Data collection methods for the prior Waves 1-8 included face-to-face interview, mail-out questionnaires, time-use diaries, Computer-Assisted Self-Interview (CASI) and Computer-Assisted Web-Interview (CAWI); while due to COVID pandemic in

2020, only CAWI was used to collect data from parents and youth in LSAC Wave 9C1. More details on the LSAC methodology, including sampling procedures and data collection techniques, is described elsewhere.^{24, 25}

Our current study included 2291 Australian youth aged 16-21 years at the time of the LSAC Wave 9C1 in 2020. Out of them, 1374 (60%) and 917 (40%) youth were from B-cohort and K-cohort, respectively. Respondents who reported on the outcome variable (COVID testing) and explanatory variables were included in the analyses; while the 'Missing', 'Don't know', 'Prefer not to say' and 'Refused' responses categories were omitted (n=667).

Measures

The LSAC Wave 9C1 online survey questionnaire included questions related to the COVID pandemic (including the coronavirus restriction period between March-May 2020) and sociodemographic characteristics among the parents and/or youths.

Outcome variable

The question related to testing for COVID included whether the youth have been tested for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) with response categories 'Yes' (coded as 1) and 'No' (coded as 0).

Explanatory variables

Based on the previous literature,^{26, 27} the following socio-demographic variables were considered as predictor variables in this study: age (16-17 years, 20-21 years), sex (Male, Female), country of birth (Australia, Overseas), remoteness (Major cities, Rural/remote areas), education (Technical/others, Secondary, University/tertiary), employment (Full-time, Part-time, Unemployed), number of household members (Alone, Two people, Three people, Four or more people), living with parents (Yes, No), currently in a relationship (Yes, No), and the SEIFA (Socio-Economic Indexes for Areas) Index of Relative Socio-Economic Advantage and Disadvantage (IRSAD) quintiles. For all areas across Australia, IRSAD quintile 1 includes the lowest 20 percent for the most disadvantaged areas, and quintile 5 contains the highest 20 percent for the most advantaged areas²⁸. Regarding family cohesion, in general, the youth were directly asked to rate the ability of family or household members to get along with each other where the responses were recorded in a 5-point scale (from Excellent to Poor).²⁹ However, for analytical purposes, we created a dichotomized variable – 'cohesion among household members' from the responses. Youth who responded 'excellent', 'very good' or 'good' were classified as 'yes' (coded as 0), while those who answered, 'fair' or 'poor' were classified as 'no' (coded as 1).

Since, the respondents of LSAC Wave 9C1 experienced the COVID pandemic in 2020, the study included COVID and coronavirus restriction period (CRP) variables between March-May 2020 in Australia.²⁵ Our study thus included the following additional variables: employment status during CRP (Employed, Unemployed), changed household composition during CRP (Yes, No), and received coronavirus supplement¹ during CRP (Yes, No) to assess whether these variables are associated with COVID testing among the respondents. The following question was directly asked to youth in W9C1 about the changes in household composition since the previous LSAC W8 in 2018 as the data were rolled forward: 'Whether the household composition changed during COVID-19 restrictions?'. Note that the regular detailed household data could not be gathered in W9C1 since the questions were meant to be asked in a face-to-face interview (which was not possible during CRP) and could not be fully converted into online inquiries.²⁵

Statistical analysis

To understand the sociodemographic predictors of COVID testing among youths, stratified by living with/without parents, initially, we computed summary statistics. Then, we conducted bivariate analyses to examine the predictor variables and their distributions over the outcome variables (COVID testing). Pearson's chi-squared test³¹ signified the strength of the bivariate associations between the characteristics and youth COVID testing, stratified by living with/without parents. Further, we employed logistic regression analysis to identify predictors of COVID testing. The results of the regression analyses were presented as adjusted odds ratios (aOR) with corresponding 95% confidence intervals (CI) after adjusting for possible confounding effects of other explanatory variables. All statistical analyses were conducted with Stata/MP 14.1, and 'svy' command³² was used to take account for of the LSAC's multi-stage clustered sampling design, and to deal with potential bias and to avoid overestimation of the significance.

Patient and public involvement

The LSAC is a population-based longitudinal study, hence no patient groups were engaged in its design or execution. To the best of our knowledge, the public was not involved in designing, recruiting, or implementation of the LSAC study. Parents received a brief health report for their children and themselves during or shortly after the assessment visit. They

¹An additional top-up payment for people on welfare during COVID-19 pandemic in Australia 30. Australia S. Coronavirus Supplement, <https://www.servicesaustralia.gov.au/individuals/services/centrelink/coronavirus-supplement> (2021, accessed 19 October 2021 2021).

agreed to participate understanding that they would not otherwise get individual results about themselves or their children.

Ethics

The LSAC has been approved by the Human Research Ethics Committee of the Australian Institute of Family Studies (AIFS) (Application number 20-09), and written informed consent was obtained for all study participants. In addition, the authorship team obtained permission from Australian Data Archive Dataverse to use LSAC data for research and publications (Reference No. 263493).

Results

Descriptive statistics of the study population are demonstrated in Table 1. Of the 2291 youth, 60% were aged between 16-17 years, more than 56% were female, 97% were born in Australia, 73% were from major cities, 57% had secondary level educational qualifications, and 57% had part-time employment. About 75% of youth had three or more household members, 84% reported cohesion among the household members, 88% of youth were living with parents and 30% were in a relationship. Nearly 70% of respondents were from the disadvantaged group (Quintile 1, Quintile 2 and Quintile 3 combined). During the coronavirus restriction period (CRP) between March-May 2020, more than 50% of youth were unemployed, the majority of the respondents (84%) did not experience changed household composition, and 82% of youth did not receive the coronavirus financial supplement.

Fig. 1 displays the percentages of youth tested for COVID, stratified by living status. Overall, 587 (26%) youth got tested for COVID. Out of which, 85% of youth were living with parents, and 15% were living without parents.

Bivariate relationships between explanatory variables and COVID testing among youths, stratified by living status are reported in Table 2. Among those who were living with their parents, more than 60% ($p=0.009$) of youth were those aged 16-17 years compared to the age group 20-21 years. Youth who were living in major cities compared to those living in rural/remote areas, and youth with changed household composition during CRP than their counterparts were also found to be factors associated with COVID testing among the young people who were living with their parents ($p<0.010$ for all). While for youths not living with parents, Table 2 also shows that remoteness and cohesion among household members were significantly associated with COVID testing ($p<0.05$ for all).

Results from logistic regression models of respondents being tested for SARS-CoV-2 testing are portrayed in Table 3. The strongest predictor of COVID testing was living in major cities regardless of whether the young person was living with parents (aOR 1.82, 95% CI: 1.27-2.60) or without parents (aOR 2.25, 95% CI: 1.20-4.20) compared to those who were living in rural/remote areas. Among those who were living with parents, changed household composition during the coronavirus restriction period (CRP) increased the probability of COVID testing (aOR 1.33, 95% CI: 0.98-1.81) compared to their counterparts. Among the respondents who were living without parents, living with multiple household members (Two people: aOR 5.92, $p < 0.001$; three people: aOR 5.34, $p < 0.001$; Four or more people: aOR 2.66, $p < 0.05$) significantly increases the likelihood of COVID testing compared to those who lived alone. Moreover, no cohesion among the household members (aOR 6.03, 95% CI: 2.40-15.16) increases the probability of getting COVID tested among the youth who were not living with parents compared to those who had a cohesive household.

Discussion

This study examined factors associated with COVID testing in adolescents and young adults. Those living with their parents were more likely to get a COVID test, and this was especially true if there had been a change in their household composition. Among those not living with their parents, youths were more likely to get a COVID test if they were living in major cities, if they had non-cohesive households, and if they were in households consisting of two or more people.

Higher COVID testing rates were observed for adolescents and young adults who were living with their parents. This may be related to parents monitoring their children for symptoms of COVID, and encouraging and taking them to testing facilities if symptoms were present.³³ Other reports have found parents are more likely to have their children tested if other family members lived at home, or if the family was in continual contact with relatives or at-risk individuals.³⁴ This could relate to parents seeking COVID testing to protect others - ultimately pro-social behaviour.³³

Changes in family composition were also associated with higher test rates when adolescents lived at home. For example, this could be attributed to family members moving to live with grandparents, parents separating, or children moving back home. This may be associated with a larger number of family members at home and therefore an increased risk of COVID entering the household.³⁴ It could also be related to family members in different homes wanting returning household members (for example, children going between families) to get tested to protect household members.³⁵ Previous research during the COVID pandemic has

shown that public health restrictions have led to negative impacts on spousal relationships,³⁶ with higher rates of domestic violence³⁷ and divorce compared to pre-pandemic.³⁸ We were not able to capture the extent to which household composition changes were related to separating parents within the coronavirus restriction period, but the Family Court of Australia did report a 39% increase in parenting-related disputes in the second month of the pandemic.³⁹ This may be attributed to parents in different homes expressing concern about the other household's adherence to social distancing and mask wearing,³⁵ or the number of people in contact with other households as reported elsewhere.^{35, 40}

Further, households with low or no cohesiveness (a measure of connectivity and caring within households) increased the testing rates for young people living outside of parental home. Low cohesiveness could mean that these youths cannot trust their household members to adequately protect themselves from COVID. In turn, their higher testing rates may be an effort to protect their families when visiting the family home, supported by parents who are mindful of the risks of COVID and recommend protective measures.⁴¹ Government promotions have framed the motivation as helping the community rather than avoiding individual risk,⁴² or using identity-based messages such as "don't be a spreader".⁴³ However, those living out of home with more secure financial footing were the least likely to have COVID tests, as indicated by the significant negative relationship between quintile 4 IRSAD and COVID testing. This could reflect possible instability in their employment position and unwillingness to risk calling in sick to isolate if testing positive.

For adolescents not living at home with their parents, factors associated with increased testing rates included living in urban areas, and the likelihood of testing increased with presence of multiple household members. Urban areas have experienced greater outbreaks of COVID⁴⁴ given increased living density, and connectivity to people.⁴⁵ In Australia, this is also because most cases (prior to June 2021) were imported from overseas and spread into the urban community from quarantine facilities based in cities. Furthermore, people living outside of urban areas may have less access to testing. Recent studies have found that the one of the most common reasons for symptomatic individuals not getting tested was not knowing where to go.^{46, 47} COVID testing rates for adolescents were lower in rural communities, and more should be done to increase these rates. Given the common combination of poverty, the higher prevalence of comorbid diseases, and poor accessibility to healthcare,⁴⁸ individuals in rural settings are likely to experience worse outcomes when COVID outbreaks occur, as was the case for the Wilcannia outbreak in rural New South Wales.⁴⁹

The majority (70%) of our study respondents were socioeconomically disadvantaged, but despite this over a quarter were tested for COVID. In the future, this age group may be dissuaded from getting tested given they may be vaccinated, asymptomatic or only mildly affected, and they may not want to isolate after testing given the potential loss of casual-work income and socialising opportunity. Some studies of predictors of COVID testing have been published in other populations. In the USA, factors predicting testing included: free testing, concern about health, and having a high number of social contacts.^{10, 50}

Interestingly, in a study using hypothetical scenarios, individuals said cost and ability to support oneself if needing to isolate were of no concern and did not influence their likelihood to get tested.¹⁰ However, in practice American Latino and Black respondents and those experiencing financial strain were disproportionately likely to indicate that resource factors would influence their decision to get tested.⁵⁰ This effect remained after controlling for being uninsured and experiencing material hardship and general financial constraints.⁵⁰ Such findings are consistent with other research^{51, 52} indicating that basic survival needs of individuals and families often outweigh other consideration in healthcare decision-making, including prosocial motives like avoiding transmission of infection to others. Further studies demonstrate that the reasons adolescents do not participate in asymptomatic COVID testing include concerns about the mental health impact of self-isolation, the impact on others if the test is positive (such as forcing others into isolation),⁵³ and thinking that it is a potential waste of resources.⁵⁴ More effective economic support for youth may assist in supporting this age group to be tested for COVID.

The generalisability of these findings may be limited to Australian young people given that the sample was not representative of the Australian overall population. Another limitation may be the difficulty in comparing these findings to other countries, for example, Australia's COVID testing rate is much higher than many other countries; 2.7 per day per 1,000 people⁵⁵ compared to the US (1.9 per 1,000), the UK (1.3 per 1,000) and India (0.654 per 1,000).⁵⁶ This may be due to Australia's relatively high national wealth (and lower inequality compared to US), as well as financial support for individuals and businesses afflicted by the pandemic.⁵⁷

Conclusion

In conclusion, adolescents and young adults living with their parents were more likely to be tested for COVID compared to those who lived outside of the family home, and changes in who lived at home during the pandemic was associated with higher testing rates. For youths not living at home, living in the city, living with two or more housemates, and for

participants reporting poor unity of households, testing rates were higher. In conclusion, while young people have been very good at getting tested for COVID, to further incentivise testing in this age group, we should consider providing financial and social support while awaiting the outcome of the test and during any isolation.

Data availability statement

The LSAC dataset is publicly available from the National Centre for Longitudinal Data (NCLD) Dataverse on request available at <https://growingupinaustralia.gov.au/data-and-documentation/accessing-lsac-data>. Authors do not have permission to share this unit record data without endorsement from the Department of Social Services and Australian Institute of Family Studies.

Ethics statements

Patient consent for publication

Not applicable.

Ethics approval and consent to participate

The LSAC has been approved by the Human Research Ethics Committee of the AIFS, and written informed consent was obtained for all study participants. In addition, the authorship team obtained permission from Australian Data Archive Dataverse to use LSAC data for research and publications.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interest statement

The authors declare that they have no competing interests.

Author's contribution

MII conceptualized the study, analyzed the data, drafted parts of the initial manuscript, reviewed and revised the manuscript. VC drafted parts of the initial manuscript, and reviewed and revised the manuscript. AM conceptualized the study, drafted parts of the initial manuscript, reviewed and revised the manuscript. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

References

1. Kerr CC, Mistry D, Stuart RM, et al. Controlling COVID-19 via test-trace-quarantine. *Nature communications* 2021; 12: 1-12.
2. Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *New England Journal of Medicine* 2021; 384: 1412-1423.
3. Tenforde MW, Patel MM, Ginde AA, et al. Effectiveness of SARS-CoV-2 mRNA vaccines for preventing Covid-19 hospitalizations in the United States. *Clin infect dis* 2021.
4. Collier A-rY, Yu J, McMahan K, et al. Differential Kinetics of Immune Responses Elicited by Covid-19 Vaccines. *New England Journal of Medicine* 2021.
5. Troiano G and Nardi A. Vaccine hesitancy in the era of COVID-19. *Public Health* 2021.
6. Ashcroft P, Lehtinen S and Bonhoeffer S. Quantifying the impact of test-trace-isolate-quarantine (TTIQ) strategies on COVID-19 transmission. *medRxiv* 2021: 2020.2012.2004.20244004.
7. Bielecki M, Patel D, Hinkelbein J, et al. Reprint of: Air travel and COVID-19 prevention in the pandemic and peri-pandemic period: A narrative review. *Travel medicine and infectious disease* 2020; 38: 101939.
8. Maguire K. Rapid antigen COVID tests a welcome relief for interstate truck drivers, but cost remains unclear. *ABC New England*, 2021.
9. Sinha IP, Harwood R, Semple MG, et al. COVID-19 infection in children. *Lancet Respir Med* 2020; 8: 446-447. 2020/04/01. DOI: 10.1016/s2213-2600(20)30152-1.
10. Thunström L, Ashworth M, Shogren JF, et al. Testing for COVID-19: Willful ignorance or selfless behavior? *Behavioural Public Policy* 2021; 5: 135-152.
11. Heath S and Kenyon L. Single young professionals and shared household living. *Journal of Youth Studies* 2001; 4: 83-100.
12. AIFS. *The Longitudinal Study of Australian Children Annual Statistical Report 2016*. 2017. Melbourne: Australian Institute of Family Studies.
13. NBC News. 70 Cases of COVID-19 at French Schools Days after Reopening. *NBC News*, 2020.
14. Sanità I and Superiore D. Apertura delle scuole e andamento dei casi confermati di SARS-CoV-2. *l'andamento in Italia: dicembre*. 2020.
15. Stein-Zamir C, Abramson N, Shoob H, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel. *Euro Surveillance* 2020; 25.
16. Eyles A, Gibbons S and Montebruno P. *Covid-19 school shutdowns: What will they do to our children's education?* 2020. London, UK: Centre for economic performance, LSE.
17. Orben A, Tomova L and Blakemore S-J. The effects of social deprivation on adolescent development and mental health. *The Lancet Child & Adolescent Health* 2020; 4: 634-640.
18. United Nations Educational and Scientific and Cultural Organization (UNESCO). *Adverse effects of school closures*. 2020.
19. Australian Institute of Health and Wellbeing. *Suicide & self-harm monitoring: The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19*. 2021. AIHW.
20. Australian Bureau of Statistics. One in two would test if they had mild COVID-19 symptoms. ABS, 2020.
21. Australian Bureau of Statistics. *Household Impacts of COVID-19 Survey*. 2021. Canberra: ABS.
22. NSW Government. COVID-19 tests by date and age range. NSW, Australia 2020.
23. Von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Bulletin of the World Health Organization* 2007; 85: 867-872.

24. Soloff C, Lawrence D and Johnstone R. *Sample design*. 2005. Australian Institute of Family Studies Melbourne.
25. Mohal J, Lansangan, C., Gasser, C., Taylor, T., Renda, J., Jessup, K., and Daraganova, G. *Growing Up in Australia: The Longitudinal Study of Australian Children – Data User Guide, Release 9C1*. June 2021 2021. Melbourne, Australia: Australian Institute of Family Studies.
26. Wu DC, Jha P, Lam T, et al. Predictors of self-reported symptoms and testing for COVID-19 in Canada using a nationally representative survey. *PloS one* 2020; 15: e0240778.
27. Murillo-Zamora E, Aguilar-Sollano F, Delgado-Enciso I, et al. Predictors of laboratory-positive COVID-19 in children and teenagers. *Public health* 2020; 189: 153-157.
28. Australia Co. *Census of Population and Housing: Socio-Economic Indexes for Areas (SEIFA), Australia*. 2013. Canberra, Australia: Australian Bureau of Statistics
29. AIHW. *National framework for protecting Australia's children indicators*. 2021. Canberra: Australian Institute of Health and Welfare.
30. Australia S. Coronavirus Supplement, <https://www.servicesaustralia.gov.au/individuals/services/centrelink/coronavirus-supplement> (2021, accessed 19 October 2021 2021).
31. Agresti A. *Categorical data analysis*. John Wiley & Sons, 2003.
32. StataCorp. *Stata survey data reference manual Release 17*. 2021. Texas, USA: StataCorp LLC.
33. Woodland L, Mowbray F, Smith L, et al. What influences whether parents recognise COVID-19 symptoms, request a test and self-isolate: A qualitative study. 2021.
34. Gaffney A, Himmelstein D and Woolhandler S. Risk for Severe COVID-19 Illness Among Teachers and Adults Living With School-Aged Children. *Annals of Internal Medicine* 2020; 173: 765-767. DOI: 10.7326/m20-5413 %m 32822221.
35. Goldberg AE, Allen KR and Smith JZ. Divorced and separated parents during the COVID-19 pandemic. *Family Process* 2021; 60: 866-887.
36. Muscat D, Ayre J, Mac O, et al. Psychological, social and financial impact of COVID-19 on culturally and linguistically diverse communities: a cross-sectional Australian study. *medRxiv* 2021: 2021.2010.2019.21265230. DOI: 10.1101/2021.10.19.21265230.
37. Lebow JL. Family in the Age of COVID-19. *Family process* 2020; 59: 309-312. 2020/05/15. DOI: 10.1111/famp.12543.
38. Prasso S. China's divorce spike is a warning to rest of locked-down world. *Bloomberg Businessweek* 2020; 31.
39. Family Court of Australia. Media release—The courts launch COVID-19 list to deal with urgent parenting disputes. 2020.
40. Frederiksen B, Gomez I, Salganicoff A, et al. Coronavirus: A look at gender differences in awareness and actions. Kaiser Family Foundation. Published Mar 20, 2020. 2020.
41. Bloomer M and Walshe C. Smiles behind the masks: A systematic review and narrative synthesis exploring how family members of seriously ill or dying patients are supported during infectious disease outbreaks. *Palliative Medicine* 2021.
42. Jordan J and Yoeli E. Don't get it or don't spread it? Comparing self-interested versus prosocially framed COVID-19 prevention messaging. *PsyArXiv Prepr* 2020.
43. Yonemitsu F, Ikeda A, Yoshimura N, et al. Warning 'Don't spread' vs. 'Don't be a spreader' to prevent the COVID-19 pandemic. *R Soc Open Sci* 2020.
44. Marziali ME, Hogg RS, Oduwole OA, et al. Predictors of COVID-19 testing rates: A cross-country comparison. *International Journal of Infectious Diseases* 2021; 104: 370-372.
45. Hamidi S, Sabouri S and Ewing R. Does density aggravate the COVID-19 pandemic? Early findings and lessons for planners. *Journal of the American Planning Association* 2020; 86: 495-509.

46. Graham MS, May A, Varsavsky T, et al. Knowledge barriers in the symptomatic-COVID-19 testing programme in the UK: an observational study. *medRxiv* 2021.
47. Bonner C, Batcup C, Ayre J, et al. Behavioural barriers to COVID-19 testing in Australia. *medRxiv* 2020.
48. Newkirk V. The coronavirus's unique threat to the south: more young people in the south seem to be dying from covid-19. *Atlantic*.
49. Green S. COVID in Wilcannia: a national disgrace we all saw coming. *The Conversation* 2021.
50. Perry BL, Aronson B, Railey AF, et al. If you build it, will they come? Social, economic, and psychological determinants of COVID-19 testing decisions. *PloS one* 2021; 16: e0252658.
51. Grossman Z and Van Der Weele JJ. Self-image and willful ignorance in social decisions. *Journal of the European Economic Association* 2017; 15: 173-217.
52. Gigerenzer G and Garcia-Retamero R. Cassandra's regret: The psychology of not wanting to know. *Psychological review* 2017; 124: 179.
53. Blake H, Knight H, Jia R, et al. Students' views towards Sars-Cov-2 mass asymptomatic testing, social distancing and self-isolation in a university setting during the COVID-19 pandemic: A qualitative study. *International journal of environmental research and public health* 2021; 18: 4182.
54. Mowbray F, Woodland L, Smith LE, et al. Is my cough a cold or covid? A qualitative study of COVID-19 symptom recognition and attitudes towards testing in the UK. *medRxiv* 2021.
55. Roser M, Ritchie. H and Ortiz-Ospina. E. Coronavirus pandemic (COVID-19), <https://ourworldindata.org/coronavirus> (2020).
56. Guardian Australia. Coronavirus Australia numbers: how many new cases are there? Covid-19 map, charts, statistics and graphs. 2020.
57. O'Sullivan D, Rahamathulla M and Pawar M. The impact and implications of COVID-19: An Australian perspective. *The International Journal of Community and Social Development* 2020; 2: 134-151.

Table 1 Sample characteristics (n=2291)

Characteristics	Total n	%
Age (in years) [Mean=18.24, SD=2.01]		
16-17	1374	60.0
20-21	917	40.0
Sex		
Male	990	43.2
Female	1301	56.8
Country of birth		
Overseas	59	2.6
Australia	2232	97.4
Remoteness		
Rural/Remote	624	27.2
Major cities	1667	72.8
Education		
Technical/Others	175	7.7
Secondary	1300	56.7
University/Tertiary	816	35.6
Employment status		
Full-time/Self-employed	152	6.6
Part-time	1320	57.6
Unemployed	819	35.8
Number of household members		
Alone	28	1.2
Two people	543	23.7
Three people	819	35.8
Four or more people	901	39.3
Cohesion among household members ¹		
No	372	16.2
Yes	1919	83.8
Living with parents		
No	276	12.1
Yes	2015	87.9
Currently in a relationship		
No	1593	69.5
Yes	698	30.5
IRSAD Quintiles ²		
Q1 (0-20%) - Most disadvantaged	556	24.3
Q2 (20-40%)	465	20.3
Q3 (40-60%)	548	23.9
Q4 (60-80%)	330	14.4
Q5 (80-100%) - Most advantaged	392	17.1
COVID related factors		
Employment status during CRP ³		
Unemployed	1215	53.0
Employed (part-time/full-time)	1076	47.0
Changed household composition ⁴ during CRP		
No		84.2
Yes	361	15.8
Received coronavirus supplement ⁵ during CRP		
No	1881	82.1
Yes	410	17.9

Characteristics	Total n	%
<p>Notes:</p> <p>¹ Cohesion - the ability of family or household members get along with each other</p> <p>² IRSAD - Index of Relative Socioeconomic Advantaged and Disadvantaged, is a general measure of both relative socioeconomic advantage and disadvantage at the area level. It uses a range of different Census variables including income, education, employment, occupation and housing characteristics.</p> <p>³ CRP - Coronavirus Restriction Period, between March and May 2020.</p> <p>⁴ Changed household composition - Whether the composition of household changed (any person added or moved out) during COVID restriction since the previous wave.</p> <p>⁵ Coronavirus supplement - Additional financial support for people on welfare during COVID pandemic from Australian Government</p>		

Table 2 Bivariate associations between factors and COVID testing, stratified by youth living with/without parents

	Study child having been tested	
	Living with parents (n=499)	Not living with parents (n=88)
Age (in years)		
16-17	310 (62.1)	8 (9.1)
20-21	189 (37.9)	80 (90.9)
p-value	0.009**	0.898
Sex		
Male	210 (42.1)	36 (40.9)
Female	289 (57.9)	52 (59.1)
p-value	0.354	0.559
Country of birth		
Overseas	10 (2.0)	4 (4.6)
Australia	489 (98.0)	84 (95.4)
p-value	0.631	0.930
Remoteness		
Rural/Remote	105 (21.1)	20 (22.7)
Major cities	394 (78.9)	68 (77.3)
p-value	0.001**	0.050*
Education		
Technical/Others	35 (7.0)	13 (14.8)
Secondary	295 (59.1)	7 (7.9)
University/Tertiary	169 (33.9)	68 (77.3)
p-value	0.051	0.883
Employment status		
Full-time/Self-employed	35 (7.0)	16 (18.2)
Part-time	294 (58.9)	45 (51.1)
Unemployed	170 (34.1)	27 (30.7)
p-value	0.168	0.247
Number of household members		
Alone	_a	9 (10.2)
Two people	104 (20.8)	51 (58.0)
Three people	201 (40.3)	18 (20.5)
Four or more people	194 (38.9)	10 (11.3)
p-value	0.107	0.691
Cohesion among household members		
No	82 (16.4)	25 (28.4)
Yes	417 (83.6)	63 (71.6)
p-value	0.620	0.016*
Currently in a relationship		
No	338 (67.7)	66 (75.0)
Yes	161 (32.3)	22 (25.0)
p-value	0.560	1.00
IRSAD Quintiles		
Q1	110 (22.0)	25 (28.4)
Q2	103 (20.6)	15 (17.1)
Q3	126 (25.3)	27 (30.7)
Q4	73 (14.6)	11 (12.5)
Q5	87 (17.8)	10 (11.4)
p-value	0.760	0.162
Employment status during CRP		
No	278 (55.7)	53 (61.4)
Yes	221 (44.3)	34 (38.6)
p-value	0.053	0.908

	Study child having been tested	
	Living with parents (n=499)	Not living with parents (n=88)
Changed household composition during CRP		
No	424 (84.9)	42 (47.7)
Yes	75 (15.1)	46 (52.3)
p-value	0.003**	0.274
Received coronavirus supplement during CRP		
No	420 (84.2)	43 (48.9)
Yes	79 (15.8)	45 (51.1)
p-value	0.173	0.278
Notes: Pearson's chi-squared test signified the bivariate relationships Level of significance considered ***p<0.001, **p<0.01, *p<0.05 ^a As the youth is living with parents, they cannot be living alone in a household		

Table 3 Logistic regression models for youth having been tested

		Youth having been tested Living with parents		Not living with parents	
		Adjusted OR	95% CI	Adjusted OR	95% CI
Age (Ref. 16-17)	20-21	1.54	0.75-3.16	2.75	0.29-25.75
Sex (Ref. Male)	Female	1.17	0.94-1.46	1.29	0.55-3.01
Country of birth (Ref. Overseas)	Australia	1.05	0.37-2.98	1.64	0.31-8.47
Remoteness (Ref. Rural/Remote)	Major cities	1.82**	1.27-2.60	2.25*	1.20-4.20
Education (Ref. Technical/Others)	Secondary	1.41	0.68-2.89	2.50	0.18-33.84
	University/Tertiary	0.92	0.62-1.39	0.60	0.18-1.95
Employment status (Ref. Full-time/Self-employed)	Part-time	0.78	0.44-1.39	0.53	0.20-1.39
	Unemployed	0.84	0.46-1.55	0.74	0.29-1.89
Number of household member (Ref. Lived alone)	Two people	- ^a	-	5.92***	2.48-14.13
	Three people	1.05	1.94-5.92	5.34***	2.67-10.66
	Four or more people	0.84	1.83-9.09	2.66*	1.13-6.27
Cohesion among household members (Ref. Yes)	No	1.12	0.82-1.34	6.03**	2.40-15.16
Currently in a relationship (Ref. No)	Yes	1.06	0.84-1.46	1.13	0.56-2.29
IRSAD Quintile (Ref. Q1)	Q2	0.85	0.51-1.40	0.51	0.15-1.70
	Q3	0.94	0.66-1.36	0.87	0.27-2.76
	Q4	0.82	0.60-1.13	0.31**	0.14-0.66
	Q5	0.69	0.44-1.09	0.62	0.26-1.48
Employment status during CRP (Ref. Unemployed)	Employed	0.81	0.57-1.07	0.87	0.41-1.84
Changed household composition during CRP (Ref. No)	Yes	1.33*	0.98-1.81	1.29	0.52-3.16
Received coronavirus supplement during CRP (Ref. No)	Yes	1.04	0.78-1.31	0.85	0.35-2.04
Level of significance: ***p<0.001, **p<0.01, *p<0.05					
^a For youth living with parents, 'Two people' used as Ref category since they cannot be living alone in a household

Figure 1 Percentages of youth been COVID tested, stratified by living with and without parents