Calorie reformulation: A systematic review and meta-analysis examining the effect of manipulating food energy density on daily energy intake and body weight

Eric Robinson¹, Mercedes Khuttan¹, India McFarland-Lesser¹, Zina Patel¹, Andrew Jones¹

¹Department of Psychology, Eleanor Rathbone Building, University of Liverpool
Liverpool, L69 7ZA, UK

Corresponding Author: Dr Eric Robinson, above address, email - eric.robinson@liv.ac.uk

Running Head: Energy density and energy intake

Registration. Registered on PROSPERO (CRD42020223973).

Data Sharing. Data described in the manuscript, code book, and analytic code is publicly
and freely available without restriction at <https://osf.io/dj4yf/>

Abstract

Background. Dietary energy density is thought to be a contributor to obesity, but the extent to which different magnitudes and types of reductions to food energy density decreases daily energy intake is unclear.

Objective. To systematically review and meta-analyse experimental studies that have examined the effect that manipulating energy density of food has on total daily energy intake.

Design. A systematic review and multi-level meta-analysis of studies on human participants that used an experimental design to manipulate the energy density of foods served and measured energy intake for a minimum of one day.

Results. Thirty-one eligible studies contributed 90 effects comparing the effect of higher vs. lower energy density of served food on daily energy intake to the primary meta-analysis. Lower energy density of food was associated with a large decrease in daily energy intake (SMD = -1.002 [95% CI: -0.745 to -1.266]). Findings were consistent across studies that did vs. did not manipulate macronutrient content to vary energy density. The relation between decreasing energy density and daily energy intake tended to be strong and linear, whereby compensation for decreases to energy density of foods (i.e. by eating more at other meals) was minimal. Meta-analysis of (n=5) studies indicated that serving lower energy dense food tended to be associated with greater weight loss than serving higher energy dense food, but this difference was not significant (-0.7kg, 95% CIs: -1.34, 0.04).

Conclusions. Decreasing the energy density of food can substantially reduce daily energy intake and may therefore be an effective public health approach to reducing population level energy intake.

Key Words: energy density; energy intake; obesity; food reformulation; low fat

Introduction

Energy density is the amount of energy contained in a given weight of food (kcal/gram). Fat (9kcal/g), protein (4kcal/g), carbohydrate (4kcal/g) and water (0kcal/g) content explain variations in food energy density (1). Although some foods tend to be higher in energy density (e.g. confectionary) than others (e.g. fruits and vegetables), even among the same type of food product there can be substantial variation in energy density between product brands (2). Increased availability of low-cost energy dense food products is likely a major contributor to higher obesity prevalence (3-5). Epidemiological data indicate that diets which are more energy dense tend to be associated with higher daily energy intake and weight gain (6-9). These findings have led to suggestions that population level approaches to reduce dietary energy density may be effective in reducing obesity (1, 10, 11). However, epidemiological studies examining energy density and risk of obesity have produced some inconsistent findings (12, 13), which may be due to the methodological challenges of accurately measuring dietary energy density and daily energy intake from self-report measures (6, 7, 13).

A number of laboratory studies have examined the causal impact that food energy density has on short-term energy intake, and it is now well established that reducing the energy density of a meal decreases acute energy consumed at that meal (10, 14). The longer-term effects of manipulating food energy density on energy intake are less well understood (14). Studies which have manipulated the energy density of some, or all food served to participants throughout the day indicate that daily energy intake tends to decrease when the energy density of served food is reduced (15-20). However, as energy density may influence the physiological processing and digestion of food (14), consumers may at least in part ‘compensate’ for reductions to food energy density by increasing consumption of other food. In line with this, there is some evidence that self-reported hunger is higher after consumption

of lower energy dense food as opposed to higher energy dense foods (21). When weight of food is held constant lower energy dense foods are also associated with greater later food intake than higher energy dense foods (22). From an applied perspective, it will be important to understand whether the method used to alter energy density of food, such as holding macronutrient composition constant vs. altering macronutrient composition (e.g. reducing % of kcals from fat), has a significant impact on the extent to which consumers compensate for changes in food energy density (19). It is also possible that the impact reducing energy density has on energy intake may be non-linear. In particular, compensatory responses in appetite may be more likely to occur for foods that are lower in energy density (i.e. < 1.75kcal/g), as opposed to more highly energy dense foods (23), as it has been suggested that humans evolutionary past leaves them poorly adapted to the recent emergence of very energy dense foods (24). Yet, these questions remain unanswered in relation to the impact that manipulating food energy density has on daily energy intake. In addition, although there is some evidence that dietary advice designed to decrease dietary energy density may benefit weight loss (1, 25), there is currently a lack of consensus on the direct impact of reformulating the energy density of food products has on body weight.

The primary aim of the present research was to systematically review and meta-analyse studies that have examined the impact that reducing energy density of served food has on daily energy intake. Secondary aims included understanding moderators of the effect that altering energy density has on daily energy intake and effects on body weight.

95

96 **Method**

PRISMA guidelines were followed (26). This review was registered on PROSPERO (CRD42020223973) and the analysis protocol was pre-registered <https://osf.io/dj4yf/> In the pre-registration we intended to review studies on both energy density and portion size in the

same report. However, prior to data extraction we updated the protocol to review portion size and energy density studies separately due to the large number of studies identified. Here we focus on energy density studies and studies on portion size are reviewed elsewhere (27).

Eligibility criteria.

Participants. Only studies sampling human participants were eligible. Studies were excluded that sampled participants who were currently undergoing any medical treatment which may influence appetite (e.g. bariatric surgery).

Intervention. Studies were required to have manipulated the energy density of food products or meals (i.e. energy content divided by weight of food served; kcal/gram) served to participants. Studies were included that manipulated the energy density of a minimum of one food/meal, and studies that manipulated energy density of up to all foods/meals served across the day were eligible. If a study only manipulated the energy density of a beverage it was deemed ineligible, as the main focus was on food energy density. However, if a study manipulated the energy density of multiple foods/meals and also extended the manipulation to accompanying drinks, it was eligible. To be eligible, studies were required to manipulate energy density by serving participants one or more varying energy densities of the same or very similar type of food/meal (e.g. lower vs. higher energy dense tomato-based pasta dish). Eligible manipulations of energy density included altering the % of energy derived from fat, protein and/or carbohydrate (e.g. standard vs. low-fat cheese). Studies that manipulated energy density through altering water content (e.g. adding water to a porridge) or substitution of lower energy dense foods (e.g. vegetables) were eligible.

Comparator. In studies with two energy density conditions the ‘comparator’ condition was the condition with the highest energy density and the condition with the lowest energy density was the ‘intervention’ condition (as public health interventions tend to aim to

decrease energy intake). It was common for studies to have multiple energy density conditions (e.g. higher vs. medium vs. lower) and all contrasts were included for use as individual contrasts (e.g. higher vs. medium, higher vs. lower, medium vs. lower).

Outcomes. To be eligible, studies had to have measured energy intake for a minimum of one day (i.e. at least 3 main meals). Measurements of energy intake that were based on an objective researcher measurement (e.g. weighing of food pre/post eating in the laboratory), participant self-reports (e.g. dietary recall) or a combination were eligible. Energy intake could be assessed under controlled laboratory settings or in real-world settings to be eligible. Measures of energy intake that were not determined by sampled participants (e.g. an infant being bottle or spoon fed) were not eligible.

Study Design. Studies using a within-subjects/repeated measures design (i.e. participants receive all energy density conditions) or a between-subjects design studies (i.e. participants receive only one energy density condition) were eligible. Some studies required participants to consume a meal/food in full (e.g. consumption of a set amount of energy density manipulated food) and these designs were eligible. Studies that ‘crossed’ energy density manipulations with another experimental factor (e.g. manipulation of both energy density of food and portion sizes in the same study) were eligible. For studies that did not manipulate energy density of all meals/foods, studies were required to measure and report energy intake at that meal(s) that energy density was manipulated for in order to be eligible (to permit quantification of the effect of the energy density manipulation independent of non-manipulated foods/meals).

Search process and article identification. The electronic databases PsycINFO, PubMed and SCOPUS (from date of inception) were searched during September-October 2020. For combinations of search terms used please refer to online supplementary material. The

reference lists of all eligible papers were searched and also contacted authors of included studies to inquire about any further eligible studies. Potential grey literature was addressed (to minimize publication bias) by searching the OSF preprint archive (includes 30 other preprint archives, including NutriXiv). Two authors independently screened and judged eligibility of all articles identified through electronic searches. One author completed the snowballing and grey literature searching approaches to identify any additional potentially eligible articles, eligibility was confirmed by a second author. Any discrepancies during eligibility assessments were discussed with a third author. Searches were re-run in October 2021 to identify any new eligible studies published between 2020-2021. None were identified.

Data extraction. The following study information was extracted by two authors independently (any discrepancies were resolved through discussion or by a third author); sampled participants (e.g. country, participant group sampled, summary information concerning sample demographics), energy density manipulation (e.g. number of foods/meals manipulated, energy density in each condition (kcal/g), nutritional composition (energy from protein, fat and carbohydrates) of energy density conditions, total number of kcals served in energy density conditions (if reported), design of study (e.g. within or between-subjects), energy intake measure (self-report vs. researcher measured, vs. mixed), whether any foods/meals had to be eaten in full (compulsory eating vs. ad libitum), number of days energy intake was measured for, energy intake information (e.g. energy intake under different conditions of energy density), whether body weight was measured before and after the different energy density conditions, and study factors related to risk of bias (see below). Authors were contacted and asked to provide details if statistical information required for analyses examining energy intake was missing (e.g. standard error not reported for energy intake under different energy density conditions).

175

176 ***Risk of bias ratings.*** A risk of bias checklist was created from existing study quality
 177 assessment tools and best practice recommendations for studying energy intake under
 178 experimental conditions (28-32), as existing bias tools (e.g. Cochrane) were not relevant to
 179 studies examining the effect of experimental manipulations of energy density on energy
 180 intake. The checklist included nine items for extraction and ‘yes’ was indicative of higher
 181 risk of bias; Was measured energy intake dependent on participant self-reporting? Did the
 182 study fail to use key participant exclusion criteria (e.g. use of medication affecting appetite)?
 183 Was any key methodological detail missing (e.g. limited information on procedures)? Was a
 184 non-random method of allocation to the different energy density condition used allocation (or
 185 was allocation method not described)? Were participants required to consume any study
 186 foods/meals in their entirety? Were demand characteristics not addressed in the study (e.g. no
 187 attempt to blind participants to study aims or measure whether differences between energy
 188 density foods were detectable)? Did the study have a small sample size (N<12 for within-
 189 subject designs)? Was the study pre-registered? Was there an absence of information on
 190 conflicts of interest or a reported relevant conflict of interest?

191

192 ***Analyses.*** The pre-registered analysis protocol and data are available online at:
 193 <https://osf.io/dj4yf/>. Deviations from planned analyses are reported in the online
 194 supplemental material.

195

196 ***Primary analyses***

197 ***Effect of energy density condition on daily energy intake.*** We first examined the effect of
 198 energy density condition on daily energy intake. Because a number of studies contributed
 199 multiple energy density comparisons (e.g. lower, medium, higher), we used multi-level meta-

analysis (33). Studies did not report on the correlation between daily energy intake under the different energy density conditions and we therefore imputed the size of this correlation based on similar studies (27) of daily energy intake ($r = 0.8$) and we conducted sensitivity analyses varying magnitude (0.6, 0.4) to examine consistency of results. Outliers were identified as effect sizes which the upper bound of their 95% confidence interval was lower than the lower bound of the meta-analysed pooled effect confidence interval of all effects or for which the lower bound of their 95% confidence interval was higher than the meta-analysed pooled effect confidence interval of all effects. Influential cases were identified as any effects with DFBETA values > 1 (indicative of a >1 change in the standard deviation of the estimated co-efficient after removal) (34) and we also conducted leave one out analyses. Egger's test for publication bias (35) and the trim and fill procedure for funnel plot asymmetry (36) were used. More detailed information is available in the online supplementary materials. If any outliers were identified, we examined the effect of removing them from the main primary meta-analysis. We also excluded them from our subsequent primary sub-group and meta-regression analyses on daily energy intake to minimize results being driven by large effects, but also examined if results were consistent when included. Dependent on outcome of interest, we either report meta-analyses as standardised mean difference (SMD), whereby SMDs of 0.2, 0.5 and 0.8 are small, moderate, and large statistical sized effects retrospectively (37) or report the mean weighted difference in energy intake (kcal) between energy density conditions to aid interpretation.

220

Sub-group and meta-regression analyses. To examine whether participant or study features moderated the effect of energy density condition on daily energy intake, we conducted a series of sub-group analyses and meta-regressions. For sub-group analyses, a-priori a minimum of $n=5$ effects per sub-group were required. We examined the effect of age group

(children vs. adult samples), sex (female vs. male vs. mixed), number of meals/foods energy density was manipulated for (all meals served vs. not), whether energy density manipulation altered macronutrient composition (% of kcals from protein vs. fat vs. carbohydrates altered between conditions vs. kept constant) and number of days energy intake was assessed for in the study (meta-regression). We also assessed risk of bias indicators for which there was sufficient variability between studies in a series of sub-groups analyses; use of random allocation (yes vs. no/not reported), energy intake measure (objective vs. reliant on self-report), whether demand characteristics were assessed (addressed vs. not addressed), conflicts of interest (statement included and no conflict vs. conflict reported or unclear). Because we found strong evidence that whether a study manipulated all foods/meals (vs did not) had a large impact on daily energy intake, we decided (unplanned) it was more appropriate to examine the relationship between absolute difference in energy density between energy density conditions (meta-regression, expressed as kcal/g) and daily energy intake for the two study types separately (see *Analyses by study type* below).

Linearity of relationship between manipulating energy density and energy intake. There was sufficient variability across studies to examine whether changes to energy density occurring at lower energy densities (e.g. reducing energy density of a relatively low energy dense food) produced smaller sized effects on daily energy intake as changes to energy density occurring at higher levels of energy density (e.g. reducing energy density of an energy dense food). We assessed this using both meta-regression (expressed as kcal/g of largest energy density condition) and in line with (23), a sub-group analysis that compared effects in which both energy density conditions < 1.75kcal/g vs. ≥ 1 energy density conditions exceeded 1.75kcal/g.

249 *Analyses by study type.* As the impact of manipulating energy density on daily energy intake
 250 differed substantially based on whether all meals/foods were manipulated vs. not, we
 251 conducted separate analyses for these two study types. For studies manipulating energy
 252 density of all food/meals, we examined whether absolute difference in energy density
 253 between energy density conditions was associated with daily energy intake (meta-regression),
 254 as well as repeating primary analyses examining number of days energy intake was assessed
 255 for and linearity of relationship between manipulating energy density and energy intake for
 256 sensitivity purposes. Next, we conducted analyses among studies that did not manipulate all
 257 foods/meals and also reported on energy intake during both manipulated meals and non-
 258 manipulated meals. We repeated the same analyses as above, as well as examining whether
 259 compulsory eating (participants required to consume one or meals eaten in full vs. ad libitum
 260 consumption) moderated results, as it was only common among this study type. We also
 261 examined the relationship between the total difference in kcals served between energy density
 262 conditions and daily energy intake using meta-regression. To further aid interpretation of
 263 ‘compensation’ effects after consuming lower vs. higher energy density foods/meals, we
 264 conducted separate meta-analyses on energy intake from manipulated foods and non-
 265 manipulated foods separately. In some instances, the manipulated meal/food was ‘fixed’ (i.e.
 266 it was compulsory for participants to eat the meal in full) and this equates to a standard error
 267 of 0, entered as 0.1 in meta-analysis. In sensitivity analyses we imputed these values as the
 268 average SE (expressed as a proportion of mean energy intake) taken from energy density
 269 manipulated meals that were not ‘fixed’, in order to ensure results were consistent.

270

271 *Body weight.* A small number of studies (n=5) reported data on change in body weight before
 272 vs. after lower and higher energy density conditions. Standard deviations were not reported
 273 and therefore we imputed this (based on average SD as a % of M weight change) from (27)

and used sensitivity analyses to examine consistency when size of SD was larger and smaller. We used generic variance inverse meta-analysis to pool change in body weight (kg). In instances where a study had multiple energy density conditions, to maximise statistical power, a-priori we only included the energy density condition contrast with the largest difference in energy density served.

Results

Summary of included studies. A total of 31 eligible studies were included in the review and meta-analysis (see Table 1). Figure 1 outlines the study selection process. Twenty-seven studies sampled adults and the remaining four studies sampled children. Most included studies were conducted in the US (n=19) and the remainder where in Europe (n=9) or Singapore (n=3). All studies used within-subjects designs to examine the effect of manipulating energy density. Energy intake was assessed between 1 and 14 days in studies and the most common study length was 1 day (n=18). Sample sizes of studies ranged from N=6 to N=95. Twenty-three studies manipulated either a single meal or a limited number of meals/food items (as opposed to all meals/foods) and the remaining (n=8) studies manipulated energy density of all food served to participants. It was most common for studies to compare the effect of two energy density conditions on daily energy intake (n=22), eight studies compared three energy density conditions and a single study had five energy density conditions. For n=9 studies, the macronutrient content (i.e. %kcal from protein, fat, carbohydrates) of food served in the different energy density conditions was held constant. For n=14 studies, energy density was manipulated by altering macronutrient content (e.g. reducing %kcal from fat) and in one study macronutrient content was held constant across two of the density conditions and differed between two energy density conditions. In n=7 studies, macronutrient information was not reported or unclear. The lowest energy density

condition in a study was 0.11–0.13kcal/g (the higher energy density condition from this study was 0.49-0.5kcal/g). The highest energy density condition in a study was 5.47kcal/g (the low energy density condition in the study was 2.53kcal/g), this study also had the largest absolute difference (2.94kcal/g). See Table 1 for individual study information.

Risk of bias indicators. Of the thirty-one included studies, a sizeable minority (n=14) used participant self-report to quantify (in part or full) daily energy intake. Only n=2/31 studies did not report use of key participant exclusion criteria, n=2/31 studies did not report key methodological information, n=10/31 used either non-random allocation to energy density condition order or did not report on allocation method, n=15/31 required participants to eat at least one meal or test food in full as part of the procedure, in n=10/31 studies demand characteristics were not addressed (e.g. measurement of participant awareness of different energy density conditions). A minority of studies (n=5/31) had a small sample size ($N < 8$), n=25/31 studies were not pre-registered and n=21/31 studies did not have a conflicts of interest statement or reported a relevant conflict. See supplementary online materials for individual study risk of bias information.

Primary analyses

Effects on daily energy intake. The multi-level meta-analysis (90 effect sizes from 31 studies) indicated that there was a large effect of energy density condition on daily energy intake ($SMD = -1.002$ [95% CI: -0.745 to -1.266], $Z = 7.54$, $p < .001$, $I^2 = 92.1\%$), whereby serving lower energy dense foods was associated with lower daily energy intake. See Figure 1. Results remained significant in sensitivity analyses varying the size of within-subjects correlation for daily energy intake (see online supplementary material). Egger's test was significant ($Z = -10.82$, $p < .001$), indicating possible publication bias, although Trim and Fill

on a single level model identified 0 studies to be filled. See online supplementary materials for funnel plot. No DFBETAs were greater than >1 , and leave-one-out analysis did not substantially influence the models ($ps < .001$). There were 28 effect sizes with confidence intervals which did not overlap the pooled analyses (outliers) and removing them from the analyses slightly reduced the pooled effect, but also the heterogeneity ($SMD = -0.872$ [95% CI: -1.001 to -0.742], $Z = 13.21$, $p < .001$, $I^2 = 60.6\%$).

Sub-group and meta-regression analyses on daily energy intake (outliers removed). Sub-group analyses comparing adult vs. child and male vs. female samples were non-significant (see online supplementary materials). Moderation analysis comparing effects for which energy density was varied by manipulating nutritional composition (28 effect sizes) vs. kept constant (21 effect sizes) was non-significant ($X^2(1) = 1.00$ $p = .318$), whereby energy density manipulations altering composition ($SMD = -0.952$ [95% CI: -0.694 to -1.209]) produced very similar effects on daily energy intake as those not altering composition ($SMD = -0.859$ [95% CI: -0.649 to -1.068]). Moderation analysis for number of meals/foods energy density was manipulated was statistically significant ($X^2(1) = 18.11$, $p < .001$). Effects in which energy density of all foods served was manipulated (9 effects, $SMD = -1.871$ [95% CI: -1.313 to -2.430]) were associated with a larger impact on daily energy intake than effects in which not all food served was manipulated (53 effects, $SMD = -0.796$ [95% CI: -0.682 to -0.910]). Meta-regression of the number of days energy intake indicated a negative but non-significant association with daily energy intake ($b = -0.039$ [95% CI: -0.080 to 0.001], $p = .060$), whereby smaller effects on daily energy intake were associated with studies measuring energy intake for longer. All analysis results remained consistent with the inclusion of outliers from the primary model, although the meta-regression on length of study became significant ($p = .018$).

349

350 *Linearity of relationship between manipulating energy density and daily energy intake*

351 *(outliers removed)*. The kcal/g of the highest energy density condition in each energy density
352 comparison was not a significant predictor of effect on daily energy intake ($b = -0.020$ [95%
353 CI: -0.129 to 0.089]) and effects for which both energy density conditions were < 1.75 kcal/g
354 vs. ≥ 1.75 kcal/g in at least one condition produced similar sized results on daily energy intake
355 ($p = .160$).

356

357 *Risk of bias indicators (outliers removed)*. Analyses examining whether effects of energy
358 density on daily energy intake were dependent on whether studies used self-report vs.
359 measured energy intake, addressed demand characteristics and conflicts of interest vs. did not
360 were all non-significant. Studies which either did not use or failed to report on random
361 allocation to energy density conditions tended to produce larger effects on daily energy intake
362 than studies which did report use of random allocation ($p = .045$), but both types of study
363 were individually significant. See online supplementary materials for results in full.

364

365 *Analyses limited to studies manipulating energy density of all foods/meals*. There was a large
366 effect of energy density condition on daily energy intake (kcal difference between higher and
367 lower energy density conditions = -855.85 [95% CI: -616.18 to -1095.52], $Z = 7.00$, $p < .001$,
368 $I^2 = 97.4\%$). See Figure 3. Removal of outliers (5 effects) slightly reduced the kcal difference
369 (-709.01 [95% CI: -602.04 to -815.97], $Z = 12.99$, $p < .001$, $I^2 = 85.4\%$). There was a
370 significant association between difference in energy density between meals and differences in
371 kcals consumed between conditions/meals ($b = -1510.70$ [95% CI: -1236.08 to -1785.33], $p <$
372 $.001$) and with outliers removed the association was smaller but remained significant ($b = -$
373 309.31 [95% CI: -115.91 to -502.71], $Z = 3.13$, $p = .002$). Length of study (number of days)

was not significantly related to effects on daily energy intake and analyses examining non-linearity were also non-significant, as in the primary analyses (see online supplementary materials).

Analyses limited to studies not manipulating energy density of all foods/meals. There was a large effect of energy density condition on daily energy intake (kcal diff = -237.84 [95% CI: -148.13 to -327.54], $Z = 5.20$, $p < .001$, $I^2 = 95.9$). See Figure 4. Twelve outliers were identified, and removal of these effects slightly reduced the kcal difference, but also the heterogeneity (-208.17 [95% CI: -160.00 to -256.37], $Z = 8.47$, $p < .001$, $I^2 = 75.5\%$). There was a significant association between difference in energy density between conditions and the difference in daily energy intake between conditions ($b = -331.86$ [95% CI: -234.62 to -429.13], $Z = 6.69$, $p < .001$) and with outliers excluded, the effect remained significant but somewhat smaller ($b = -104.50$ [95% CI: -12.03 to 196.98], $Z = 2.21$, $p = .027$). Length of study was not significantly related to effects on daily energy intake. For analyses examining potential non-linearity, results were largely consistent with the primary analyses. See online supplementary materials.

Difference in total kcals served and daily energy intake. Among studies that did not manipulate energy density of all foods/meals, there was a significant association between differences in kcals served between energy density conditions and difference in daily energy intake ($b = -0.774$ [95% CI: -0.644 to -0.905], $Z = 11.64$, $p < .001$), whereby a 100kcal difference in energy served (due to energy density manipulation) was predictive of a 77kcal difference in daily energy intake. Results remained significant with removal of outliers. See online supplementary materials.

Energy intake during manipulated energy density meals vs. later in the day. For studies that provided complete data (i.e. Ms and SDs) on both energy intake from energy density manipulated foods/meals and energy intake from subsequent non-manipulated foods/meals (16 effects from 7 studies), the difference in kcals consumed from manipulated meals between higher and lower energy density conditions was - 330.78kcals ([95% CI: -224.27 to - 437.29), $Z = 6.09$, $p < .001$, $I^2 = 100\%$) and similar in sensitivity analyses that varied size of SD for manipulated meals that required compulsory eating [-326.40 ([95% CI:-222.53 to - 431.31]. There was a small increase in kcals consumed after consuming lower vs. high energy dense food, but this increase was not statistically significant (kcals = 35.08 [95% CI: - 28.32 to 98.48], $Z = 0.28$, $p = .278$, $I^2 = 95.15$).

Body weight. Pooled across the five studies that provided data on weight change, weight loss tended to be greater in lower compared to higher energy dense conditions, but this difference was not statistically significant, kg change = -0.69 [95% CI: -1.43 to 0.04]. See Figure 5. Results were similar in sensitivity analyses. See online supplementary materials.

Discussion

Serving lower energy dense (vs. higher energy density) foods significantly reduced daily energy intake and this effect was statistically large. Studies with the most pronounced differences in energy density produced the largest changes to daily energy intake (i.e. a dose-dependent response) and studies which manipulated the energy density of all foods served produced larger effects on daily energy intake than studies that did not manipulate all foods. The impact energy density had on daily energy intake was similar among males vs. females and in adults vs. children. Studies tended to manipulate energy density by either altering macronutrient content (e.g. by reducing %kcals from fat) or by holding macronutrient content

(e.g. by increasing water content) and both manipulations produced similar sized effects on daily energy intake. Due to the available data, we were unable to examine whether distinct macronutrient manipulations (e.g. replacing fat with protein as opposed to carbohydrate) affected daily energy intake differentially. However, over and above absolute changes to energy density, any nutrient specific effects may be relatively subtle (14). Due to the satiety-providing effects of protein, decreasing energy by increasing protein content may have a more pronounced effect on daily energy intake (38, 39), but evidence is mixed (40, 41).

We found consistent evidence that the relationship between energy density and daily energy intake was strong and linear. Consistent with this, among studies that manipulated energy density for some but not all food served, analyses suggested that for every 100 fewer kcals of food served to participants (due to reduced energy density), daily energy intake was reduced by approximately 77kcals. Furthermore, in these studies participants consumed approximately 326kcals fewer during the lower (vs. higher) energy dense meals but increased their later ad-libitum energy intake (from non-manipulated foods) by a non-significant 35kcals. Therefore, unlike manipulation of food product portion size for which more substantial compensation appears to occur (37), there is minimal evidence of energy intake compensation in response to manipulations of food energy density. An implication of the present findings is that public health policies which reduce energy density of food being sold (e.g. through voluntary industry reformulation or mandatory action) are likely to be more effective in reducing daily energy intake than policies which target portion size alone.

A limitation of included studies was their relatively short duration (between 1 and 14 days). We found some inconsistent evidence that the length of time energy intake was measured for moderated findings, whereby effects of energy density on daily energy intake were smaller among studies with longer duration in our main analysis. However, the statistical significance of this effect was dependent on the exclusion of outliers from analyses

and this association was not observed when studies that manipulated all foods (vs. did not) were analysed separately. This finding may indicate that over time consumers learn about the energy density of food served and adapt their food intake, but this adaptation is only partial. A small sub-set of studies examined change in body weight and although after being served lower vs. energy dense foods participants tended to lose more weight (1.4kg difference), this difference was not statistically significant. Although previous studies that have directed participants to reduce energy density through dietary advice provide evidence for significant changes to body weight (1, 25), the effect of reformulating the energy density of foods on body weight therefore remains less clear. Future research will therefore be needed examining the effect that manipulations of energy density have on body weight in order to understand whether mass reformulation of the energy density of food products is likely to benefit population level obesity.

Contrary to suggestions that humans may be more sensitive to changes in energy density to less energy dense foods (23, 24), we found no evidence that the impact of reducing energy density of food served was non-linear in nature; studies comparing two low energy density conditions (e.g. 1.1kcal/g vs. 0.8kcal/g) produced similar sized effects to studies comparing more energy dense foods (e.g. 2.6kcal/g vs. 2.3kcal/g). However, the majority of studies examined lower food energy densities, as opposed to ‘highly’ energy dense foods (i.e. ≥ 4 kcal/g) and it may be the case that differences would be observed for the latter. Further research directly addressing this question will now be important because public health approaches would presumably target reformulation of highly energy dense foods, as opposed to food products that are already relatively low in energy density.

There are strengths and limitations to the present research. We followed best practice guidelines for systematic review methodology and attempted to identify eligible published and unpublished articles using a combination of supplementary methods including grey

474 literature searching and contact authors of eligible articles. Methodological quality of
475 included studies was variable, but studies tended to be well-reported, few were of very small
476 sample size and most study designs addressed demand characteristics. We assessed whether a
477 range of potential risk of bias indicators affected results in sub-group analyses and found little
478 convincing evidence that risk of bias indicators predicted study outcomes. As discussed,
479 study durations were relatively short and the artificial nature of the laboratory settings used in
480 most studies increases confidence in experimental control but at the expense of ecological
481 validity (32, 42). It may be the case that alterations to food energy density would be
482 associated with greater compensation outside of the laboratory when concerns about social
483 desirability are reduced and/or a wider selection of food is available (32, 43, 44), which
484 would result in smaller effects on daily energy intake and body weight. It is also important to
485 note that a number of included studies allowed participants to consume foods and meals
486 outside of the laboratory and later self-report this intake, and in these studies the effect of
487 manipulating energy density of food served in the laboratory on daily energy intake was still
488 sizeable. As noted, the relatively short duration of studies is a limitation and it may be that
489 over longer time periods, the post-ingestive consequences of lower energy density foods
490 would result in dietary learning. However, it is not clear how long foods would need to be
491 consumed for, as in one study repeated daily exposures of higher vs. lower energy density
492 versions of the same product for 5 days produced no evidence of dietary learning (45). It
493 should also be noted that we detected evidence of funnel plot asymmetry which may be
494 indicative of publication bias. However, this appears to have been largely caused by there
495 being a number of studies that had particularly large manipulations to energy density which
496 would be expected to cause large decreases in daily energy intake and therefore contribute to
497 asymmetry. A final limitation was that we were only able to examine a small number of
498 participant characteristics in moderation analyses (sex, age) and it may be the case that there

are other characteristics (e.g. BMI, socioeconomic status) or participant traits (e.g. satiety responsiveness) that moderate the effect reducing energy density has on daily energy intake.

Conclusions. Experimental studies indicate that decreasing energy density of food products has a strong and largely linear effect on daily energy intake, although effects on body weight are less clear and warrant further study. Reformulation of the energy density of food products may be an effective public health approach to reducing population level energy intake.

Studies included in review. Refs 15-20, 46-70.

Acknowledgements: N/A.

Funding: No external funding.

Conflicts of Interest: All authors report no conflicts of interest. ER has previously received funding from the American Beverage Association and Unilever for projects unrelated to the present research.

Author Contributions: ER designed the research, conducted the research, had primary responsible for the final content and wrote the paper. AJ designed the research, conducted the research, analysed data and wrote the paper. IML, MK and ZP conducted the research.

Figures

Figure 1. Systematic review study search and eligibility flowchart

Figure 2. Forest plot of all studies included in primary meta-analysis

Figure 3. Forest plot for analyses limited to studies which manipulated energy density for all foods served

Figure 4. Forest plot for analyses limited to studies which manipulated energy density of some but not all foods served

Figure 5. Body weight meta-analysis forest plot

524

References

- 525 1. Rolls BJ. Dietary energy density: Applying behavioural science to weight
526 management. *Nutr Bull.* 2017;42(3):246-53.
- 527 2. Hardman CA, Ferriday D, Kyle L, Rogers PJ, Brunstrom JM. So Many Brands and
528 Varieties to Choose from: Does This Compromise the Control of Food Intake in Humans?
529 *PLOS ONE.* 2015;10(4):e0125869.
- 530 3. Drewnowski A, Darmon N. The economics of obesity: dietary energy density and
531 energy cost. *Am J Clin Nutr.* 2005;82(1 Suppl):265s-73s.
- 532 4. Drewnowski A. Obesity and the food environment: dietary energy density and diet
533 costs. *Am J Prev Med.* 2004;27(3 Suppl):154-62.
- 534 5. Prentice AM, Jebb SA. Fast foods, energy density and obesity: a possible mechanistic
535 link. *Obesity Reviews.* 2003;4(4):187-94.
- 536 6. Rouhani MH, Haghighatdoost F, Surkan PJ, Azadbakht L. Associations between
537 dietary energy density and obesity: A systematic review and meta-analysis of observational
538 studies. *Nutrition.* 2016;32(10):1037-47.
- 539 7. Stelmach-Mardas M, Rodacki T, Dobrowolska-Iwanek J, Brzozowska A, Walkowiak
540 J, Wojtanowska-Krosniak A, et al. Link between Food Energy Density and Body Weight
541 Changes in Obese Adults. *Nutrients.* 2016;8(4):229.
- 542 8. Ledikwe JH, Blanck HM, Kettel Khan L, Serdula MK, Seymour JD, Tohill BC, et al.
543 Dietary energy density is associated with energy intake and weight status in US adults. *Am J*
544 *Clin Nutr.* 2006;83(6):1362-8.
- 545 9. Karl JP, Roberts SB. Energy Density, Energy Intake, and Body Weight Regulation in
546 Adults. *Advances in Nutrition.* 2014;5(6):835-50.
- 547 10. Rolls BJ. The relationship between dietary energy density and energy intake.
548 *Physiology & Behavior.* 2009;97(5):609-15.
- 549 11. Mendoza JA, Drewnowski A, Christakis DA. Dietary Energy Density Is Associated
550 With Obesity and the Metabolic Syndrome in U.S. Adults. *Diabetes Care.* 2007;30(4):974-9.
- 551 12. Du H, van der A DL, Ginder V, Jebb SA, Forouhi NG, Wareham NJ, et al. Dietary
552 Energy Density in Relation to Subsequent Changes of Weight and Waist Circumference in
553 European Men and Women. *PLOS ONE.* 2009;4(4):e5339.
- 554 13. Gomez-Bruton A, Arenaza L, Medrano M, Mora-Gonzalez J, Cadenas-Sanchez C,
555 Migueles JH, et al. Associations of dietary energy density with body composition and
556 cardiometabolic risk in children with overweight and obesity: role of energy density
557 calculations, under-reporting energy intake and physical activity. *British Journal of Nutrition.*
558 2019;121(9):1057-68.
- 559 14. Stubbs J, Ferres S, Horgan G. Energy density of foods: effects on energy intake.
560 *Critical reviews in food science and nutrition.* 2000;40(6):481-515.
- 561 15. Stubbs R, Johnstone A, Harbron C, Reid C. Covert manipulation of energy density of
562 high carbohydrate diets in 'pseudo free-living' humans. *International journal of obesity.*
563 1998;22(9):885-92.
- 564 16. Stubbs R, Johnstone A, O'Reilly L, Barton K, Reid C. The effect of covertly
565 manipulating the energy density of mixed diets on ad libitum food intake in 'pseudo free-
566 living' humans. *International Journal of Obesity.* 1998;22(10):980-7.
- 567 17. Stubbs R, Ritz P, Coward WA, Prentice AM. Covert manipulation of the ratio of
568 dietary fat to carbohydrate and energy density: effect on food intake and energy balance in
569 free-living men eating ad libitum. *Am J Clin Nutr.* 1995;62(2):330-7.
- 570 18. Stubbs RJ, Harbron CG, Murgatroyd PR, Prentice AM. Covert manipulation of
571 dietary fat and energy density: effect on substrate flux and food intake in men eating ad
572 libitum. *Am J Clin Nutr.* 1995;62(2):316-29.

- 573 19. Williams RA, Roe LS, Rolls BJ. Comparison of three methods to reduce energy
574 density. Effects on daily energy intake. *Appetite*. 2013;66:75-83.
- 575 20. Leahy KE, Birch LL, Rolls BJ. Reducing the energy density of multiple meals
576 decreases the energy intake of preschool-age children. *Am J Clin Nutr*. 2008;88(6):1459-68.
- 577 21. Moosavian SP, Haghighatdoost F. Dietary energy density and appetite: A systematic
578 review and meta-analysis of clinical trials. *Nutrition*. 2020;69:110551.
- 579 22. Rouhani MH, Surkan PJ, Azadbakht L. The effect of preload/meal energy density on
580 energy intake in a subsequent meal: A systematic review and meta-analysis. *Eat Behav*.
581 2017;26:6-15.
- 582 23. Brunstrom JM, Drake ACL, Forde CG, Rogers PJ. Undervalued and ignored: Are
583 humans poorly adapted to energy-dense foods? *Appetite*. 2018;120:589-95.
- 584 24. Brunstrom JM, Cheon BK. Do humans still forage in an obesogenic environment?
585 Mechanisms and implications for weight maintenance. *Physiology & Behavior*.
586 2018;193:261-7.
- 587 25. Ello-Martin JA, Roe LS, Ledikwe JH, Beach AM, Rolls BJ. Dietary energy density in
588 the treatment of obesity: a year-long trial comparing 2 weight-loss diets. *Am J Clin Nutr*.
589 2007;85(6):1465-77.
- 590 26. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for
591 systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine*.
592 2009;6(7):e1000097.
- 593 27. Robinson E, McFarland-Lesser I, Patel Z, Jones A. Downsizing food: A systematic
594 review and meta-analysis examining the effect of reducing served food portion sizes on daily
595 energy intake and body weight. *medRxiv*. 2021:2021.09.22.21263961.
- 596 28. Blundell J, De Graaf C, Hulshof T, Jebb S, Livingstone B, Lluch A, et al. Appetite
597 control: methodological aspects of the evaluation of foods. *Obesity reviews*. 2010;11(3):251-
598 70.
- 599 29. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The
600 Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *Bmj*. 2011;343.
- 601 30. Robinson E, Bevelander K, Field M, Jones A. Methodological and reporting quality in
602 laboratory studies of human eating behavior. *Appetite*. 2018;125:486-91.
- 603 31. Kersbergen I, Whitelock V, Haynes A, Schroor M, Robinson E. Hypothesis
604 awareness as a demand characteristic in laboratory-based eating behaviour research: An
605 experimental study. *Appetite*. 2019;141:104318.
- 606 32. Livingstone MBE, Robson PJ, Welch RW, Burns AA, Burrows MS, McCormack C.
607 Methodological issues in the assessment of satiety. *Näringsforskning*. 2000;44(1):98-103.
- 608 33. Pastor DA, Lazowski RA. On the Multilevel Nature of Meta-Analysis: A Tutorial,
609 Comparison of Software Programs, and Discussion of Analytic Choices. *Multivariate*
610 *Behavioral Research*. 2018;53(1):74-89.
- 611 34. Viechtbauer W, Cheung MWL. Outlier and influence diagnostics for meta-analysis.
612 *Research synthesis methods*. 2010;1(2):112-25.
- 613 35. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a
614 simple, graphical test. *Bmj*. 1997;315(7109):629-34.
- 615 36. Duval S, Tweedie R. Trim and fill: a simple funnel plot-based method of testing and
616 adjusting for publication bias in meta-analysis. *Biometrics*. 2000;56(2):455-63.
- 617 37. Cohen J. A power primer. *Psychological bulletin*. 1992;112(1):155.
- 618 38. Martens EA, Lemmens SG, Westerterp-Plantenga MS. Protein leverage affects energy
619 intake of high-protein diets in humans. *Am J Clin Nutr*. 2012;97(1):86-93.
- 620 39. Gosby AK, Conigrave AD, Lau NS, Iglesias MA, Hall RM, Jebb SA, et al. Testing
621 Protein Leverage in Lean Humans: A Randomised Controlled Experimental Study. *PLOS*
622 *ONE*. 2011;6(10):e25929.

- 623 40. Blatt AD, Roe LS, Rolls BJ. Increasing the protein content of meals and its effect on
624 daily energy intake. *Journal of the American Dietetic Association*. 2011;111(2):290-4.
- 625 41. Raben A, Agerholm-Larsen L, Flint A, Holst JJ, Astrup A. Meals with similar energy
626 densities but rich in protein, fat, carbohydrate, or alcohol have different effects on energy
627 expenditure and substrate metabolism but not on appetite and energy intake. *Am J Clin Nutr*.
628 2003;77(1):91-100.
- 629 42. Meiselman HL. Methodology and theory in human eating research. *Appetite*.
630 1992;19(1):49-55.
- 631 43. Robinson E, Hardman CA, Halford JC, Jones A. Eating under observation: a
632 systematic review and meta-analysis of the effect that heightened awareness of observation
633 has on laboratory measured energy intake. *Am J Clin Nutr*. 2015;102(2):324-37.
- 634 44. Robinson E, Kersbergen I, Brunstrom JM, Field M. I'm watching you. Awareness that
635 food consumption is being monitored is a demand characteristic in eating-behaviour
636 experiments. *Appetite*. 2014;83:19-25.
- 637 45. Hogenkamp PS, Brunstrom JM, Stafleu A, Mars M, de Graaf C. Expected satiation
638 after repeated consumption of low- or high-energy-dense soup. *British Journal of Nutrition*.
639 2012;108(1):182-90.
- 640 46. Bell EA, Castellanos VH, Pelkman CL, Thorwart ML, Rolls BJ. Energy density of
641 foods affects energy intake in normal-weight women. *Am J Clin Nutr*. 1998;67(3):412-20.
- 642 47. Bell EA, Rolls BJ. Energy density of foods affects energy intake across multiple
643 levels of fat content in lean and obese women. *Am J Clin Nutr*. 2001;73(6):1010-8.
- 644 48. Blatt AD, Roe LS, Rolls BJ. Hidden vegetables: an effective strategy to reduce energy
645 intake and increase vegetable intake in adults. *Am J Clin Nutr*. 2011;93(4):756-63.
- 646 49. Blatt AD, Williams RA, Roe LS, Rolls BJ. Effects of energy content and energy
647 density of pre-portioned entrées on energy intake. *Obesity (Silver Spring)*. 2012;20(10):2010-
648 8.
- 649 50. Buckland NJ, Camidge D, Croden F, Lavin JH, Stubbs RJ, Hetherington MM, et al. A
650 Low Energy-Dense Diet in the Context of a Weight-Management Program Affects Appetite
651 Control in Overweight and Obese Women. *J Nutr*. 2018;148(5):798-806.
- 652 51. Caputo FA, Mattes R. Human dietary responses to covert manipulations of energy,
653 fat, and carbohydrate in a midday meal. *Am J Clin Nutr*. 1992;36-43.
- 654 52. Devitt AA, Mattes RD. Effects of food unit size and energy density on intake in
655 humans. *Appetite*. 2004;42(2):213-20.
- 656 53. Duncan KH, Bacon JA, Weinsier RL. The effects of high and low energy density
657 diets on satiety, energy intake, and eating time of obese and nonobese subjects. *Am J Clin*
658 *Nutr*. 1983;37(5):763-7.
- 659 54. Foltin RW, Fischman MW, Moran TH, Rolls BJ, Kelly TH. Caloric compensation for
660 lunches varying in fat and carbohydrate content by humans in a residential laboratory. *Am J*
661 *Clin Nutr*. 1990;52(6):969-80.
- 662 55. Gray RW, French SJ, Robinson TM, Yeomans MR. Dissociation of the effects of
663 preload volume and energy content on subjective appetite and food intake. *Physiology &*
664 *Behavior*. 2002;76(1):57-64.
- 665 56. Kral TVE, Roe L, Rolls BJ. Does nutrition information about the energy density of
666 meals affect food intake in normal-weight women? *Appetite*. 2002;39.
- 667 57. Kral TV, Roe LS, Rolls BJ. Combined effects of energy density and portion size on
668 energy intake in women. *Am J Clin Nutr*. 2004;79(6):962-8.
- 669 58. Kral TVE, Moore RH, Chittams J, O'Malley L, Jones E, Quinn RJ, et al. Caloric
670 compensation and appetite control in children of different weight status and predisposition to
671 obesity. *Appetite*. 2020;151.

59. Mazlan N, Horgan G, Stubbs RJ. Energy density and weight of food effect short-term caloric compensation in men. *Physiology & Behavior*. 2006;87(4):679-86.
60. McCrickerd K, Lim CM, Leong C, Chia-Ming E, Forde C. Texture-Based Differences in Eating Rate Reduce the Impact of Increased Energy Density and Large Portions on Meal Size in Adults. *journal of nutrition Ingestive Behavior and Neurosciences*. 2017.
61. Miller DL. Effect of fat-free potato chips with and without nutrition labels on fat and energy intakes. 1998.
62. Pritchard SJ, Davidson I, Jones J, Bannerman E. A randomised trial of the impact of energy density and texture of a meal on food and energy intake, satiation, satiety, appetite and palatability responses in healthy adults. *Clin Nutr*. 2014;33(5):768-75.
63. Rolls BJ, Roe LS, Meengs JS. Reductions in portion size and energy density of foods are additive and lead to sustained decreases in energy intake. *Am J Clin Nutr*. 2006;83(1):11-7.
64. Shide DJ, Rolls BJ. Information about the fat content of preloads influences energy intake in healthy women. *J Am Diet Assoc*. 1995;95(9):993-8.
65. Silver HJ, Dietrich MS, Castellanos VH. Increased energy density of the home-delivered lunch meal improves 24-hour nutrient intakes in older adults. *J Am Diet Assoc*. 2008;108(12):2084-9.
66. Smethers AD, Roe LS, Sanchez CE, Zuraikat FM, Keller KL, Rolls BJ. Both increases and decreases in energy density lead to sustained changes in preschool children's energy intake over 5 days. *Physiology & Behavior*. 2019;204:210-8.
67. Spill MK, Birch LL, Roe LS, Rolls BJ. Hiding vegetables to reduce energy density: an effective strategy to increase children's vegetable intake and reduce energy intake. *Am J Clin Nutr*. 2011;94(3):735-41.
68. Tey SL, Chia-Ming E, Forde C. Impact of dose-response calorie reduction or supplementation of a covertly manipulated lunchtime meal on energy compensation. *Physiol Behav*. 2016:15-21.
69. Tey SL, Salleh N, Henry CJ, Forde CG. Effects of Consuming Preloads with Different Energy Density and Taste Quality on Energy Intake and Postprandial Blood Glucose. *Nutrients*. 2018;10(2).
70. Westerterp-Plantenga MS, Wijckmans-Duijsens NE, ten Hoor F, Weststrate JA. Effect of replacement of fat by nonabsorbable fat (sucrose polyester) in meals or snacks as a function of dietary restraint. *Physiol Behav*. 1997;61(6):939-47.

Tables

Table 1. Summary information on included studies

Study	Country and sample	Sample characteristics	Child or adult sample	Study design information	Number of participants	Meals/foods ED manipulated for all meals/foods?	ED conditions	ED manipulation
Bell, 1998 (46)	US University students and local community	BMI: M = 23 Gender: 18F	Adult	Within-subjects Researcher measured EI EI measured for 2 days	N = 18	No: Lunch entrée, dinner entrée, evening snack only	<u>Lower ED</u> 0.8 kcal/g <u>Medium ED</u> 1.1 kcal/g <u>Higher ED</u> 1.3 kcal/g	Nutritional composition matched
Bell, 2001 (47)	US University students and local community	BMI: M = 28 Gender: 36F	Adult	Within-subjects Researcher measured EI EI measured for 1 day	N = 36	No: Breakfast entrée, lunch entrée, evening snack only	<u>Lower ED</u> 1.26-1.31 kcal/g <u>Higher ED</u> 1.79-1.81 kcal/g	Nutritional composition not matched
Blatt, 2011 (48)	US University students and local community	BMI: M = 24 Gender: 20M, 21F	Adult	Within-subjects Researcher measured EI EI measured for 1 day	N = 41	No: Breakfast entrée, lunch entrees, dinner entrée only	<u>Lower ED</u> 1.99 kcal/g <u>Medium ED</u> 2.25 kcal/ <u>Higher ED</u> 2.64 kcal/g	Nutritional composition matched
Blatt, 2012 (49)	US University students and local community	BMI: M = 24 Gender: 28M, 40F	Adult	Within-subjects Researcher measured EI EI measured for 1 day	N = 68	No: Breakfast entrée, lunch entrees, dinner entrée only	<u>Lower ED</u> 1 kcal/g <u>Higher ED</u> 1.6 kcal/g	Nutritional composition matched
Buckland,	UK	BMI:	Adult	Within-subjects	N = 77	Yes	<u>Lower ED</u>	Nutritional composition not

2018 (50)	Participants in weight loss trial	M = 34 Gender: 77F		Self-reported and researcher measured EI EI measured for 1 day			0.8 kcal/g <u>Higher ED</u> 2.5 kcal/g	matched
Caputo, 1992 (51)	US Healthy/normal weight sample	BMI: M = N/R Gender: 8M, 8F	Adult	Within-subjects Self-reported and researcher measured EI EI measured for 5 days	N = 8	No: Lunch only	<u>Lower ED</u> 0.45 kcal/g 1.19 kcal/g <u>Higher ED</u> 1.2 kcal/g 3.9 kcal/g	Nutritional composition not matched
Devitt, 2004 (52)	US University students and local community	BMI: M = 25 Gender: 11M, 9F	Adult	Within-subjects Researcher measured EI EI measured for 4 days	N = 20	Yes	<u>Lower ED</u> 1.49 kcal/g <u>Higher ED</u> 2.6 kcal/g	Nutritional composition not matched
Duncan, 1983 (53)	US Sample N/R	BMI: M = N/R Gender: 10M, 10F	Adult	Within-subjects Researcher measured EI EI measured for 5 days	N = 20	Yes	<u>Lower ED</u> 0.7 kcal/g <u>Higher ED</u> 1.5 kcal/g	Nutritional composition N/R
Foltin, 1983 (54)	US Sample N/R	BMI: M = N/R Gender: 6M	Adult	Within-subjects Self-reported and researcher measured EI EI measured for 3 days	N = 6	No: Lunch only	<u>Lower ED</u> 0.4 kcal/g <u>Higher ED</u> 0.8-0.9 kcal/g	Nutritional composition not matched
Gray, 2002 (55)	England University students and staff	BMI: M = 23 Gender: 20M	Adult	Within-subjects Self-reported and researcher measured EI	N = 20	No: Starter (preload) at lunch only	<u>Lower ED</u> 0.33 kcal/g <u>Higher ED</u> 1 kcal/g	Nutritional composition N/R

				EI measured for 1 day				
Kral, 2002 (56)	US University students and local community	BMI: M = 22 Gender: 40F	Adult	Within-subjects Researcher measured EI EI measured for 1 day	N = 40	No: Breakfast entrée, lunch entrée and dinner entree only	<u>Lower ED</u> 1.25 kcal/g <u>Medium ED</u> 1.5 kcal/g <u>Higher ED</u> 1.75 kcal/g	Nutritional composition matched
Kral, 2004 (57)	US University students and local community	BMI: M = 24 Gender: 39F	Adult	Within-subjects Researcher measured EI EI measured for 1 day	N = 39	No: Lunch entrée only	<u>Lower ED</u> 1.25 kcal/g <u>Higher ED</u> 1.75 kcal/g	Nutritional composition matched
Kral, 2020 (58)	US Local community	BMI percentiles ranging from 43-95 Gender: 29M, 46F	Children	Within-subjects Self-reported and researcher measured EI EI measured for 1 day	N = 75	No: Breakfast only	<u>Lower ED</u> 1 kcal/g <u>Higher ED</u> 1.6 kcal/g	Nutritional composition matched
Leahy, 2008 (20)	US Pre-school children at university day care centre	BMI percentile: M=71 Gender: 10M, 16F	Children	Within-subjects Researcher measured EI EI measured for 2 day	N = 26	No: Breakfast, lunch, afternoon snack only	<u>Lower ED</u> 0.91 kcal/g <u>Higher ED</u> 1.23 kcal/g	Nutritional composition N/R
Mazlan, 2006 (59)	Scotland Local community	BMI: M = 23 Gender: 16M	Adult	Within-subjects Self-reported and researcher measured EI EI measured for 1 day	N = 16	No: Breakfast, dessert snack only	<u>Lower ED</u> 0.96 kcal/g <u>Higher ED</u> 1.94 kcal/g	Nutritional composition matched
McCrickerd, 2017 (60)	Singapore	BMI: M = 22	Adult	Within-subjects	N = 62	No: Breakfast, only	<u>Lower ED</u> 0.57 kcal/g	Nutritional composition not matched

	University students and staff	Gender: 30M, 31F		Self-reported and researcher measured EI EI measured for 1 day			<u>Higher ED</u> 1.01 kcal/g	
Miller, 1998 (61)	US University students, staff and local community	BMI: M = 20-26 Gender: 51M, 44F	Adult	Within-subjects Self-reported and researcher measured EI EI measured for 10 days	N = 95	No: Snack food only	<u>Lower ED</u> 2.53 kcal/g <u>Higher ED</u> 5.47 kcal/g	Nutritional composition N/R
Pritchard, 2014 (62)	UK University students, staff and local community	BMI: M = 24 Gender: 10M, 23F	Adult	Within-subjects Self-reported and researcher measured EI EI measured for 1 day	N = 33	No: Lunch only	<u>Lower ED</u> 1 kcal/g <u>Higher ED</u> 1.4 kcal/g	Nutritional composition not matched
Rolls, 2006 (63)	US Local community	BMI: M = 22 Gender: 24F	Adult	Within-subjects Researcher measured EI EI measured for 2 days	N = 24	Yes	<u>Lower ED</u> 1.61 kcal/g <u>Higher ED</u> 2.11 kcal/g	Nutritional composition not matched
Shide, 1995 (64)	US Local community	BMI: M = 24 Gender: 48F	Adult	Within-subjects Self-reported and researcher measured EI EI measured for 1 day	N = 48	No: Lunch food only	<u>Lower ED</u> 0.46 kcal/g <u>Higher ED</u> 1.02 kcal/g	Nutritional composition not matched
Silver, 2008 (65)	US Adults aged over 60 in care	BMI: M = 24 Gender:	Adult	Within-subjects Self-reported and researcher measured	N = 45	No: Lunch only	<u>Lower ED</u> 1.1 kcal/g <u>Higher ED</u>	Nutritional composition N/R

	setting	14M, 31F		EI EI measured for 1 day			2.2 kcal/g	
Smethers, 2019 (66)	US Children in childcare centre	BMI percentile: M = 60 Gender: 26M, 23F	Children	Within-subjects Self-reported and researcher measured EI EI measured for 5 days	N = 49	No: Breakfast, lunch entrée, dinner entrée and afternoon snack only	<u>Lower ED</u> 1.47 kcal/g <u>Medium ED</u> 1.82 kcal/ <u>Higher ED</u> 2.19 kcal/g	Nutritional composition N/R
Spill, 2011 (67)	US Children in day care centre	BMI percentile: M = 57 Gender: 18M, 21F	Children	Within-subjects Researcher measured EI EI measured for 1 day	N = 41	No: Breakfast entrée, lunch entrees, dinner entrée, evening snack only only	<u>Lower ED</u> 1.46 kcal/g <u>Medium ED</u> 1.56 kcal/ <u>Higher ED</u> 1.95 kcal/g	Nutritional composition N/R
Stubbs, 1995a (18)	UK Local community	BMI: M = N/R Gender: 6M	Adults	Within-subjects Researcher measured EI EI measured for 7 days	N = 6	Yes	<u>Lower ED</u> 1.15 kcal/g <u>Medium ED</u> 1.34 kcal/ <u>Higher ED</u> 1.68 kcal/g	Nutritional composition not matched
Stubbs, 1995b (17)	UK Local community	BMI: M = N/R Gender: 7M	Adults	Within-subjects Researcher measured EI EI measured for 14 days	N = 7	Yes	<u>Lower ED</u> 1.15 kcal/g <u>Medium ED</u> 1.34 kcal/ <u>Higher ED</u> 1.68 kcal/g	Nutritional composition not matched
Stubbs, 1998a (15)	UK Local	BMI: M = 22	Adults	Within-subjects Researcher measured	N = 6	Yes	<u>Lower ED</u> 0.89 kcal/g	Nutritional composition matched

	community	Gender: 6M		EI EI measured for 14 days			<u>Medium ED</u> 1.31 kcal/ <u>Higher ED</u> 1.76 kcal/g	
Stubbs, 1998b (16)	UK (Scotland) Local community	BMI: M = 22 Gender: 6M	Adults	Within-subjects Participant self- reported and researcher measured EI EI measured for 14 days	N = 6	Yes	<u>Lower ED</u> 0.85 kcal/g <u>Higher ED</u> 1.50 kcal/g	Nutritional composition matched
Tey, 2016 (68)	Singapore University campus	BMI: M = 22 Gender: 27M	Adults	Within-subjects Participant self- reported and researcher measured EI EI measured for 1 day	N = 27	No: Lunch only	<u>Very low ED</u> 0.27 kcal/g <u>Low ED</u> 0.48 kcal/g <u>Medium ED</u> 0.95 kcal/g <u>High ED</u> 1.37 kcal/g <u>Very high ED</u> 1.81 kcal/g	Nutritional composition not matched
Tey, 2018 (69)	Singapore University campus	BMI: M = 22 Gender: 32M	Adults	Within-subjects Participant self- reported and researcher measured EI EI measured for 1 day	N = 32	No: Lunch food only	<u>Lower ED</u> 0.11-0.13 kcal/g <u>Higher ED</u> 0.49-0.5 kcal/g	Nutritional composition not matched
Westerstep, 1997(70)	Netherlands	BMI: M = 22 & 27	Adults	Within-subjects	N = 36	No: Breakfast food and lunch only	<u>Lower ED</u> 1.21 kcal/g	Nutritional composition not matched

	University campus and local community	Gender: 36F		Participant self-reported and researcher measured EI EI measured for 1 day			<u>Higher ED</u> 1.78 kcal/g	
Williams, 2013 (19)	US University campus and local community	BMI: M = 26 Gender: 30M, 29F	Adults	Within-subjects Researcher measured EI EI measured for 1 day	N = 59	No: Breakfast entrée, lunch entrée, dinner entrée only	<u>Lower ED</u> 1.44 kcal/g <u>Higher ED</u> 1.8 kcal/g	Nutritional composition matched for one comparison (water content manipulation), not matched for other comparisons (e.g. fat content manipulation)

711

712

713

714

715

716

717

718

Records identified through
electronic database searches
(n=6,917)

Records title and abstract
screened to list duplicates
(removed) (n=273)

▼

Records title and
abstract screened (n=6,644)

▼

Eligible articles from
electronic searches
(n=214)

▼

Additional eligible articles
from other sources (n=32)

▼

Articles included in meta-analysis
(n = 246)

Eligible based on title and
abstract (n=5042)

Articles excluded (n=4928)
Not experimental study (n=4)
Energy intake measured > 1 day (n=60)
Unstable participants (n=3)
No comparison available (n=7)
Study design not permit assessment of
TD on energy intake (n=28)
Data not available (n=4)

Articles excluded (n=45)
No direct TD measurement/insufficient
information on int-differences
between conditions (n=37)
Trial not randomized (n=3)
Energy intake measured > 1 day (n=2)
Data not reported as accessible (n=2)