

Supplementary Table 2. Lesion overlap with FC networks, clusters associated with GDS scores, and white matter tracts

	<i>n</i>	jaccard index				voxels			
		mean	<i>SD</i>	min	max	mean	<i>SD</i>	min	max
NAC-seeded									
correlated	16	0.0063	0.0155	0	0.071	534.57	1339.2	0	6440
anticorrelated	3	0.0002	0.0009	0	0.005	6.52	28.13	0	170
GDS cluster	12	0.0104	0.027	0	0.107	236.11	620.834	0	2389
WM tracts	6	0.0002	0.0008	0	0.004	2.17	7.68	0	43
Amy-seeded									
correlated	24	0.0078	0.0132	0	0.051	909.5	1527.07	0	6078
anticorrelated	4	0.0021	0.0078	0	0.045	104.25	400.04	0	2397
dIPFC-seeded									
correlated	10	0.0011	0.0035	0	0.016	327.73	1009.26	0	4469
anticorrelated	9	0.001	0.004	0	0.025	59.41	254.76	0	1655
DMN									
correlated	10	0.002	0.0059	0	0.027	588.32	1711.6	0	8317
anticorrelated	11	0.0023	0.0072	0	0.038	259.05	845.36	0	4783
GDS cluster	4	0.0003	0.0013	0	0.007	31.75	151.33	0	868
WM tracts	4	0.0004	0.0009	0	0.005	1.58	3.9	0	21
V1-seeded									
correlated	8	0.0128	0.0365	0	0.174	2380.89	6876.92	0	33397
anticorrelated	6	0.001	0.0035	0	0.019	126.82	453.49	0	2592

Note. FC = functional connectivity; GDS = geriatric depression scale; SD = standard deviation; NAc = nucleus accumbens; Amy = amygdala; dIPFC = dorsolateral prefrontal-cortex; DMN = default mode network; V1 = primary auditory cortex.