Development of a Decision Support Tool for Acute Appendicitis

Joshua E. Rosen, MD, MHS,^{1,2,3} Joshua M. Liao, MD, MSc,^{3,4} David R. Flum, MD, MPH,^{1,2,3} Jonathan E. Kohler, MD, MA, Giana H. Davidson, MD, MPH^{1,2,6}

¹Surgical Outcomes Research Center, University of Washington, Seattle, Washington

²Department of Surgery, University of Washington, Seattle, Washington

³Decision Science Group, University of Washington, Seattle, Washington

⁴Department of Medicine, University of Washington, Seattle, Washington

⁵Division of Pediatric General, Thoracic and Fetal Surgery, Department of Surgery, University of California – Davis, Sacramento, California

⁶Department of Health Systems and Population Health, University of Washington, Seattle, Washington

Study Funding

This work was supported by the National Institute of Diabetes and Digestive and Kidney Diseases [grant number T32DK070555]; and a generous gift from Marty and Linda Ellison. The development of aspects of the decision support tool was funded by the Patient Centered Outcomes Research Institute. The funding sources had no role in the study design, collection, analysis, or interpretation of data, in the writing of the manuscript or in the decision to submit for publication.

Corresponding Author:

Joshua E. Rosen, MD, MHS

Department of Surgery

University of Washington

1959 NE Pacific Street, Box 356410

Seattle, WA 98195

jerosen@uw.edu

Abstract

Background: Multiple randomized controlled trials have shown that it is safe and effective to treat appendicitis with antibiotics or surgery. There are no tools available to assist surgeons and their patients in choosing the optimal treatment for each individual patient. Here we describe the development of a new decisions support tool (DST) for acute appendicitis and place it in the context of international guidelines for decision aid development.

Methods: The stakeholder engagement and development process for the DST is described. The DST and its development process are placed in the context of the International Patient Decision Aid Standards (IPDAS) and the DEVELOPTOOLS checklist for a user-centered design process.

Results: A diverse group of over 60 stakeholders were involved in the needs-assessment, development, and evaluation of the DST. The development process met 11/11 of the scored items on the DEVELOPTOOLS checklist. Of the 34 applicable IPDAS items, the current version of the DST meets 31 of them including 6/6 qualifying criteria, 6/6 certification criteria, and 18/22 quality criteria.

Conclusions: The novel appendicitis DST was developed with the input of multiple stakeholders. The development process and the tool itself complies with best practices recommended by the IPDAS.

Introduction

Appendicitis is one of the most common acute surgical conditions in the United States and has been largely treated with appendectomy. However, accumulating data from 9 randomized controlled trials has indicated that treatment with antibiotics with surgery if there is progression of symptoms is a safe option that will allow many patients to avoid an urgent surgical procedure.(1–9) Furthermore, data from the recent Comparison of Drugs and Antibiotic Medicines for the Treatment of Acute Appendicitis (CODA) trial(1) has shown that up to 7 in 10 patients may avoid an urgent operation and patient reported quality of life measures are similar at 30 and 90 days for participants in both treatment arms. However, there are important differences with respect to healthcare utilization, days of work missed, and risk of various complications that may influence patient decision-making and surgeon willingness to discuss and offer both treatment options.

The decision of which treatment to pursue, surgery or antibiotic therapy for acute appendicitis, requires careful consideration of multiple data points across different outcome domains in the context of the unique preferences and values of individual patients. This type of individualized patient-centered decision making may be particularly difficult for acute conditions in the emergency department given both the time-pressured nature of the decision, pain and anxiety experienced by the patient, and the fact that clinicians and patients have no prior established relationship. Furthermore, the priorities of patients and surgeons may be discordant. For example, surgeons may be most concerned about recurrence whereas patients may be most concerned about surgical pain or missing work.(10) Based on the feedback of multiple stakeholder groups involved in the CODA trial, we set out to develop a decision support tool (DST) (appyornot.org) to inform patients about their treatment options and assist patients in the emergency setting making this treatment decision. This article will describe the development process of the tool and place it specifically within the published framework of the recently published DEVELOPTOOLS checklist.(11) We also outline how the tool complies with the International Patient Decision Aid Standards (IPDAS) guidelines for a patient decision aid.(12,13)

Intended Users and Setting

The intended users of this decision support tool are adult patients with imaging-confirmed acute appendicitis and their caregivers and the clinicians who care for them. The typical setting for use would be in the emergency department on an electronic device (either patient-owned or hospital-provided) in collaboration with the surgical consultation. Notable features of this setting include: 1) patients will typically be experiencing acute abdominal pain and may also be experiencing nausea, vomiting, or other distressing symptoms; 2) patients will typically not have a long-standing relationship with the clinicians they interact with and will often be meeting them for the first time during this encounter; 3) there is a time-pressured nature to the environment, both because appendicitis is an acute condition that requires an urgent treatment decision (that cannot be delayed and discussed at a second appointment for example) and because the emergency department often cares for a high-volume of patients and must focus on patient flow into and out of the department.

Purpose of the Tool and Needs Assessment

Over the past number of years there has been increasing calls for enhancing the use of SDM around surgical procedures.(14,15) While shared decision making (SDM) has been embraced in many fields of surgery, emergency general surgery has for a long time been seen as a particularly difficult environment in which to implement it given that patients are often in acute distress and decisions are made in a time-pressured environment. The mounting evidence for an antibiotics-first approach to treating appendicitis, has resulted in profound interest in methods to facilitate strong shared decision making for this disease process.(14) While there have been efforts to build better tools to facilitate these discussions for pediatric patients,(16,17) there has not been any tools developed to facilitate the integration of adult clinical trial data into shared decision making. This is particularly important, as the mounting evidence for antibiotic treatments may meet criteria under the Affordable Care Act requiring shared decision making.(14,15)

While the most recent data from the CODA trial has shown similar outcomes for surgery and antibiotics for overall quality of life and time to symptom resolution, the treatments do differ on a number of important outcomes that may be important to patients. These outcome dimensions include days of missed work, risk of needing future surgery, and risk of missing an occult malignancy.(1) Early discussions with patient and clinician stakeholder groups acknowledged the complexity these results may pose for effective SDM and identified the need to develop tools and procedures to facilitate it. Stakeholders recognized the variation in patient experiences and practice environments across the country and world, and the inherent variability in treatment discussions that this may lead to. They also recognized that patients and surgeons may value certain outcomes and define treatment success differently.(10) They identified key attributes of any appendicitis DST as including: effective transmission of knowledge in both video and text formats, facilitation of outcomes comparison between treatments, exercises to clarify the needs and values of individual patients and how they map to the different treatments and their outcomes, and finally the DST must be useable under the constraints of the typical emergency general surgery environment described above.

Main Stakeholder Representation Groups

Figure 1 shows the main groups involved in the development of the decision support tool. The CODA trial included key stakeholders through the research cycle from conception to dissemination(18,19): a Patient Advisory Board (6 patient advisors), a Clinical Advisory Group made up of 43 surgeons and emergency medicine physicians, and a National Stakeholder Advisory Group representing professional societies, payors, and policymakers (8 members). The detailed makeup of these groups and their formation has been described previously.(19) All three of these groups served as key sources of opinions and feedback throughout the development process.

In addition to the main stakeholder groups involved in the CODA trial, we employed several other groups in the development of this decision aid. We used online pilot-testing in people at-risk for appendicitis

(i.e., no prior appendicitis, typical age distribution from the CODA trial) using Amazon's Mechanical Turk (MTurk) crowdsourcing platform.(10,20) We engaged with non-healthcare provider content experts at our institution in the fields of cognitive and behavioral psychology, behavioral economics, and marketing to design both the decision aid elements and the testing framework used in our online pilot tests (4 people overall). We solicited opinions from a diverse group of clinical providers including acute care, surgery residents, emergency medicine physicians, and emergency department nurses. Finally, our core research team developing the decision aid included acute care surgeons, surgery residents, and experts in behavioral economics and educational video development.

Figure 1: Main groups involved in the development of the appendicitis DST

Development Process

The appendicitis DST was developed through an iterative process involving representatives of the the stakeholder groups. During analysis of the trial results,(1) multiple focus groups were held with patient and physician stakeholders focusing on the interpretation of the trial results and planning for dissemination and implementation strategies. Stakeholders provided input on the purpose and goals of an

appendicitis DST and the concept of a DST consisting of an informational video followed by an interactive decision aid was solidified. Furthermore, CODA stakeholder groups identified the key outcome data points for inclusion in the decision aid and video.

The informational video was developed based on a prior video that had been created to facilitate a more uniform informed consent process for participation in the CODA trial.⁽²¹⁾ The video script underwent multiple rounds of development and review by patient and clinician takeholders, as well as review by healthcare communications experts. Finally, the videos were professionally shot and edited and final videos were reviewed by key stakeholders prior to inclusion in the DST.

The interactive patient decision aid was developed by using the needs assessment specified by the CODA advisory groups to complete a literature search identifying best practices in the key areas of information communication and values clarification. We also consulted with non-clinicina experts as described above. We then performed five rounds of iterative and sequential testing of our information delivery practices using over 1,300 subjects on MTurk. After developing an initial prototype based on this iterative testing, we solicited feedback from CODA stakeholder groups, and the additional clinical stakeholders identified in Figure 1.

Description of Decision Aid

The DST consists of two main components, an informational video, and an interactive decision aid. Patients first view a professionally produced video in either English or Spanish that describes the pathophysiology of appendicitis, reviews evidence-based treatment options (e.g., antibiotics and surgery), and reviews information on the expected clinical course. Following the video, patients view the interactive decision aid which reviews clinical outcomes comparing antibiotics and surgery followed by a values-clarification exercise where patients rate the importance of various outcomes and see how they

align with each treatment option. Finally, a summary screen is provided that can be used to help facilitate discussion of treatment options with the clinical team and patient.

Figure 2: Example screenshots from decision aid a) video selection screen where users can select the informational video they would like to watch; b) data-presentation screen demonstrating the graphical format of data presentation involving point estimates and graphical confidence intervals; c) users can input what outcomes are most important to them; d) the tool categorizes those outcomes as favoring either antibiotics or surgery to aid in patient decision making.

Assessing the Development Process – The DEVELOPTOOLS Checklist

The IPDAS group provides a basis for a recommended systematic development process and a recently published update incorporates principles of user-centered design to create the DEVELOPTOOLS checklist(11) for reporting the design and development process for patient decision aids. In Table 1 we

outline the components of the development process for the appendicitis DST using the DEVELOPTOOLS checklist.

Table 1: DEVELOPTOOLS checklist and evidence items from the Appendicitis DST

<u>Checklist Item</u>	<u>Evidence from Appendicitis DST</u>	<u>Scoring</u> <u>(1 =</u> <u>Present,</u> <u>0 =</u> <u>Absent)</u>
Pre-prototype Involvement		
Were potential users involved in any steps to help understand users and their needs?	Yes, members of the CODA patient advisory board, clinical advisory group and national stakeholder advisory group (see above for numbers) were involved. emergency department nurses and surgery residents were involved.	1
Were potential users involved in any steps of designing, developing, and/or refining a prototype?	Yes, see Figure 1 and Stakeholder section above for all groups involved in prototype development.	1
Iterative Responsiveness		
Were potential users involved in any steps to evaluate prototypes of the tool or a final version of the tool?	Yes. We used online crowd-sourcing via MTurk (over 1200 subjects), as well as the clinical and patient advisors described in Figure 1. Furthermore, we are actively trialing the pilot version of the tool with patients in the emergency department at our institution.	1
Were potential users asked their opinions of prototypes of the tool or a final version of the tool in any way?	Yes, see Figure 1 and stakeholder section above.	1
Were potential users observed using the tool in any way?	Yes, surgeons and patients are currently being observed using the tool in the emergency department.	1
Did the development process have 3 or more iterative cycles? (Definition of a cycle is that your team developed something and showed it to at least one person outside of the team before making changes in response to their feedback)	Yes, there was 5 cycles of online development studies and at least 3 more cycles involving feedback from the other clinician and patient stakeholder in Figure 1.	1
Were changes between iterative cycles explicitly reported in any way?	Yes in internal logs	1
Other Expert Involvement		
Were health professionals asked their opinion of the tool at any point?	Yes the clinical advisory group as well as additional acute care surgery faculty, surgery residents, and emergency medicine physicians were asked about the tool.	1
Were health professionals consulted before a first prototype was developed?	Yes, as described above the clinical advisory group was consulted prior to prototype development.	1
Were health professionals consulted between initial and final prototypes?	Yes the clinical groups in Figure 1 were consulted at multiple points.	1
Was an expert panel involved?	Yes, as described in the stakeholder section above.	1
Additional Elements		
Was a formal advisory panel of users involved?	Yes, see(19) for information on the assembly of stakeholder panels.	N/A
Were users, health professionals, and other relevant stakeholders involved as members of the research team?	Yes, acute care surgeons and marketing/behavioral economics experts were research team members. Furthermore, the CODA patient advisors are considered to be members of the research team and contributed to the results dissemination discussions.	N/A
Were members of populations marginalized by social norms and policies involved?	Unable to answer, under-represented minorities were under-recruited to the patient advisory board.	N/A
How many users and health professionals were involved in total and of each type?	See stakeholder section and Figure 1 above.	N/A
Does the tool have a defined purpose?	Yes, the tool's purpose is to provide consistent information about treatment options for appendicitis and to help patients decide between surgical and	N/A

	antibiotic treatment	
Is the tool intended to be used in a particular context?	Yes, the tool is intended to be used in the emergency setting.	N/A
Were any methods used to facilitate sharing of perspectives between groups?	Yes, stakeholder meetings often were cross-disciplinary and both structured and open ended questions were asked of participants. Deidentified summary of discussion points were also shared.	N/A
Were users involved from the outset of the project?	Yes, patient and clinician users have been involved from the outset of the CODA project	N/A
Were translations and cultural adaptation used to render the patient decision aid available to users across languages and cultures?	Yes, for the videos we have multiple actors from different cultural backgrounds as well as a Spanish language option. The decision aid is translated into Spanish. Additional languages including American Sign Language are planned.	N/A

Assessment of Decision Support Tool by IPDAS Guidelines

The IPDAS provides a framework for the evaluation of patient decision aids. Version 4.0 published in 2013 specifies a set of minimum criteria broken up into three categories: qualifying criteria (defining if a tool is a decision aid), certification criteria (defining essential facets of a decision aid), and quality criteria (those that will enhance a decision aid but are not essential).(12) Table 2 below shows the 34 applicable criteria in IPDAS 4.0 (9 of the standard 44 criteria are for decision aids involving the use and interpretation of medical tests, 1 additional criterion is not applicable to the appendicitis DST) and whether or not the appendicitis DST meets them in its current form. Overall 6/6 qualifying criteria are met, 6/6 certification criteria are met, and 18/22 quality criteria are met, with three currently in progress.

Table 2: IPDAS 4.0 criteria and scoring of appendicitis DST

<u>Dimension</u>	<u>Criteria</u>	<u>Type of Criteria</u>	<u>Met by Appendicitis DST?</u>
Information	The patient decision aid describes the health condition or problem (treatment, procedure, or investigation) for which the index decision is required.	Qualifying	Yes
	The patient decision aid explicitly states the decision that needs to be considered (index decision).		Yes
	The patient decision aid describes the options available for the index decision.		Yes
	The patient decision aid describes the positive features (benefits or advantages) of each option.		Yes
	The patient decision aid describes the negative features		Yes

	(harms, side effects, or disadvantages) of each option.		
	The patient decision aid shows the negative and positive features of options with equal detail (e.g., using similar fonts, sequence, presentation of statistical information).	Certification	Yes
	The patient decision aid describes the natural course of the health condition or problem, if no action is taken (when appropriate).	Quality	NA
	The patient decision aid makes it possible to compare the positive and negative features of the available options.		Yes
Probabilities	The patient decision aid provides information about outcome probabilities associated with the options (i.e., the likely consequences of decisions).	Quality	Yes
	The patient decision aid specifies the defined group (reference class) of patients for whom the outcome probabilities apply.		Yes
	The patient decision aid specifies the event rates for the outcome probabilities		Yes
	The patient decision aid allows the user to compare outcome probabilities across options using the same time period (when feasible).		Yes
	The patient decision aid allows the user to compare outcome probabilities across options using the same denominator (when feasible).		Yes
	The patient decision aid provides more than 1 way of viewing the probabilities (e.g., words, numbers, and diagrams).		Yes
Values	The patient decision aid describes what it is like to experience the consequences of the options (e.g., physical, psychological, social).	Qualifying	Yes
	The patient decision aid asks patients to think about which positive and negative features of the options matter most to them (implicitly or explicitly).	Quality	Yes
Guidance	The patient decision aid provides a step-by-step way to make a decision.	Quality	Yes
	The patient decision aid includes tools like worksheets or lists of questions to use when discussing options with a practitioner		Yes
Development	The development process included a needs assessment with clients or patients.	Quality	Yes
	The development process included a needs assessment with health professionals.		Yes
	The development process included review by clients/patients not involved in producing the decision support intervention.		Yes
	The development process included review by professionals not involved in producing the decision support intervention.		Yes
	The patient decision aid was field tested with patients who were facing the decision.		In Progress*
	The patient decision aid was field tested with practitioners who counsel patients who face the decision.		In Progress*
Evidence	The patient decision aid (or associated documentation) provides citations to the evidence selected.	Certification	Yes
	The patient decision aid (or associated documentation) provides a production or publication date.		Yes

	The patient decision aid (or associated documentation) provides information about the update policy.		Yes
	The patient decision aid provides information about the levels of uncertainty around event or outcome probabilities (e.g., by giving a range or by using phrases such as “our best estimate is . . .”).		Yes
	The patient decision aid (or associated documentation) describes how research evidence was selected or synthesized.	Quality	Yes
	The patient decision aid (or associated documentation) describes the quality of the research evidence used.		Yes
Disclosure	The patient decision aid (or associated documentation) provides information about the funding source used for development.	Certification	Yes
	The patient decision aid includes authors’/ developers’ credentials or qualifications.	Quality	Yes
Plain Language	The patient decision aid (or associated documentation) reports readability levels (using 1 or more of the available scales).	Quality	No
Evaluation	There is evidence that the patient decision aid improves the match between the preferences of the informed patient and the option that is chosen.	Quality	In Progress*
	There is evidence that the patient decision aid helps patients improve their knowledge about options’ features.		Yes

*A field-test of the decision aid in at-risk users has been completed. A field test in users facing the healthcare decision is currently ongoing and will be reported in the future

Conclusions

In this article we have described the development process for a new decision aid for acute appendicitis.

The process involved input from multiple clinician and patient stakeholders and adhered to principles of user-centered and evidence-based design as described in the DEVELOPTOOLS checklist. The resulting decision support tool complies with the IPDAS 4.0 criteria. To our knowledge this is one of the first DSTs specifically developed for use in adult emergency general surgery patients. This may serve as a roadmap to more efficiently develop patient-centered tools for SDM for emergency surgical patients.

References

1. Collaborative C, Flum DR, Davidson GH, Monsell SE, Shapiro NI, Odom SR, et al. A Randomized Trial Comparing Antibiotics with Appendectomy for Appendicitis. *New Engl J Med*. 2020;383(20):1907–19.

2. O'Leary DP, Walsh SM, Bolger J, Baban C, Humphreys H, O'Grady S, et al. A Randomised Clinical Trial Evaluating the Efficacy and Quality of Life of Antibiotic Only Treatment of Acute Uncomplicated Appendicitis: Results of the COMMA trial. *Ann Surg.* 2021;Publish Ahead of Print.
3. Salminen P, Tuominen R, Paaanen H, Rautio T, Nordström P, Aarnio M, et al. Five-Year Follow-up of Antibiotic Therapy for Uncomplicated Acute Appendicitis in the APPAC Randomized Clinical Trial. *Jama.* 2018;320(12):1259–65.
4. Eriksson S, Granström L. Randomized controlled trial of appendicectomy versus antibiotic therapy for acute appendicitis. *Brit J Surg.* 1995;82(2):166–9.
5. Styrd J, Eriksson S, Nilsson I, Ahlberg G, Haapaniemi S, Neovius G, et al. Appendectomy versus Antibiotic Treatment in Acute Appendicitis. A Prospective Multicenter Randomized Controlled Trial. *World J Surg.* 2006;30(6):1033.
6. Turhan AN, Kapan S, Kütükçü E, Yiğitbaş H, Hatipoğlu S, Aygün E. Comparison of operative and non operative management of acute appendicitis. *Ulusal Travma Ve Acil Cerrahi Dergisi Turkish J Trauma Emerg Surg Tjtes.* 2009;15(5):459–62.
7. Vons C, Barry C, Maitre S, Pautrat K, Leconte M, Costaglioli B, et al. Amoxicillin plus clavulanic acid versus appendicectomy for treatment of acute uncomplicated appendicitis: an open-label, non-inferiority, randomised controlled trial. *Lancet.* 2011;377(9777):1573–9.
8. Hansson J, Körner U, Khorram-Manesh A, Solberg A, Lundholm K. Randomized clinical trial of antibiotic therapy versus appendicectomy as primary treatment of acute appendicitis in unselected patients. *Brit J Surg.* 2009;96(5):473–81.
9. Talan DA, Saltzman DJ, Mower WR, Krishnadasan A, Jude CM, Amii R, et al. Antibiotics-First Versus Surgery for Appendicitis: A US Pilot Randomized Controlled Trial Allowing Outpatient Antibiotic Management. *Ann Emerg Med.* 2017;70(1):1-11.e9.
10. Rosen JE, Agrawal N, Flum DR, Liao JM. Willingness to undergo antibiotic treatment of acute appendicitis based on risk of treatment failure. *Brit J Surg.* 2021;znab280-.
11. Witteman HO, Maki KG, Vaissou G, Finderup J, Lewis KB, Steffensen KD, et al. Systematic Development of Patient Decision Aids: An Update from the IPDAS Collaboration. *Med Decis Making.* 2021;0272989X2110141.
12. Hoffmann TC, Bakhit M, Durand M-A, Perestelo-Pérez L, Saunders C, Brito JP. Basing Information on Comprehensive, Critically Appraised, and Up-to-Date Syntheses of the Scientific Evidence: An Update from the International Patient Decision Aid Standards. *Med Decis Making.* 2021;0272989X2199662.
13. Joseph-Williams N, Newcombe R, Politi M, Durand M-A, Sivell S, Stacey D, et al. Toward Minimum Standards for Certifying Patient Decision Aids. *Med Decis Making.* 2013;34(6):699–710.
14. Telem DA. Shared Decision Making in Uncomplicated Appendicitis: It Is Time to Include Nonoperative Management. *Jama.* 2016;315(8):811–2.

15. Lee EO, Emanuel EJ. Shared Decision Making to Improve Care and Reduce Costs. *New Engl J Medicine*. 2013;368(1):6–8.
16. Minneci PC, Mahida JB, Lodwick DL, Sulkowski JP, Nacion KM, Cooper JN, et al. Effectiveness of Patient Choice in Nonoperative vs Surgical Management of Pediatric Uncomplicated Acute Appendicitis. *Jama Surg*. 2015;151(5):1–8.
17. Minneci PC, Cooper JN, Leonhart K, Nacion K, Sulkowski J, Porter K, et al. Effects of a Patient Activation Tool on Decision Making Between Surgery and Nonoperative Management for Pediatric Appendicitis. *Jama Netw Open*. 2019;2(6):e195009.
18. Ehlers AP, Davidson GH, Bizzell BJ, Guiden MK, Skopin E, Flum DR, et al. Engaging Stakeholders in Surgical Research: The Design of a Pragmatic Clinical Trial to Study Management of Acute Appendicitis. *Jama Surg*. 2016;151(6):580–2.
19. Ehlers AP, Davidson GH, Deeney K, Talan DA, Flum DR, Lavalley DC. Methods for Incorporating Stakeholder Engagement into Clinical Trial Design. *Egms Generating Évid Methods Improv Patient Outcomes*. 2017;5(1):4.
20. Rosen JE, Agrawal N, Flum DR, Liao JM. Verbal Descriptions of the Probability of Treatment Complications Lead to High Variability in Risk Perceptions: A Survey Study. *Ann Surg*. 2021;Publish Ahead of Print.
21. Davidson GH, Flum DR, Talan DA, Kessler LG, Lavalley DC, Bizzell BJ, et al. Comparison of Outcomes of antibiotic Drugs and Appendectomy (CODA) trial: a protocol for the pragmatic randomised study of appendicitis treatment. *Bmj Open*. 2017;7(11):e016117.