Manuscript Title: Dengue Fever Distribution, Climate and Visitors: A Study from Badung District of Bali, Indonesia

Authors:

Mochamad Rizal Maulana¹, Ririh Yudhastuti^{1,2}, Muhammad Farid Dimjati Lusno^{1,2}, Yudied Agung Mirasa³, Setya Haksama⁴, Zida Husnina^{1,2*}

¹Department of Environmental Health, Faculty of Public Health, Universitas Airlangga Surabaya, Indonesia

²Research Center for Tropical Diseases, Infectious Diseases and Herbal Medicine, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia.

³Center for Disease Control and Prevention (BBTKLPP) Surabaya, Indonesia

⁴Department of Health Administration and Policy, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

*Corresponding Author

Authors' contact details:

1. Mochamad Rizal Maulana¹ (email: mochamad.rizal.maulana-2018@fkm.unair.ac.id)
2. Yudhastuti, Ririh, Dr. DMV., MSc.^{1,2} (email: ririh.unair@gmail.com)
3. Lusno, Muhammad Farid D. dr., MKL.^{1,2} (email: faridlusno@fkm.unair.ac.id)
4. Mirasa, Yudied Agung, Dr., SKM., MKes.³ (email: ymirasa@gmail.com)
5. Haksama, Setya Dr.drg. MKes. (email: setyahaksama@fkm.unair.ac.id)
<https://orcid.org/0000-0002-9080-3442>
6. Husnina, Zida, MPH.^{1,2} (email: zida.husnina@fkm.unair.ac.id) <http://orcid.org/0000-0003-3703-870X>

Corresponding Author:

Name : Zida Husnina, MPH.

Address : Department of Environmental Health, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia.

Phone numbers: +62 31 592 0948

Facsimile numbers: +62 31 592 0948

E-mail address: zida.husnina@fkm.unair.ac.id

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

ABSTRACT:

As a national and international tourism destination, Badung district has recorded the highest dengue fever (DF) in Bali Province with multiple suspected influencing factors. This research presents the distribution of dengue fever (DF) incidence in Badung district and analyses its association with climate and visitors. The monthly data of DF surveillance, climate and reported number of visitors during January 2013 – December 2017 was analysed using Poisson Regression. 10689 new DF cases were notified from January 2013 to December 2017 while 2016 was recorded as the heaviest DF incidence. Monthly DF cases have positive association with average temperature, precipitation, humidity and local visitors. While negative association was shown between DF cases with foreign visitors. This study underlines the urgency to integrate climate and tourism for dengue fever control. Further study is recommended to take both factors together with population mobility at finer spatial and time unit of analysis.

Keyword: *Dengue Fever (DF), Climate, Badung, Visitors, Bali Indonesia*

1. Introduction

The incidence of Dengue Fever (DF) around the world increased in recent decades (Waggoner et al. 2016). The world health organization (WHO) recorded an increasing number of DF cases

15 fold over the last two decades, from 505,430 cases in 2000 to 2,400,138 cases in 2010 and 3,312,040 in 2015. The mortality number increased from 960 in 2000 to more than 4032 in 2015 (WHO 2019). It is estimated 390 million dengue infections occur annually with 96 million cases were manifesting clinical severity (Bhatt et al. 2013). The infection impacted 3.9 billion population in 129 countries at risk, whereas 70 percent of the actual burdens is located in Asia (Brady et al. 2012). Where this region was detected to have profound synchronous dengue fever infection pattern since 1990's (Van Panhuis et al. 2015).

Dengue fever is caused by *Flaviviridae* family viruses that are categorized into four types (DENV-1, DENV-2, DENV-3, and DENV-4), that all of these strains are exist in Indonesia. The World Health Organization (WHO) classifies Indonesia as one of the endemic countries due to the high periodic DF outbreaks. The DF IR in Indonesia was fluctuated during 2010 – 2017. The lowest IR recorded was 27.67 per 100,000 population (in 2011) and reached the peak 78.85 per 100,000 population in 2017. In 2018, DF incidence was reported by 85.65 percent of all districts in Indonesia. Regions with the highest DF contracted districts were Papua, North Maluku, Bali, Banten, Central Java, Yogyakarta, Jakarta, Jambi, and Riau. Up to 2018, the national IR was IR 24.73/100,000 population with Case Fatality Rates (CFR) was 0.71. This numbers are lower than 2017 with IR 26.10/ 100,000 population with CFR 0.72 percent (MOH 2018).

Bali Province is an endemic are to DF for years. In 2016, the disease transmission emerged in all districts with 21,668 new cases (IR 544.3/100,000 population) and 63 mortalities (CFR 0.29 percent). Badung is the district with the highest number of cases (3,998 cases), followed by Buleleng and Gianyar (3,787 and 3,673 new cases respectively) (PHO 2016). This district has various tourists destinations and has the largest visitors in the Island of Bali. In

2013 as many as 438 thousand local tourists and 3.3 million foreign tourists were recorded to have had visited Badung, and it is expected to increase annually (The Government 2018). This condition was indicated could increase the risk of DF transmission abroad (Yoshikawa and Kusriastuti 2013).

Climate affects the pattern of infectious diseases because vector and viruses sensitivity for any change that directly or indirectly affect the onset of a disease. Some studies have revealed that there is a significant association between DHF with climatic conditions including air temperature, humidity, and precipitation due to increasing breeding place for of *Aedes aegypti* as primary vector of dengue virus (Colón-González et al. 2013, Fatmawati and Sulistyawati 2019, Xiang et al. 2017). Thus, investigation on how the climate variables impact to dengue incidence in Badung could enrich the discussion around this issue.

One of the social demographics and population factors is tourism that have long been studied as the contributing factor to the transmission of infectious disease, as reported by study in South Korea that international travellers from South East Asia are associated with the DF occurrence during 2006 – 2015 (Cho et al. 2018). Another dengue case found from traveller who came from Saudi Arabia whose type-2 virus strain clustered with isolates of Singapore and India (Matsui et al. 2019). The impact of mobility and tourism not only exacerbated local transmission but also have driven global spread of DF, for instance the research in 2018 found significant association between the DF incidence in Bali with the imported cases in Australia (Xu et al. 2018).

Despite the outgrowing research around dengue transmission and the high burden of the disease in Badung district, little is known about the distribution characteristics and the

association with driving forces both from climate and visitors. Most of the current research examined the association of dengue events with climate factors at one certain time. Such as the study in Denpasar City examined the association of climate and DHF in 2010-2015 (Wiradarma and Somia 2015) another study did comprehensive analysis between DF cases and climate for the entire of Bali Island, however has not taking into account for other factors such as socio economic, transportation and population dynamics (Dhewantara et al. 2019).

Aside from the substantial information presented by these studies, further analysis of DF in relation with climate and visitors at longer observation period could comprehend the recommendation for mitigation and control of DF. In this research, the authors did exploration for the pattern of DF incidence and the explanatory variables (climate and visitors) in Badung district and identify the association between them.

2. Materials and Methods

2.1 Study design

This research is ecological study with population level as unit analysis. Yearly data from 2013 to 2019 (7 years) at sub-district level was used for descriptive choropleth maps and monthly data for the period of January 2013 – December 2017 (60 months) at district level was used for statistical analysis.

2.2 Study Area

Badung district is located between 08°14'20"- 08°50'48" South latitude, and 115°05'00"-115°26'16" East longitude, sized 418.5 Kilometers square of lowland area with elevation up to 2075 meters above sea level. Administratively, the district is consisted of 6 sub districts (Petang, Abiansemal, Mengwi, Kuta Utara, Kuta and Kuta Selatan). As tropical area, Badung District has only rainy season and dry season. The rainfall is ranging between 2.0 mm

– 622.8 mm. While, the maximum temperature is 26.6 °C – 31.2 °C, the minimum temperature is 25.4 °C - 24.1 °C. The humidity is ranging from 78 percent (around August) – 84 percent (around June).

2.3 Variables and Data Sources

The outcome variable in this study is number of DF cases obtained from monthly province surveillance data since January 2013 until December 2017. Dengue fever (DF) is defined as diagnosed by physician with symptoms of 2 – 7 days fever accompanied by manifestations of bleeding, decreased platelets, and hemoconcentration. The confirmation of DF case is supported by Tourniquet test, thrombocytopenia (platelets $\leq$ 100000/mm³), increased hematocrit $\geq$ 20 percent, and the increase of the specific IgM antibodies dengue virus in blood tests in the laboratoris (WHO 2011). The independent variables are climate factors (temperature, precipitation and humidity) gathered from Meteorology, Climatology and Geophysics Bureau (BMKG) and number of recorded visitors (local and foreign) were extracted from Badung Statistics Bureau (BPS). The data for producing descriptive mapping was extracted from Badung District health profile 2013 – 2019 that provided DF yearly data at sub-district level categorized based on the gender of reported patients.

2.4 Data Analysis

2.4.1 Spatial Visualisation

Choropleth maps were generated to provide geographical distribution of DF incidence rates (IR) across 6 sub-districts for 7 years (2013 – 2019). The incidence rates (IR) was calculated by dividing total cases with the total population of each district and each observed year. The value was classified using natural breaks due to small spatial units. The maps were overlaid with the presentation of gender distribution of reported cases for each sub-districts. The production of maps used ArcGIS 10.4 software by ESRI.

Statistical Descriptive and Time Series Decomposition

The descriptive analysis was performed by presenting the variables in distribution frequency table and box plots in order to describe the fluctuation of mean values of each variable in each consecutive years. The DF incidence rates (IR) data was decomposed using multiplicative approach to examine the trend, seasonal and random pattern. The decomposition graph was generated using R statistical software version 4.0.2.

2.4.2 Analysis with Poisson Regression

Uni-variate analysis of Poisson regression was performed to examine the association of DF cases as outcome variable with each of explanatory variables (air temperature, temperature, and precipitation, local and foreign visitors). The analysis was performed for multiple lags variables (1-3 months) to explore the most significant association and to choose variables to be included into the next analysis. The detection for Variance Inflation Factor (VIF) was conducted to examine any multicollinearity the analysis was conducted using STATA 14 software by Stata Corp.

The Multivariate Poisson regression was adopted to examine the association of monthly cases with the selected explanatory variables simultaneously.

$$\theta_i \sim \text{Poisson}(\mu_i)$$

$$\text{Log } \theta_i = \alpha + (\beta_1 \times \text{temperature}_i) + (\beta_2 \times \text{precipitation}_i) + (\beta_3 \times \text{number of local visitors}_i)$$

$$+ (\beta_4 \times \text{number of foreign visitors}_i)$$

θ_{ij} is the number of DF cases in Badung district in month i, while α is the intercept, β_1 , β_2 , β_3 and β_4 are the poisson regression coefficients for the four explanatory variables (temperature, precipitation, number of local visitors and number of foreign visitors). The goodness and significance of the association was evaluated by looking at p value (<0.05), 95% confidence interval (CI) and Akaike Information Criterion (AIC).

Ethics

This study was approved by Ethical Committee for Health Research, Faculty of Public Health, Universitas Airlangga, Indonesia, registered on October 1st, 2019 with reference number: 208/EA/KEPK/2019.

3. Results

3.1 Geographics Distribution

In 2016 was recorded as the heaviest DF incidence rates in Badung District where the highest IRs were located in Abiensemal and Mengwi (almost reached 1000 cases/ 100000 population) followed by Kuta Selatan Sub-district. The year of 2015 was recorded as the second highest IR (reached above 500 cases/100000 population) where 2 sub-districts (Mengwi and Kuta Selatan) were the largest transmission. Even though during 2017 and 2018 the IRs were tend to decrease for all 6 sub-districts in Badung District and landed at the lowest in 2018 (below 100 cases/ 100000 population for entire sub-districts). It is however showed darkening IR for Mengwi sub-districts in 2019. Overall, since 2013 – 2019, the areas that frequently experiencing high IR were Mengwi and Kuta Selatan, while the lowest IR were Petang and Kuta district [Figure 1].

The gender proportion of cases showed that during 7 years in 6 sub-districts were dominated by male patients, except in 2015 and 2018 for Abiensemal sub-disrict when the female patients were higher than male. However the rest of 5 illustrated the male patients were higher or slightly equal to female patient [Figure 1]. The detail table distribution of DF IR and gender proportion available in appendix 2.

3.2 Descriptive Statistics

During 60 months of observation (January 2013 – December 2017) there were 10,689 new DF cases. The maximum number of cases occurred in April 2016 and the minimum occurred in

August 2017, with average of incidence rates almost 30 cases per 100000 Population. The highest temperature was recorded in November 2015 and the lowest was in August 2015 with average 27°C. The precipitation reached its peak in December 2017 (above 620 mm), while the lowest was in October 2015 (0 mm) with average 154.1 mm. The highest humidity was recorded in February 2016 and the lowest was in September 2014 with average value 79.98 percent. The highest number of local visitors was above 94 thousands recorded in November 2016 with average was nearly 50 thousands each month. While the highest foreign visitors was close to 600 thousands as recorded in August 2017 with average above 359 thousand every month. The detail statistic description is available in Table 1.

The decomposition plot of DF time series data demonstrated an increasing trend that reached its highest on mid year of 2016 before it was sloping down thereafter. The plot also illustrated profound seasonal shape with consistent peaks around early months and the lowest around October - November for every year [Figure.2].

The box plots demonstrate the mean values distribution of variables in each 5 consecutive years. The highest mean of DF cases and incidence rates (IR) were reported in 2016. This pattern is similar to temperature variable, as the mean temperature in 2016 was reported as the highest. The rest of climatic variable have different pattern, whereas the mean of humidity in 2017 indicated as the highest, while the mean of precipitation in 2013 showed as the highest. The highest mean value of visitors was appeared in 2016 (local visitors) and 2017 (foreign visitors) [Figure. 3].

3.3 Analysis and Model Construction with Poisson Regression

The association between outcome and each of explanatory variables for multiple lags (0- 3 months lag) have been examined and all of them have significant result except for humidity at lag 3 months. Based on the AIC, *p value* and 95% CI there were 5 variable selected: temperature lag 3 months, precipitation lag 2 months, humidity lag 2 months, local visitors lag 1 month and foreign visitors lag 3 months. The detection of multicollinearity using Variance Inflation Factor (VIF) showed there is no value exceeding 4 means there is no multicollinearity among selected variables [Appendix 1].

DF cases is significantly associated with temperatures, the number of foreign tourists in the previous 3 months and local tourist in the previous 1 month. The DF Cases is also significantly associated with the precipitation and humidity occurring in the previous 2 months. DF cases increases by 5.909×10^{-1} (95% CI: $5.610 \times 10^{-1} : 6.209 \times 10^{-1}$), by 5.779×10^{-4} (95%CI: $3.882 \times 10^{-4} : 7.676 \times 10^{-4}$), 1.434×10^{-2} (95% CI: $0.303 \times 10^{-2} : 2.565 \times 10^{-2}$) and by 7.40×10^{-6} (95%CI: $5.88 \times 10^{-6} : 8.91 \times 10^{-6}$) for each increment 1°C of temperature (3 months lag), 1 mm of precipitation (2 months lag), 1% humidity (2 months lag) and additional 1 local visitor (1 month lag) respectively. However, the DF cases decreased by 2.18×10^{-6} (95% CI: $-2.50 \times 10^{-6} : -1.87 \times 10^{-6}$) for every additional 1 foreign visitors in 3 months lag [Table 2].

4. Discussion

Climatic factors are found to have significant association with dengue cases in Badung District. Many possible mechanisms between climate and dengue were proposed by previous studies. Rising temperatures due to climate change led to shorter incubation period of mosquitoes, it impacted on the rapid and faster reproduction of the dengue vectors. Furthermore, optimal air temperature can accelerate mosquitos' growth and the incubation period of the DENV virus

(Huang et al. 2013, Tjaden et al. 2013, Zhang et al. 2016). Temperature ranges 25-27°C is optimal for the multiplication of mosquito vectors that it could change the pattern of interaction between infection diseases agent and human as the host (Goindin et al. 2015) as examined by previous studies about dengue fever transmission and environmental climate factors in Indonesia (Hasanah and Susanna 2018).

The optimum air humidity ranges from 70-80 percent which makes mosquitoes mobility more aggressive and increases the risk of viral transmission and the incidence of DHF disease. This association was signified by several studies (Astuti et al. 2019, Hasanah and Susanna 2018). This finding is also in line with the current study result in Badung district. However there was one study in South Sulawesi, Indonesia found otherwise stating that there is no significant influence of humidity on the lowering or increased incidence of DHF (Tosepu et al. 2018). The discrepancy is probably due to humidity variation in different topography of observed areas, moreover the location of South Sulawesi is closer to equator than Badung, Bali that it could drives different humidity characteristics and impacts.

Precipitation is one of the affecting factors for DF case in Badung district. The exploratory analysis for 0 – 3 months lags showed significant positive association. Similar result was reported from Neighbour country Thailand that precipitation together with temperature were indicated as the primary factors to drive dengue cases incidence (Langkulsen et al. 2020) as the presence of suitable precipitation could affects the humidity and initiates of mosquito-breeding place, as *Aedes aegypti* mosquitoes have high durability against high precipitation intensity . This finding also agrees with the study in Kerala, India that dengue occurrence during 2010 – 2014 was significantly associated with various lags of climatic factors (temperature, rainfall and humidity) and it showed spatial and temporal clustered

(Valson and Soman 2017). However, the different result was reported by study in Singapore that found significant negative association between weekly precipitation and dengue incidence (Koh et al. 2018). The difference might be caused by the different time unit analysis as Badung study used monthly data while Singapore study used weekly data.

The number of local and foreign tourists is the influencing factor of DF case in Badung district. The result implies the association between the local and foreign visitors to the incidence of DF. This findings agrees with the previous study stated that dengue transmission was accounted for 2 percent of all of diseases experienced by visitors returned from endemic areas such us Southeast Asia (Wilder-Smith 2012). This result is also supported by the research in South Korea to found that in majority of imported DF cases were correlated with overseas travellers (Cho et al. 2018).

Aside from the geographical originis of Indonesia as tropical area located in the equator that it is conducive to support the breeding of *Aedes aegypti* as a vector of the dengue virus. Dengue transmission in Indonesia could be exacerbating by increasing population mobility and uncontrolled urbanization flows. Study in 2010 concluded that mobility of local and foreign visitors to Bali was significantly contributed for the DF transmission, in particularly densely populated of Badung district (Ishikawa and Shimogawara 2019). It is supported by another study in Japan that the dengue case (genotype I DENV-1) found in Okinawa in 2014 was confirmed as imported from Bali, Indonesia (Saito et al. 2016). These findings magnify the urgency of strengthening the monitoring and protection of international border starting from the arrival. Study in Yogyakarta reported that the breeding sites for *Aedes spp* is higher in airport leads to the potential for DF outbreak (Satoto et al. 2018). Nevertheless, the local

residents and foreign visitors have been proven to have significant contribution to involve in dengue fever reduction program (Yoshikawa and Kusriastuti 2013).

The highly seasonal pattern of DF incidence found in Badung district agrees with the previous study of entire Bali Island (Dhewantara et al. 2019) also similar to the pattern of DF cases found in Timor Leste located in Timor Island (Wangdi et al. 2018). As the neighbors' Island of Bali, this phenomena is most likely caused by the similar climate condition of southern part of equator that leads to the similar DF incidence pattern. The seasonal pattern was coincided with the climatic dynamics, for instance the peaks always occurred around early months, and this condition is concurrent with the high intensity of reported precipitation in Badung that occurred around January – March. While, the lowest curve occurred around August – November that is coincided with the lowest reported humidity in this district.

The maps show that Abiansemal and Mengwi sub district frequently have highest dengue incidence from 2013 – 2019. According to Badung health profile, that these two sub-districts have the highest number of household: 39,663 and 22,124 respectively (PHO 2019). It could drive domestic transmission setting of DF incidences. While among 6 sub districts, Petang was consistently has the lowest IR during 7 years. It could be due to its elevation is the highest in Badung district around 2,075 meters above sea level, as reported by Badung Bureau of Statistics (BPS 2016) that lead to the unfavourable weather, temperature and humidity for mosquitoes vectors' proliferation. The DF cases were dominated by male patients in almost all districts during 2013 – 2019, most probably because in Badung district in majority male works as informal workers close to their house where water containers for mosquitoes breeding sites are available.

The present study have numbers of areas to be improved such as the inconsistent range of data and spatial units of analysis. For statistical analysis only covered 60 months (January 2013 – December 2017) at only at district level due to limited access to surveillance data. While for the visual geographical distribution, even though it covered 7 years (2013 – 2019) period however the spatial unit was only limited to 6 sub-districts that made it impossible to explore any further spatial pattern and analysis such as cluster and association. This study also has not integrated others potential risk variables into analysis such as health system and service accessibility, vector and larvae density, local population behaviour, population movement, transportation, interventions and others factors that directly and indirectly impacts on DF distribution.

Aside from the limitations stated, this study try to provide analysis that integrating both environmental (climate) and social determinant (visitors) in determining the risk for dengue spread. The result of this study could be the baseline information for developing comprehensive analysis on the impact of human movement on the emerging of infectious diseases.

5. Conclusion

The climate and tourism factors are the influencing factors for DF transmission dynamics in Badung District, therefore it need to be integrated for determining for DF monitoring and mitigation intervention in densely populated destination area.

Acknowledgement

The authors thanks to the Diseases Prevention and Control Division of Bali Province Health Office, Bureau of Meteorology, Climatology, and Geophysics (BMKG) and Badung

Regency Statistics Bureau (BPS) that have supported the research and article by providing the data.

References

- Astuti, E. P., Dhewantara, P. W., Prasetyowati, H., Ipa, M., Herawati, C. & Hendrayana, K. 2019. Paediatric dengue infection in Cirebon, Indonesia: a temporal and spatial analysis of notified dengue incidence to inform surveillance. *Parasites & vectors*, 12, 186. <https://doi.org/10.1186/s13071-019-3446-3> DOI: 10.1186/s13071-019-3446-3
- Bhatt, S., Gething, P. W., Brady, O. J., Messina, J. P., Farlow, A. W., Moyes, C. L., Drake, J. M., Brownstein, J. S., Hoen, A. G. & Sankoh, O. 2013. The global distribution and burden of dengue. *Nature*, 496, 504-507. <https://doi.org/10.1038/nature12060> DOI: 10.1038/nature12060
- Bps. 2016. *Social and Demography* [Online]. Badung Districts Bureau of Statistics. Available: <https://badungkab.bps.go.id/indicator/153/64/1/luas-wilayah-kabupaten-badung-dan-ketinggiannya-dari-permukaan-laut-menurut-kecamatan.html> [Accessed 30 November 2020].
- Brady, O. J., Gething, P. W., Bhatt, S., Messina, J. P., Brownstein, J. S., Hoen, A. G., Moyes, C. L., Farlow, A. W., Scott, T. W. & Hay, S. I. 2012. Refining the global spatial limits of dengue virus transmission by evidence-based consensus. *PLoS Negl Trop Dis*, 6, e1760. <https://doi.org/10.1371/journal.pntd.0001760> DOI: 10.1371/journal.pntd.0001760
- Cho, K.-H., Park, S.-Y., Lee, W.-C., Lee, M.-J. & Lee, J.-B. 2018. International Travel and Exotic Dengue Fever in South Korea from 2006 to 2015. *Japanese journal of infectious diseases*, 71, 378-381. <https://doi.org/10.7883/yoken.JJID.2017.553> DOI: 10.7883/yoken.JJID.2017.553
- Colón-González, F. J., Fezzi, C., Lake, I. R. & Hunter, P. R. 2013. The effects of weather and climate change on dengue. *PLoS Negl Trop Dis*, 7, e2503. <https://doi.org/10.1371/journal.pntd.0002503> DOI: 10.1371/journal.pntd.0002503
- Dhewantara, P. W., Marina, R., Puspita, T., Ariati, Y., Purwanto, E., Hananto, M., Hu, W. & Magalhaes, R. J. S. 2019. Spatial and temporal variation of dengue incidence in the island of Bali, Indonesia: An ecological study. *Travel medicine and infectious disease*, 101437. <https://doi.org/10.1016/j.tmaid.2019.06.008> DOI: 10.1016/j.tmaid.2019.06.008
- Fatmawati, F. & Sulistyawati, S. 2019. CLIMATE CHANGE AND DENGUE IN INDONESIA: A SYSTEMATIC REVIEW. *Epidemiology and Society Health Review (ESHR)*, 1, 29-40
- Goindin, D., Delannay, C., Ramdini, C., Gustave, J. & Fouque, F. 2015. Parity and longevity of *Aedes aegypti* according to temperatures in controlled conditions and consequences on dengue transmission risks. *PLoS One*, 10, e0135489. <https://doi.org/10.1371/journal.pone.0135489> DOI: 10.1371/journal.pone.0135489
- Hasanah & Susanna, D. 2018. Weather Implication for Dengue Fever in Jakarta, Indonesia 2008-2016. *KnE Life Sciences*, 184–192-184–192. <https://doi.org/10.18502/cls.v4i10.3719> DOI: 10.18502/cls.v4i10.3719
- Huang, X., Williams, G., Clements, A. C. & Hu, W. 2013. Imported dengue cases, weather variation and autochthonous dengue incidence in Cairns, Australia. *PLoS One*, 8, e81887. <https://doi.org/10.1371/journal.pone.0081887> DOI: 10.1371/journal.pone.0081887
- Ishikawa, H. & Shimogawara, R. 2019. Risk assessment of dengue autochthonous infections in Tokyo in summer, especially in the period of the 2020 Olympic Games. *Japanese Journal of Infectious Diseases*, JJID. 2019.094. <https://doi.org/10.7883/yoken.JJID.2019.094> DOI: 10.7883/yoken.JJID.2019.094

- Koh, Y.-M., Spindler, R., Sandgren, M. & Jiang, J. 2018. A model comparison algorithm for increased forecast accuracy of dengue fever incidence in Singapore and the auxiliary role of total precipitation information. *International journal of environmental health research*, 28, 535-552
- Langkulsen, U., Promsakha Na Sakolnakhon, K. & James, N. 2020. Climate change and dengue risk in central region of Thailand. *International journal of environmental health research*, 30, 327-335
- Matsui, T., Kinoshita, N., Maeki, T., Kutsuna, S., Nakamura, K., Nakamoto, T., Ishikane, M., Tajima, S., Kato, F. & Taniguchi, S. 2019. Dengue virus type 2 infection in a traveler returning from Saudi Arabia to Japan. *Japanese Journal of Infectious Diseases*, JJID. 2018.537.
<https://doi.org/10.7883/yoken.JJID.2018.537> DOI: 10.7883/yoken.JJID.2018.537
- Moh 2018. Indonesia Health Profile 2018. Jakarta: Ministry of Health.
- Pho 2019. Health Profile of Badung District 2019. In: (Pho), P. H. O. (ed.). The Government of Badung District.
- Pho, B. 2016. Health Profile of Bali Province 2016. In: (Pho), B. P. H. O. (ed.). Denpasar: Bali Province Health Office (PHO).
- Saito, M., Tamayose, M., Miyagi, K., Takaragawa, H., Tateyama, M., Tadano, M. & Fujita, J. 2016. Serologic and virologic studies of an imported dengue case occurring in 2014 in Okinawa, Japan. *Japanese journal of infectious diseases*, 69, 60-65.
<https://doi.org/10.7883/yoken.JJID.2015.061> DOI: 10.7883/yoken.JJID.2015.061
- Satoto, T. B. T., Listyantanto, A., Agustjahjani, S. D., Josef, H. K. & Widartono, B. S. 2018. Vertical transmission of dengue virus in the Yogyakarta airport area. *Environmental health and preventive medicine*, 23, 1-7. <https://doi.org/10.1186/s12199-018-0711-6> DOI: 10.1186/s12199-018-0711-6
- The Government, O. B. R. 2018. Badung Regency in Numbers 2018. Badung, Bali: The Government of Badung Regency.
- Tjaden, N. B., Thomas, S. M., Fischer, D. & Beierkuhnlein, C. 2013. Extrinsic incubation period of dengue: knowledge, backlog, and applications of temperature dependence. *PLoS neglected tropical diseases*, 7, e2207. <https://doi.org/10.1371/journal.pntd.0002207> DOI: 10.1371/journal.pntd.0002207.g002
- Tosepu, R., Tantrakarnapa, K., Nakhapakorn, K. & Worakhunpiset, S. 2018. Climate variability and dengue hemorrhagic fever in Southeast Sulawesi Province, Indonesia. *Environmental Science and Pollution Research*, 25, 14944-14952. <https://doi.org/10.1007/s11356-018-1528-y> DOI: 10.1007/s11356-018-1528-y
- Valson, J. S. & Soman, B. 2017. Spatiotemporal clustering of dengue cases in Thiruvananthapuram district, Kerala. *Indian Journal of Public Health*, 61, 74. DOI: 10.4103/ijph.IJPH_26_16. DOI: 10.4103/ijph.IJPH_26_16
- Van Panhuis, W. G., Choisy, M., Xiong, X., Chok, N. S., Akarasewi, P., Iamsirithaworn, S., Lam, S. K., Chong, C. K., Lam, F. C. & Phommasak, B. 2015. Region-wide synchrony and traveling waves of dengue across eight countries in Southeast Asia. *Proceedings of the National Academy of Sciences*, 112, 13069-13074
- Waggoner, J. J., Gresh, L., Vargas, M. J., Ballesteros, G., Tellez, Y., Soda, K. J., Sahoo, M. K., Nuñez, A., Balmaseda, A. & Harris, E. 2016. Viremia and clinical presentation in Nicaraguan patients infected with Zika virus, chikungunya virus, and dengue virus. *Clinical Infectious Diseases*, ciw589. <https://doi.org/10.1093/cid/ciw589> DOI: 10.1093/cid/ciw589
- Wangdi, K., Clements, A. C., Du, T. & Nery, S. V. 2018. Spatial and temporal patterns of dengue infections in Timor-Leste, 2005–2013. *Parasites & Vectors*, 11, 9.
<https://parasitesandvectors.biomedcentral.com/articles/10.1186/s13071-017-2588-4> DOI: 10.1186/s13071-017-2588-4
- Who 2011. *Comprehensive Guidelines for Prevention and Control of Dengue and Dengue Haemorrhagic Fever*, India, World Health Organization Regional Officer for South East Asia.

- Who. 2019. *Dengue and severe dengue* [Online]. Geneva: World Health Organization. Available: <https://www.who.int/en/news-room/fact-sheets/detail/dengue-and-severe-dengue>. [Accessed 16 January 2020].
- Wilder-Smith, A. 2012. Dengue infections in travellers. *Paediatrics and international child health*, 32, 28-32
- Wiradarma, I. G. a. B. A. & Somia, I. K. A. 2015. Hubungan antara perubahan iklim dengan jumlah kasus demam dengue (DD) di Denpasar, Bali tahun 2010-2015. <https://isainsmedis.id/index.php/ism/article/viewFile/423/353>.DOI: 10.15562/ism.v10i3.423
- Xiang, J., Hansen, A., Liu, Q., Liu, X., Tong, M. X., Sun, Y., Cameron, S., Hanson-Easey, S., Han, G.-S. & Williams, C. 2017. Association between dengue fever incidence and meteorological factors in Guangzhou, China, 2005–2014. *Environmental research*, 153, 17-26. <https://doi.org/10.1016/j.envres.2016.11.009>.DOI: 10.1016/j.envres.2016.11.009
- Xu, Z., Lau, C. L., Zhou, X., Fuimaono, S., Magalhães, R. J. S. & Graves, P. M. 2018. The extensive networks of frequent population mobility in the Samoan Islands and their implications for infectious disease transmission. *Scientific reports*, 8, 1-11. <https://doi.org/10.1038/s41598-018-28081-x>.DOI: 10.1038/s41598-018-28081-x
- Yoshikawa, M. J. & Kusriastuti, R. 2013. Surge of dengue virus infection and chikungunya Fever in bali in 2010: the burden of mosquito-borne infectious diseases in a tourist destination. *Tropical medicine and health*, 2011-05. <https://doi.org/10.2149/tmh.2011-05>.DOI: 10.2149/tmh.2011-05
- Zhang, Y., Wang, T., Liu, K., Xia, Y., Lu, Y., Jing, Q., Yang, Z., Hu, W. & Lu, J. 2016. Developing a time series predictive model for dengue in Zhongshan, China based on weather and Guangzhou dengue surveillance data. *PLoS neglected tropical diseases*, 10, e0004473. <https://doi.org/10.1371/journal.pntd.0004473>.DOI: 10.1371/journal.pntd.0004473

TABLES AND FIGURES

Figure 1: Geographical Distribution of Dengue Incidence Rates (IR) in 6 Sub-Districts in Badung 2013 – 2019

Table 1. Descriptive Frequency Distribution of Monthly Variables and Covariates 2013-2017

Variable and Covariates	Median	Mean	Min	Max	Standar Deviation
Case	140,5	178,15	27	796	149,25
IR	23.1	29,26	4,2	128,2	24,23
Temperature	27.5	27,41	25,6	29	0,81
Precipitation	102.5	154,09	0	620,1	157,1962
Humidity	80	79,98	75	85	2,213
Foreign Visitors	344,510.5	359047,80	232935	593.337	84372,31
Local Visitors	47,682.5	48893,60	15.224	94.847	16761,33

Figure 2: Decomposition of DF (IR/ 100000 Population) time series data in Badung District

Figure 3: The monthly average of case, IR, temperature, precipitation, humidity, local and foreign visitors

Table 2. Result of Multivariate preliminary analysis of selected variables using poisson regression

Model with Selected Variables	Coefficient	P value	Confidence Interval(95% CI)
α (Intercept)	-11.98101	0,000	-13.19878 : -10.76324
Average Temperature lag3	0.59095	0,000	0.5610301 : 0.6208698
Average Precipitation lag2	0.0005779	0,000	0.0003882 : 0.0007676
Average Humidity lag2	0.0143432	0,013	0.0030307 : 0.0256557
Local Visistors lag1	7.40×10^{-6}	0,000	5.88×10^{-6} : 8.91×10^{-6}
Foreign Visitors lag3	-2.18×10^{-6}	0,000	-2.50×10^{-6} : -1.87×10^{-6}

APPENDIX

Appendix 1: Result of Univariate preliminary analysis of selected variables using poisson regression and VIF identification using multivariate regression

Variables	VIF	IRR	95% CI	P value (poisson)	AIC	BIC
Without lag						
Average Temperature	1,14	1.332228	1.300285 : 1.364955	0,000	6198.693	6202.881
Average Precipitation	1,88	0.9995045	0.9993775 : 0.9996316	0,000	6684.254	6688.443
Average Humidity	1,80	1.044416	1.035398 : 1.053513	0,000	6647.66	6651.849
Local Visistors	1,18	1.00001	1.000008 : 1.000011	0,000	6452.497	6456.686
Foreign Visitors	1,18	0.9999994	0.9999992 : 0.9999996	0,000	6717.178	6721.366
With lag1						
Average Temperature lag1	1,16	1.581323	1.542961 : 1.620639	0,000	5257.154	5261.309
Average Precipitation lag1	1,83	1.001175	1.001053 : 1.001296	0,000	6293.349	6297.504
Average Humidity lag1	1,75	1.098213	1.088593 : 1.107919	0,000	6197.537	6201.692
Local Visistors lag1*	1,18	1.000011	1.000010 : 1.000012	0,000	6257.705	6261.86
Foreign Visitors lag1	1,17	0.9999984	0.9999981 : 0.9999986	0,000	6445.778	6449.933
With lag2						
Average Temperature lag2	1,15	1.774695	1.731259 : 1.819220	0,000	4456.405	4460.526
Average Precipitation lag2*	1,77	1.002338	1.002221 : 1.002455	0,000	5190.697	5194.818
Average Humidity lag2*	1,70	1.100248	1.090501 : 1.110083	0,000	6154.343	6158.463
Local Visistors lag2	1,18	1.00001	1.000009 : 1.000011	0,000	6271.487	6275.608
Foreign Visitors lag2	1,17	0.9999978	0.9999975 : 0.999998	0,000	6263.738	6267.858
With lag3						
Average Temperature lag3*	1,17	2.084046	2.032162 : 2.137255	0,000	3128.901	3132.987
Average Precipitation lag3	1,77	1.002119	1.002001 : 1.002238	0,000	5442.098	5446.184
Average Humidity lag3	1,70	0.9962442	0.9874804 : 1.005086	0.404	6577.703	6581.789
Local Visistors lag3	1,19	1.00001	1.000009 : 1.000011	0,000	6261.01	6265.096
Foreign Visitors lag3*	1,19	0.9999973	0.9999971 : 0.9999976	0,000	6105.728	6109.814
With selected lag						
Average Temperature lag3	1,51	2.084046	2.032162 : 2.137255	0,000	3128.901	3132.987
Average Precipitation lag2	2,07	1.002338	1.002221 : 1.002455	0,000	5190.697	5194.818
Average Humidity lag2	1,74	1.100248	1.090501 : 1.110083	0,000	6154.343	6158.463
Local Visistors lag1	1,38	1.000011	1.000010 : 1.000012	0,000	6257.705	6261.86
Foreign Visitors lag3	1,20	0.9999973	0.9999971 : 0.9999976	0,000	6105.728	6109.814

Appendix 2: Distribution of DF in 6 Sub Districts in Badung from 2013 - 2019

Sub-Districts	2013			2014			2015			2016			2017			2018			2019		
	Male (%)	Female (%)	IR/100000 Population	Male (%)	Female (%)	IR/100000 Population	Male (%)	Female (%)	IR/100000 Population	Male (%)	Female (%)	IR/100000 Population	Male (%)	Female (%)	IR/100000 Population	Male (%)	Female (%)	IR/100000 Population	Male (%)	Female (%)	IR/100000 Population
Abiansemal	52.90	47.10	343.95	53.96	46.04	292.70	40.08	59.92	271.76	51.96	48.04	923.53	54.68	45.32	221.50	46.30	53.70	58.67	54.55	45.45	142.95
Kuta	55.96	44.04	114.65	47.62	52.38	107.50	60.17	39.83	117.65	57.32	42.68	159.58	62.22	60.00	42.75	62.50	37.50	44.58	57.50	42.50	145.32
Kuta Selatan	52.09	47.91	195.50	54.41	45.59	491.90	54.36	45.64	563.75	57.26	42.74	794.23	60.20	39.80	126.76	62.50	37.50	33.98	55.89	44.11	173.85
Kuta Utara	54.68	45.32	294.70	58.67	41.33	187.80	55.45	44.55	259.65	53.44	46.56	354.00	57.89	42.11	130.40	54.29	45.71	51.96	54.92	45.08	190.72
Mengwi	50.20	49.80	584.76	56.22	43.54	326.30	53.80	46.20	489.62	57.79	42.21	952.78	52.58	47.42	222.10	56.25	43.75	97.02	58.97	41.03	306.52
Petang	62.86	37.14	133.79	63.89	36.11	138.00	58.82	41.18	130.87	51.69	48.31	343.50	56.67	43.33	116.10	60.00	40.00	38.82	66.67	33.33	58.32