1 Genomic epidemiology and longitudinal sampling of ward wastewater
2 environments and patients reveals complexity of the transmission dynamics of
3 *bla*_{KPC}-carbapenemase-producing Enterobacterales in a hospital setting

4
5 Stoesser N^{a, b, *}, George R^{c, *}, Aiken Z^c, Phan HTT^d, Lipworth S^a, Wyllie DH^e, Quan
6 TP^a, Mathers AJ^f, De Maio N^g, Seale AC^h, Eyre DW^{a, b}, Vaughan A^a, Swann J^a, Peto
7 TEA^{a, i}, Crook DW^{a, b, i}, Cawthorne J^c, Dodgson A^c, Walker AS^{a, b, i}, and the TRACE
8 Investigators' Group

9
10 ^a Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

11 ^b NIHR Health Protection Research Unit in Healthcare Associated Infections and
12 Antimicrobial Resistance at University of Oxford in partnership with Public Health
13 England, Oxford, United Kingdom

14 ^c Manchester University NHS Foundation Trust, Manchester, United Kingdom

15 ^d University of Southampton, Southampton, United Kingdom

16 ^e United Kingdom Health Security Agency, United Kingdom

17 ^f University of Virginia, Charlottesville, United States of America

18 ^g EMBL-European Bioinformatics Institute, Cambridge, United Kingdom

19 ^h University of Warwick, Coventry, United Kingdom

20 ⁱ NIHR Oxford Biomedical Research Centre, Oxford University Hospitals NHS
21 Foundation Trust, John Radcliffe Hospital, Oxford, United Kingdom

22
23 *Contributed equally
24

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

25 **Keywords:** Carbapenemase-producing Enterobacterales, genomic epidemiology,
 26 *bla*_{KPC}, wastewater sites, healthcare-associated
 27
 28 **Corresponding author:** Nicole Stoesser
 29 **Corresponding address:** Department of Microbiology (Research), Level 7, John
 30 Radcliffe Hospital, Headley Way, Headington, OX3 9DU
 31 **Affiliation:** Nuffield Department of Medicine, University of Oxford, Oxford UK and
 32 Oxford University Hospitals NHS Foundation Trust, Oxford, UK
 33 **e-mail:** nicole.stoesser@ndm.ox.ac.uk

ABSTRACT

Background

Healthcare-associated wastewater reservoirs and asymptomatic gastrointestinal patient colonisation by carbapenemase-producing Enterobacterales (CPE) can be important in nosocomial CPE dissemination and infection. We characterised these niches and within-niche diversity in a *bla*_{KPC}-associated CPE (KPC-E) endemic healthcare setting, to better understand transmission potential.

Methods

We systematically sampled wastewater sites and patients across three units (six wards) over 6-12 months in 2016 in a KPC-E endemic hospital. We used Illumina sequencing to characterise up to five isolates per sample. Recombination-adjusted phylogenies were used to define genetically related strains; assembly and mapping-based typing approaches were used to characterise antimicrobial resistance gene and insertion sequence profiles, and *Tn4401* types/target site sequences. The wider accessory genome was evaluated in a subset of the largest clusters, and those crossing niches.

Findings

Wastewater site KPC-E-positivity was substantial (101/349 sites [28.9%] positive, 319/4,488 [7.1%] sampling events positive); 183/4,425 (4.1%) of patients were CPE culture-positive over the same timeframe. 13 species and 109 KPC-E strains were observed across niches, and 24% of wastewater and 26% of patient KPC-E-positive samples harboured ≥ 1 strain. Most diversity was explained by the individual niche, suggesting highly localised factors are important in selection and spread. *Tn4401*+target site sequence (TSS) diversity was greater in wastewater sites ($p < 0.001$), suggesting these might favour *Tn4401*-associated transposition/evolution

and dissemination. Shower/bath and sluice/mop-associated sites were more likely to be KPC-E-positive (Adjusted Odds Ratio [95% CI]: 2.69 [1.44-5.01], $p=0.0019$ and 2.60 [1.04-6.52], $p=0.0410$, respectively). Different strains had different transmission and *bla*_{KPC} dissemination dynamics.

Interpretation

There may be substantial KPC-E colonisation of wastewater sites and patients in KPC-E-endemic healthcare settings. Niche-specific factors, and different strains with different transmission dynamics influence carbapenemase gene dissemination. New transmission models incorporating complex, multi-level dynamics are needed to better quantify CPE dissemination to inform interventions and reduce transmission.

Funding

This study was supported by the National Institute for Health Research, UK.

RESEARCH IN CONTEXT

Evidence before this study

We searched PubMed to identify previous studies using whole genome sequencing (WGS) to characterise in-hospital dissemination and persistence of carbapenemase-producing Enterobacterales (CPEs) or the presence of CPEs in hospital wastewater reservoirs. In our search (01/Jan/1996-30/Sep/2021) we used the terms (((CPE) OR (KPC)) AND (healthcare-associated OR nosocomial OR hospital-associated)) AND (genom*) AND ((environment OR patient))). We had no restrictions on language. We found sixty-six studies, but these were limited by: (i) evaluating single species or lineages only (n=34), (ii) including patients or environmental outbreaks only, and not both (n=23), (iii) sampling only small numbers (n<50) of patients or environmental sites (n=26), or (iv) performing no evaluation of within-niche diversity (n=64). 13 studies were not relevant (e.g. investigating non-hospital locations, *ex vivo* or *in vivo* evolution), two studies did not use WGS, and one study was not primary research.

Added value of this study

We provide a comprehensive assessment of healthcare-associated CPE dissemination and persistence in hospital wastewater reservoirs through systematic sampling of a large number of environmental sites (n=349 sites, n=4,488 samples) and patients (n=4,425) over 6-12 months in a single hospital. For all CPE-positive environmental samples (n=319) and a subset of CPE-positive patient samples (n=97/399 [24.3%] samples from 76/183 [41.5%] CPE-positive patients), we sampled up to five CPE isolates per sample to capture within-niche diversity, and used short-read Illumina WGS (n=1,732 isolates successfully sequenced) to characterise within-niche diversity, changes in colonisation over time, and genetic overlap between patient and environmental niches.

96 **Implications of all the available evidence**

97 Through a detailed and resource-intensive study we captured the dynamics of seeding
 98 and dissemination of important carbapenemase genes amongst patients and
 99 environmental reservoirs within a hospital. We found differential colonisation and
 100 dissemination dynamics for different species and lineages within and between niches.
 101 Improved approaches incorporating variable within-niche diversity, accessory
 102 distances and horizontal gene transfer in transmission evaluations are required to
 103 better understand CPE dissemination and persistence, in order to direct interventions
 104 limiting transmission.

INTRODUCTION

Carbapenemase-producing Enterobacterales (CPE) are a global health threat¹, and commonly associated with multi-drug resistance, rendering treatment of CPE infections difficult. Major transmissible global carbapenemases include the metallo-beta-lactamases (*bla*_{NDM}, *bla*_{VIM}, *bla*_{IMP}), some oxacillinases (*bla*_{OXA-48/48-like} variants), and the *Klebsiella pneumoniae* carbapenemase (KPC, encoded by *bla*_{KPC})². Intra- and inter-species horizontal transfer of these genes is facilitated by mobile genetic elements such as transposons and plasmids, resulting in rapid dissemination of carbapenem resistance^{3, 4}.

CPE have been isolated from human and animal gastrointestinal tracts, and from sewage, rivers and sink drains⁵⁻⁷. Recent studies have highlighted that hospital wastewater sites may act as a CPE reservoir⁵, but the diversity within these niches, genetic overlap with patient isolates and likely directionality and rates of transmission remain to be elucidated. Studies evaluating the contribution of patient-to-patient transmission as the sole explanatory factor explaining transmission have generally been unable to robustly explain the majority of transmission events, suggesting that patient, staff and/or environmental reservoirs remain insufficiently sampled to account for the diversity observed⁸. Few studies have considered within-niche diversity of strains by sampling multiple isolates per individual and/or site⁹⁻¹¹.

The Manchester University NHS Foundation Trust (MFT; Manchester, UK), has experienced on-going transmission and patient acquisition of multi-species CPE, predominantly *bla*_{KPC}-positive Enterobacterales (KPC-E), since 2009. Following a KPC-*Escherichia coli* outbreak on two cardiac wards in the Manchester Royal

Infirmiry (MRI; part of MFT), and subsequent ward closure and plumbing infrastructure replacement¹², we undertook a prospective study systematically sampling all wastewater sites on six wards for 6-12 months alongside patient rectal screening/clinical sampling, and used an anonymised electronic database to characterise patient admission, ward movement, sampling profiles and culture results. Multiple colonies were isolated from KPC-E-positive samples to define within-niche diversity, and whole genome sequencing (Illumina) of individual isolates was used as to characterise this diversity and consider modes of evolution and transmission.

METHODS

Study setting and epidemiological data

MFT is a large, regional referral centre in northwest England, UK, managing >10,000 patients/year. In response to the regional emergence of KPC-E¹³, the Trust implemented an extensive Infection Prevention and Control (IPC) programme, consistent with UK CPE guidelines (published in 2014)¹⁴. Despite this, in April 2015 there was a large *bla*_{KPC}-*E. coli* outbreak in the cardiac unit¹², leading to its closure (September 2015-January 2016) and complete refurbishment, including plumbing replacement. Subsequently systematic, prospective wastewater site sampling was undertaken (described further below).

To evaluate patient-level microbiological data and hospital stays, MFT electronic bacteriology records were linked, based on NHS numbers, to patient administration data, and anonymised. For this study, we analysed anonymised patient data from 1st January 2016-31st December 2016 inclusive, and included only those patients with exposure to any one of the three units (i.e. six study wards) during that time.

155

156 *Environmental sampling and laboratory processing*

157 All sink/drain/wastewater sites on six wards in three units were sampled: the cardiac
158 unit (wards 3 [W3] and 4 [W4]); the geratology unit (wards 45 [W45] and 46 [W46]);
159 and the acute medicine unit (wards AM1 and AM2) (Table S1 shows sampling site
160 designations). W3/W4 wastewater sites were sampled fortnightly on rotation from 8th
161 January 2016-28th December 2016. W45/W46 and AM1/AM2 wastewater sites were
162 similarly sampled, but from 18th July 2016-31st December 2016. Wastewater site
163 types are described in Appendix Methods.

164

165 Environmental sampling was carried out by aspirating ~20mls of wastewater from
166 sink P-traps, shower drains or toilets and performing enrichment-based culture (see
167 Appendix methods). Multiplex, real-time, quantitative PCR (for *bla*_{NDM}, *bla*_{KPC},
168 *bla*_{OXA-48}¹²) was performed on broths following incubation. 10µl of any *bla*_{KPC}-
169 positive-broth were streaked onto CPE-selective agar plates (Chromid CARBA) and
170 re-incubated (aerobically, 37°C).

171

172 *Patient sampling and laboratory processing*

173 Routinely collected patient clinical samples were processed using standard operating
174 procedures in line with UK Standards¹⁵. Rectal swabs were screened for CPE using
175 the Cepheid Xpert Carba-R assay or an in-house multiplex PCR (identifying *bla*_{KPC},
176 *bla*_{NDM}, and *bla*_{OXA-48}) from November 2014. Species identification of isolates was
177 performed by MALDI-ToF (Bruker Biotyper, Bruker MD); antimicrobial
178 susceptibility testing was performed as per EUCAST guidelines¹⁶.

179

Analyses of within-niche diversity

To evaluate within-niche diversity in the environment and patients, up to five different Enterobacterales colonies were individually sub-cultured; *bla*_{KPC}-positivity in sub-cultures was re-confirmed by PCR and species identification performed by MALDI-ToF prior to storage in nutrient broth at -80°C.

DNA extraction and isolate sequencing

DNA was extracted from frozen sub-culture stocks with the QuickGene kit, as per the manufacturer's instructions, with an additional mechanical lysis step post-chemical lysis (6m/s at 40secs x2 FastPrep, MPBio). Isolates were sequenced on the Illumina HiSeq 4000 generating 150bp paired-end reads.

Sequence read processing and analysis

Sequencing reads were initially trimmed/quality controlled¹⁷; species identification was performed with Kraken¹⁸. If at least 50 sequenced isolates were of a single species, reads were mapped to species-specific references (Appendix Methods). Variant calling and filtering was performed, and consensus fasta sequences of variant calls with respect to the reference generated. Recombination-adjusted phylogenies by species were created from core chromosomal single nucleotide variants (SNVs)(padded to the length of the reference genome prior to analysis) using IQtree¹⁹ (flags: -m GTR+G -blmin 0.00000001 -t PARS) and then ClonalFrameML²⁰ (default parameters). A distance of ≤ 400 SNVs was used as a threshold to define strains (threshold used similar to²¹; Fig.S2A-E for most common species identified). For detailed sub-strain analysis, sub-strains were defined on the basis of accessory

204 genome clusters characterised using Panaroo (default parameters including --clean-
205 mode strict²²) and clustered using the R heatmap package.

206

207 Isolates were assembled using SPAdes (v3.6)²³ with default parameters. Multi-locus
208 sequence types were derived using BLASTn to relevant species-specific databases
209 hosted at <https://pubmlst.org> (Achtman 7-gene MLST for *E. coli*). AMR genes were
210 identified using ARIBA²⁴ with the CARD database²⁵. *bla*_{KPC} genes, any *Tn4401*
211 transposon context and 5-bp target site sequences reflecting *Tn4401* insertion
212 signatures were characterised using TETyper²⁶, using a *Tn4401b* reference sequence,
213 a 5bp flank length and the default structural and SNV profiles.

214

215 For the analysis of *K. pneumoniae* strain 9 (see Results), putative transmission
216 networks were inferred first using SCOTTI²⁷ (Appendix Methods), and then
217 heuristically, linking isolates identified as clusters on the basis of accessory
218 component profiles and geographic/temporal overlap (Appendix Methods).

219

220 Sequence data have been deposited in NCBI's GenBank (BioProject accession
221 PRJNA768622).

222

223 *Statistical analyses and data visualisation*

224 Descriptive statistics were calculated in R v3.6.2 or STATA v16.1. To conduct
225 permutational analyses of variance, we used the *adonis* function from the vegan R
226 package (v2.5-7)²⁸ on a matrix of pairwise Gower distances based on genetic features
227 (i.e. composite of species-strain type, AMR gene profile, IS profile, and
228 *Tn4401*+flanking sequence profiles), which was calculated using the *daisy* function in

the R cluster package. To evaluate the impact of unit location or environmental site type on the odds of CPE positivity, we used logistic regression with robust standard errors clustered by environmental site sampled, using the R rms package (v6.2-0)²⁹. Data visualisation was using the ggplot2³⁰ and scatterpie³¹ (v.0.1.5) R packages, and BioRender (www.biorender.com).

Ethical approvals

This study was performed as part of a Trust board-approved and UK Health Security Agency (formerly Public Health England) long-term outbreak response, and ethics approval was not required under NHS governance arrangements; all analyses were undertaken using anonymised data. This was reviewed and agreed by the Joint Research Office of the Oxford University Hospitals NHS Foundation Trust and the University of Oxford (who oversee research governance, ethics and assurance for our institution), and who deemed that neither sponsorship nor research ethics review was required. Of note, sample and patient numbers represented in the figures/supplementary data are study-specific labels that are not patient-identifiable.

RESULTS

Environmental sampling revealed high prevalence of wastewater site KPC-E colonisation

349 sites across the six study wards (i.e. three units) were sampled a total of 4,488 times (Fig.1): Of these, 23/76 (30%) sites and 111/1,564 (7.1%) sampling events on the cardiac unit were positive for KPC-E, 38/129 (29%) sites and 86/1,309 (6.6%) sampling events on the acute medicine unit and 40/144 [28%] and 122/1,615 (7.6%) on the geratology unit (Fig.2, top three panels). 248/349 (71.1%) sites were never

identified as KPC-E-positive (Fig.S3); 7/349 (2.0%) sites appeared more likely to be persistently KPC-E positive (≥ 10 consecutive KPC-E-positive sampling events) and interestingly five of these were medicines/treatment room handwash basin drains likely only accessible to healthcare staff (A59, F45, F113, T43, T44; Fig.S4). The proportion of positive sites did not vary significantly over time (Fig.2, bottom panel). Two utility room sink drains on the cardiac unit (across W3, W4) were positive for KPC-E a day after W3 reopened to patient admissions (11th January 2016) following closure and complete plumbing replacement over three months; thereafter the unit was rapidly re-colonised (Fig.2)¹². Shower/bath drain sites and sluices/sluice sinks/mop sinks were significantly more likely to be positive for KPC-E than others (Adjusted Odds Ratio [AOR] (95% CI): 2.69 (1.44-5.01), $p=0.0019$ and 2.60 (1.04-6.52), $p=0.041$, respectively), and toilet water sampling less likely to be positive (AOR (95% CI): 0.28 (0.10-0.78), $p=0.015$); there was no evidence of a unit-dependent difference (Table 1).

Approximately one in twenty patients was affected by CPE, with most cases reflecting rectal colonisation with KPC-producing K. pneumoniae, E. coli or E. cloacae

Of 2,172 Enterobacterales positive-samples (obtained from 943 patients) and 2,000 with an associated carbapenem susceptibility result (172 carbapenem susceptibility results missing), 399/2,000 (20.0%) were carbapenem-resistant Enterobacterales (CRE; i.e. had intermediate or resistant susceptibilities) (Figs.1, S5). For 344/431 (79.8%) CRE isolates cultured from these 399 CRE-positive samples carbapenemase PCR data were available: All contained at least one known carbapenemase resistance gene, and 274 (79.7%) were KPC-E (Table S2, henceforth denoted CPE). These CPEs were cultured from 183 patients (4.1% of all patients overall, 8.4% of all patients with

an Enterobacterales-positive culture) during the study period (median=1, IQR: 1-2, range: 1-17 CPE isolates). Amongst these patients, CPE-positive samples represented a median of 10% (IQR: 5-20%) of all samples taken and cultured.

The majority of confirmed CPE-positive samples were rectal screens (n=318/399 [79.7%]); others were non-screening clinical specimens (81/399 [20.3%] including blood cultures [n=4], urine [45], respiratory [20], wound/intra-abdominal [12]) (Table S3). 16/183 individuals were positive only on clinical culture (9%), 141 only on rectal screen (77%), and 26 (14%) on both. By routine laboratory methods, at least 16 species and eight genera were represented amongst CPEs, with *K. pneumoniae*, *E. coli* and *E. cloacae* complex predominating (364/431 [84.5%]). Notably, the proportions of carbapenem-resistant cases within these top species varied (n=239/464 [51%] vs 88/1,083 [8%] vs 37/86 [43%] respectively; Fisher's exact test, p<0.001, Fig.S5).

The proportion of patient CPE-positive cultures was lower than that of environmental site samples, typically ~3% (Fig.3). Occasional peaks in prevalence were observed, consistent with outbreaks (e.g. week 31, acute medicine unit). CPE prevalence amongst patient cultures on the refurbished cardiology ward was 0% (97.5% CI: 0-0.02) for the first 10 weeks post-reopening.

Structuring of diversity by niche, with most diversity explained by the individual niche

Overall, amongst successfully sequenced isolates (n=1,732) across patients and environmental reservoirs, at least 13 Enterobacterales species harboured *bla*_{KPC}, with 109 unique strains represented (and a further ten species not characterised at the strain

level because of small numbers, see Methods). Of the 109 strains, 61 (56%) were found only in the environment, 27 (24%) only in patients and 21 (19%) in both patients and the environment (Fig.4); the median strain cluster size was 5 isolates (range: 1-122), and the median number of unique niches (i.e. unique patients or wastewater sites) affected by a strain was 2 (range: 1-23) (Fig.4A, Fig.4B). A much greater proportion of all identified *Tn4401*/TSS types was observed in the environment versus patients (118/121 [98%] of types versus 9/121 [7%] respectively; Fig.S6; Fisher's exact test, $p < 0.001$), and all the isolates with >1 set of TSSs were seen in environmental sites, consistent with these representing a more favourable niche for *Tn4401*-associated transposition/evolution and dissemination.

For colonised niches (i.e. patient or environmental), using a metric of genomic diversity based on strain type, AMR gene profile, IS profile, and *Tn4401*+ 5bp target site sequence profiles, most variance was explained by the specific individual niche sampled ($R^2=46.0\%$; $p < 0.001$), versus small but significant contributions made by reservoir type (patient vs specific environmental niche; $R^2=5.9\%$; $p < 0.001$) and unit location ($R^2=3.0\%$; $p < 0.001$). For the fifteen examples where exactly the same strain, AMR gene profile, IS profile, plasmid profile and *Tn4401*/TSS type were seen in patient and environmental niches, the temporal/geographic relationship of these suggested: patient $>$ environment transmission in 14/31 (45%) events, environment $>$ patient transmission in 2/31 (7%) events, environment $>$ environment transmission in 4/31 (13%) events, and patient $>$ patient transmission in 11/31 (35%) events (test of equality of proportions, $p < 0.001$).

328 *Environmental and patient niches harbour diverse KPC-E within single samples and*
329 *over time*

330 Environmental sites showed evidence of major strain-level diversity (Fig.5): in the
331 101/349 sites from which KPC-E was successfully sequenced at any timepoint,
332 78/319 (24.4%) KPC-E-positive samples had >1 KPC-E strain, namely 52/319
333 (16.3%) with two strains, 16/319 (5.0%) with three, 9/319 (2.8%) with four and 1/319
334 (0.3%) with five strains. Over the study period, colonised environmental sites had a
335 median of 3 (range: 1-17, IQR: 2-5) KPC-E strains. 49/101 sites (49%) were
336 apparently colonised by a single KPC-E strain only, of which 38 were at single
337 sampling-points, with 11 sites showing persistent colonisation (≥ 2 timepoints positive
338 with the same strain).

339

340 For patient samples, based on routine laboratory microbiology, evidence of mixed-
341 species KPC-E colonisation/infection was found in 34/399 (9%) samples from 27/183
342 (15%) patients; these were predominantly *E. coli*/*K. pneumoniae* mixtures (n=21
343 samples; maximum two species identified). To evaluate within-host diversity with
344 genomic rather than species-level resolution, in a subset of 76 patients, strains were
345 sequenced from at least one CPE-positive sample during the study period; for 97
346 samples, multiple colonies were picked. In total 463 sequenced colonies from 97
347 samples were available for evaluation of within-patient diversity; 25/97 samples
348 (26%) had more than one strain per sample, namely 17/97 samples (18%) with two
349 KPC-E strains, 6/97 (6%) with three strains and 2/97 (2%) with four strains (Fig.5) -
350 i.e. considerably more diversity identified than based on routine laboratory species
351 identification data (26% vs 9% samples; Fisher's exact test, $p < 0.001$). Longitudinal
352 switches in strain composition of KPC-E-positive samples was common, occurring in

353 11/21 (52%) KPC-E-positive longitudinal patient samples and 108/202 (53%) KPC-
354 E-positive longitudinal wastewater site samples.

355

356 *Phylogenetic analysis reveals distinct population biology for different KPC-E strains*

357 To investigate different modes of *bla*_{KPC} dissemination, we characterised in detail
358 several clusters, including the largest environmental-only cluster (*E. cloacae* strain 10
359 [n=122 isolates]), a large cluster involving patients and the environment (*K.*
360 *pneumoniae* strain 9 [n=116 isolates], and a cluster in which signatures of Tn4401
361 transposition appeared highly frequent (*K. pneumoniae* strain 11 [n=106 isolates]).

362

363 *Rapid churn of diversity in restricted environmental niches observed at the*

364 *chromosome, plasmid, IS, and AMR gene level - E. cloacae strain 10*

365 *E. cloacae* strain 10 (eclo10, ST32) appeared a restricted but persistent coloniser of
366 three co-located sink drains on a geratology ward (in a drug room clinical handwash
367 basin drain [F45] and utility sink drain [F46], and a separate treatment room clinical
368 handwash basin drain [F47]) over ~5 months (18/Jul/2016-20/Dec/2016) (Fig.6A). At
369 the core chromosomal level, strains appeared largely structured by site, with 7 SNVs
370 separating the majority of isolates in F45 from those in F47, and 14 SNVs those in
371 F46 from F45 (Fig.6A). There was evidence of rapid churn in AMR gene and IS
372 content (111 and 36 different profiles respectively; Fig.6A), and all isolates in F47
373 had acquired an additional set of IncHI2/HI2A replicons on a background of FIB/FII,
374 except for 1954277_1_4, consistent with plasmid gain/loss (Fig.6B). Although the
375 Tn4401 type and flanking sequences were largely identical (Tn4401a-ATTGA-
376 ATTGA in 105/122 (86%) isolates across sinks), there was evidence of unique

additional *Tn4401* deletions occurring in multiple isolates at four distinct timepoints in the three sites (Fig.6C).

Rapid patient-patient transmission - K. pneumoniae strain 9

K. pneumoniae strain 9 (kpne9, ST252) was highly related (Fig.S7; maximum 6 SNVs across strain, 68 isolates within 0 SNVs of each other), and spread amongst 21 patients across three wards on two units (acute medicine AM1 and AM2 and geratology W46), with a dense outbreak in Aug 2016, most consistent, at least initially, with direct/indirect patient-patient transmission (Fig.7). *bla_{KPC}* was consistently nested in *Tn4401*-ATTGA-ATTGA, except for one isolate which acquired a *Tn4401*-associated mutation (isolate: 2216698_14, C4620T), and another in which the right target site sequence (TSS) could not be identified (2035791_11).

Amongst the 106 isolates there was evidence of four transient plasmid replicon acquisitions on a background of the stable presence of a set of IncFIB/IncFII replicons (Fig.S7). Transmission probabilities inferred by SCOTTI were weak (see Appendix; Fig.S8), as previously⁸, but using accessory cluster profiling to attempt further discrimination amongst strains suggested that there were several major transmission clusters (Fig.S9), with some individuals involved in several of these (Fig.7). A single shower drain site was involved (A13, serving a particular bay with four beds), which was likely a seeded bystander from affected patients, but may have contributed to transmission of sub-strain 3g (Fig.7).

Cross-unit dissemination with evidence of significant Tn4401-associated bla_{KPC}

transposition - K. pneumoniae strain 11

Given that Tn4401 transposition is associated with *bla*_{KPC} dissemination²¹, we also evaluated whether the number of different Tn4401-TSS types was different across strains, noting that the observed frequency of different Tn4401-TSS combinations was higher in *K. pneumoniae* strain 11 (kpne11, ST11) than in other strains (Fig.S10). Kpne11 represented two sub-clusters, separated by ~40SNVs (Fig.7, designated “cluster 1” and “cluster 2”), both of which were isolated from patients and wastewater sites across ward settings, including the cardiac unit after plumbing replacement. Within both clusters of isolates there were ten different Tn4401a flanking signatures observed, only one of which was seen in the other >1700 isolates sequenced in the study (Tn4401a-1_ATTGA_ATTGA, the most common flanking signature), and therefore most strongly consistent with the occurrence of multiple transposition events within this *K. pneumoniae* strain. Similarly plasmid replicon gain/loss events were observed (Col, IncN, IncR and IncFIA); all isolates contained IncFIB and IncFII replicons.

DISCUSSION

This work highlights the substantial CPE diversity and flux that can occur in both human and wastewater niches in healthcare settings. Many CPE strains have unique features likely to make generalising results from analyses of transmission, evolution and gene sharing of *bla*_{KPC} amongst these species very challenging. Although some sites are persistently colonised with individual strains over a period of months, many appear to be only transiently colonised. Sites particularly associated with CPE colonisation included sluices, sluice sinks, mop sinks and shower/bath drains, supported by other studies⁵, and consistent with these representing a major interface between human faeces and premise plumbing. Toilets were less likely to be CPE-

426 positive, perhaps because of regular flushing or particular cleaning protocols;
 427 consistent with this, several studies have demonstrated that daily bleach application
 428 decontaminated wastewater sites^{32, 33}.
 429
 430 At the strain-level, likely transmissions occurred from patient-patient, environment-
 431 patient, patient-environment and environment-environment; however, our study
 432 highlights the difficulty in accurately tracking strains across niches, and the detailed
 433 and dense sampling effort required to understand within-sample diversity and
 434 longitudinal flux. This difficulty has also been demonstrated in other recent work
 435 investigating *bla*_{OXA-48}, where only patient-based sampling was undertaken and most
 436 *bla*_{OXA-48}-associated transmission events either remained non-delineated or were only
 437 weakly supported by joint phylogenetic/epidemiological inference⁸. Our study clearly
 438 shows that different species and lineages represent different epidemiological risks and
 439 have their own behaviours, and a focus on single, clonal strains as mediators of AMR
 440 gene transmission is not relevant in this and potentially many other settings where
 441 important AMR genes have become endemic.
 442
 443 The importance of evaluating asymptomatic CPE carriage, characterising within-host
 444 diversity of strains, considering rapid acquisition/loss of strains over short
 445 timeframes, and including environmental sites in transmission networks has been
 446 partly demonstrated in previous studies investigating ESBL-*E. coli*^{10, 34} and
 447 *Enterococcus* spp.³⁵, but generally these features are not jointly investigated because
 448 doing so is highly resource-intensive. However, without this effort, such as in studies
 449 which consider only single or infrequent sampling timepoints, and do not investigate

asymptomatic colonisation or consider within-site diversity³⁶, our study highlights that relevant events may easily be missed.

There are several limitations to this work. We undertook IPC-led real-time collection of patient isolates; sample and data capture for this process was not 24/7, and we therefore only evaluated a subset of all the patient-associated CPE isolates. We were not resourced to perform long-read sequencing to enable plasmid sequence reconstruction, and plasmid replicon or *bla*_{KPC}-contig profiling is relatively low-resolution²¹; we therefore avoided a detailed plasmid analysis which would be highly relevant to evaluating horizontal *bla*_{KPC} transmission. However, even with relatively low-resolution approaches, the diversity in *bla*_{KPC} contexts was substantial, supporting high rates of HGT. A further limitation is the absence of metadata capturing relevant selection pressures in each niche, including cleaning protocols for environmental reservoirs and drug prescription/antimicrobial stewardship data for patients. Although we only investigated one healthcare setting, the polyspecies, polyclonal phenomenon we observed has been seen in other centres and with other carbapenemase genes^{4,8}. We focused only on Enterobacterales and may have missed wider HGT of carbapenemase genes to other bacterial orders. In line with this, at least 23 *Pseudomonas* spp., five *Stenotrophomonas maltophilia* and two *Acinetobacter* spp. that were PCR-positive for *bla*_{KPC} or *bla*_{NDM} were inadvertently picked up on rectal screening.

Further optimisation of methods to evaluate multi-level genetic transmission and within-sample diversity without individual colony-level characterisation is needed. Holistic approaches using fully reconstructed chromosome/plasmid assemblies to

475 quantify the diverse types of genomic transmission (including clonal transmission,
476 horizontal transfer at the gene, transposon and plasmid-level) are also required to
477 better understand what facilitates emergence, selection and persistence of different
478 lineages. This information will improve our understanding of how to intervene to
479 limit the transmission of drug-resistant Enterobacterales in healthcare settings, and
480 avoid incorrect inference.

Declaration of interests

DWE has received lecture fees from Gilead, outside the submitted work.

Acknowledgements

We are grateful to the microbiology staff and infection prevention and control teams at Manchester University NHS Foundation Trust, and to Alvaro San Millán Ricardo León-Sampedro for helpful discussion.

This study was funded by the National Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance (NIHR200915), a partnership between the UK Health Security Agency (UKHSA) and the University of Oxford, and was supported by the NIHR Oxford Biomedical Research Centre (BRC). The computational aspects of this research were funded from the NIHR Oxford BRC with additional support from the Wellcome Trust Core Award Grant Number 203141/Z/16/Z. The views expressed are those of the author(s) and not necessarily those of the NHS, NIHR, UKHSA or the Department of Health and Social Care. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

DWE is a Robertson Foundation Fellow and an NIHR Oxford Biomedical Research Centre Senior Fellow. NS is an Oxford Martin Fellow and an NIHR Oxford Biomedical Research Centre Senior Fellow. DWC, TEAP and ASW are NIHR Senior Investigators.

The Transmission of Carbapenemase-producing Enterobacteriaceae (TRACE)
Investigators' Group is made up of the following (listed alphabetically, includes
several of the authors also listed by name in the main author list): Zoie Aiken,
Oluwafemi Akinremi, Aiysha Ali, Julie Cawthorne, Paul Cleary, Derrick W. Crook,
Valerie Decraene, Andrew Dodgson, Michel Doumith, Matthew J. Ellington, Ryan
George, John Grimshaw, Malcolm Guiver, Robert Hill, Katie L. Hopkins, Rachel
Jones, Cheryl Lenney, Amy J. Mathers, Ashley McEwan, Ginny Moore, Andrew
Mumford, Mark Neilson, Sarah Neilson, Tim E.A. Peto, Hang T.T. Phan, Mark
Regan, Anna C. Seale, Nicole Stoesser, Jay Turner-Gardner, Vicky Watts, A. Sarah
Walker, Jimmy Walker, David H. Wyllie, William Welfare and Neil Woodford.

REFERENCES

1. Munoz-Price LS, Poirel L, Bonomo RA et al. Clinical epidemiology of the
global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis* 2013;
13: 785-96.
2. Logan LK, Weinstein RA. The Epidemiology of Carbapenem-Resistant
Enterobacteriaceae: The Impact and Evolution of a Global Menace. *J Infect Dis* 2017;
215: S28-S36.
3. Nordmann P, Dortet L, Poirel L. Carbapenem resistance in
Enterobacteriaceae: here is the storm! *Trends Mol Med* 2012; **18**: 263-72.
4. Mathers AJ, Cox HL, Kitchel B et al. Molecular dissection of an outbreak of
carbapenem-resistant enterobacteriaceae reveals Intergenous KPC carbapenemase
transmission through a promiscuous plasmid. *MBio* 2011; **2**: e00204-11.

- 529 5. Kizny Gordon AE, Mathers AJ, Cheong EY et al. Is the hospital water
530 environment a reservoir for carbapenem-resistant organisms causing hospital-acquired
531 infections? A systematic review of the literature. *Clin Infect Dis* 2017.
- 532 6. Bleichenbacher S, Stevens MJA, Zurfluh K et al. Environmental
533 dissemination of carbapenemase-producing Enterobacteriaceae in rivers in
534 Switzerland. *Environ Pollut* 2020; **265**.
- 535 7. Cahill N, O'Connor L, Mahon B et al. Hospital effluent: A reservoir for
536 carbapenemase-producing Enterobacterales? *The Science of the total environment*
537 2019; **672**: 618-24.
- 538 8. León-Sampedro R, DelaFuente J, Díaz-Agero C et al. Pervasive transmission
539 of a carbapenem resistance plasmid in the gut microbiota of hospitalized patients. *Nat*
540 *Microbiol* 2021; **6**: 606-16.
- 541 9. Weingarten RA, Johnson RC, Conlan S et al. Genomic Analysis of Hospital
542 Plumbing Reveals Diverse Reservoir of Bacterial Plasmids Conferring Carbapenem
543 Resistance. *MBio* 2018; **9**.
- 544 10. Stoesser N, Sheppard AE, Moore CE et al. Extensive Within-Host Diversity in
545 Fecally Carried Extended-Spectrum-Beta-Lactamase-Producing Escherichia coli
546 Isolates: Implications for Transmission Analyses. *J Clin Microbiol* 2015; **53**: 2122-
547 31.
- 548 11. Ludden C, Coll F, Gouliouris T et al. Defining nosocomial transmission of
549 Escherichia coli and antimicrobial resistance genes: a genomic surveillance study. *The*
550 *Lancet Microbe* 2021; **2**: e472-e80.
- 551 12. Decraene V, Phan HTT, George R et al. A Large, Refractory Nosocomial
552 Outbreak of Klebsiella pneumoniae Carbapenemase-Producing Escherichia coli

553 Demonstrates Carbapenemase Gene Outbreaks Involving Sink Sites Require Novel
554 Approaches to Infection Control. *Antimicrob Agents Chemother* 2018; **62**.
555 13. Public Health England. Carbapenemase-producing Enterobacteriaceae:
556 laboratory confirmed cases, 2003 to 2015. London: Public Health England, 2016.
557 14. Public Health England. Carbapenemase-producing Enterobacteriaceae: early
558 detection, management and control toolkit for acute trusts. 2014.
559 15. UK Standards for Microbiology Investigations.
560 [https://www.gov.uk/government/collections/standards-for-microbiology-](https://www.gov.uk/government/collections/standards-for-microbiology-investigations-smi)
561 [investigations-smi](https://www.gov.uk/government/collections/standards-for-microbiology-investigations-smi). Last updated: 1st July 2021.
562 16. The European Committee on Antimicrobial Susceptibility Testing. Breakpoint
563 tables for interpretation of MICs and zone diameters. 2017.
564 17. Stoesser N, Giess A, Batty EM et al. Genome sequencing of an extended
565 series of NDM-producing *Klebsiella pneumoniae* isolates from neonatal infections in
566 a Nepali hospital characterizes the extent of community- versus hospital-associated
567 transmission in an endemic setting. *Antimicrob Agents Chemother* 2014; **58**: 7347-57.
568 18. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence
569 classification using exact alignments. *Genome Biol* 2014; **15**: R46.
570 19. Nguyen LT, Schmidt HA, von Haeseler A et al. IQ-TREE: a fast and effective
571 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*
572 2015; **32**: 268-74.
573 20. Didelot X, Wilson DJ. ClonalFrameML: efficient inference of recombination
574 in whole bacterial genomes. *PLoS Comput Biol* 2015; **11**: e1004041.
575 21. Sheppard AE, Stoesser N, Wilson DJ et al. Nested Russian Doll-Like Genetic
576 Mobility Drives Rapid Dissemination of the Carbapenem Resistance Gene blaKPC.
577 *Antimicrob Agents Chemother* 2016; **60**: 3767-78.

578 22. Tonkin-Hill G, MacAlasdair N, Ruis C et al. Producing polished prokaryotic
579 pangenomes with the Panaroo pipeline. *Genome Biol* 2020; **21**: 180.

580 23. Bankevich A, Nurk S, Antipov D et al. SPAdes: a new genome assembly
581 algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012; **19**: 455-
582 77.

583 24. Hunt M, Mather AE, Sánchez-Busó L et al. ARIBA: rapid antimicrobial
584 resistance genotyping directly from sequencing reads. *Microb Genom* 2017; **3**:
585 e000131.

586 25. Jia B, Raphenya AR, Alcock B et al. CARD 2017: expansion and model-
587 centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids*
588 *Res* 2017; **45**: D566-D73.

589 26. Sheppard AE, Stoesser N, German-Mesner I et al. TETyper: a bioinformatic
590 pipeline for classifying variation and genetic contexts of transposable elements from
591 short-read whole-genome sequencing data. *Microb Genom* 2018; **4**.

592 27. De Maio N, Wu CH, Wilson DJ. SCOTTI: Efficient Reconstruction of
593 Transmission within Outbreaks with the Structured Coalescent. *PLoS Comput Biol*
594 2016; **12**: e1005130.

595 28. Oksanen JB, Blanchet, FG, Friendly M, Kindt R, Legendre P, McGlinn D,
596 Minchin PR, O'Hara RB, Simpson GL, Solymos P, Henry M, Stevens H, Szoecs E,
597 Wagner H. *vegan: Community Ecology Package*. 2020.

598 29. Harrell Jr FE. *rms: Regression Modeling Strategies*. 2021.

599 30. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. 2016.

600 31. Yu G. *scatterpie: Scatter Pie Plot*. 2020.

601 32. Valentin AS, Santos SD, Goube F et al. A prospective multicentre surveillance
602 study to investigate the risk associated with contaminated sinks in the intensive care
603 unit. *Clin Microbiol Infect* 2021; **27**: 1347.e9-.e14.

604 33. Heireman L, Hamerlinck H, Vandendriessche S et al. Toilet drain water as a
605 potential source of hospital room-to-room transmission of carbapenemase-producing
606 *Klebsiella pneumoniae*. *The Journal of hospital infection* 2020; **106**: 232-9.

607 34. Kantele A, Kuenzli E, Dunn SJ et al. Dynamics of intestinal multidrug-
608 resistant bacteria colonisation contracted by visitors to a high-endemic setting: a
609 prospective, daily, real-time sampling study. *The Lancet Microbe* 2021; **2**: e151-e8.

610 35. Gouliouris T, Coll F, Ludden C et al. Quantifying acquisition and transmission
611 of *Enterococcus faecium* using genomic surveillance. *Nat Microbiol* 2021; **6**: 103-11.

612 36. Qiao F, Wei L, Feng Y et al. Handwashing Sink Contamination and
613 Carbapenem-resistant *Klebsiella* Infection in the Intensive Care Unit: A Prospective
614 Multicenter Study. *Clin Infect Dis* 2020; **71**: S379-s85.

615

616 TABLES

617 **Table 1. Association of site type and unit location with *bla*_{KPC}-Enterobacterales**
 618 **(KPC-E) positivity in environmental samples.** Outputs of a multivariable logistic
 619 regression model with robust standard errors clustered by environmental site type and
 620 unit location; no other variables were considered in the model. P-values <0.05 were
 621 considered significant and are highlighted in bold. No interactions with an interaction
 622 Wald p <0.05 were observed.

623

Environmental site	Number of sites	Number of sampling events KPC-E-positive (% of total sampling events)	Number of sampling events KPC-E-negative	OR (95% CI)	p
Environmental site type					
Clinical handwash basin drain	96	79 (5.9%)	1270	Reference	
Handwash basin drain	85	52 (4.6%)	1077	0.99 (0.51-1.93)	0.9735
Shower/bath drain	59	100 (19.2%)	421	2.69 (1.44-5.01)	0.0019
Sluice/sluice	16	47 (17.9%)	215	2.60 (1.04-	0.0410

sink/mop sink				6.52)	
Toilet	74	5 (0.5%)	931	0.28 (0.10-0.78)	0.0152
Utility sink	19	36 (12.4%)	255	1.32 (0.47-3.68)	0.5965
Environmental site location					
Acute medicine unit	129	86 (6.6%)	1223	Reference	
Cardiac unit	76	111 (7.0%)	1453	1.07 (0.57-2.00)	0.8412
Geratology unit	144	122 (7.6%)	1493	0.88 (0.52-1.48)	0.6243

FIGURE LEGENDS

Figure 1. Sampling and sequencing flowchart for the study.

Figure 2. *bla*_{KPC}-Enterobacterales (KPC-E)-positive environmental sites by week in 2016 and hospital unit. The number of environmental site samples performed and KPC-E-positive samples by week of sampling, stratified by unit (top three panels), and the proportion of KPC-E-positive sites by week of sampling, again stratified by unit (bottom panel).

Figure 3. Carbapenem-susceptible Enterobacterales (CSE) and carbapenem-resistant-Enterobacterales (CRE) patient cultures by week in 2016 and hospital unit. Top three panels show counts of Enterobacterales culture-positive samples stratified by unit and carbapenem susceptibility; bottom panel shows the proportion of all cultures taken that were CRE, again stratified by unit.

Figure 4. Number of unique patient/environmental niches colonised by common *bla*_{KPC} Enterobacterales species/strains. A) Includes species/strains identified in both patients and wastewater niches during the study (if >50 isolates of a species were identified [see Methods]). Orange circles denote wastewater niches, black circles unique patients. B) Includes species/strains identified only in patients (left panel) and only in wastewater sites (right panel). For *Raoultella ornithinolytica*, six strains were identified and these were observed in wastewater sites only (data not shown).

Figure 5. Top panel: Within-patient KPC-E diversity. Pies reflect the proportion of strains within a sample that were identified using whole genome sequencing, with

649 evaluation of diversity at the species-strain level. “cfre” = *Citrobacter freundii*, “eclo”
650 = *Enterobacter cloacae*, “ecol” = *Escherichia coli*, “evul” = *Escherichia vulnificus*,
651 “koxy” = *Klebsiella oxytoca*, “kpne” = *Klebsiella pneumoniae*, “lecl” = *Leclercia*
652 *adecarboxylata*. Numbers after species abbreviations denote unique strains. Bottom
653 panel: Within-wastewater sample KPC-E diversity. Pies reflect the proportion of
654 strains within a sample that were identified using whole genome sequencing, with
655 evaluation of diversity at the species-strain level. “aher” = *Escherichia hermannii*,
656 “cfar” = *Citrobacter farmeri*, “easb” = *Enterobacter asburiae*, “pant” = *Pantoea* spp.,
657 “plur” = *Pluralibacter gergoviae*; other assignments as for top panel.

658
659 **Figure 6.** A) Recombination-adjusted phylogeny for *E. cloacae* strain 10. Main
660 annotations as described in the legends as part of the figure. In the panel to the right
661 of the phylogeny, AMR profile, IS profile, plasmid replicon profile and Tn4401/target
662 site sequences (TSSs) are represented, with different colours distinguishing different
663 versions of each. Three isolates in F47 and one isolate in F46 clustered with the F45-
664 associated isolates; sampling dates suggest that these represented transmission events
665 from F45. The scale bar is in SNVs. B) Represents plasmid replicon profiles in the
666 isolates obtained from the environmental site samples over time, and C) similarly
667 represents the Tn4401/TSS profiles with additional deletions observed annotated as
668 an overlay for isolates typed as containing the “unknown-1_ATTGA_ATTGA
669 variants” using TETyper.

670
671 **Figure 7.** Putative transmission network for sub-lineages of *K. pneumoniae* strain 9,
672 defined by accessory genome clustering (clusters 1, 2, 3a-h). Horizontal bars
673 represent admission episodes to study wards for each patient (pale blue=W45,

geratology; pale pink=AM1, acute medicine; pale purple=AM2, acute medicine; A13 represents the only shower site involved. Empty black triangles denote CPE-negative rectal screens on patients. Arrows denote possible transmission links (dashed arrows denote events that have no geographic overlap or could be associated with two links based on negative screen, geographic location and timing). For clarity, arrows are not drawn for clusters 3g and 3h, which affected the most patients - these are shown as an inset panel at the bottom right.

681

Figure 8. Left-hand panel shows a recombination-corrected phylogeny for *K. pneumoniae* strain 11, designating the two identified clusters, 1 and 2, and annotated with isolate names, unit, collection date, niche, and Tn4401-TSS type. Right-hand panels reflect the plasmid replicon and Tn4401-TSS diversity within sequenced isolates in given niches over time for each chromosomally genetically-related cluster, highlighting features consistent with clonal transmission, Tn4401 transposition events within isolates (Tn4401a, change in inner dot colour with consistent outer ring colour), deletions in Tn4401a (to “unknown-a”), plasmid replicon loss/gain events (change in outer ring colour).

Wastewater sites cultured

n=349

Clinical handwash basin drains n=96

Handwash basin drains n=85

Shower/bath drains n=59

Sluices/slucie sink/mop sinks n=16

Utility sinks n=19

Toilets n=74

Wastewater samples

Total cultures - n=4,488

CRE-positive -n=319 (7.1%)

Patients cultured

n=4,425

Patients ever CRE-positive 183 (4.1%)

Patient samples

Total cultures n=41,203

Culture-positive n=5,154 (12.5%)

Enterobacterales-positive n= 2,172
(2,000 with carbapenem susceptibility
results) (42.1%)

CRE-positive n=399 (20.0% of
Enterobacterales; 1.0% of total)

Total isolates sequenced
n=1452

Excluded

n=177

Not Enterobacterales
Mixed/poor quality
Replicates

Total wastewater isolate sequences available
for analysis
n=1275

Enterobacter cloacae - n=536 isolates

Klebsiella pneumoniae - n=463 isolates

Citrobacter freundii - n=133 isolates

Raoultella ornithinolytica - n=63 isolates

Enterobacter asburiae - n=15 isolates

Citrobacter farmeri - n=14 isolates

Klebsiella oxytoca - n=13 isolates

Escherichia coli - n=12 isolates

Pluralibacter gergoviae - n=10 isolates

Leclercia adecarboxylata - n=9 isolates

Kluyvera intermedia - n=6 isolates

Pantoea spp. - n=1 isolate

Total isolates sequenced
n=482

Excluded

n=25

Not Enterobacterales
Mixed/poor quality
Replicates

Total patient isolate sequences available for
analysis
n=457

Klebsiella pneumoniae - n=329 isolates

Escherichia coli - n=64 isolates

Enterobacter cloacae - n=28 isolates

Klebsiella oxytoca - n=22 isolates

Escherichia vulneris - n=7 isolates

Citrobacter freundii - n=4 isolates

Leclercia adecarboxylata - n=3 isolates

A.

B.

Species/strain

Species/strain

A
E. cloacae strain 10
Geratology ward

Plasmid replicon profile

a IncFIB|IncFII
a IncFIB|IncFII|IncH2|IncHI2A

tn4401type

a Tn4401a-1_ATTGA_ATTGA
a unknown-1_ATTGA_ATTGA

Sample collection date