1 Manuscript draft

2 **Full title:** Differences in bacterial colonization and mucosal responses between
3 high and low SES children in Indonesia

4 **Running title:** SES alters mucosal responses to bacteria

5

6 Authors:

7 Marloes M.A.R. van Dorst^{1*}, Shohreh Azimi^{1*}, Sitti Wahyuni², Aldian I. Amaruddin^{1,2}, Erliyani

8 Sartono¹, Linda J. Wammes³, Maria Yazdanbakhsh¹, Simon P. Jochems^{1#}

9

10 ¹ Department of Parasitology, Leiden University Medical Center, Leiden, The Netherlands

11 ² Department of Parasitology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

12 ³ Department of Medical Microbiology, Leiden University Medical Center, Leiden, the Netherlands

13

14 *contributed equally

15 # correspondence to s.p.jochems@lumc.nl

16

17 Abstract:

18 **BACKGROUND.** Nasopharyngeal carriage of pathogenic bacteria precedes invasive disease and
19 higher rates are found in low socioeconomic-status (SES) settings. Local immune responses are
20 important for controlling colonization, but whether SES affects these responses is currently
21 unknown.

22 **OBJECTIVE.** Examining bacterial colonization and cytokine response in nasal mucosa of children from
23 high and low SES.

24 **METHODS.** Twenty-five cytokines were measured in nasal fluid. qPCR was performed to determine
25 carriage and density of *Haemophilus influenzae* (H. influenzae), *Streptococcus pneumoniae* (S.
26 pneumoniae), *Moraxella catarrhalis* (M. catarrhalis) and *Staphylococcus aureus* (S. aureus).

27 **RESULTS.** The densities of H. influenzae and S. pneumoniae were increased in low compared to the
28 high SES ($p=0.006$, $p=0.026$), with respectively 6 and 67 times higher median densities. Densities of
29 H. influenzae and S. pneumoniae were positively associated with levels of IL-1beta ($p=0.002$,
30 $p=0.008$) and IL-6 ($p<0.001$, $p=0.006$). After correcting for bacterial density, IL-6 levels were
31 increased in colonized children from high compared to low SES for both H. influenzae and S.
32 pneumoniae (both $p=0.039$).

33 **CONCLUSION.** Increased density of H. influenzae and S. pneumoniae was observed in low SES
34 children, while IL-6 levels associated with colonization were reduced in these children, indicating
35 that immune responses to bacterial colonization were altered by SES.

36 **Key words:** *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Moraxella catarrhalis*,
37 *Staphylococcus aureus*, IL-1beta, IL-6, IL-1RA, socio-economic status, upper respiratory tract
38 colonization.

39 **Word count abstract text:** 200 words

40 **Number of insertions:** 5 figures

41 Introduction

42 Lower respiratory tract infections (LRTIs) remain the leading cause of mortality amongst children
43 under five, accounting for 2.4 million deaths in 2016 most of them in low- and middle income
44 countries (LMICs) (1, 2).

45

46 A number infectious agents can establish LRTI, prevalent bacterial causes include *Streptococcus*
47 *pneumoniae* (*S. pneumoniae*), *Haemophilus influenzae* (*H. influenzae*), *Moraxella catarrhalis* (*M.*
48 *catarrhalis*) and *Staphylococcus aureus* (*S. aureus*). Nasopharyngeal carriage of these bacteria is
49 common in infants and colonization by these bacteria is considered an essential first step in
50 development of disease (3-5). Moreover, nasopharyngeal carriage and increased densities of *S.*
51 *pneumoniae*, *H. influenzae* and *M. catarrhalis* have been associated with pneumonia in Tanzanian
52 children and increased *H. influenzae* densities were found among children with (very) severe
53 pneumonia in seven LMICs (6, 7).
54
55 Control of nasopharyngeal bacterial colonization is mediated by local immune responses and
56 impaired innate cytokine responses have been associated with persistent bacterial colonization. A
57 human inoculation study showed that clearance of nasal *S. aureus* requires an upregulation of
58 chemokines, growth factors and inflammatory cytokines and that a low IL-1RA/IL-1beta ratio
59 associates with *S. aureus* persistence (8). Furthermore, an increased prevalence of nasopharyngeal
60 colonization with *M. catarrhalis* was observed infants with anemia, while the pro-inflammatory
61 cytokine responses were reduced upon *in vitro* stimulation (9). Finally, murine models show that IL-1
62 signaling is essential for clearance of *S. pneumoniae* and suggest that reduced IL-1 responses might
63 be permissive for persistent colonization during infancy (10).
64
65 Increased carriage rates have been observed in various settings and are associated with lower socio-
66 economic status (SES). High nasopharyngeal bacterial colonization rates have been observed in low-
67 income countries (11), rural areas (12) and indigenous populations (4). Furthermore, a study in Israel
68 showed that SES rather than ethnicity drove these differences in pneumococcal carriage (13). Finally,
69 socio-economic factors were identified as risk factors for carriage of potential pathogenic bacteria in
70 Indonesian children (14).

71

Since increased bacterial colonization has been observed in low SES settings and local immune responses play an important role in the control of the bacterial colonization, reduced mucosal immune responses in low SES populations could play a role. Studies comparing immune responses between socio-economic settings are scarce and often focus on the systemic immune system. To our knowledge, no studies have compared the immune responses to bacterial colonization between different socio-economic settings. Moreover, most studies that found an association between SES and colonization have been performed in a heterogenous population. Therefore, this study aimed to examine the cytokine response to nasopharyngeal bacterial colonization of children from high and low SES schools in a single urban area.

81

82 **Methods:**

83 **Data collection:**

The data was obtained as part of a larger study into the effects of SES in school-aged children in Makassar, Indonesia (15). This cross-sectional study included children attending two primary schools in Makassar, the capital of South Sulawesi. These schools were selected based on the socio-economic background of the children attending the school. One of the schools is attended by children from High SES families, a privately funded school, whereas the other school is fully government funded and predominantly attended by low SES children. Both schools are located in the center of Makassar, 2km apart.

91

Samples were collected during September-October 2019. Informed written consent was obtained from primary caregivers. All children with written consent were included in the study, except for children using antihistamines/corticosteroids. Ethical approval was obtained from the Health Research Ethical Committee, Faculty of Medicine, Hasanuddin University (No:703/H4.6.4.5.31/PP36/2019). Socio-demographic information was obtained from all participants via questionnaires.

98

99 **Nasosorption samples preparation**

100 Nasosorption samples were collected by inserting an absorptive matrix strip (Nasosorption™, Hunt
101 Developments) into one nostril and pushed against the nasal lining for 30-60 seconds (16). These
102 strips were then placed on ice until storage at -80°C within 8 hours after collection. The nasosorption
103 samples were eluted by adding 100µL sterilized PBS with 1%BSA+0.05%Triton-X100 to the filter and
104 spun down at 3600xg for 10 minutes at 4°C. The supernatant was transferred to a new tube and
105 centrifuged at 16000xg, 4°C, 10 minutes. The supernatant was moved to a new tube for cytokine
106 analysis and the pellet was stored at -20°C until DNA extraction.

107

108 **Cytokine concentration measurement**

109 Of the supernatant, 12 µl was used to measure the concentration of 25 cytokines, by the Human
110 Cytokine 25-plex ProcartaPlex Panel (Invitrogen, ThermoFisher No:EXP250-12166-901) according to
111 manufacturer's instructions, but additional washing steps and alcohol flushes were performed to
112 account for the muckiness. The samples were measured by the Luminex200 device at normal RP1
113 target and concentrations were obtained by using the xPonent3.1 software in pg/mL. Children for
114 who >90% of the cytokines could not be measured in their nasosorption sample were excluded from
115 further analysis. This was the case for 15% of the children, 6 high and 9 low SES children.

116

117 **Generation of standard curves**

118 The DNA concentrations of *S. aureus* (ATCC43300), *H. influenzae* (ATCC49766), *S. pneumoniae* 6305
119 and clinical isolate of *M. catarrhalis* were determined spectrophotometrically with the
120 NanoDrop1000 (Thermo scientific). The copy numbers of template were calculated using the
121 genome length computed with URI Genomics & Sequencing Center
122 (<http://cels.uri.edu/gsc/cndna.html>). A standard curve was generated of 10-fold serial dilution
123 scheme ranging from 10⁷-10¹ copy/mL

124

125 **Bacterial colonization measurement**

126 To determine nasal bacterial colonization, qPCR was performed on the bacterial pellet. DNA
 127 extraction was performed using magnetic beads, as previously published (17). To determine the
 128 colonization of *S. pneumoniae* and *H. influenzae*, qPCR was performed using the *LytA* and *IgA1* gene
 129 respectively, as previously described (17, 18). qPCR for the *copB* gene was performed to determine
 130 colonization of *M. catarrhalis* and for *S. aureus* the *nuc* gene was used as previously described (6).
 131 Using the standard curves, the genome length and Avogadro's number, the number of genome
 132 copies/ μ L was determined in the samples per bacterium. Details regarding the primers and probes
 133 utilized in this study can be found in supplementary (**TableS1**).

134

135 **Statistical analysis**

136 The standardized z-scores of body mass index (z-BMI) were determined according to the WHO
 137 guidelines (19). To obtain approximately normally distributed data, cytokine concentrations and
 138 bacterial loads were $\log_{10}$ -transformed. A Student's t-test was performed for continuous data and
 139 for binary and categorical data Pearson's Chi-square was used. Pearson's correlation was used to
 140 examine the correlation of two continuous variables. To investigate the effect of SES on the bacterial
 141 loads and to estimate the effect of bacterial density and SES on cytokine levels, a regression model
 142 was performed. For all regression models in this study age, sex and z-BMI were considered *a priori*
 143 confounders and adjustment was performed accordingly. To identify the main drivers of the
 144 correlation between bacterial density and cytokine concentrations, a canonical correlation analysis
 145 was performed by using Wilk's Lambdas including the density of the four bacteria and the
 146 concentrations of all widely measurable cytokines. The two bacteria and two cytokines that showed
 147 the largest coefficient in the canonical correlation analysis were selected for further analysis. P-
 148 values smaller than 0.05 were considered statistically significant. Data was analyzed and visualized
 149 using RStudio and R software.

150

151 Results:

152 Bacterial density of *H. influenzae* and *S. pneumoniae* increased in low SES

153 A total of 98 school-aged children were included, 50 children attended the low SES school and 48
154 children the high SES school (**TableS2**). There was no significant difference in age or sex, but z-BMI
155 and parental income were between lower in low SES compared to high SES children. The carriage
156 rate was highest for *M. catarrhalis*, lowest for *S. aureus* and a combination of two or three bacteria
157 was most common. Neither the carriage rates of the bacteria nor the number of co-colonizing
158 bacteria differed between high and low SES.

159 Although the carriage rates and combinations did not differ between high and low SES, differences in
160 the densities of bacteria were observed (**Figure1**). The median bacterial load of *H. influenzae* was
161 59,075 copies per nasosorption sample in low and 9,804 in high SES, indicating a 6-times higher
162 median density. For *S. pneumoniae*, the median bacterial load was 97,422 in the low and 1,446 in
163 the high SES, corresponding to a 67-times higher median density. No significant differences were
164 observed in the bacterial loads of *M. catarrhalis* and *S. aureus* between high and low SES. To further
165 examine the relation between the bacterial load and SES, a multivariate regression analysis was
166 performed adjusted for *a priori* confounders. These results showed that SES significantly affected the
167 *H. influenzae* ($\beta_{SES}=-1.19$, $p=0.006$) and *S. pneumoniae* density ($\beta_{SES}=-1.25$, $p=0.026$).

168

169 Correlation between densities of the different bacteria

170 Multiple studies have shown that microbial species interact with each other and associations
171 between different species have been described (20, 21). Correlation analysis showed that the *H.*
172 *influenzae* density positively correlated with the *M. catarrhalis* density ($\beta=0.53$, $p<0.001$) and the
173 bacterial load of *S. pneumoniae* negatively associated with *S. aureus* ($\beta=-0.50$, $p=0.04$) (**Figure2**).

Furthermore, there was a trend for positive association between *H. influenzae* and *S. pneumoniae* density ($\beta=0.34$, $p=0.07$). No association was found between the other combinations of bacteria.

IL-1beta concentrations are increased in the low SES compared to the high SES.

To understand whether impaired mucosal responses might be responsible for the higher densities of colonized bacteria in low SES children, the concentration of 25 cytokines in nasal fluid were analyzed using a multiplex assay. Nine of these cytokines were detectable in the majority of samples: IL-18, IL-1alpha, IL-1beta, IL-1RA, IL-27, IL-4, IL-6, IL-7 and TNF-alpha. The other cytokines were below the limit of reliable detection in most samples (**TableS3**). Comparison of the cytokine levels between high and low SES showed that IL-1beta levels were significantly increased in the high SES, with a mean concentration of 338.8 in low and 102.3 in high SES after adjusting for confounders ($\beta_{SES}=-0.518$, $p=0.048$) (**Figure3, TableS4**).

IL-6 concentrations are increased in high SES children colonized by *H. influenzae* or *S. pneumoniae*

Similarly to PCA analysis, canonical correlation allows the integration of two multi-dimensional datasets to find common drivers of variation maximizing covariance between the datasets, allowing us to compare bacteria and cytokines on a global level (22). The densities of *H. influenzae* and *S. pneumoniae* and IL-1beta and IL-6 concentrations were identified as the main drivers of the correlation between bacteria and cytokines (**TableS5**). Indeed both *H. influenzae* and *S. pneumoniae* densities positively associated with IL-1beta and IL-6 levels (**Figure4A-D**). To quantify these correlations, a regression model including *a priori* confounders was used. The results showed that *H. influenzae* and *S. pneumoniae* densities are positively associated with IL-1beta ($\beta_{bacteria} = 0.294$, $p=0.002$, $\beta_{bacteria}=0.221$, $p=0.008$, respectively) and IL-6 ($\beta_{bacteria}=0.190$, $p<0.001$ and $\beta_{bacteria}=0.136$, $p=0.006$) (**TableS6**). After adjusting for *a priori* confounders and the bacterial densities, IL-6 levels were increased in high compared to low SES in children colonized by *H. influenzae* ($\beta_{SES}=0.360$, $p=0.039$) and *S. pneumoniae* ($\beta_{SES}=0.366$, $p=0.039$), indicating that IL-6 levels were increased in high

compared to low SES at any given density of these bacteria. To examine whether SES affected the strength and direction of this association, a regression model including the interaction between these bacterial density and cytokine levels was performed, which showed that this was not significant (**TableS7**). Thus, children from both high and low SES had dose-dependent cytokine responses to *H. influenzae* or *S. pneumoniae* colonization, but these responses were stronger in high SES children for IL-6.

IL-1RA and IL-1beta response to *S. aureus* carriage differs between high and low SES

Since a limited number of children were colonized with *S. aureus* and the presence and density was not associated with that of the other bacteria, responses to colonization by *S. aureus* were not likely to be identified by the canonical correlation analysis. Therefore, the relation between *S. aureus* carriage status and cytokine concentrations was further examined, whereby IL-1 cytokines were of main interest since these cytokines were shown previously to be important for *S. aureus* control (8). In the low SES, the concentration of IL-1RA seemed to be higher in *S. aureus* carriers compared to non-carriers, whereas in high SES IL-1RA levels were lower in the carriers compared to non-carriers (**Figure5A**). A similar pattern was observed for IL-1beta (**Figure5B**). To quantify these observations, we performed a regression analysis including the *a priori* confounders, showing a significant difference between high and low SES for IL-1RA ($\beta_{\text{interaction}}=-0.700$, $p=0.007$), and a similar trend for IL-1beta ($\beta_{\text{interaction}}=-0.700$, $p=0.059$) (**TableS8**). Since IL-1beta and IL-1RA compete for the same receptor and the activity of IL-1beta is affected by the levels of IL-1RA, the ratio between IL-1RA and IL-1beta was determined (**Figure5C**). The interaction between *S. aureus* carriage and SES was not seen for the ratio between these cytokines ($\beta_{\text{interaction}}=0.074$, $p=0.859$), but the IL-1RA/IL-1beta ratio tended to be higher in high compared to low SES ($\beta_{\text{SES}}=0.465$, $p=0.078$) (**TableS8**). The pattern observed for IL-1beta and IL-1RA did not exist for other IL-1 cytokines, including IL-1alpha (**FigureS1**).

Discussion

This is the first study evaluating the nasal cytokine response in relation to bacterial colonization in high and low SES children. This study showed increased densities of *H. influenzae* and *S. pneumoniae* in low compared to high SES. The densities of these bacteria were positively associated with IL-1 β and IL-6 levels. After correcting for bacterial density, IL-6 levels were increased in high SES, indicating that IL-6 levels were higher at any given density of *H. influenzae* and *S. pneumoniae* in high compared to low SES. Furthermore, increased levels of IL-1RA were found in *S. aureus* carriers compared to non-carriers in low SES, whereas in high SES IL-1RA were higher in the non-carriers. A similar pattern was observed for IL-1 β but not for other cytokines. The IL-1RA/IL-1 β ratio was not affected by *S. aureus* carrier status, but tended to be increased in high compared to low SES. SES has been identified as risk factor for increased nasopharyngeal carriage in a number of studies (4, 11-14). A previous study in Indonesia found increased densities of *S. aureus* in children living in semi-rural compared to urban areas. Furthermore, they identified family income as a risk factor for pneumococcal carriage densities and showed that low maternal education was associated with increased *H. influenzae* densities (14). The results of the current study are in line with these previous findings, however in contrast to the study Fadlyana *et al.* we included children from different SES within one urban center, thereby providing unique insights into the role of SES on nasopharyngeal bacterial carriage and densities in children.

Without taking bacterial colonization status into account, only IL-1 β was significantly different between high and low SES, while there was a large dynamic range in cytokine levels observed, indicating that local factors are important for driving cytokine levels at the mucosa. Indeed, densities of *H. influenzae* and *S. pneumoniae* were positively associated with IL-1 β and IL-6 levels, highlighting the importance of the activation of the IL-1 signaling pathway in response to bacterial colonization. The importance of the IL-1 cytokine signaling in the clearance of *S. pneumoniae* has been shown in mice and reduced IL-1 responses have been suggested to be permissive for persistent

colonization during infancy (10, 23). Moreover, high levels of IL-6 and IL-1beta were found in response to Non-typeable H. influenzae in individuals suffering from chronic suppurative lung disease (24).

After adjusting for bacterial densities, increased IL-6 levels were observed in children colonized by H. influenzae or S. pneumoniae from high compared low SES, whereas this was not observed for IL-1beta. This might be explained by the role of IL-6 in the IL-1 signaling pathway and its capacity to drive promote the differentiation of Th17-cells. Whereas IL-1beta is primarily initiates the IL-1 signaling, IL-6 production is one of the numerous downstream effects of the IL-1 signaling pathway. Furthermore, high levels of IL-6 are known to promote differentiation of CD4⁺-cells to form Th17-cells, a cell type that has been shown essential for the clearance of S. pneumoniae in mice (25). A study analyzing the Th17-cells and cytokines in adenoidal tissue from S. pneumoniae-positive and S. pneumoniae-negative children found an increased number of Th17-cells and higher levels of IL-17A and IL-6 in S. pneumoniae-negative compared to S. pneumoniae-positive children (26). In the current study increased IL-6 levels were observed in the high SES, whereas the density of H. influenzae and S. pneumoniae was reduced, supporting the role of IL-6 in the control of these bacteria. The IL-17A levels were below the level of detection in most samples collected in this study, however very low levels of IL-17A might be enough to activate local T-cells and therefore be involved in the bacterial clearance.

In the current study in the low SES increased IL-1beta and IL-1RA levels were observed in S. aureus carriers compared to non-carriers, whereas in the high SES decreased levels were found in carriers. The ratio between IL-1RA and IL-1beta was not affected by the carrier status but tended to be decreased in low compared to high SES after adjusting for S. aureus carriage. A human nasal inoculation with S. aureus showed that IL-1beta was upregulated after inoculation in individuals able to clear the bacteria compared to individuals, but found no difference in IL-1RA levels. Furthermore, they showed that IL-1RA/IL-1beta ratio was significantly decreased in individuals with persistent S.

aureus and proposed this ratio as metric for the clearance of *S. aureus* in the nasal mucosa rather than the expression of individual cytokines (8). In the current study, the IL-1RA/IL-1beta ratio tended to be increased in high compared to low SES, indicating that the response to *S. aureus* in low SES children might be reduced. No significant differences in the IL-1RA/IL-1beta ratio between carriers and non-carriers were found, however, it should be kept in mind that the duration of the colonization was not measured in the current study and some of the carriers would be able to clear *S. aureus* over time. To elucidate the relation between SES and *S. aureus* persistence, longitudinal studies are thus needed.

A factor that can affect the bacterial colonization and cytokine responses and which can be impacted by SES is the vaccination status against pneumococcal and *H. influenzae* type B (Hib) diseases. The children in our study were not vaccinated through their national childhood vaccination program. However, in a private clinic on payment additional vaccines can be administered, which predominantly involves the Hib vaccine. Although vaccination status might differ between SES its effect on the bacterial load detected here is probably limited, since the (sero)types prevalent in Indonesia only partially match with the available vaccines. A cross-sectional study amongst 302 young children in Indonesia in 2016, thus after Hib vaccine was incorporated in the national program but before the pneumococcal conjugate vaccine (PCV13) was introduced, showed a carriage rate of 27.5% of *H. influenzae*, but none of these isolates was type B (27). Moreover, before the introduction of Hib vaccination it was shown that Hib vaccination would not have an effect on the pneumonia incidence in Indonesia whereas it in Africa and South America significantly did, indicating that in Indonesia other respiratory pathogens are responsible for LRTIs (28). Moreover, study by Dunne *et al.* showed that only 46% of the pneumococcal isolates obtained from Indonesian children are covered by the PCV13 (27). While location-specific serotype circulation is of course possible, one can expect that the isolates found in our study are also (sero)types that will only partly be covered by the vaccine, which limits the effect of differences in vaccination status between high and low SES on the results.

One of the limitations of the current study is that only four pathogenic bacteria were measured in a cross-sectional manner, whereas it is known that viral coinfections and microbiota in general are also important in shaping responses. Moreover, future studies should assess changes and responses longitudinally. In addition, while school-going children are important drivers of transmission within communities, most childhood pneumonia occurs before the age of 5, therefore it would also be of interest to study mucosal immune responses in infants in future studies.

To conclude, the results in the current study indicate that the local immune response to nasopharyngeal bacterial colonization is altered by SES. Since nasopharyngeal carriage of these bacteria precedes disease and local immune responses are important in controlling potential pathogenic bacteria elucidating the relation between the bacterial colonization and local cytokine response is an important first step in understanding the bacteria-host relationship. Insights in this relationship are important for the development of vaccine strategies and treatment options and will eventually lead to reduced LRTIs worldwide.

Conflict of interest statement

The authors declare that they have no conflict of interest.

Funding statement

This work was supported by the PhD fellowship of the Leiden University Medical Center in the Netherlands. The sponsors of the study had no role in study design, data collection, data analysis, data interpretation or writing of the report.

Presentation of the work

This work has not been previously presented.

326 Corresponding author information

327 Dr. Simon P. Jochems

328 Department of Parasitology, Leiden University Medical Center

329 Albinusdreef 2, 2333 ZA Leiden, The Netherlands.

330 Tel: +31 71 5265072

331

332 Acknowledgements

333 The authors would like to thank Koen Stam and Mikhael Manurung for their advice on the statistical

334 analysis and all participants involved in this study as well as all students of the LUMC and

335 Hasanuddin University for their efforts on sample collection in participating primary schools in

336 Makassar. We thank Hasanuddin University Medical Research Center (HUM-RC) for laboratory

337 facilities during sample collections, and Hasanuddin University for their support.

338

339 Author contributions

340 Study design: SJ, MY, ES

341 Local organization of the study: SW, AA, MM, ES

342 Sample collection: MD, AA, MM, SJ, SW

343 Experimental measurements: MD, SA

344 Data analysis: MD, SA, SJ

345 Data interpretation: MD, SA, SJ

346 Writing manuscript: MD, SJ, MY

347 Sources:

348 1. Collaborators GBDCoD. Global, regional, and national age-sex specific mortality for 264 causes of

349 death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*.

350 2017;390:1151-1210.

2. Collaborators GBDLRI. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory infections in 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Infect Dis.* 2018;18:1191-1210.
3. Simell B, Auranen K, Kayhty H, et al. The fundamental link between pneumococcal carriage and disease. *Expert Rev Vaccines.* 2012;11:841-855.
4. Mackenzie GA, Leach AJ, Carapetis JR, Fisher J, Morris PS. Epidemiology of nasopharyngeal carriage of respiratory bacterial pathogens in children and adults: cross-sectional surveys in a population with high rates of pneumococcal disease. *Bmc Infect Dis.* 2010;10.
5. von Eiff C, Becker K, Machka K, Stammer H, Peters G, Grp S. Nasal carriage as a source of *Staphylococcus aureus* bacteremia. *New Engl J Med.* 2001;344:11-16.
6. Chochua S, D'Acremont V, Hanke C, et al. Increased Nasopharyngeal Density and Concurrent Carriage of *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Moraxella catarrhalis* Are Associated with Pneumonia in Febrile Children. *PLoS One.* 2016;11:e0167725.
7. Park DE, Baggett HC, Howie SRC, et al. Colonization Density of the Upper Respiratory Tract as a Predictor of Pneumonia-*Haemophilus influenzae*, *Moraxella catarrhalis*, *Staphylococcus aureus*, and *Pneumocystis jirovecii*. *Clin Infect Dis.* 2017;64:S328-S336.
8. Cole AL, Muthukrishnan G, Chong C, et al. Host innate inflammatory factors and staphylococcal protein A influence the duration of human *Staphylococcus aureus* nasal carriage. *Mucosal Immunol.* 2016;9:1537-1548.
9. Liao SL, Hsu SY, Lai SH, et al. Infant anemia is associated with reduced TLR-stimulated cytokine responses and increased nasopharyngeal colonization with *Moxarella catarrhalis*. *Sci Rep-Uk.* 2018;8.
10. Kuipers K, Lokken KL, Zangari T, Boyer MA, Shin S, Weiser JN. Age-related differences in IL-1 signaling and capsule serotype affect persistence of *Streptococcus pneumoniae* colonization. *Plos Pathog.* 2018;14.

375 11. Adegbola RA, DeAntonio R, Hill PC, et al. Carriage of *Streptococcus pneumoniae* and Other
376 Respiratory Bacterial Pathogens in Low and Lower-Middle Income Countries: A Systematic Review
377 and Meta-Analysis. *Plos One*. 2014;9.

378 12. Hanieh S, Hamaluba M, Kelly DF, et al. *Streptococcus pneumoniae* Carriage Prevalence in Nepal:
379 Evaluation of a Method for Delayed Transport of Samples from Remote Regions and Implications for
380 Vaccine Implementation. *Plos One*. 2014;9.

381 13. Dagan R, Juergens C, Trammel J, et al. Modeling pneumococcal nasopharyngeal acquisition as a
382 function of anticapsular serum antibody concentrations after pneumococcal conjugate vaccine
383 administration. *Vaccine*. 2016;34:4313-4320.

384 14. Fadlyana E, Dunne EM, Rusmil K, et al. Risk factors associated with nasopharyngeal carriage and
385 density of *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, and
386 *Staphylococcus aureus* in young children living in Indonesia. *Pneumonia*. 2018;10.

387 15. Wahyuni S, van Dorst M, Amaruddin AI, et al. The relationship between malnutrition and TH 2
388 immune markers: a study in school-aged children of different socio-economic backgrounds in
389 Makassar, Indonesia. *Trop Med Int Health*. 2021;26:195-203.

390 16. Thwaites RS, Jarvis HC, Singh N, et al. Absorption of Nasal and Bronchial Fluids: Precision
391 Sampling of the Human Respiratory Mucosa and Laboratory Processing of Samples. *J Vis Exp*. 2018.

392 17. Nikolaou E, Blizard A, Pojar S, et al. Minimally Invasive Nasal Sampling in Children Offers Accurate
393 Pneumococcal Colonization Detection. *Pediatric Infectious Disease Journal*. 2019;38:1147-1149.

394 18. Olwage CP, Adrian PV, Nunes MC, Madhi SA. Evaluation of the association of pneumococcal
395 conjugate vaccine immunization and density of nasopharyngeal bacterial colonization using a
396 multiplex quantitative polymerase chain reaction assay. *Vaccine*. 2018;36:3278-3285.

397 19. de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO
398 growth reference for school-aged children and adolescents. *B World Health Organ*. 2007;85:660-
399 667.

- 400 20. Margolis E, Yates A, Levin BR. The ecology of nasal colonization of *Streptococcus pneumoniae*,
401 *Haemophilus influenzae* and *Staphylococcus aureus*: the role of competition and interactions with
402 host's immune response. *BMC Microbiol.* 2010;10:59.
- 403 21. Chien YW, Vidal JE, Grijalva CG, et al. Density interactions among *Streptococcus pneumoniae*,
404 *Haemophilus influenzae* and *Staphylococcus aureus* in the nasopharynx of young Peruvian children.
405 *Pediatr Infect Dis J.* 2013;32:72-77.
- 406 22. Gonzalez I, Dejean S, Martin PGP, Baccini A. CCA: An R package to extend canonical correlation
407 analysis. *J Stat Softw.* 2008;23:1-14.
- 408 23. Lemon JK, Miller MR, Weiser JN. Sensing of Interleukin-1 Cytokines during *Streptococcus*
409 *pneumoniae* Colonization Contributes to Macrophage Recruitment and Bacterial Clearance. *Infect*
410 *Immun.* 2015;83:3204-3212.
- 411 24. Pizzutto SJ, Upham JW, Yerkovich ST, Chang AB. High Pulmonary Levels of IL-6 and IL-1 beta in
412 Children with Chronic Suppurative Lung Disease Are Associated with Low Systemic IFN-gamma
413 Production in Response to Non-Typeable *Haemophilus influenzae*. *Plos One.* 2015;10.
- 414 25. Zhang Z, Clarke TB, Weiser JN. Cellular effectors mediating Th17-dependent clearance of
415 pneumococcal colonization in mice. *J Clin Invest.* 2009;119:1899-1909.
- 416 26. Jiang XL, Zhang GL, Yang T, et al. Association of Pneumococcal Carriage and Expression of
417 Foxp3+Regulatory T Cells and Th17 Cells in the Adenoids of Children. *Respiration.* 2015;90:25-32.
- 418 27. Dunne EM, Murad C, Sudigdoadi S, et al. Carriage of *Streptococcus pneumoniae*, *Haemophilus*
419 *influenzae*, *Moraxella catarrhalis*, and *Staphylococcus aureus* in Indonesian children: A cross-
420 sectional study. *PLoS One.* 2018;13:e0195098.
- 421 28. Gessner BD, Sutanto A, Linehan M, et al. Incidences of vaccine-preventable *Haemophilus*
422 *influenzae* type b pneumonia and meningitis in Indonesian children: hamlet-randomised vaccine-
423 probe trial. *Lancet.* 2005;365:43-52.

424

Figure legends

Figure 1: Bacterial loads in high and low SES school-aged children.

All children for which bacterial colonization could be determined were included, thus n = 90 of which 46 low SES (blue) and 44 high SES (red) children. The boxes represent the interquartile range and the line within represents the median. The whiskers represent the 1.5 IQ of the upper and lower quartile bacterial load in each group. P-values are depicted above the corresponding boxplot and are derived with a multivariate regression model, correcting for age, sex and z-BMI. HI: *Hemophilus influenzae*, MC: *Moraxella catarrhalis*, SA: *Staphylococcus aureus*, SES: Socio-economic status, SPN : *Streptococcus pneumoniae*.

Figure 2: Correlation between the loads of different bacteria

All children for which bacterial colonization could be determined were included, thus n = 90. Left lower panel are the bivariate scatterplot of $\log_{10}$ (bacterial loads) of the different bacteria with a regression line in red and for each plot the correlation (Pearson's correlation) and as stars the significance level of this correlation '***' for p-value < 0.001, '**' for p-value ≤ 0.01, '*' for p-value ≤ 0.05 and 'ns' for p ≤ 0.10.

Figure 3: Nasal cytokine levels in high and low SES school-aged children.

All children for which cytokines could be measured were included, thus n = 79 of which 39 low SES (blue) and 40 high SES (red) children. The boxes represent the interquartile range and the line within represents the median. Individuals are depicted by circles. The whiskers represent the 1.5 IQ of the upper and lower quartile. SES: Socio-economic status.

Figure 4: Association between *H. influenzae* (A and B) or *S. pneumoniae* concentration (C and D) and IL-1beta and IL-6 concentration in high and low SES school-aged children. For all models with *H. influenzae* only children colonized are included thus n = 49, which includes 27 high SES children and 22 low SES children. For

all models with *S. pneumoniae* only children colonized are included thus n = 36, which includes 15 high SES children and 21 low SES children.

Figure 5: The IL-1RA and IL-1Beta concentrations in *S. Aureus* carrier and non-carrier in high and low SES. All children for which both cytokines and bacterial colonization could be measured were included, thus n = 73 of which 36 low SES and 37 high SES children. The boxes represent the interquartile range and the line within represents the median. The whiskers represent the 1.5 IQ of the upper and lower quartile concentration of the cytokine. SES: Socio-economic status.

Supplementary figure S1: IL-1 α concentration in *S. aureus* carrier and non-carrier in high and low SES. All children for which both cytokines and bacterial colonization could be measured were included, thus n = 73 of which 36 low SES and 37 high SES children. SES: Socio-economic status.

Figure 1: Bacterial loads in high and low SES school-aged children.

All children for which bacterial colonization could be determined were included, thus n = 90 of which 46 low SES (blue) and 44 high SES (red) children. The boxes represent the interquartile range and the line within represents the median. The whiskers represent the 1.5 IQ of the upper and lower quartile bacterial load in each group. P-values are depicted above the corresponding boxplot and are derived with a multivariate regression model, correcting for age, sex and z-BMI. HI: *Hemophilus influenzae*, MC: *Moraxella catarrhalis*, SA: *Staphylococcus aureus*, SES: Socio-economic status, SPN: *Streptococcus pneumoniae*.

Figure 2: Correlation between the loads of different bacteria

All children for which bacterial colonization could be determined were included, thus $n = 90$. Left lower panel are the bivariate scatterplot of $\log_{10}(\text{bacterial loads})$ of the different bacteria with a regression line in red and for each plot the correlation (Pearson's correlation) and as stars the significance level of this correlation '***' for $p\text{-value} < 0.001$, '**' for $p\text{-value} \leq 0.01$, '*' for $p\text{-value} \leq 0.05$ and '.' for $p \leq 0.10$.

Figure 3: Nasal cytokine levels in high and low SES school-aged children.

All children for which cytokines could be measured were included, thus n = 79 of which 39 low SES (blue) and 40 high SES (red) children. The boxes represent the interquartile range and the line within represents the median. Individuals are depicted by circles. The whiskers represent the 1.5 IQ of the upper and lower quartile. SES: Socio-economic status.

Figure 4: Association between *H. influenzae* (A and B) or *S. pneumoniae* concentration (C and D) and IL-1beta and IL-6 concentration in high and low SES school-aged children. For all models with *H. influenzae* only children colonized are included thus n = 49, which includes 27 high SES children and 22 low SES children. For all models with *S. pneumoniae* only children colonized are included thus n = 36, which includes 15 high SES children and 21 low SES children.

A

B

C

Figure 5: The IL-1RA and IL-1Beta concentrations in *S. Aureus* carrier and non-carrier in high and low SES. All children for which both cytokines and bacterial colonization could be measured were included, thus n = 73 of which 36 low SES and 37 high SES children. The boxes represent the interquartile range and the line within represents the median. The whiskers represent the 1.5 IQ of the upper and lower quartile concentration of the cytokine. SES: Socio-economic status.