

Supplementary Table 1. Characteristics of Study Participants with Functional Seizures.

Age group (years)	Sex	Illness Duration	Psychiatric Diagnoses	CNS-Active Medications
41-45	F	15 years	PD, AG, BP II, MD, PTSD, INS, ICD, SI, SHB	Valproate, Quetiapine
41-45	F	13 years	PD, AG, SAD, HME, MD, ICD, INS, PSY, SI	Pregabalin, Quetiapine, Risperidone, Venlafaxine
26-30	F	8 years	GAD, INS, SI	-
21-25	F	less than a year	MD	-
41-45	F	2 years	INS	-
26-30	F	9 years	PTSD, MD	-
46-50	M	less than a year	PD, SAD, MD, PDD	Quetiapine, Zopiclone
26-30	F	14 years	MD, PTSD, INS	-
46-50	M	3 years	PDD, AUD, INS	Trimipramine, Pregabalin
21-25	F	2 years	GAD, MD, PTSD, INS, PSY, SHB	-
21-25	F	1.5 years	AG, SP, GAD, MD, INS, SHB	Rotigotine

41-45	M	3 years	PD, SP, GAD, MD	Paroxetine
36-40	F	1.5 years	PD, AG, SP, PDD, PTSD, ED, INS	Clonazepam
26-30	F	4 years	GAD, INS	-
21-25	M	less than a year	-	-
51-55	M	4 years	GAD, MD, PDD, ICD, SSD, AUD, INS, SI	-
56-60	F	8 years	PD, GAD, MD, PDD, PTSD, INS	Trazodone, Amitriptyline
21-25	F	less than a year	SP, BP I, MD, SHB, PTSD, ICD, PSY, SI	Levetiracetam
26-30	F	15 years	PD, AG, SAD, MD, OCD, PTSD, ICD, SD, PSY, SI, SHB	Valproate, Levetiracetam, Stiripentol, Perampanel, Pregabalin, Clobazam, Clonazepam, Oxycodone
31-35	F	1 year	AG, SP, GAD, MD, OCD, PTSD, INS, SD, SI	Lamotrigine, Mirtazapine

Psychiatric diagnoses are based on the structured psychiatric interview (Mini-DIPS) administered during this study. Abbreviations: CNS=central nervous system, F=Female, M=Male, PD=panic disorder, AG=agoraphobia, BP=bipolar disorder, MD=major depression, PTSD=post-traumatic stress disorder, INS=insomnia, ICD=impulse control disorder, SI=suicide ideation, SHB=self-harming behavior, SAD=social anxiety disorder, HME=hypomanic episode, PSY=psychotic episodes, GAD=generalized anxiety disorder, PDD=persistent depressive disorder, AUD=alcohol use disorder, SP=specific phobia, ED=eating disorder, SSD=somatic symptom disorder, OCD=obsessive compulsive behavior, SD=sexual dysfunction. In patients with Mini-DIPS diagnoses of panic disorder (PD), seizure semiology was distinct from panic attack symptoms