

Supplementary Table 4. Full list of PANTHER Pathways associated with the genes with high spatial similarity to GM volume correlations for different analyses.

a) Full list of PANTHER Pathways associated with the genes with high spatial similarity to gray matter volume correlation with SDQ-20 Cluster 2, surviving to FDR correction (genes listed in Supplementary Table 4a)

PANTHER Pathways	Fold Enrichment	p (FDR)
5HT3 type receptor mediated signaling	32.36	0.02
Beta3 adrenergic receptor signaling	30.74	0.03
5HT4 type receptor mediated signaling	26.35	0.02
Opioid prodynorphin	25.62	0.02
Opioid proopiomelanocortin	24.92	0.01
Opioid proenkephalin	24.92	0.01
Histamine H2 receptor mediated signaling	24.59	0.03
Beta2 adrenergic receptor signaling	19.62	0.02
Beta1 adrenergic receptor signaling	19.62	0.01
5HT1 type receptor mediated signaling	19.21	0.01
Metabotropic glutamate receptor group II	18.82	0.01
Ionotropic glutamate receptor	18.44	0.01
Dopamine receptor mediated signaling	15.63	0.02
Oxytocin receptor mediated signaling	15.37	0.02
Thyrotropin-releasing hormone receptor signaling	14.87	0.02
5HT2 type receptor mediated signaling	13.36	0.02
Metabotropic glutamate receptor group III	13.17	0.02
Heterotrimeric G-protein signaling-Gi alpha and Gs alpha mediated	7.41	0.02

b) Full list of PANTHER Pathways associated with the genes with high spatial similarity to gray matter volume correlation with TEC Emotional neglect Cluster 2, surviving to FDR correction (genes listed in Supplementary Table 4b)

PANTHER pathways	Fold Enrichment	p (FDR)
Metabotropic glutamate receptor group III	21.63	0.00
Heterotrimeric G-protein signaling-Gq alpha and Go alpha mediated	12.21	0.01
Wnt signaling	6.69	0.01

c) Full list of PANTHER Pathways associated with the genes with high spatial similarity to gray matter volume correlation with TEC Sexual trauma Cluster 1, surviving to FDR correction (genes listed in Supplementary Table 4c)

PANTHER pathways	Fold Enrichment	p (FDR)
5HT1 type receptor mediated signaling	35.76	0.02
5HT4 type receptor mediated signaling	32.69	0.05
GABA-B receptor II signaling	31.78	0.05
Opioid prodynorphin	31.78	0.04
Opioid proopiomelanocortin	30.92	0.04
Opioid proenkephalin	30.92	0.04
Histamine H1 receptor mediated signaling	26.00	0.04
Beta2 adrenergic receptor signaling	24.34	0.05
Beta1 adrenergic receptor signaling	24.34	0.04
Metabotropic glutamate receptor group II	23.35	0.04
Ionotropic glutamate receptor	22.88	0.04
Dopamine receptor mediated signaling	19.39	0.05
Oxytocin receptor mediated signaling	19.07	0.05
Thyrotropin-releasing hormone receptor signaling	18.45	0.05
Heterotrimeric G-protein signaling-Gi alpha and Gs alpha mediated	13.79	0.02

d) Full list of PANTHER Pathways associated with the genes overlapping between SDQ-20 Cluster 2 and TEC Sexual Trauma Cluster 1, surviving to FDR correction (genes listed in Supplementary Table 4d)

PANTHER Pathways	Fold Enrichment	p (FDR)
Beta3 adrenergic receptor signaling	62.41	0.04
Enkephalin release	53.49	0.04
5HT4 type receptor mediated signaling	53.49	0.03
GABA-B receptor II signaling	52.01	0.02
Opioid prodynorphin	52.01	0.02
Opioid proopiomelanocortin	50.60	0.02
Opioid proenkephalin	50.60	0.01
Histamine H1 receptor mediated signaling	42.55	0.02
Beta2 adrenergic receptor signaling	39.84	0.02
Beta1 adrenergic receptor signaling	39.84	0.02
5HT1 type receptor mediated signaling	39.01	0.02
Metabotropic glutamate receptor group II	38.21	0.02
Dopamine receptor mediated signaling	31.73	0.02
Oxytocin receptor mediated signaling	31.20	0.02
Thyrotropin-releasing hormone receptor signaling	30.20	0.02
5HT2 type receptor mediated signaling	27.13	0.02

Linking gene expression patterns and brain morphometry to trauma and symptom severity in patients with functional seizures (Jungilligens et al.)

Heterotrimeric G-protein signaling -Gi alpha and Gs alpha mediated	16.92	0.02
Inflammation mediated by chemokine and cytokine signaling	11.01	0.02