***De novo* emergence of a remdesivir resistance mutation during treatment of persistent SARS-CoV-2 infection in an immunocompromised patient: A case report**

Shiv Gandhi^{*1†}, Jonathan Klein^{*2}, Alexander Robertson^{3*}, Mario A. Peña-Hernández^{*2}, Michelle J Lin⁴, Pavitra Roychoudhury⁴, Peiwen Lu², John Fournier¹, David Ferguson⁵, Shah A. Mohamed Bakhash⁴, M. Catherine Muenker³, Ariktha Srivathsan³, Elsio A. Wunder Jr³, Nicholas Kerantzas⁶, Wenshuai Wang⁷, Brett Lindenbach⁸, Anna Pyle^{7,9,11}, Craig B. Wilen^{2,6}, Onyema Ogbuagu¹, Alexander L. Greninger^{4,10}, Akiko Iwasaki^{2,3,7,11}, Wade L. Schulz^{5,6}, and Albert I. Ko^{1,3†}

¹Department of Medicine, Section of Infectious Diseases, Yale University School of Medicine, New Haven, CT, USA

²Department of Immunobiology, Yale University School of Medicine, New Haven, CT, USA

³Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT, USA

⁴Department of Laboratory Medicine, University of Washington School of Medicine, Seattle, Washington, USA

⁵Center for Outcomes Research and Evaluation, Yale New Haven Hospital, New Haven, CT, USA

⁶Department of Laboratory Medicine, Yale School of Medicine, New Haven, CT, USA

⁷Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT, USA

⁸Department of Microbial Pathogenesis, Yale School of Medicine, New Haven, CT, USA

⁹Department of Chemistry, Yale University, New Haven, CT, USA

¹⁰Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA

¹¹Howard Hughes Medical Institute, Chevy Chase, MD, USA.

*These authors contributed equally

Corresponding Authors[†]: S.G. (shiv.gandhi@yale.edu), A.I.K. (albert.ko@yale.edu)

Abstract

SARS-CoV-2 remdesivir resistance mutations have been generated *in vitro* but have not been reported in patients receiving treatment with the antiviral agent. We present a case of an immunocompromised patient with acquired B-cell deficiency who developed an indolent, protracted course of SARS-CoV-2 infection. Remdesivir therapy alleviated symptoms and produced a transient virologic response, but her course was complicated by recrudescence of high-grade viral shedding. Whole genome sequencing identified a mutation, E802D, in the nsp12 RNA-dependent RNA polymerase, which was not present in pre-treatment specimens. *In vitro* experiments demonstrated that the mutation conferred a ~6-fold increase in remdesivir IC₅₀ but resulted in a fitness cost in the absence of remdesivir. Sustained clinical and virologic response was achieved after treatment with casirivimab-imdevimab. Although the fitness cost observed *in vitro* may limit the risk posed by E802D, this case illustrates the importance of monitoring for remdesivir resistance and the potential benefit of combinatorial therapies in immunocompromised patients with SARS-CoV-2 infection.

Text

Remdesivir (RDV), an adenosine nucleoside analog that interferes with the SARS-CoV-2 RNA polymerase, nsp12, when converted to its active form¹, has been widely used in the treatment of patients hospitalized with COVID-19. *In vitro* selection experiments have generated mutations at residue 802 of nps12, which confer resistance to RDV². These mutations have been detected in genome sequences deposited in GISAID², but their emergence has not been demonstrated in patients receiving RDV therapy. Selection for RDV resistance is a specific concern for immunocompromised patients who may receive multiple courses of the antiviral agent for persistent SARS-CoV-2 infection³⁻⁵. Here, we report a case of an immunocompromised patient

with an indolent, protracted course of SARS-CoV-2 infection from whom a RDV resistance mutation, E802D, was identified during recrudescence of viral shedding following treatment with the antiviral agent.

A woman in her 70s with Stage IV Non-Hodgkin's lymphoma (NHL) had completed a course of rituximab and bendamustine in March 2019, which was complicated by lymphocytopenia and hypogammaglobulinemia. The NHL was in remission when she developed a SARS-CoV-2 infection in May 2020, which presented as the acute onset of fever, anosmia, cough and rhinorrhea (Figure 1a) and was confirmed by reverse transcriptase-polymerase chain reaction (RT-PCR) on day 0 (Figure 1b). The patient was hospitalized twice in the following two months for persistent fever and new onset neutropenia and anemia (Figure 1c), during which she had low SARS-CoV-2 RT-PCR cycle threshold (Ct) values from nasopharyngeal specimens (Ct 20.5 and 17.1 on days 17 and 36, respectively, Figure 1b), undetectable serum anti-SARS-CoV-2 IgG antibody (Abbott), and elevated serum inflammatory markers (Figure 1c). Chest computed tomography (CT) revealed bilateral ground glass opacities (day 38, Figure 1d), but she did not experience dyspnea, hypoxemia or recurrence of respiratory symptoms during her hospitalizations and subsequent course of illness. The workup for other sources of fever, which included a bone marrow biopsy, was unrevealing. CD19⁺ B-cells were not identified in the bone marrow biopsy or by flow-cytometry of peripheral blood mononuclear cells (PBMC). The patient was initiated on treatment with filgrastim 1-2 times per week and monthly intravenous immunoglobulin (IVIG) to manage her neutropenia and hypogammaglobulinemia, respectively, and both were continued after hospital discharge on day 51.

The patient continued to have daily fevers, except for a 30-day period of defervescence (days 103-132), refractory neutropenia, anemia, and positive RT-PCR test results with Ct values <25

when she was hospitalized at our institution on day 145. Her chest CT on day 140 was notable for increased opacities compared with prior examinations (Figure 1d). The patient initiated a 10-day course of RDV (days 148-157), which precipitated a resolution of her fever (day 149, Figure 1a), normalization of CRP (day 156, Figure 1c) and improvement in the opacities on chest CT (day 162, Figure 1d). An initial virologic response to therapy was observed, with an increase in RT-PCR Ct values and decreasing numbers of plaque forming units (PFU) in viral cultures and lower proportions of SARS-CoV-2 subgenomic RNA (sgRNA), measured in amplicon sequencing data, in her respiratory tract specimens by day 152 (Figure 2a).

However, a recrudescence of viral shedding occurred during and after RDV therapy, as evidenced by decreasing RT-PCR Ct values which reached 18 on day 160 and increasing PFU and sgRNA in respiratory tract specimens on day 156 (Figure 2a). The patient received an 8g infusion of casirivimab-imdevimab^{6,7} on day 163 after expanded access for the use of the monoclonal antibody (mAb) therapy was approved. A rapid and sustained virologic response was observed with undetectable cultured virus, low sgRNA and negative NP RNA in specimens by days 164, 166 and 217, respectively. The patient's anosmia resolved roughly 17 days after administration of casirivimab-imdevimab (day 180, Figure 1a). During the following five-month period (days 160-292), the patient did not have recurrence of COVID-19 related symptoms and had a chest CT which showed minimal opacities (day 170, Figure 1d), and had high anti-SARS-CoV-2 S1 protein IgG titers (Figure 2a). Peripheral blood neutrophil counts, hemoglobin levels and serum inflammatory markers normalized during convalescence (Figure 1c).

To identify SARS-CoV-2 mutations that arose during the course of illness, we longitudinally sampled multiple sources of the patient's tissues and secretions (Supplemental Table 1) and performed WGS on an Illumina NextSeq platform using the Swift SARS-CoV-2 multiplex

amplicon sequencing panel⁸. Viral N1 or N2 RNA was detected in 19 of 21 nasopharyngeal, 10 of 10 saliva, 10 of 10 stool and 0 of 10 whole blood specimens. Among the 39 samples with detectable N1 or N2 RNA, we assembled 27 whole SARS-CoV-2 genomes from nasopharyngeal (12), saliva (9) and stool (6) specimens. One of the nasopharyngeal specimens was obtained from the initial phase of illness (day 36, Figure 2a). Phylogenetic analysis found that sequenced genomes belonged to a single lineage (Pango B1) within Nextstrain clade 20C (Figure S1), indicating that viral genomes identified during the course of illness were derived from intra-host diversification following infection with a single strain.

Analysis of assembled viral genomes identified a mutation, E802D, in nsp12, whose detection in patient specimens was temporally associated with RDV therapy (Figure 2b). E802D was not identified at an allele frequency above 1% in either the specimen obtained during the initial phase of illness (day 36) or in specimens collected during the first 5 days (days 148-152) of RDV therapy. The mutation was first detected 7 days after initiation of RDV therapy (day 155) and accounted for 23% and 96% of the allele frequency in nasopharyngeal and saliva specimens, respectively, by day 160. The E802 residue of nsp12 resides in the palm sub-domain which contains a portion of the residues that comprise the active site of the SARS-CoV-2 RNA polymerase^{1,2,9}. The E802 residue participates in an electrostatic network with D804 and K807 which stabilizes the loop involved with binding to the nascent RNA (Figure 2c, S2a). E802D has been identified in an *in vitro* RDV resistance selection experiment and was found to confer a ~2.5-fold increase in IC₅₀ to the drug². Together, the temporality of E802D emergence in the patient, its location in nsp12 and the *in vitro* identification of the same mutation with a RDV resistance phenotype support the plausibility that that RDV treatment of the patient selected for

variants with the E802D mutation, which in turn contributed to the observed rebound viral shedding.

To validate the resistance phenotype of E802D, we engineered this mutation and an E802A mutation, which had been also shown to confer RDV resistance², into an infectious molecular clone of SARS-CoV-2/WA01 (icSARS-CoV-2-mNG), which expresses the mNeon Green reporter and is ORF7a depleted¹⁰. Replication kinetics of the E802D and E802A mutants in Vero-E6 cells revealed decreased viral replication relative to parental icSARS-CoV-2-mNG (Figure 2d), suggesting that these mutations may impart a fitness cost. In the presence of high concentrations of RDV (5 μ M), the E802D mutant replicated to higher titers than the parental virus (Figure S2b), confirming that this mutation confers resistance to RDV. RDV dose-response curves demonstrated that E802D and E802A mutants had significant increases in IC₅₀ values (4.2 μ M and 2.7 μ M, respectively) relative to parental icSARS-CoV-2-mNG (0.7 μ M) (Figure 2e). Assessment of RDV cytotoxicity demonstrated limited impact on cell death rates below 10 μ M, indicating that the RDV resistance phenotype conferred by mutations at E802 may occur at physiologically relevant concentrations (Fig S2c).

These findings suggest that RDV resistance mediated by substitutions at residue 802 in nsp12 may lead to therapeutic failure in the setting of an immunocompromised host. We speculate that mutations at 802 distort the active site in a way that either enables the enzyme to exclude RDV or alleviates the steric clash mediated by S861, thereby allowing the enzyme to escape RDV-mediated chain termination^{1,9}. Further biochemical and structural characterization will be needed to delineate the mechanism by which substitutions at residue E802 confer resistance to RDV.

The therapeutic impact of RDV resistance mutations at E802 may be attenuated by a fitness cost imparted by these mutations, which we observed *in vitro*. Although RDV therapy has been

widely administered to patients during the pandemic, E802D and all substitutions at residue 802 have been found in 131 and 297, respectively, of the 4.8M genome sequences obtained from patient isolates in the GISAID database (www.gisaid.org; accessed 11/07/21). Consistent with the observation of a fitness cost, the allele frequency of E802D in nasopharyngeal specimens from the patient decreased from 22.6% to <1% between the period (day 160-162) after completion of RDV therapy and prior to casirivimab-imdevimab administration (Figure 2b). However, E802 allele frequencies were significantly higher (50-96% at days 158-167) in saliva specimens during this period and did not decline until casirivimab-imdevimab was administered. An explanation for these discordant findings is unclear and may relate to the differential compartmentalization of variants within the respiratory tract or the stochastic nature of sampling the nasopharyngeal compartment as opposed to saliva where virus from a wider geographic distribution may pool. Given that saliva is infrequently sampled, RDV resistance mediated by E802 mutations may be under-detected. Furthermore, analysis of the 131 genomic sequences that have been found in GISAID to contain the E802D mutation identified two instances where sequences were geographically and temporally clustered, suggesting that local transmission of E802D variants may have occurred (Figure S3). Continued surveillance will thus be needed to elaborate whether the occurrence of E802D, as well as other substitutions at this residue, will be limited because of an underlying fitness cost or alternatively, emerge to pose a broader risk for RDV resistance.

We did not identify nsp12 mutations at sites other than residue 802 which were temporally associated with RDV therapy. A mutation at D484Y has been reported in a post-treatment sample from an immunocompromised patient who failed RDV therapy¹¹, but the RDV resistance phenotype associated with this mutation, whose location is distal to the active site, has not been

confirmed. We identified two additional mutations, A504V in nsp14 (exonuclease) and I115L in nsp15 (endoRNase), whose allele frequencies increased during and after RDV treatment (Figure S4). However, A504V in nsp14 was present (allele frequency 85.1%) in a specimen obtained during her initial phase of illness (day 36). Although I115L in nsp15 was not detected in the early-phase specimen, the plausibility of a mutation in the endoRNase gene which confers RDV resistance is less clear.

The case highlights additional issues for the management of SARS-CoV-2 infection in immunocompromised patients. Firstly, the course of infection was notable for being indolent despite a six-month period of sustained high-grade viral shedding and underlying lung involvement. We identified six mutations, including a frameshift mutation in ORF3a (N257fs), which were present throughout the course of illness prior to RDV treatment (Figure S5). Yet none provides a clear explanation for an attenuated phenotype. Longitudinal profiling of PBMC demonstrated profound depletion of circulating B-cell populations and a T-cell phenotype consistent with chronic antigen exposure as evidenced by the increased activation (CD38+, HLA-DR+) and exhaustion (PD-1, Tim-3) markers (Figure S6). Given reports that maladaptive immune responses contribute to the risk of severe COVID-19^{12,13}, we speculate that the compromised immune response was unable to control viral load yet conversely, protected the patient from developing severe complications.

Secondly, SARS-CoV-2 infection was associated with the development of severe refractory neutropenia and moderate anemia in the patient (Figure 1c). While we are unable to discern if the cytopenia was due to direct or indirect effects of infection, SARS-CoV-2 has been detected in the bone marrow of an immunocompromised patient who developed pancytopenia during

infection¹⁴. Our patient's cytopenia resolved after casirivimab-imdevimab therapy, suggesting a causal link between SARS-CoV-2 infection and cytopenia in this setting.

Finally, the case illustrates the potential benefit of mAb therapy for immunocompromised patients with persistent SARS-CoV-2 infection. Prior to initiation of mAb therapy, two non-synonymous mutations, H655Y (Figure S7a) and P812L (Figure S7b), emerged which were not present in the initial specimen (day 36), indicating that spike protein continues to evolve and adapt to the host in the absence of selective pressure from an adaptive immune response. One week after casirivimab-imdevimab was administered (day 170), we identified a new spike protein mutation, A348S in the RBD domain, which persisted to day 177 (Figure S7c). WGS was not performed on specimens collected after day 177 because of low amounts or undetectable viral RNA. A348S is not proximal to the mAb binding sites (Figure S7d)¹⁵ and was not identified in a comprehensive analysis of resistance mutations⁶. In the setting of a negligible cellular immune response, mAb therapy cleared viral shedding, abrogated the residual complaint of anosmia, and resolved the blood dyscrasias.

In summary, we identified the *de novo* emergence of a RDV-resistance mutation, E802D, following initiation of RDV in an immunocompromised patient with persistent SARS-CoV-2 infection. While the finding is limited to a single case and requires confirmation of its generalizability in larger patient populations, it suggests that RDV may impart a selective pressure *in vivo* to drive evolution of the virus. E802D is associated with a fitness cost *in vitro*, which may limit the broader impact of this mutation on the development of secondary resistance during treatment and the risk for primary resistance through transmission of resistant variants. Yet, our findings underscore the importance of immunocompromised hosts with uncontrolled viral replication as a source of genetic diversification^{3,4,16–18} and selection of mutations that may

potentially impart adverse consequences for antiviral therapy. Enhanced genomic surveillance of immunocompromised patients may thus be warranted. As observed in this case, initiation of anti-SARS-CoV-2 mAb may serve as a therapeutic option to achieve rapid and sustained virologic responses and improved clinical outcomes in immunocompromised patients.

Online Methods

Regulatory and ethical approvals

Regulatory approval to administer an 8g dose of casirivimab/imdevimab was obtained from the U.S. Food and Drug Administration under investigational new drug application no. 153390.

This study was also approved by Yale Human Research Protection Program Institutional Review Boards (FWA00002571, protocol ID 2000027690, 2000029277). Informed consent was obtained from both the patient and healthcare workers.

RNA extraction and PCR

RNA was extracted on a Qiagen Bio Robot (Qiagen, Hilden, Germany) using the QIAamp Virus Bio Robot MDx Kit (Qiagen) following manufacturer's instructions. RT-PCR was performed using AgPath-ID One-Step RT-PCR kit (Life Technologies, Carlsbad, CA, USA) with CDC primers and probes for N1/N2 following protocol and concentrations defined previously¹⁹. Each 25 µL of reaction mix included 2X RT-PCR buffer, 25X enzyme mix, primers-probes, and 5 µL of the extracted RNA. Cycle parameters on the ABI 7500 real-time PCR system (Applied Biosystems) were 10 min at 48°C for reverse transcription, 10 min of inactivation at 95°C, 40 cycles of 15 sec at 95°C, and 45 sec at 60°C. Internal control (EXO), positive, and negative (water) control were included on every run.

Library preparation and sequencing

Using 11 µL of extracted RNA, single strand complementary DNA (sscDNA) was synthesized using SuperScript IV First-Strand Synthesis System (Thermofisher). Sequencing libraries were prepared using the Swift Normalase Amplicon Panel (SNAP) for SARS-CoV-2 (Swift

Biosciences, Ann Arbor, MI, USA) as described previously⁸. Libraries were quantified using fluorometric methods (Quant-iT dsDNA high sensitivity kit, Life Technologies, Carlsbad, CA, USA), normalized, and sequenced on the Illumina NextSeq 500 or 2000 instruments (Illumina, San Diego, CA, USA) using 2x150 reads.

Analysis of sequencing data

Reads were processed using a custom bioinformatic pipeline (https://github.com/greninger-lab/covid_swift_pipeline), as described⁸. Briefly, raw FASTQs were trimmed of adapter content and low-quality reads, then aligned to the NCBI reference Wuhan-1 (NC_045512). PCR primers were soft-clipped with Primerclip (<https://github.com/swiftbiosciences/primerclip>) and a consensus genome was generated from this alignment using bcftools v1.9²⁰. Variants were examined longitudinally with a modified Python script adapted from LAVA (<https://github.com/greninger-lab/lava>).

A k-mer based approach was used to identify reads corresponding to subgenomic RNAs. After removing adapters and low-quality regions, junction-spanning reads corresponding to individual sgRNAs were filtered using BBduk (<https://sourceforge.net/projects/bbmap/>) against a custom FASTA file composed of 30 nucleotides of the 5' leader sequence and 30 nucleotides of the respective 3' junctions of downstream ORFs²¹, allowing an edit distance of 1. In order to avoid calling of genomic reads, we constrained matches to those that had 10 consecutive matches of 31-mers.

Isolation of patient plasma and PBMCs

As described previously, patient whole blood was collected in sodium heparin-coated vacutainers and gently agitating at room temperature until sample pick-up by IMPACT team members. All blood was processed on the day of collection. Plasma samples were collected after centrifugation of whole blood at 400g for 10 minutes at room temperature (RT) without brake. The undiluted serum was then transferred to 15-ml polypropylene conical tubes, and aliquoted and stored at -80°C for subsequent analysis. PBMCs were isolated using Histopaque (Sigma-Aldrich, #10771-500ML) density gradient centrifugation in a biosafety level 2+ facility. After isolation of undiluted serum, blood was diluted 1:1 in room temperature PBS, layered over Histopaque in a SepMate tube (StemCell Technologies; #85460) and centrifuged for 10 minutes at 1,200g. The PBMC layer was isolated according to the manufacturer's instructions. Cells were washed twice with PBS before counting. Pelleted cells were briefly treated with an ACK lysis buffer for 2 minutes and then counted. Percentage viability was estimated using standard Trypan blue staining and an automated cell counter (Thermo-Fisher, #AMQAX1000).

Flow cytometry

As described previously²², antibody clones and vendors used for flow cytometry analysis were as follows: BB515 anti-hHLA-DR (G46-6) (1:400) (BD Biosciences), BV605 anti-hCD3 (UCHT1) (1:300) (BioLegend), BV785 anti-hCD4 (SK3) (1:200) (BioLegend), APCFire750 or PE-Cy7 or BV711 anti-hCD8 (SK1) (1:200) (BioLegend), BV421 anti-hCCR7 (G043H7) (1:50) (BioLegend), AlexaFluor 700 anti-hCD45RA (HI100) (1:200) (BD Biosciences), PE anti-hPD1 (EH12.2H7) (1:200) (BioLegend), APC anti-hTIM3 (F38-2E2) (1:50) (BioLegend), BV711 anti-hCD38 (HIT2) (1:200) (BioLegend), BB700 anti-hCXCR5 (RF8B2) (1:50) (BD Biosciences), PE-Cy7 anti-hCD127 (HIL-7R-M21) (1:50) (BioLegend) and PE-CF594 anti-hCD25 (BC96)

(1:200) (BD Biosciences). In brief, freshly isolated PBMCs were plated at $1-2 \times 10^6$ cells per well in a 96-well U-bottom plate. Cells were resuspended in Live/Dead Fixable Aqua (ThermoFisher) for 20 min at 4 °C. Following a wash, cells were blocked with Human TruStan FcX (BioLegend) for 10 min at RT. Cocktails of desired staining antibodies were added directly to this mixture for 30 min at RT. For reinfection stains, cells were first washed and supernatant aspirated; then to each cell pellet a cocktail of secondary markers was added for 30 min at 4 °C. Prior to analysis, cells were washed and resuspended in 100 µl 4% PFA for 30 min at 4 °C. Following this incubation, cells were washed and prepared for analysis on an Attune NXT (ThermoFisher). Data were analysed using FlowJo software version 10.6 software (Tree Star). The specific sets of markers used to identify each subset of cells are summarized in ²⁰Figure S6.

SARS-CoV-2 specific-antibody measurements

ELISAs were performed as previously described²³. Triton X-100 and RNase A were added to serum samples at final concentrations of 0.5% and 0.5mg/ml respectively and incubated at room temperature (RT) for 30 minutes. 96-well MaxiSorp plates (Thermo Scientific #442404) were coated with 50 µl/well of recombinant SARS Cov-2 S1 protein (ACROBiosystems #S1N-C52H3-100ug) at a concentration of 2 µg/ml in PBS and were incubated overnight at 4 °C. The coating buffer was removed, and plates were incubated for 1h at RT with 200 µl of blocking solution (PBS with 0.1% Tween-20, 3% milk powder). Serum was diluted 1:50 in dilution solution (PBS with 0.1% Tween-20, 1% milk powder) and 100 µl of diluted serum was added for two hours at RT. Plates were washed three times with PBS-T (PBS with 0.1% Tween-20) and 50 µl of HRP anti-Human IgG Antibody (GenScript #A00166, 1:5,000) diluted in dilution solution added to each well. After 1 h of incubation at RT, plates were washed three times with PBS-T.

Plates were developed with 100 µl of TMB Substrate Reagent Set (BD Biosciences #555214) and the reaction was stopped after 15 min by the addition of 2 N sulfuric acid. Plates were read at a wavelength of 450 nm and 570nm.

Cell lines and virus

As previously described²², Vero E6 kidney epithelial cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 1% sodium pyruvate (NEAA) and 5% fetal bovine serum (FBS) at 37°C and 5% CO₂. The cell line was obtained from the ATCC and has been tested negative for contamination with mycoplasma. SARS-CoV-2, strain USA-WA1/2020, was obtained from BEI Resources (#NR-52281) and was amplified in Vero E6 cells. Cells were infected at a MOI 0.01 for four three days to generate a working stock and after incubation the supernatant was clarified by centrifugation (450g × 5min) and filtered through a 0.45-micron filter. The pelleted virus was then resuspended in PBS then aliquoted for storage at – 80°C. Viral titers were measured by standard plaque assay using Vero E6 cells. All experiments were performed in a biosafety level 3 with the Yale Environmental Health and Safety office approval.

Virus titration from nasopharyngeal swabs.

SARS-CoV-2 titers from longitudinal nasopharyngeal swabs were assessed by conventional plaque assay. Nasopharyngeal swabs were collected and swabs stored and frozen in universal viral transport media (VTM) prior to plaquing. Each nasopharyngeal sample was ten-fold serially diluted (dilution range from 1:10 to 1:1000000). Diluted NP samples were added on a monolayer of Vero E6 cells. After 1 hour of infection, DMEM (2% FBS), and 1.2% of Avicel (Dupont, 9004346) were overlaid. 72 hours post inoculation, media was removed and 4%

Paraformaldehyde added for 1 hour. Cells were stained with 0.1% of crystal violet to visualize SARS-CoV-2 plaques. Plaques were counted by single operator. Each sample was tested in duplicate.

Site-directed mutagenesis and full length icSARS-CoV-2 mNG E802D and E802A production.

The reverse genetics system used to generate E802D and E802A SARS-CoV-2 mutants is described at length elsewhere¹⁰. The icSARS-CoV-2 mNG (mNeonGreen) backbone previously reported was employed to construct both full-length E802D and E802A mutant viruses utilizing the following primer pairs: forward (E802D): *gttggactgaTactgaccttactaaaggac*, reverse (E802D): *aggtcagtAtcagtcaccaacattttgcttc*; and forward (E802A): *gttggactgCgactgaccttactaaaggac*, reverse (E802A): *aggtcagtcGcagtcaccaacattttgcttc*. Site-directed mutagenesis was performed using Pfu Turbo Polymerase (Agilent) according to manufacturer's instructions. E802D and E802A mutations were confirmed in icSARS-CoV-2 mNG backbone by Sanger sequencing using the primer: *gatgatactctctgacgatg*.

Steps to produce full-length infectious virus were performed as described previously¹⁰, with the following minor modifications: Plasmid propagation and digestion were performed as described and excision of DNA bands from the gel was done by using a Blue-Light transilluminator (Accuris). Next, equal molar amounts of the seven plasmids were reduced by half relative to the prior reports to perform the full-length genomic ligation: F1 (.305 µg), F2 (.325 µg), F3 (.375 µg), F4 (.47 µg) were ligated using T4 DNA ligase with a final volume of 50 µl. Plasmids F5 (.375 µg), F6 (.36 µg) and F7 (.3 µg) were incubated with T4 DNase Ligase in a volume of 40 µl.

The primary reactions were incubated at 4°C for 16 h. After this, the two reactions were mixed and volumed up to 100 µl by adding the corresponding additional volume of ligation reagents. The secondary ligation was incubated for 18 h at 4°C. The full-length genomes were purified by using the Genomic DNA Clean and Concentrator kit from Zymo Research according to manufacturer's instructions. Successful ligation products were visualized using a 0.8% agarose gel and standard gel imager.

Next, full-length *in vitro* RNA transcription was also performed with additional modifications. First, 500 ng of each ligation product was employed. Reactions were performed using the mMESSAGE mMACHINE T7 Transcription Kit (Thermo Scientific) with a cap analog-to-GTP ratio of 1:1. Incubation of the reaction was done at 32°C for 12 h. Following DNase treatment was performed at 37°C for 15 min. Next, viral RNA was purified by using the RNA Clean and Concentrator kit from Zymo Research according to the manufacturer's instructions. Successful transcription products were visualized using a 0.8% agarose gel and standard gel imager.

RNA transcripts were next electroporated into BHK-N expressing cells (courtesy of Dr. Brett Lindenbach). 15 µg of each mutant transcript was electroporated into BHK cells using the Electro Square Porator Device (BTX). Electroporated cells were seeded into a T75 flask containing 1:2 ratio of BHK-N:Vero-E6 cells. Complete monolayer CPE was evident at 72 h post-electroporation and supernatant was centrifuged and .22 µm filtered. Vero-E6 were infected with 500 µL of each of the clarified supernatants. Supernatants were collected for each mutant virus at 48 hours post infection and tittered as described previously. Infectious clones are available with MTA and appropriate biosafety approvals.

Sequencing and variant calling

RNA was extracted from SARS-CoV-2 mutants using the MagMAX viral/pathogen nucleic acid isolation kit and tested with the multiplexed RT-qPCR variant assay²⁴. Using a standard dilution series, Ct values were converted to RNA copies per μL . Extracted RNA was diluted to 1 million copies per μL and used as input into the COVIDSeq Test RUO version for library preparation, in duplicate. Pooled libraries were sequenced on the NovaSeq (paired-end 150) at the Yale Center for Genome Analysis. Single nucleotide variants were called at a minimum frequency threshold of 0.02 and minimum read depth of 400X using iVar (version 1.3.1)²⁵, and filtered between duplicate replicates. Introduction of the E802D and E802A was confirmed at a minimum frequency of 0.82-0.9 in initial stocks.

Replication kinetics of icSARS-CoV-2 mutants.

Isolates from initial stocks were expanded and used to infect Vero-E6 cells with 0.01 MOI of each E802D and E802A mutant virus to assess their replication kinetics. Supernatants were collected at 8, 24 and 48 hours post-infection. Titers for each time point were assessed by conventional plaque assay as described above. Each virus was assessed in triplicate.

Remdesivir resistance assay.

Each icSARS-CoV-2 mutant virus (E802D and E802A) was cultured in the presence of increasing concentrations of RDV, ranging from 0.39 to 40 μM . Infections were performed using Vero-E6 in 384-well plates containing the corresponding RDV dilutions in phenol-red free DMEM media with 5% FBS and incubated at 37°C for 4 hours. Next, cells were infected with

0.01 MOI of each mutant virus and infected cell frequencies were measured at 48 hours post-infection by mNeonGreen expression by high content imaging (Cytation 5, BioTek)²⁶. Cell culture supernatants were also collected for plaque assay. Cell viability in uninfected cells was assessed at 72 h post-infection using the CellTiter-Glo kit (Promega) according to the manufacturer's instructions.

Neutralization assay

As described previously²⁷, patient sera was isolated from whole blood and heat treated for 30 minutes at 56 °C. Sixfold serially diluted plasma, from 1:10 to 1:2430 were incubated with SARS-CoV-2 for 1 h at 37 °C. The mixture was subsequently incubated with Vero E6 cells in a 6-well plate for 1 hour for infection. Cells were overlaid with DMEM supplemented NaHCO₃, 2% FBS, and 0.6% Avicel mixture²⁶. Plaques were resolved at 40h post infection by fixing in 4% formaldehyde for 1h followed by staining in 0.5% crystal violet. All experiments were performed in parallel with healthy, uninfected control sera to establish degree of protection as reflected in difference in plaque counts.

Phylogenetic analysis

Phylogenetic trees were generated using the Nextstrain software package²⁸. 326 background sequences from CT, USA collected April 1st-September 30th 2020 were downloaded from GISAID (Supplemental Table 3). The tree was rooted using the Wuhan reference strain. Genome sequences which contain the E802D mutation were identified from the GISAID database (accessed on 10/31/21; Supplemental Table 4) and were compared to sequences in the global Auspice dataset.

Data availability

Sequencing data are available under NCBI BioProject no. PRJNA774781 (Supplementary Table 2).

Acknowledgements

The authors thank Dr. Michael Jacobs, Dr. Sanjay Bhagani and Dr. David Lowe at the Royal Free Hospital for their advice regarding the management of the patient, Dr. Jennifer Hamilton, David Stein, Dr. Wendy Kampman and Dr. David M. Weinreich at Regeneron Pharmaceuticals, Inc. for their guidance on the use of monoclonal antibody therapy and laboratory evaluations, Nathaniel Price at Yale School of Medicine for his assistance in identifying data and samples for the analyses, and Dr. Chantal Vogels, Dr. Joseph Fauver, Mallery Breban and Dr. Nathan Grubaugh at the Yale School of Public Health for sequencing infectious clones and providing advice on the analysis and interpretation of the genomic data. Casirivimab and imdevimab was provided by Regeneron Pharmaceuticals, Inc. for the purpose of emergency compassionate use treatment. We also thank the health care staff at Yale New Haven Hospital, which include Dr. Jensa Morris, Dr. Anne Spichler, Dr. Frederick Altice and Dr. Nikhil Seval, who cared for the patient during her hospital course and provided key input, and most of all, the patient for her equanimity and contributions to this study.

This study was supported by the Beatrice Kleinberg Neuwirth Fund; the Sendas Family Fund, Yale School of Public Health; and Department of Internal Medicine at the Yale School of Medicine. SG was supported by a grant from the National Institutes of Health (2T32AI007517-

21A1). C.B.W. was supported by Burroughs Welcome Fund, Ludwig Family Foundation, and Mathers Foundation.

Contributions

J.K., A.R., and M.P.H contributed to data collection, analysis, and writing. M.L., P.R., J.F., D. F., C.V., M.C.M., A.S., E.W, N.K., P.L, W.W., A.P. contributed to data collection and analysis. B.L. contributed to study design. O.O., A.G, C.B.W, A.I., and W.L.S. conceived of the study and contributed to study design, interpretation, and editing of the manuscript. S.G. and A.I.K conceived of the study and contributed to study design, data interpretation, writing, and regulatory approvals.

Competing Interests

ALG reports institutional central testing contracts from Abbott and research grants from Merck and Gilead, outside of the proposed work. AIK received consulting fees from Tata Sons and is recipient of grants on COVID-19 from Merck, Regeneron and Serimmune all of which are outside the submitted work. OO received consulting fees from Gilead and ViiV, as well as research support and honoraria from Gilead, outside of the submitted work. WLS is a consultant for Hugo Health, founder of Refactor Health and is recipient of grants on COVID-19 from Merck and Regeneron, all of which are outside the submitted work.

References

1. Kokic, G. *et al.* Mechanism of SARS-CoV-2 polymerase stalling by remdesivir. *Nat Commun* 12, 279 (2021).

2. Szemiel, A. M. *et al.* In vitro selection of Remdesivir resistance suggests evolutionary predictability of SARS-CoV-2. *Plos Pathog* 17, e1009929 (2021).
3. Collaboration, T. C.-N. B. C.-19 *et al.* SARS-CoV-2 evolution during treatment of chronic infection. *Nature* 1–10 (2021) doi:10.1038/s41586-021-03291-y.
4. Choi, B. *et al.* Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host. *New Engl J Med* 383, 2291–2293 (2020).
5. Collaboration, C.-N. C.-19 B. *et al.* Treatment of COVID-19 with remdesivir in the absence of humoral immunity: a case report. *Nat Commun* 11, 6385 (2020).
6. Baum, A. *et al.* Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. *Science* 369, 1014–1018 (2020).
7. Baum, A. *et al.* REGN-COV2 antibodies prevent and treat SARS-CoV-2 infection in rhesus macaques and hamsters. *Science* eabe2402 (2020) doi:10.1126/science.abe2402.
8. Addetia, A. *et al.* Sensitive Recovery of Complete SARS-CoV-2 Genomes from Clinical Samples by Use of Swift Biosciences’ SARS-CoV-2 Multiplex Amplicon Sequencing Panel. *J Clin Microbiol* 59, (2020).
9. Yin, W. *et al.* Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-CoV-2 by remdesivir. *Science* 368, 1499–1504 (2020).
10. Xie, X. *et al.* Engineering SARS-CoV-2 using a reverse genetic system. *Nat Protoc* 16, 1761–1784 (2021).
11. Martinot, M. *et al.* Remdesivir failure with SARS-CoV-2 RNA-dependent RNA-polymerase mutation in a B-cell immunodeficient patient with protracted Covid-19. *Clin Infect Dis* ciaa1474- (2020) doi:10.1093/cid/ciaa1474.
12. Wang, E. Y. *et al.* Diverse Functional Autoantibodies in Patients with COVID-19. doi:10.1101/2020.12.10.20247205.
13. Wilk, A. J. *et al.* A single-cell atlas of the peripheral immune response in patients with severe COVID-19. *Nat Med* 26, 1070–1076 (2020).
14. Issa, N., Lacassin, F. & Camou, F. First case of persistent pancytopenia associated with SARS-CoV-2 bone marrow infiltration in an immunocompromised patient. *Ann Oncol* 31, 1418–1419 (2020).
15. Hansen, J. *et al.* Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science* 369, 1010–1014 (2020).

16. Drouin, A. C. *et al.* Successful Clearance of 300 Day SARS-CoV-2 Infection in a Subject with B-Cell Depletion Associated Prolonged (B-DEAP) COVID by REGEN-COV Anti-Spike Monoclonal Antibody Cocktail. *Viruses* 13, 1202 (2021).
17. Avanzato, V. A. *et al.* Case Study: Prolonged infectious SARS-CoV-2 shedding from an asymptomatic immunocompromised cancer patient. *Cell* (2020) doi:10.1016/j.cell.2020.10.049.
18. Clark, S. A. *et al.* SARS-CoV-2 evolution in an immunocompromised host reveals shared neutralization escape mechanisms. *Cell* (2021) doi:10.1016/j.cell.2021.03.027.
19. Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Eurosurveillance* 25, 2000045 (2020).
20. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* 27, 2987–2993 (2011).
21. Alexandersen, S., Chamings, A. & Bhatta, T. R. SARS-CoV-2 genomic and subgenomic RNAs in diagnostic samples are not an indicator of active replication. *Nat Commun* 11, 6059 (2020).
22. Lucas, C. *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* 584, 463–469 (2020).
23. Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat Med* 26, 1033–1036 (2020).
24. Vogels, C. B. F. *et al.* Multiplex qPCR discriminates variants of concern to enhance global surveillance of SARS-CoV-2. *Plos Biol* 19, e3001236 (2021).
25. Grubaugh, N. D. *et al.* An amplicon-based sequencing framework for accurately measuring intrahost virus diversity using PrimalSeq and iVar. *Genome Biol* 20, 8 (2019).
26. Wei, J. *et al.* Genome-wide CRISPR Screens Reveal Host Factors Critical for SARS-CoV-2 Infection. *Cell* 184, 76-91.e13 (2021).
27. Lucas, C. *et al.* Delayed production of neutralizing antibodies correlates with fatal COVID-19. *Nat Med* 27, 1178–1186 (2021).
28. Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* 34, 4121–4123 (2018).

Figures

Fig. 1: Clinical course of the immunocompromised patient with persistent SARS-CoV-2 infection

Timeline of a) patient symptoms and hospitalizations, b) SARS-CoV-2 N1 RT-PCR Ct values and c) clinical laboratory parameters from time of laboratory diagnosis of SARS-CoV-2 infection (day 0) to end of the follow-up period (day 292) and d) computed tomography (CT) scans of chest at indicated days after time of initial diagnosis. The timing of remdesivir and

casirivimab-imdevimab are shown as grey and light blue shading, respectively. RT-PCR results that were positive but performed on assays that did not generate a Ct value are denoted by the open circle in panel b. The timing of filgrastim treatments are denoted by green diamonds in panel b. IsCRP values were converted to hsCRP values using a correction factor of 9.2. Ground-glass opacities marked by white arrows in panel d. Abbreviations: Real-time polymerase chain reaction (RT-PCR); Cycle threshold (Ct); Absolute neutrophil count (ANC); Absolute lymphocyte count (ALC); Hemoglobin (Hgb); high-sensitivity C-reactive protein (hsCRP).

Fig. 2: *de novo* emergence of remdesivir resistance mutation during and following treatment with the antiviral agent

a) Anti-SARS-CoV-2 Spike Protein IgG ELISA endpoint titers (1st panel), RT-PCR Ct values (2nd panel), subgenomic RNA (sgRNA, 3rd panel), and plaque forming units (PFU) on viral

culture (4th panel) during the course of illness which include the period when remdesivir (grey shading) and casirivimab/imdevimab (light blue shading) were administered. b) Allele frequencies of E802D in nsp12 as ascertained by whole genome sequencing. c) Crystal structure (PDB:7BV2) depicting nsp12 E802 (dark blue), its putative hydrogen bonds (yellow dashed lines) within the palm domain (yellow), and the residues (green) that interact with the replicating RNA molecule (orange). Remdesivir monophosphate (RMP) is depicted in light blue. d) Viral growth curves from icSARS-CoV-2 mNG WT, E802D (patient), and E802A (control) nsp12 mutants on a ORF7a depleted backbone. Significance was assessed by unpaired two sample t-tests (** $p < 0.01$) in experiments of three biological replicates. e) SARS-CoV-2 inhibition by RDV assessed at 48 h post-infection (0.01 multiplicity of infection) as determined by quantitative image analysis of percentage of cells expressing mNG. Vertical lines indicate IC_{50} for WT (black) and E802D (red). Shading represents 95% confidence intervals of the IC_{50} estimate. Figure is representative of four biological replicates.

Supplemental Material

Fig. S1: Phylogenetic analysis of SARS-CoV-2 genomes from patient samples and contemporaneous samples from Connecticut, USA.

Nextstrain maximum likelihood analysis of patient whole genome samples (blue) in comparison to 283 contemporaneous local samples (grey) obtained from GISAID (10/30/21; Supplemental Table 3). Samples are annotated by type (nasopharyngeal (NP), saliva (SL), or stool (ST)) and day from diagnosis. Asterisk denotes consensus genomes that contain nsp12 E802D. Divergence from root reference genome (Wuhan-Hu-1) by nucleic acid changes.

Fig. S2: Validation of a remdesivir resistance mutation

a) Structure depicting E802 in relationship to nsp12 (PDB:7BV2) b) Viral titers (PFU) assessed under 5 μM RDV treatment. Significance was assessed by an ANOVA (* p < 0.05) with 3 biological replicates. c) Cell viability experiment with RDV performed using CellTiter Glo. Significance of infected cells with SARS-CoV-2 WT (top panel) or E802D/A (bottom panel) was assessed by two-way ANOVA corrected for multiple comparisons (**** p < 0.0001) with 3 biological replicates.

Fig. S3: Global analysis of genomes with nsp12 E802D mutations.

Nextstrain maximum likelihood analysis of 131 genomes containing the E802D mutation in comparison to the global representative Auspice dataset. Genomes with the E802D mutation (denoted by circles) are distributed on various PANGO lineages (color coded according to the legend). The four patient genomes harboring the mutation are denoted by the arrow. Two genetically, geographically, and temporally linked networks are depicted in detail.

Fig S4: Identification of additional mutations temporally associated with remdesivir treatment

Longitudinal allele frequencies of A504V in nsp14 (a) and I115L (b) by whole genome sequencing of in relationship to RDV and casirivimab/imdevimab

SNP	Protein	AA	NP 36	NP 148	SL 148
G521T	leader protein	V86F			x
G569A	leader protein	E102K		x	
C1059T	nsp2	T85I	x	x	x
A4229G	nsp3	T504A		x	
C4230G	nsp3	T504R			x
C5178T	nsp3	T820I	x		x
G5180A	nsp3	D821N		x	
C9344T	nsp4	L264F			x
C9870T	nsp4	T439M		x	
G9929A	nsp4	D459N			x
A10323G	3C-like_proteinase	K90R		x	
G11083T	nsp6	L37F			x
A12519G	nsp8	D143G		x	
C12525T	nsp8	T145I			x
A12759G	nsp9	D25G		x	
A12759C	nsp9	D25A			x
C14408T	RdRp	P323L	x	x	x
C19550T	3'-to-5' exonuclease	A504V	x		
A23403G	surface glycoprotein	D614G	x	x	x
C23525T	surface glycoprotein	H655Y			x
C23997T	surface glycoprotein	P812L		x	
C25511T	ORF3a_protein	S40L			x
G25563T	ORF3a_protein	Q57H	x	x	x
C25714T	ORF3a_protein	L108F		x	
G25912T	ORF3a_protein	G174C		x	
A26162del	ORF3a_protein	N257fs	x	x	x
T26767C	membrane glycoprotein	I82T		x	
A28281G	nucleocapsid	D3G		x	
T28829C	nucleocapsid	S186P			x
C28472A	nucleocapsid	P67T	x	x	x
G29151A	nucleocapsid	R293K	x		
A29567G	ORF10_protein	I4V			x

Fig. S5: Variants identified in persistently infected host

Non-synonymous (NS) mutations identified at days 36 (NP) and 148 (NP and saliva) of infection. During early infection (day 36), whole genome sequencing identified 9 NS mutations from reference strain Wuhan-1 (NC_045512). Six of those persisted through day 148 (green background). Abbreviations: single nucleotide polymorphism (SNP); amino acid (AA); nasopharyngeal (NP); saliva (SL)

Fig. S6: Serial lymphocyte profiling reveals T-cell exhaustion

T-cell profiling of uninfected healthcare workers (HCW), patient's with moderate and severe SARS-CoV-2 infection. a) Global, CD4⁺, and CD8⁺ T-cells. CD4⁺ (b) and CD8⁺ (c) subsets demonstrates a phenotype consistent with T-cell exhaustion. For all boxplots, the centre is drawn through the median of the measurement, while the lower and upper bounds of the box correspond to the first and third percentile. Whiskers beyond these points denote 1.5×the interquartile range. P values were determined by two-sided, Wilcoxon rank-sum test. ****p < .0001 ***p < .001 **p < .01. *p < .05. Population definitions: naïve CD4/CD8 subset: CD4/CD8+CD45RA+CD127+CCR7+PD-1-; activated CD4/CD8: CD4/CD8+CD38+HLA-DR+;effector memory CD4/CD8:CD4/CD8CD45RA-CD127+CCR7-; exhaustion CD4/CD8: CD4/CD8CD45RA-Tim-3+PD-1+; effTreg: CD4+CD45RA-CD127-CD25hiHLA-DR+. Gating strategy depicted in Fig S8.

c

Source	Day	D80Y	L141fs	A348S	D614G	V635G	H655Y	V656I	P812L	G1124V
NP	36	<5	<5	<5	100	<5	<5	<5	<5	<5
NP	148	<5	<5	<5	99.9	<5	<5	<5	96.9	<5
SL	148	<5	<5	<5	99.9	<5	97.9	<5	<5	<5
NP	149	<5	<5	<5	99.9	<5	99.5	<5	<5	<5
SL	149	<5	<5	<5	99.9	5.6	79.6	<5	20.7	<5
SL	152	<5	<5	8.8	99.8	<5	74.1	<5	<5	<5
NP	155	<5	<5	<5	99.8	<5	36.1	7.9	69	<5
NP	156	<5	<5	<5	99.8	<5	42.6	<5	54.2	<5
SL	158	<5	<5	<5	99.9	6.2	61.1	<5	20.9	<5
NP	160	<5	<5	<5	99.8	<5	58.9	<5	42.3	<5
NP	162	62.9	<5	<5	99.8	<5	35.4	24.3	<5	65.5
NP	163	13.9	<5	<5	99.8	<5	86.6	59.2	<5	15.8
SL	163	<5	<5	<5	100	5.3	91.5	11.3	10.4	<5
NP	164	17.9	12.6	<5	99.9	<5	<5	<5	75.7	17.3
NP	166	<5	<5	<5	100	<5	13.1	13.2	86.3	8.6
SL	167	<5	<5	<5	100	5.4	99.3	7.4	<5	<5
SL	170	<5	5.6	75	100	5.4	21.6	<5	<5	<5
NP	177	<5	<5	86.5	100	<5	<5	<5	17.9	<5

Fig. S7: Evolution of Spike protein in patient samples.

a) H655Y and b) P812L allele frequency over time. c) Allele frequencies of non-synonymous Spike protein mutations identified on more than one occasion by whole genome sequencing. Allele frequencies >50% are denoted by green background. d) Crystal structure of RBD (PDB:6XDG) depicting residue A348 in relationship to the imdevimab (heavy chain: yellow; light chain: red) and casirivimab (heavy chain: green; light chain: blue) binding sites. Abbreviations: nasopharyngeal (NP); saliva (SL); receptor binding domain (RBD).

Figure S8: Gating strategy for flow cytometry

Gating strategies are shown for the key T cell populations described in Figure S6. The T cell surface staining gating strategy to identify CD8 & CD4 T cells, naïve T cells, TCR-activated T cells, exhaustion T cells, effector memory T cells (Tem), and effector regulatory T cells (eff Treg) are depicted.

Days after diagnosis	Nasopharyngeal specimen				Saliva specimen			Stool specimen			Whole blood specimen		
	N1 RT-PCR Ct	N2 RT-PCR Ct	RP RT-PCR Ct	Viral Culture PFU/ml	N1 RT-PCR Ct	N2 RT-PCR Ct	RP RT-PCR Ct	N1 RT-PCR Ct	N2 RT-PCR Ct	RP RT-PCR Ct	N1 RT-PCR Ct	N2 RT-PCR Ct	RP RT-PCR Ct
36	21.8	21.5	-										
148	26.1	25.4	32.9		22.9	22.2	22.3						
149	19.2	18.0	24.0	41.7	29.4	29.8	21.8	30.0	31.6	36.6	ND	ND	36.0
152	28.3	27.7	27.8	12.5	28.9	29.0	22.1	33.2	32.9	34.4			
155	26.1	25.6	27.6										
156	26.2	26.1	28.2	1741.7	28.0	27.6	23.3	32.8	ND	33.7			
157											ND	ND	26.9
158	24.2	23.7	34.5	16666.7	30.0	30.8	23.0	32.4	ND	33.1	ND	ND	27.9
159					22.1	21.2	23.5	31.7	ND	36.4	ND	ND	28.2
160	23.2	22.7	27.0								ND	ND	29.2
162	23.1	22.7	28.7								ND	ND	27.1
163	22.5	22.1	28.3	533.3	27.1	27.2	22.1	32.7	33.9	34.5	ND	ND	30.6
164	27.3	27.1	28.5	0.0									
166	35.5	36.1	29.3	0.0									
167	36.2	36.9	30.7		28.0	27.9	23.1	34.1	41.1	31.5	ND	ND	30.1
168	39.5	38.7	28.2	0.0									
170	36.7	40.2	26.9	0.0	31.4	32.4	22.4	34.1	37.7	33.0	ND	ND	30.0
171	39.4	41.0	30.9										
176	39.7	ND	28.0	0.0	33.9	36.9	22.3	35.0	ND	32.5	ND	ND	29.8
177	33.3	33.7	26.2										
189											ND	ND	28.0
202	ND	ND	33.7								ND	ND	30.3
210	39.8	44.3	29.1	0.0									
217	35.9	37.1	24.4	0.0							ND	ND	28.1
245								34.5	ND	31.9	ND	ND	29.1

Supplementary Table 1: Serial patient samples from various tissues and secretions

Catalogue of patient samples collected for research during the course of illness. N1, N2, and RP cycle threshold values are reported for each sample. WGS was performed on the bolded samples (accession numbers in table S2). Days of RDV treatment are denoted by grey shading and casirivimab/imdevimab was administered on day 163 (blue shading). Note that NP Ct values depicted in Figures 1,2 were obtained from clinical platforms and are not shown here.

Abbreviations: ND=Not Detected

Sample Name	SRA Accession	GenBank Accession
INP-0411-NP-148	SRR16576689	OL307721
INP-0411-NP-149	SRR16576688	OL307722
INP-0411-NP-152*	SRR16576677	
INP-0411-NP-155	SRR16576668	OL307723
INP-0411-NP-156	SRR16576667	OL307724
INP-0411-NP-158	SRR16576666	OL307725
INP-0411-NP-160	SRR16576665	OL307726
INP-0411-NP-162	SRR16576664	OL307727
INP-0411-NP-163	SRR16576663	OL307728
INP-0411-NP-164	SRR16576662	OL307729
INP-0411-NP-166	SRR16576687	OL307720
INP-0411-NP-177	SRR16576686	OL307730
INP-0411-NP-36	SRR16576685	OL307731
INP-0411-SL-148	SRR16576684	OL307732
INP-0411-SL-149	SRR16576683	OL307733
INP-0411-SL-152	SRR16576682	OL307734
INP-0411-SL-156	SRR16576681	OL307735
INP-0411-SL-158	SRR16576680	OL307736
INP-0411-SL-159	SRR16576679	OL307737
INP-0411-SL-163	SRR16576678	OL307738
INP-0411-SL-167	SRR16576676	OL307739
INP-0411-SL-170	SRR16576675	OL307740
INP-0411-ST-149	SRR16576674	OL307741
INP-0411-ST-152	SRR16576673	OL307742
INP-0411-ST-156	SRR16576672	OL307743
INP-0411-ST-158	SRR16576671	OL307744
INP-0411-ST-159	SRR16576670	OL307745
INP-0411-ST-163	SRR16576669	OL307746

*depth was sufficient to calculate allele frequency at position E802
in nsp12 but coverage was insufficient to generate consensus genome

Supplemental Table 2: Accession numbers for submitted samples

We gratefully acknowledge the following Authors from the Originating laboratories responsible for obtaining the specimens, as well as the Submitting laboratories where the genome data were generated and shared via GISAID, on which this research is based.

All Submitters of data may be contacted directly via www.gisaid.org

Authors are sorted alphabetically.

Accession ID	Originating Laboratory	Submitting Laboratory	Authors	
EPI_ISL_3354875	CT-Dr. Katherine A. Kelley State Public Health Lab	Centers for Disease Control and Prevention Division of Viral Diseases, Pathogen Discovery	Alex Burgin; Ben Rambo-Martin; Clinton Paden; Dakota Howard; Dave Wentworth; Dhwani Batra; Jasmine Padilla; Justin Lee; Krista Queen; Kristen Knipe; Kristine Lacey; Mark Burroughs; Matthew Schmerer; Meghan Bentz; Mili Sheth; Peter Cook; Sam Shepard; Sarah Nobles; Suxiang Tong; Vivien Dugan; Yvette Unoarumhi	
EPI_ISL_454642, EPI_ISL_454645	CT-Dr. Katherine A. Kelley State Public Health Lab	Pathogen Discovery, Respiratory Viruses Branch, Division of Viral Diseases, Centers for Disease Control and Prevention	Anna Uehara; Bettina Bankamp; Clinton R. Paden; Haibin Wang; Jing Zhang; Krista Queen; Suxiang Tong; Yan Li; Ying Tao; Zachary Weiner	
EPI_ISL_729735, EPI_ISL_729736, EPI_ISL_729741, EPI_ISL_729743, EPI_ISL_729744, EPI_ISL_729745, EPI_ISL_729746, EPI_ISL_729747, EPI_ISL_729748, EPI_ISL_729749, EPI_ISL_729750, EPI_ISL_729752, EPI_ISL_729753, EPI_ISL_729754, EPI_ISL_729755, EPI_ISL_729756, EPI_ISL_729758, EPI_ISL_729759, EPI_ISL_729760, EPI_ISL_729765, EPI_ISL_729772, EPI_ISL_729773, EPI_ISL_729778, EPI_ISL_729779, EPI_ISL_729780, EPI_ISL_729786, EPI_ISL_729789, EPI_ISL_729792	see above	Connecticut Department of Health	Grubaugh Lab - Yale School of Public Health Adam Moore; Akiko Iwasaki; Albert Ko; Alice Lu; Allison Nelson; Anderson Brito; Anne Wyllie; Annie Watkins; Arnaud Casanovas; Catherine Muenker; Chaney Kalinich; Chantal Vogels; Charles Dela Cruz; Isabel Ott; Joseph Fauver; Maria Tokuyama; Mary Petrone; Nathan Grubaugh; Patrick Wong; Peiwen Lu; Richard Martinello; Saad Omer; Shelli Farhadian; Tara Alpert	
EPI_ISL_527738, EPI_ISL_527761, EPI_ISL_527762, EPI_ISL_527763, EPI_ISL_527764, EPI_ISL_527765, EPI_ISL_527766, EPI_ISL_527767, EPI_ISL_527768, EPI_ISL_527769, EPI_ISL_527770, EPI_ISL_527771, EPI_ISL_527772, EPI_ISL_527773, EPI_ISL_527774, EPI_ISL_527775, EPI_ISL_527776, EPI_ISL_527777, EPI_ISL_527778, EPI_ISL_527779, EPI_ISL_527780, EPI_ISL_527781, EPI_ISL_527782, EPI_ISL_527783, EPI_ISL_527784, EPI_ISL_527785	see above	Connecticut State Department of Public Health	Grubaugh Lab - Yale School of Public Health Albert Ko; Anderson Brito; Anthony Muyombwe; Chantal Vogels; Jafar Razeq; Joseph Fauver; Mary Petrone; Nathan Grubaugh; Tara Alpert	
EPI_ISL_738284, EPI_ISL_738285, EPI_ISL_738289, EPI_ISL_738298, EPI_ISL_738299, EPI_ISL_738300, EPI_ISL_738301, EPI_ISL_738302, EPI_ISL_738307, EPI_ISL_738308, EPI_ISL_738309, EPI_ISL_738310, EPI_ISL_738311, EPI_ISL_738312, EPI_ISL_1017585, EPI_ISL_1017589, EPI_ISL_1017592, EPI_ISL_1017593, EPI_ISL_1017602, EPI_ISL_1017604, EPI_ISL_1017627, EPI_ISL_1017628, EPI_ISL_1017636, EPI_ISL_1017637, EPI_ISL_1017639, EPI_ISL_1017643, EPI_ISL_1017652, EPI_ISL_1017675, EPI_ISL_1067646, EPI_ISL_1067647, EPI_ISL_1067686, EPI_ISL_1067687, EPI_ISL_1080455, EPI_ISL_1080462, EPI_ISL_1091839, EPI_ISL_1091869, EPI_ISL_1239847	see above	Connecticut Veterans' Affairs Hospital	Grubaugh Lab - Yale School of Public Health Chantal Vogels; Danielle Plank; Ellen Foxman; Isabel Ott; Joseph Fauver; Mary Petrone; Nathan Grubaugh; Shaali Gupta; Tara Alpert	
EPI_ISL_802136	MSHS Clinical Microbiology Laboratories	MSHS Pathogen Surveillance Program	Adolfo Garcia-Sastre; Adriana van de Guchte; Ajay Obia; Alberto Paniz-Mondolfi; Ana S. Gonzalez-Reiche; Andrew Kasarskis; Angela Amoako; Ashley S. Salimangon; Betsaida Salom Melo; Brenny Albuquerque; Brianne Ciferri; Charles Gleason; Deena R. Altman; Denise Jurczynszak; Elena Hirsch; Emilia Mia Sordillo; Emily Ferrer; Gintaras Deikus; Giulio Kleiner; Gopi Patel; Hala Alshammary; Harm van Bakel; Irina Oussenko; Jayeeta Dutta; Juan Soto; Katherine Beach; Kathryn Twyman; Kayla Russo; Komal Srivastava; Levy Sominsky; Mahmoud Awawda; Marta Luksha; Matthew M. Hernandez; Melissa Gitman; Michael D. Nowak; Mitchell J. Sullivan; Nancy Francoeur; Rachel Chermet; Robert Sebra; Sarah Schaefer; Shellee Fabre; Shwetha Hara Sridhar; Viviana Simon; Ying-Chih Wang; Zenab Khan	
EPI_ISL_430401	NYU Langone Health	Departments of Pathology and Medicine, New York University School of Medicine	Adriana Heguy; Amy Rapkiewicz; Andre M. Ribeiro-dos-Santos; Andrew Lytle; Antonio Serrano; Brendan Belovarac; Christian Marier; Dacia Dimartino; Emily Guzman; Gael Westby; George Jour; Guomiao Shen; Iman Osman; Jared Pinnell; John Cadley; John Chen; Lawrence H. Lin; Ludovic Boyard; Margaret Black; Maria Agüero-Rosenfeld; Marie Samanovic-Golden; Mark J. Mulligan; Matija Snuderl; Matthew T. Maurano; Megan Hogan; Nick Vulpescu; Paolo Cotzia; Paul Zapple; Peter Meyn; Raquel Ordonez Ciriza; Raven Luther; Sitharam Ramaswami; Tatyana Gindin; Theodore Vougiouklakis; Vanessa Raabe; Xiaojun Feng; Yutong Zhang	
EPI_ISL_937036, EPI_ISL_937037	Quest Diagnostics	Quest Diagnostics	Anderson, B.; D.F.; Gerasimova, A.; Hua, M.; K.E.; Kagan; Lacbawan, F.; Liu Y.; Livingston; Owen, R.; R.M.; Rosenthal; S.H.; Shalhout	
EPI_ISL_429640, EPI_ISL_429641, EPI_ISL_429642, EPI_ISL_429645, EPI_ISL_429646, EPI_ISL_430907, EPI_ISL_430908, EPI_ISL_430917, EPI_ISL_430919, EPI_ISL_430921, EPI_ISL_430922, EPI_ISL_430923, EPI_ISL_430925, EPI_ISL_430926, EPI_ISL_430927, EPI_ISL_430960, EPI_ISL_430961, EPI_ISL_430962, EPI_ISL_430963, EPI_ISL_430971, EPI_ISL_430972, EPI_ISL_430973, EPI_ISL_437865, EPI_ISL_437866, EPI_ISL_437867, EPI_ISL_437868, EPI_ISL_437869, EPI_ISL_437870, EPI_ISL_437871, EPI_ISL_437872, EPI_ISL_461408, EPI_ISL_461409, EPI_ISL_461411, EPI_ISL_461412, EPI_ISL_461414, EPI_ISL_461416, EPI_ISL_461417, EPI_ISL_461420, EPI_ISL_461435, EPI_ISL_461441, EPI_ISL_570070, EPI_ISL_570076, EPI_ISL_570092, EPI_ISL_570093, EPI_ISL_570094, EPI_ISL_570095, EPI_ISL_570104, EPI_ISL_570107, EPI_ISL_570109, EPI_ISL_570130, EPI_ISL_570131, EPI_ISL_570132, EPI_ISL_570133, EPI_ISL_570135, EPI_ISL_570137, EPI_ISL_570141, EPI_ISL_570145, EPI_ISL_570146, EPI_ISL_570148, EPI_ISL_570166, EPI_ISL_570178, EPI_ISL_570179, EPI_ISL_570180, EPI_ISL_570182, EPI_ISL_570184, EPI_ISL_570185	see above	UW Virology Lab	Alexander Greninger; Amin Addetta; Hong Xie; Keith Jerome; Keith R Jerome; Lasata Shrestha; Meei-Li Huang; Pavitra Roychoudhury; Truong Nguyen; Victoria M Rachleff	
EPI_ISL_484885	University of Wisconsin-Madison AIDS Vaccine Research Laboratories	University of Wisconsin-Madison AIDS Vaccine Research Laboratories	Gage Moreno; Katarina Braun; et al. AIDS Vaccine Research Laboratories	
EPI_ISL_1168354, EPI_ISL_1168356, EPI_ISL_1168357, EPI_ISL_1168363, EPI_ISL_1168368, EPI_ISL_1168377, EPI_ISL_1168384, EPI_ISL_1168391, EPI_ISL_1168396, EPI_ISL_1168406, EPI_ISL_1168426, EPI_ISL_1168427, EPI_ISL_1168428, EPI_ISL_1168431, EPI_ISL_1168435, EPI_ISL_1168437	see above	VA Connecticut Healthcare System	Grubaugh Lab - Yale School of Public Health Chantal Vogels; Danielle Plank; Isabel Ott; Joseph Fauver; Mary Petrone; Nathan Grubaugh; Shaali Gupta; Tara Alpert	
EPI_ISL_428384, EPI_ISL_428385, EPI_ISL_428386, EPI_ISL_428389, EPI_ISL_428391, EPI_ISL_428392, EPI_ISL_428393, EPI_ISL_428394, EPI_ISL_428395, EPI_ISL_428396, EPI_ISL_428397, EPI_ISL_428398, EPI_ISL_428399, EPI_ISL_428401, EPI_ISL_428402, EPI_ISL_428403, EPI_ISL_428404, EPI_ISL_428405, EPI_ISL_428746, EPI_ISL_428747, EPI_ISL_428748, EPI_ISL_428749, EPI_ISL_428750, EPI_ISL_428751, EPI_ISL_428752, EPI_ISL_431081, EPI_ISL_431082, EPI_ISL_431083, EPI_ISL_431084, EPI_ISL_431085, EPI_ISL_431086, EPI_ISL_431087, EPI_ISL_431088, EPI_ISL_431089, EPI_ISL_431090, EPI_ISL_431091, EPI_ISL_431092, EPI_ISL_431093, EPI_ISL_431094, EPI_ISL_431095, EPI_ISL_431096, EPI_ISL_435702, EPI_ISL_435703, EPI_ISL_435704, EPI_ISL_435705, EPI_ISL_435706, EPI_ISL_435707, EPI_ISL_435708, EPI_ISL_452143, EPI_ISL_452144, EPI_ISL_452145, EPI_ISL_452146, EPI_ISL_452147, EPI_ISL_594133, EPI_ISL_730052	see above	Yale COVID-19 Biorepository	Grubaugh Lab - Yale School of Public Health Adam Moore; Akiko Iwasaki; Albert Ko; Alice Lu; Allison Nelson; Anderson Brito; Anne Wyllie; Annie Watkins; Arnaud Casanovas; Catherine Muenker; Chaney Kalinich; Chantal Vogels; Charles Dela Cruz; Cole Jensen; Isabel Ott; Joseph Fauver; Maria Tokuyama; Mary Petrone; Nathan Grubaugh; Patrick Wong; Peiwen Lu; Richard Martinello; Saad Omer; Shelli Farhadian; Tara Alpert	
EPI_ISL_463167, EPI_ISL_463168, EPI_ISL_463169, EPI_ISL_463170, EPI_ISL_463171, EPI_ISL_463172, EPI_ISL_463173, EPI_ISL_463174, EPI_ISL_469171, EPI_ISL_469172, EPI_ISL_469173, EPI_ISL_469174, EPI_ISL_469175, EPI_ISL_469176, EPI_ISL_469177, EPI_ISL_469178, EPI_ISL_469179, EPI_ISL_469180, EPI_ISL_469181, EPI_ISL_469182, EPI_ISL_469183, EPI_ISL_469184, EPI_ISL_469185, EPI_ISL_469186, EPI_ISL_469187, EPI_ISL_469188, EPI_ISL_469189, EPI_ISL_469190, EPI_ISL_469191, EPI_ISL_469192, EPI_ISL_469193, EPI_ISL_469194, EPI_ISL_469195, EPI_ISL_469196, EPI_ISL_469197, EPI_ISL_469198, EPI_ISL_469199, EPI_ISL_469200, EPI_ISL_469201, EPI_ISL_469202, EPI_ISL_469203, EPI_ISL_469204, EPI_ISL_469205, EPI_ISL_469206, EPI_ISL_469207, EPI_ISL_469208, EPI_ISL_476547, EPI_ISL_476548, EPI_ISL_476549, EPI_ISL_476550, EPI_ISL_476551, EPI_ISL_476552, EPI_ISL_476553, EPI_ISL_476554, EPI_ISL_476555, EPI_ISL_476556, EPI_ISL_476557, EPI_ISL_730062, EPI_ISL_730063, EPI_ISL_730064, EPI_ISL_730065, EPI_ISL_730066, EPI_ISL_730067, EPI_ISL_730068, EPI_ISL_730069, EPI_ISL_730070, EPI_ISL_730071, EPI_ISL_730072, EPI_ISL_730073, EPI_ISL_730074, EPI_ISL_730075, EPI_ISL_730076, EPI_ISL_730077, EPI_ISL_730078, EPI_ISL_730079, EPI_ISL_730080, EPI_ISL_730081, EPI_ISL_730082, EPI_ISL_730083, EPI_ISL_730084	see above	Yale Clinical Virology Laboratory	Grubaugh Lab - Yale School of Public Health Adam Moore; Akiko Iwasaki; Albert Ko; Alice Lu; Allison Nelson; Anderson Brito; Anne Wyllie; Annie Watkins; Arnaud Casanovas; Catherine Muenker; Chaney Kalinich; Chantal Vogels; Charles Dela Cruz; Cole Jensen; Isabel Ott; Joseph Fauver; Maria Tokuyama; Mary Petrone; Nathan Grubaugh; Patrick Wong; Peiwen Lu; Richard Martinello; Saad Omer; Shelli Farhadian; Tara Alpert	
EPI_ISL_738290, EPI_ISL_738291, EPI_ISL_738292, EPI_ISL_738293, EPI_ISL_738294, EPI_ISL_738295, EPI_ISL_738296, EPI_ISL_738297	see above	Yale Pathology Lab	Grubaugh Lab - Yale School of Public Health Chantal Vogels; Danielle Plank; Ellen Foxman; Isabel Ott; Joseph Fauver; Mary Petrone; Nathan Grubaugh; Shaali Gupta; Tara Alpert	

Supplemental Table 3: GISAID acknowledgments for local genomes used to generate Figure S1

We gratefully acknowledge the following Authors from the Originating laboratories responsible for obtaining the specimens, as well as the Submitting laboratories where the genome data were generated and shared via GISAID, on which this research is based.

All Submitters of data may be contacted directly via www.gisaid.org

Authors are sorted alphabetically.

Accession ID	Originating Laboratory	Submitting Laboratory	Authors
EPI_ISL_4361721, EPI_ISL_4362029	Aegis Sciences Corporation	Centers for Disease Control and Prevention Division of Viral Diseases, Pathogen Discovery	Alec Vest; Benjamin Rambo-Martin; Christopher Gulvick; Clinton Paden; Cyndi Clark; Dakota Howard; Dhvani Batra; Dillon Nall; Duncan MacCannell; Erisa Sula; Ethan Sanders; Holly Houdeshell; Jason Caravas; Kristine Lacek; Matthew Hardison; Matthew Schmerer; Ola Kvalvaag; Patrick Campbell; Peter Cook; Rob Case; Scott Sammons; Shatavia Morrison; Shaun Westlund; Tymeckia Kendall; Victoria Caban Figueroa; Vikramsinha Ghorpade; Yvette Unoarumhi
EPI_ISL_2102156	CHUV	Laboratory of genomics and metagenomics	Claire Bertelli; Damien Jacot; Gilbert Greub; Sébastien Aebys; Trestan Pillonel
EPI_ISL_3473275	CNR Virus des Infections Respiratoires - France SUD	CNR Virus des Infections Respiratoires - France SUD	Antonin Bal; Bruno Lina; Gregory Destras; Gwendolynne Burfin; Hadrien Regue; Laurence Josset; Martine Valette; Quentin Semanas
EPI_ISL_2170967	CS DE NIPOA	Instituto Butantan / Mendelics	Antonio Jorge Martins; Bianca Cechetto Carlos, Mendelics; Bibiana Santos; Claudia Renata dos Santos Barros; Clintia Bittar; David Schlesinger; Hemocentro Ribeirão Preto; Simone Kashima; Debora Botequilo Moretti; Elaine Cristina Marqueze; Elaine Vieira dos Santos; Elisangela Chicaroni Mattos; Erika Freitas; Evandra Strazza Rodrigues; Felipe Allan da Silva da Costa; Flavia Aburjaile; Fábio Sossai Possebon; Guilherme Campos; Guilherme Targino Valente; Heidge Fukumasu; USP-Botucatu; Rejane Maria Tommasini Grotto; Helena Lage Ferreira; Instituto Butantan; Dimas Tadeu Covas; Jardelina de Souza Todao Bernardino; Jayme A. Souza-Neto; Jessika Cristina Chagas Lesbon; Jorge A. Petroll Marchesi; José Salvatore Leister Patané; João Paulo Kitajima; João Pessoa Araújo Jr.; Leila Sabrina Ullmann; Loyze Paola Oliveira de Lima; Luiz Aurelio de Campos Crispin, Centro de Genômica Funcional da ESALQ; Luiz Lehmann Coutinho; Luiz Carlos Junior de Alcantara; Livia Sacchetto; Maisa C. Pereira Parra; Maria Carolina Elias; Marta Giovanetti; Marília Moraes; Mauricio Lacerda Nogueira, Prefeitura de Sao Paulo; Melissa Palmieri; Patricia Akemi Assato; Paula Rahal; Paulo Inacio da Costa; Rafael dos Santos Bezerra; Raquel de Lello Rocha Campos Cassano, NGS Soluções Genômicas; Pilar Drummond Sampaio Corrêa Mariani, FZEA-USP Pirassununga; Mirele Daiana Poletti; Raul Machado Neto; Ricardo Augusto Brassaloti; Ricardo Haddad; Rodrigo Tocantins Calado, FAMERP-SJRP; Cecília Artico Banho; Sandra Coccuzzo Sampaio; Svetoslav Nanev Slavov; Vagner Fonseca; Vincent Louis Viala
EPI_ISL_1138787	Centro Hospitalar do Baixo Vouga (CHBV)	Institute of Biomedicine (IBIMED), Universidade de Aveiro	Gabriela Moura; Miguel Pinheiro and Manuel Santos; Patricia Arinto; Sofia Marques
EPI_ISL_862176, EPI_ISL_862177	Charité Universitätsmedizin Berlin, Institut für Virologie/Labor Berlin	Charité Universitätsmedizin Berlin, Institut für Virologie	Barbara Mühlemann; Christian Drosten; Julia Schneider; Julia Tesch; Jörn Beheim-Schwarzbach; Talitha Veith; Terry Jones; Tobias Blecker; Victor M Corman
EPI_ISL_1156362	Chemische und Veterinäruntersuchungsamt MünsterLand-Emscher-Lippe	Robert Koch Institute	
EPI_ISL_3339857, EPI_ISL_3463152, EPI_ISL_3463851, EPI_ISL_3707915, EPI_ISL_3708049, EPI_ISL_3770479, EPI_ISL_3770513, EPI_ISL_3806078, EPI_ISL_3904028, EPI_ISL_3904237, EPI_ISL_3904270, EPI_ISL_3917497, EPI_ISL_4007212, EPI_ISL_4008134, EPI_ISL_4355696	see above	Statens Serum Institut Bioinformatics and Microbial Genomics	Danish Covid-19 Genome Consortium
EPI_ISL_1654424	Department of Public Health Microbiology Ljubljana, National Laboratory for Health, Environment and Food	Department of Public Health Microbiology Ljubljana, National Laboratory for Health, Environment and Food	José Gonçalves; Katarina Proscenc; Martin Bosilj; Metka Paragi; Natasa Berginc; Tom Koritnik
EPI_ISL_759609, EPI_ISL_793664	Department of Virus and Microbiological Special Diagnostics, Statens Serum Institut, Copenhagen, Denmark	Albertsen Lab, Department of Chemistry and Bioscience, Aalborg University, Denmark	Danish Covid-19 Genome Consortium
EPI_ISL_666605	Dept. of Microbiology and Infection Control, Akershus University Hospital HF	Dept. of Microbiology and Infection Control, Akershus University Hospital HF	Alexander Hesselberg Løvestad; Hege Vangstein Aamot; Nina Handal; Ole Herman Ambur; Silje Bakken Jørgensen
EPI_ISL_1457742, EPI_ISL_2672226, EPI_ISL_3257488	Dutch COVID-19 response team	National Institute for Public Health and the Environment (RIVM)	Adam Meijer; AnneMarie van den Brandt; Annelies Kroneman; Bas van der Veer; Chantal Reusken; Dennis Schmitz; Dirk Eggink; Eunice Then; Florian Zwagemaker; Harry Vennema; James Groot; Jeroen Cremer; Jolienke Hardeman; Karim Hajji; Kim Freriks; Linda van de Nes; Lisa Wijsman; Lynn Aarts; Melissa van Tuil; Robert Kohl; Rianne Jaarsma; Sanne Bos; Sharon van den Brink; Sjoerd Kulling; on behalf of the national COVID-19 response team
EPI_ISL_4636562	Genetica Molecular and Subdepartamento de Virologia ISP Chile	Instituto de Salud Publica de Chile	Andres Castillo; Barbara Parra; Constanza Campano; Gisselle Barra; Javier Tognarelli; Jorge Fernandez; Karen Orostica; Loredana Arata; Patricia Bustos; Rodrigo Fasce; Soledad Ulloa
EPI_ISL_2301853	Greek Genome Center, Biomedical Research Foundation of the Academy of Athens (BRFAA)	Greek Genome Center, Biomedical Research Foundation of the Academy of Athens (BRFAA)	Dimitrios Thanos; Emmanouil Athanasiadis; Giannis Vatsellas; Katerina Zoi; Theodoros Loupis
EPI_ISL_2187692	HLAGYN - Laboratorio de Imunologia de Transplantes de Goias	HLAGYN - Laboratorio de Imunologia de Transplantes de Goias	Alessandro Leonardo Alvares Magalhães; Daniel Ferreira de Sousa; Erika Lopes Rocha Batista; Fernando Antonio Vinhal dos Santos; Frederico Rodrigues Vinhal; Lucas Carlos Gomes Pereira; Sabrina Sara Moreira Duarte
EPI_ISL_1581735, EPI_ISL_1907121, EPI_ISL_4346271, EPI_ISL_4962483	Helix/Illumina	Centers for Disease Control and Prevention Division of Viral Diseases, Pathogen Discovery	Adrian Paskey; Alexandre Bolze; Ary Ascencio; Benjamin Rambo-Martin; Brad Sickler; Charlotte Rivera-Garcia; Christine Tran; Christopher Gulvick; Christine Tran; Clinton Paden; Clinton R. Paden; Dakota Howard; Darlene Wagner; David Becker; Dhvani Batra; Duncan MacCannell; Efrén Sandoval; Eileen De Feo; Eileen de Feo; Elizabeth Cirulli; Eric Allen; Geraint Levan; James Lu; Jan Antico; Jason Caravas; Jason Nguyen; Jimmy Ramirez; Jingtao Liu; Kara Moser; Kelly Barrett; Kelly Schiabor Barrett; Kim Gietzen; Kristine Lacek; Magnus Isaksson; Marc Laurent; Matthew Schmerer; Matthew Tolentino; Nicole L. Washington; Nicole Washington; Peter Cook; Peter W. Cook; Phil Febbo; Ryan Cho; Scott Sammons; Shannon Wickline; Shatavia Morrison; Sherry Wang; Simon White; Tyler Cassens; William Lee; Yvette Unoarumhi
EPI_ISL_5070788	Houston Methodist Hospital	Houston Methodist Hospital	Ilya J. Finkelstein; James J. Davis; Jessica Cambric; Jimmy Gollihar; Kristina Reppond; Layne Pruitt; Madison N. Shyer; Marcus Nguyen; Matthew Ojeda Saavedra; Paul A. Christensen; Prasanti Yerramilli; Randall J. Olsen; Robert Olson; Ryan Gadd; S. Wesley Long; Sishir Subedi; and James M. Musser
EPI_ISL_2152548	Hôpital Cochin	Department of Virology, Henri Mondor University Hospital, Assistance Publique Hôpitaux de Paris, Université Paris-Est Créteil, INSERM U955	Alexandre Soulier; Christophe Rodriguez; Elisabeth Trawinski; Guillaume Gricourt; Jean-Michel Pawlotsky; Melissa N'Debi; Slim Fourati; Vanessa Demontant
EPI_ISL_1819243	I.R.C.C.S. "S. De Bellis" - Ente Ospedaliero	Istituto Zooprofilattico Sperimentale della Puglia e della Basilicata	Bianco A.; Capozzi L.; Del Sambro L.; Lippolis A.; Notarricola M.; Parisi A.; Simone D.
EPI_ISL_1678325, EPI_ISL_1678326	ID Bureau of Laboratories	Centers for Disease Control and Prevention Division of Viral Diseases, Pathogen Discovery	Alison Laufer Halpin; Ben L. Rambo-Martin; Clinton R. Paden; Dakota Howard; Darlene Wagner; Dave Wentworth; Dhvani Batra; Jasmine Padilla; Justin Lee; Katie Dillon; Krista Queen; Kristen Knipe; Kristine Lacek; Mark Burroughs; Matthew Schmerer; Mili Sheth; Peter Cook; Sam Shepard; Sarah Nobles; Shoshona Le; Suixiang Tong; Vivien Dugan; Yvette Unoarumhi
EPI_ISL_3275089, EPI_ISL_3275095	Idaho Bureau of Laboratories	Center for Global Health, University of New Mexico Health Sciences Center	Christopher Ball; Darrell Dinwiddie; Daryl Domman; Kurt Schwalm; Matthew Burns; Robert Voermans; Valerie Morley
EPI_ISL_1219920, EPI_ISL_3543382, EPI_ISL_3756841, EPI_ISL_3922840, EPI_ISL_3923187	Idaho Bureau of Laboratories	Idaho Bureau of Laboratories	"R. Beukelman; Aimee Ceniseros; Christian Loera; Christopher Ball"; Matthew Charles Burns; Robert L. Voermans
EPI_ISL_1964532, EPI_ISL_3039233, EPI_ISL_3039248	Institute of Microbiology and Immunology, Faculty of Medicine, University of Ljubljana	Institute of Microbiology and Immunology, Faculty of Medicine, University of Ljubljana	Alen Suljić; Andraž Celar; Dominika Šturm; Doroteja Vaj; Mario Poljak; Matic Brvar; Miša Korva; Patricija Pozvek; Samo Zakotnik; Tatjana Avšič – Županc; Tomaž Mark Zorec; Špela Pleh
EPI_ISL_4862793	Integrated Covid Hub North East	Wellcome Sanger Institute for the COVID-19 Genomics UK (COG-UK) Consortium	Cordelia Langford; David K. Jackson; Dominic Kwiatkowski; Ewan Harrison; Ian Johnston; Integrated Covid Hub North East and Alex Alderton; Jeffrey Barrett; John Sillitoe on behalf of the Wellcome Sanger Institute COVID-19 Surveillance Team; Roberto Amato; Sonia Goncalves
EPI_ISL_1689555	Lab voor klinische biologie	Lab voor klinische biologie	Bruno Verhasselt; Hannelore Hamerlinck; Marija Janevica

EPI_ISL_1140857, EPI_ISL_1141807	Labor Dr. Wisplinghoff - Berlin	Robert Koch Institute	
EPI_ISL_1643477	Labor LÄbeck bzw. LaborÄrztliche Gemeinschaftspraxis LÄbeck	Robert Koch Institute	
EPI_ISL_1218124, EPI_ISL_1282942	Labor ZOTZKLIMAS; MVZ Düsseldorf-Centrum	Robert Koch Institute	
EPI_ISL_5617461	Laboratorio Nacional de Salud, Ministerio de Salud Publica y Asistencia Social	Genomics and Proteomics Departament, Gorgas Memorial Institute For Health Studies	Alexander A Martinez; Ambar Moreno; Claudia Estrada; Claudia Gonzalez V; César Roberto Conde Pereira; Jessica Gondola; Leyda Abrego; Marlene Castillo; Melissa Gaitan; Oris Chavarria
EPI_ISL_4635252, EPI_ISL_5504867	Laboratorio de Referencia Nacional de Virus Respiratorios, Centro Nacional de Salud Publica, Instituto Nacional de Salud Peru.	Laboratorio de Referencia Nacional de Virus Respiratorios, Centro Nacional de Salud Publica, Instituto Nacional de Salud Peru.	Alicia Nuñez Llanos; Carlos Padilla Rojas; Henri Bailon Calderon; Iris Silva Molina; Joseph Huayra Niqueen; Lely Solari Zepa; Luis Barcena Flores; Marco Galarza Perez; Nancy Rojas Serrano; Nieves Sevilla Castañeda; Omar Caceres Rey; Orson Mestanza Millones; Princesa Medrano Alhuay; Priscila Lope Pari; Sandra Morales Ruiz; Sara Gordillo Vilchez; Steve Acedo Lazo; Veronica Hurtado Vela; Victor Jimenez Vasquez; Wendy Lizarraga Olivares
EPI_ISL_2044785, EPI_ISL_2927081	Laboratory Corporation of America	Centers for Disease Control and Prevention Division of Viral Diseases, Pathogen Discovery	Adrian Paskey; Amanda Douglas; Amanda Suchanek; Andrea Throop; Ayla Burns; Benjamin Rambo-Martin; Bobbi Croy; Brian Krueger; Brian Norvell; Christopher Gulvick; Christos Petropoulos; Clinton R. Paden; Craig Lukasik; Dakota Howard; Darlene Wagner; Debbie Boles; Dhwani Batra; Duncan MacCannell; Eyad Almasri; Goran Stevovic; Howard Engler; Hrushikesh Deshmukh; Jake Humphrey; Jana Schroth; Jason Caravas; Joe Voshell; John Pruitt; Jonathan Maltzer; Jonathan Williams; Kara Moser; Kimberly Wagner; Lax Iyer; Lisa Pfeiffer; Lyndon Tilson; Manoj Jain; Marcia Eisenberg; Mary Ann Cristobal; Mary Williamson; Matthew Robinson; Matthew Schnerer; Michael Levandoski; Mike Sapeta; Minda Nye; Minoo Agarwal; Mohan Kalli; Nuthawin Charoensri; Oren Cohen; Peter W. Cook; Prashant Gupta; Qian Zeng; Rama Ghatti; Scott Parker; Scott Ryan; Scott Sammons; Shatavia Morrison; Stanley Letovsky; Steven Ragan; Suresh Babu Selvaraju; Susan Countryman; Susan Hicks; Suzanne Dale; Thomas Urban; Tim Kuphal; Tricia Zwiefelhofer; Vincent Drouillon; Yvette Unarumhi
EPI_ISL_1504634, EPI_ISL_2393531, EPI_ISL_2569439, EPI_ISL_4099792, EPI_ISL_4189290	see above	Wellcome Sanger Institute for the COVID-19 Genomics UK (COG-UK) Consortium	Cordelia Langford; David K. Jackson; Dominic Kwiatkowski; Ewan Harrison; Ian Johnston; Jacquelyn Wynn; Jeffrey Barrett; John Sillitoe on behalf of the Wellcome Sanger Institute COVID-19 Surveillance Team; Mairead Hyland; Roberto Amato; Sonia Goncalves; The Lighthouse Lab in Alderley Park and Alex Alderton
EPI_ISL_822325, EPI_ISL_2513334, EPI_ISL_4289982, EPI_ISL_4683454	Lighthouse Lab in Glasgow	Wellcome Sanger Institute for the COVID-19 Genomics UK (COG-UK) Consortium	Anna Dominiczak and Alex Alderton; Carol Clugston; Cordelia Langford; David Gray; David K. Jackson; Dominic Kwiatkowski; Ewan Harrison; Harper VanSteenhouse; Ian Johnston; Jeffrey Barrett; John Sillitoe on behalf of the Wellcome Sanger Institute COVID-19 Surveillance Team; Roberto Amato; Sonia Goncalves; Yumi Kasai
EPI_ISL_993421, EPI_ISL_2118914, EPI_ISL_2236023, EPI_ISL_3763330	Lighthouse Lab in Milton Keynes	Wellcome Sanger Institute for the COVID-19 Genomics UK (COG-UK) Consortium	Cordelia Langford; David K. Jackson; Dominic Kwiatkowski; Ewan Harrison; Ian Johnston; Jeffrey Barrett; John Sillitoe on behalf of the Wellcome Sanger Institute COVID-19 Surveillance Team; Roberto Amato; Sonia Goncalves; The Lighthouse Lab in Milton Keynes and Alex Alderton
EPI_ISL_1280209	Limbach - MVZ Labor EVELD & Kollegen Essen	Robert Koch Institute	
EPI_ISL_1153475	MVZ Dr. Eberhard & Partner Dortmund	Robert Koch Institute	
EPI_ISL_1218607, EPI_ISL_1439643, EPI_ISL_1439663, EPI_ISL_1439670, EPI_ISL_1440873, EPI_ISL_1440888, EPI_ISL_1440919	see above	Robert Koch Institute	
EPI_ISL_1347837	Medical Laboratories Duesseldorf	Center of Medical Microbiology, Virology, and Hospital Hygiene, University of Dueseldorf	Alexander Dikhey; Andreas Walker; Angelika Holmer; Christian Lange; Daniel Strolow; Jessica Nicolai; Jörg Timm; Klaus Pfeffer; Lisanna Hülse; Malte Kohns Vasconcelos; Maximilian Damagnez; Nadine Lübke; Tobias Wienemann; Torsten Houwaart
EPI_ISL_1527767	Michigan Department of Health and Human Services, Bureau of Laboratories	Michigan Department of Health and Human Services, Bureau of Laboratories	Blankenship HM; Riner D; Soehnlen MK
EPI_ISL_5015439, EPI_ISL_5015483	Nastavni zavod za javno zdravstvo Splitsko-Dalmatinske županije	Hrvatski zavod za javno zdravstvo	Irena Tabain; Ivana Ferencak
EPI_ISL_1915533	National Institute of Health Research and Development	National Institute of Health Research and Development	Agustiniingsih; Arie Ardiansyah Nugraha; Fauzul Muna; Hana Apsari Pawestri; Hartanti Dian Ikawati; Herna; Holy Arif Wibowo; Irene Lorida Indalao; Kartika Dewi Puspa; Kindi Adam; Krisna Nur Andriana Pangesti; Natalie Laurencia Kipuw; Nelly Pusparandi; Ni Ketut Susilarini; Nike Susanti; Nurika Hariastuti; Reni Herman; Rinin Ramadhany; Subangit; Tati FEBriyanti; Triyani Soekarso; Uily Afri Nikmah; Vivi Setiawaty.; Yuni Rukminiati
EPI_ISL_1911219	Naval Infectious Diseases Diagnostic Laboratory	Naval Medical Research Center Biological Defense Research Directorate	Andrea Luquette; Andrew Bennett; Bishwo Adhikari; Catherine Arnold; Francisco Malagon Bautista; Gregory Rice; Kimberly Bishop-Lilly; Kyle Long; Lindsay Giang; Logan Voeghtly; Megan Schilling; Michael Deschenes; Regina Cer; Victor Sugiharto
EPI_ISL_1196461	New Mexico Department of Health Scientific Laboratory	New Mexico Department of Health Scientific Laboratory	Anastacia Griego-Fisher; D'eldra Malone; Ellie Johnson; Jennifer Benoit
EPI_ISL_3903716	Nicolae Testemitanu State University of Medicine and Pharmacy	International Centre for Genetic Engineering and Biotechnology (ICGEB) and ARGO Open Lab Platform for Genome Sequencing	Buzunili L; Dal Monego S; Groppa S; Licastro D; Marcello A; Orsini E; Paraschiv A; Todiras M; Ulinici M
EPI_ISL_612110	Northumbria University / South Tees Hospitals NHS Foundation Trust / North Cumbria Integrated Care NHS Foundation Trust / North Tees and Hartlepool NHS Foundation Trust / Newcastle Hospitals NHS Foundation Trust	COVID-19 Genomics UK (COG-UK) Consortium	Andrew Nelson; Brendan Payne; Clive Graham; Darren L Smith; Debra Padgett; Edward Barton; Emma Swindells; Garren Scott; Gary Black; Gary Eltringham; Giles S Holt; Greg R Young; Jane Greenaway; Jennifer Collins; John Allan; Joshua Loh; Lynn Dover; Matthew Bashton; Mohammad A Tariq; Paul Baker; Sarah Essex; Steve Liggett; Wen C Yew; Yusri Taha
EPI_ISL_2373016	Northwestern Memorial Hospital	Northwestern University - Ozer Lab	Chad J. Achenbach; Chao Qi; Egon A. Ozer; Judd F. Hultquist; Lacy M. Simons; Lawrence J. Jennings; Michael G. Ison; Ramon Lorenzo-Redondo; Taylor J. Dean
EPI_ISL_5294882	Originating lab: Wales Specialist Virology Centre Sequencing lab: Pathogen Genomics Unit	Public Health Wales Microbiology Cardiff Wales Specialist Virology Centre	Alec Birchley; Alexander Adams; Amy Gaskin; Angela Marchbank; Bree Gatica-Wilcox; Catherine Moore; Jason Coombes; Joanne Watkins; Joel Southgate; Johnathan Evans; Laura Gifford; Lauren Gilbert; Lee Graham; Malorie Perry; Matthew Bull; Nicole Pacchiarini; Sally Corden; Sara Kumziene-Summerhayes; Sara Rey; Sarah Taylor; Simon Cottrell; Sophie Jones; Tom Connor
EPI_ISL_2037330	Pandemic Response Lab - NYC	Pandemic Response Lab, R&D	Cybill del Castillo; Dylan Law; Haiping Hao; Henry Lee; Jon Laurent; Katharine Nelson; Melissa Hopkins; Michael Hammerling; Pradeep Bugga; Shinyoung Clair Kang; Sol Rey; William Ward
EPI_ISL_1281983, EPI_ISL_1435362	Procomcure Biotech Germany GmbH	Robert Koch Institute	
EPI_ISL_2253727	Randox Laboratories	Wellcome Sanger Institute for the COVID-19 Genomics UK (COG-UK) Consortium	Cordelia Langford; David K. Jackson; Dominic Kwiatkowski; Ewan Harrison; Ian Johnston; Jeffrey Barrett; John Sillitoe on behalf of the Wellcome Sanger Institute COVID-19 Surveillance Team; Randox Laboratories and Alex Alderton; Roberto Amato; Sonia Goncalves
EPI_ISL_5041500, EPI_ISL_5281579	Respiratory Virus Unit, Microbiology Services Colindale, Public Health England	COVID-19 Genomics UK (COG-UK) Consortium	PHE Covid Sequencing Team
EPI_ISL_2776550	SARS-CoV-2 testing team, National Institute of Infectious Diseases	Pathogen Genomics Center, National Institute of Infectious Diseases	Hazuka Y Furihata; Hiromizu Takahashi; Kentaro Itokawa; Makoto Kuroda; Masanori Hashino; Masumichi Saito; Naomi Nojiri; Nozomu Hanaoka; Rina Tanaka; Tsuguto Fujimoto; Tsuyoshi Sekizuka
EPI_ISL_2112492	SYNLAB Jena Oncoscreen	Robert Koch Institute	
EPI_ISL_1729353, EPI_ISL_1851074, EPI_ISL_2110293	SYNLAB MVZ Berlin	Robert Koch Institute	
EPI_ISL_2130345	SYNLAB MVZ Leverkusen	Robert Koch Institute	
EPI_ISL_2934764, EPI_ISL_2934792, EPI_ISL_2934796	Servicio de Microbiología Clínica (Complejo Hospitalario de Navarra, Pamplona)	Centro de Secuenciación NASERTIC	Ana Miqueleiz; Ana Navascués; Carmen Ezpeleta Baquedano
EPI_ISL_2383298	Servicio de Microbiología, Hospital General	SeqCOVID-SPAIN	Maria Dolores Tirado Balaguer and SeqCOVID-SPAIN consortium; Rosario Moreno Muñoz

EPI_ISL_4518949, EPI_ISL_4519084	Universitario de Castellón Sharp HealthCare Laboratory	consortiumIBV(CSIC) Andersen Lab at Scripps Research	Art Mendoza; Cathy Woelfle; Jacquelyn Berumen; Liam McGinnis; Omid Bakhtar; SEARCH Alliance San Diego with Aaron Harding
EPI_ISL_1770548	State Laboratories Division, Hawaii State Department of Health	State Laboratories Division, Hawaii State Department of Health	Ayana Garnet; Drew Kuwazaki; Edward Desmond; Pamela O'Brien; Razvan Sultana
EPI_ISL_1899466, EPI_ISL_2419503	Swedish national genomic surveillance program of SARS-CoV-2	The Public Health Agency of Sweden	Alma Brolund; Maria Lind Karlberg; Maximilian Riess; Swedish national genomic surveillance program of SARS-CoV-2
EPI_ISL_3655956	University of Florida Health Pathology Laboratories	Salemi Lab, University of Florida	Cash MN; Lauzardo M; Magalis BR; Mavian C; Riva A; Salemi M; Tagliamonte M
EPI_ISL_2292512	Utah Public Health Laboratory	Utah Public Health Laboratory	Erin L. Young; Kelly F. Oakeson; Tara Gallagher
EPI_ISL_456551	Victorian Infectious Diseases Reference Laboratory (VIDRL)	Microbiological Diagnostic Unit Public Health Laboratory and Victorian Infectious Diseases Reference Laboratory, Doherty Institute	Cathy L.; Druce J.; Salt, M.; Schultz M.; Seemann T.; Sherry, N.
EPI_ISL_2662416	Viollier AG	Department of Biosystems Science and Engineering, ETH Zürich	Andrea Patrizia Salzmann; Chaoran Chen; Christiane Beckmann; Christoph Noppen; Henriette Kurth; Ivan Topolsky; Kim Philipp Jablonski; Lara Fuhrmann; Maurice Redondo; Niko Beerenwinkel; Olivier Kobel; Sarah Nadeau; Tanja Stadler
EPI_ISL_3115359	Virology Laboratory, University of Maiduguri Teaching Hospital	Africa Centre for Excellence for Genomics of Infectious Diseases (ACEGID), Redeemer's University	A.T.; Abechi; Ajogbasile; Akano; C.A.; C.T.; Eromon; F.V.; Folarin, O.; Happi; I.B.; J.N.; J.J.; K.O.; Kayode; Nosamiefan, I.; Oguzie; Olawoye; Olumade; Oluniyi; P.E.; P.S.; T.J.; Ugwu; Uwanibe
EPI_ISL_473537	West of Scotland Specialist Virology Centre, NHSGGC / MRC-University of Glasgow Centre for Virus Research	COVID-19 Genomics UK (COG-UK) Consortium	Alasdair MacLean; Alice Broos; Ana da Silva Filipe; Antonia Ho; Daniel Mair; David L. Robertson; Elihu Aranday-Cortes; Emma Thomson; James Shepherd; Jenna Nichols; Joseph Hughes; Kathy Li; Kathy Smollett; Kirstyn Bruncker; Kyriaki Nomikou; Lily Tong; Marc Niebel; Natasha Jesudason; Natasha Johnson; Patawee Asamaphan; Rajiv Shah; Richard Orton; Rory Gunson; Sarah McDonald; Sreenu Vattipally; Stephen Carmichael; Yasmin Parr
EPI_ISL_1447611, EPI_ISL_2296455	Yale Clinical Virology Lab	Grubaugh Lab • Yale School of Public Health	Anderson Brito; Annie Watkins; Chaney Kalinich; Chantal Vogels; Isabel Ott; Jessica Rothman; Joseph Fauver; Mallory Breban; Marie L. Landry; Mary Petrone; Nathan Grubaugh; Tara Alpert
EPI_ISL_1727175	amedes MVZ Hannover	Robert Koch Institute	
EPI_ISL_1156939	amedes MVZ für Laboratoriumsmedizin Rhein/Ruhr	Robert Koch Institute	

Supplemental Table 4: GISAID acknowledgments for genomes containing nsp12 E802D used to generate Figure S3