

Appendix

Table S1. Model Parameters and Data Sources.

Symbol	Description	Values	Sources
Ω_t^c	Number of travelers from SA to region c at time t	Daily mobility	Facebook Data for Good (1)
Υ_t^c	Number of flight seats from SA to region c on day t	Daily flight seats	Number of aircraft flying from SA to c in the OpenSky dataset (2); number of seats per aircraft (3).
κ	Proportion of available seats on flights from SA that are occupied in November, 2021	Scaling factor	The ratio of the total number of travelers (1) and the total number of flight seats (2) from SA to the 30 focal regions (2) in October 2021.
p_{sym}	Proportion of infections that are symptomatic	57%	Ref. (4), assuming Omicron has the same symptomatic proportion as the wildtype.
dH_t^{SA}	Number of COVID-19 hospital admissions in SA at time t	Daily admissions	Ref. (5)
$dI_{\text{presym},t}^{\text{SA}}$	Number of new pre-symptomatic variant infections in SA at time t	Daily cases	Estimated
$dI_{\text{asym},t}^{\text{SA}}$	Number of new asymptomatic variant infections in SA at time t	Daily cases	Estimated
ξ_t^{SA}	Prevalence of asymptomatic variant cases as a percentage of SA population at time t	Daily prevalence	Estimated
γ_t^c	Rate of variant introductions from SA to region c on day t	Daily rate	Estimated
dH_t^c	Number of COVID-19 hospital admissions in region c at time t	Daily admissions	Refs. (6,7)
$dI_{\text{presym},t}^c$	Number of new pre-symptomatic variant infections in region c at time t	Daily cases	Estimated
$dI_{\text{asym},t}^c$	Number of new asymptomatic variant	Daily cases	Estimated

	infections in region c at time t		
r	Proportion of symptomatic cases that are hospitalized	7.5/124	Ref. (8)
ω_t	Proportion of the Omicron variant among new SARS-CoV-2 cases at time t	Proportion of sequenced SARS-CoV-2 specimens in Africa (weekly data)	Ref. (9)
D_h	Expected delay from infection to hospital admission	10 days	5 days from infection to symptom onset (10); 5 days from symptom onset to hospital admission (11,12)
D_{presym}	Expected incubation period between infection and symptom onset for symptomatic cases	5 days	Ref. (10)
$D_{inf,s}$	Expected time from infection to recovery, for symptomatic cases	11 days	Sum of expected 5-day incubation (10) and expected 6-day symptomatic (11) period
$D_{inf,a}$	Expected time from infection to recovery, for asymptomatic cases	11 days	Ref. (11)
N_{SA}	Population of SA	59.31 million (2019)	Ref. (13)
N_c	Population of region c	2019 population estimates	Ref. (14)

Data

To estimate the daily number of passengers traveling between SA and other countries, we obtained daily between-country mobility data from Facebook Data for Good (1). Based on geolocation data from Facebook users that enabled the ‘location history’ feature, Facebook reconstructed the travel volume of anonymized passengers between different countries. Facebook reports the mobility flows for travel connections from SA to other countries with at least 1000 daily recorded passengers in the raw dataset. We obtained data for ~93 thousand passengers originating from SA to the other 30 non-African regions between November 1 and November 28, 2021. Out of the 840 region-day combinations (30 regions/28 days), 836 were missing data because of volumes below 1000.

The nomenclature system for SARS-CoV-2 genetic lineages is established by GISAID (15). The GISAID database includes ~200 sequenced SARS-CoV-2 specimens from Africa per week during the study period (9). Using this data, we estimate the relative frequency of Omicron per week and assume that proportion is the same each day of a given week.

Methods

Estimating missing daily travel volumes from South Africa

For regions with missing data at time t , we first estimate the daily flow Ω_t^c from SA as given by

$$F_t^c = \kappa \Upsilon_t^c$$

where κ is the estimated proportion of aircraft seats occupied and Υ_t^c is the number of available seats on flights from SA to region c . We then scale so that all estimated travel volumes fall between 1 and 1000 passengers, as given by

$$\Omega_t^c = F_t^c \cdot \frac{1000}{\max_j(F_t^j)}.$$

Risk of COVID-19 variant introduction via infected travelers from South Africa

To estimate the probability of Omicron introductions from SA, we first estimate the prevalence of pre-symptomatic and asymptomatic cases in SA and then use mobility data to estimate the likelihood that infectious cases traveled to regions across the world.

Our notation and parameter values are provided in **Table S1**. Briefly, we assume that the infected cases will develop symptoms with probability p_{sym} and hence remain asymptomatic with probability $1 - p_{\text{sym}}$ throughout their infection. The length of the

infectious period is $D_{\text{inf},a}$ and $D_{\text{inf},s}$ days for asymptomatic and symptomatic cases, respectively. Symptomatic cases will develop symptoms after an incubation period of D_{presym} days. Let r be the proportion of symptomatic cases who require hospitalization with a delay of D_h days after infection. Let dH_t^{SA} denote the number of new COVID-19 hospital admissions, and ω_t the proportion of COVID-19 cases infected by the Omicron variant among all sequenced SARS-CoV-2 specimens in SA on day t . We assume that the numbers of new pre-symptomatic and asymptomatic infections caused by the Omicron variant in SA on day t are given by:

$$\begin{aligned} dI_{\text{presym},t}^{\text{SA}} &= \frac{dH_{t+D_h}^{\text{SA}} \omega_t}{r} \\ dI_{\text{asym},t}^{\text{SA}} &= dI_{\text{presym},t}^{\text{SA}} \left(\frac{1-p_{\text{sym}}}{p_{\text{sym}}} \right). \end{aligned}$$

Then, the numbers of pre-symptomatic and asymptomatic cases in SA on day t are given by

$$\begin{aligned} I_{\text{presym},t}^{\text{SA}} &= \sum_{i=t-D_{\text{presym}}}^{t-1} dI_{\text{presym},i}^{\text{SA}} \\ I_{\text{asym},t}^{\text{SA}} &= \sum_{i=t-D_{\text{inf},a}}^{t-1} dI_{\text{asym},i}^{\text{SA}} \end{aligned}$$

The prevalence of pre-symptomatic and asymptomatic cases infected by the Omicron variant as a proportion of the SA population is given by

$$\xi_t^{\text{SA}} = \frac{I_{\text{presym},t}^{\text{SA}} + I_{\text{asym},t}^{\text{SA}}}{N_{\text{SA}}}$$

where N_{SA} is the population size of SA.

We assume that the proportion of travelers from SA that leave while infected is equal to the overall prevalence of pre-symptomatic and asymptomatic infections on a given day.

Then the rate of case introductions from SA to a region c on day t is approximated by

$$\gamma_t^c = \xi_t^{\text{SA}} \cdot \Omega_t^c,$$

where Ω_t^c is the estimated number of travelers from SA to region c on day t . We assume that

symptomatic cases do not travel during their symptomatic period.

Assuming that the introduction of Omicron cases from SA to each region c is essentially a non-homogeneous Poisson process (16–18), we estimate the probability of at least one Omicron case being introduced from SA to each region c by time t (starting at time t_0) as

$$1 - \exp\left(-\sum_{i=t_0}^t \gamma_i^c\right).$$

References

1. Home - Facebook Data for Good [Internet]. [cited 2021 Jan 2]. Available from: <https://dataforgood.fb.com/>
2. Strohmeier M, Olive X, Lübke J, Schäfer M, Lenders V. Crowdsourced air traffic data from the OpenSky Network 2019–2020. *Earth Syst Sci Data*. 2021 Feb 11;13(2):357–66.
3. Aéroports de Montréal. Reference table – Number of seats per aircraft type [Internet]. Admtl. 2021 [cited 2021 Dec 2]. Available from: https://www.admtl.com/sites/default/files/2021/2021_Appendix_6-Number-of-seats-per-aircraft-type.pdf
4. Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, et al. Spread of SARS-CoV-2 in the Icelandic Population. *N Engl J Med*. 2020 Jun 11;382(24):2302–15.
5. DAILY HOSPITAL SURVEILLANCE (DATCOV) REPORT - NICD [Internet]. 2020 [cited 2021 Dec 2]. Available from: <https://www.nicd.ac.za/diseases-a-z-index/disease-index-covid-19/surveillance-reports/daily-hospital-surveillance-datcov-report/>
6. Official UK Coronavirus Dashboard [Internet]. [cited 2020 Dec 29]. Available from: <https://coronavirus.data.gov.uk/details/healthcare>
7. Coronavirus (COVID-19) Hospitalizations [Internet]. [cited 2021 Jan 3]. Available from: <https://ourworldindata.org/covid-hospitalizations>
8. Centers for Disease Control and Prevention. Estimated Disease Burden of COVID-19 [Internet]. 2021 [cited 2021 Dec 2]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/burden.html>
9. GISAID - hCov19 Variants [Internet]. [cited 2021 Dec 2]. Available from: <https://www.gisaid.org/hcov19-variants/>
10. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20–28 January 2020.

Eurosurveillance. 2020 Feb 6;25(5):2000062.

11. Aleta A, Martín-Corral D, Pastore Y Piontti A, Ajelli M, Litvinova M, Chinazzi M, et al. Modelling the impact of testing, contact tracing and household quarantine on second waves of COVID-19. *Nat Hum Behav.* 2020 Aug 5;4(9):964–71.
12. ISARIC. International Severe Acute Respiratory and Emerging Infections Consortium (ISARIC). COVID-19 Report: 19 [Internet]. 2020 [cited 2021 Feb 1]. Available from: https://media.tghn.org/medialibrary/2020/05/ISARIC_Data_Platform_COVID-19_Report_19MAY20.pdf
13. Population of South Africa 1800-2020 [Internet]. [cited 2021 Dec 2]. Available from: <https://www.statista.com/statistics/1067083/population-south-africa-historical/>
14. World Population Prospects - Population Division - United Nations [Internet]. [cited 2021 Feb 3]. Available from: <https://population.un.org/wpp/Download/Standard/Population/>
15. Tracking SARS-CoV-2 variants [Internet]. [cited 2021 Dec 6]. Available from: <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>
16. Wang L, Wu JT. Characterizing the dynamics underlying global spread of epidemics. *Nat Commun.* 2018 Jan 15;9(1):218.
17. Scalia Tomba G, Wallinga J. A simple explanation for the low impact of border control as a countermeasure to the spread of an infectious disease. *Math Biosci.* 2008 Jul;214(1-2):70–2.
18. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet.* 2020 Feb 29;395(10225):689–97.