

Supplementary Equations

December 2, 2021

2-compartment PK model for bamlanivimab and etesevimab

$$\begin{aligned}\frac{dAb_{1c}}{dt} &= -Ab_{1c} \cdot \frac{CL_{Ab_{1c}}}{V_{Ab_{1c}}} - Q_a \cdot \frac{Ab_{1c}}{V_{Ab_{1c}}} + Q_a \cdot \frac{Ab_{1p}}{V_{Ab_{1p}}} \\ \frac{dAb_{1p}}{dt} &= Q_a \cdot \frac{Ab_{1c}}{V_{Ab_{1c}}} - Q_a \cdot \frac{Ab_{1p}}{V_{Ab_{1p}}} \\ \frac{dAb_{2c}}{dt} &= -Ab_{2c} \cdot \frac{CL_{Ab_{2c}}}{V_{Ab_{2c}}} - Q_b \cdot \frac{Ab_{2c}}{V_{Ab_{2c}}} + Q_b \cdot \frac{Ab_{2p}}{V_{Ab_{2p}}} \\ \frac{dAb_{2p}}{dt} &= Q_b \cdot \frac{Ab_{2c}}{V_{Ab_{2c}}} - Q_b \cdot \frac{Ab_{2p}}{V_{Ab_{2p}}}\end{aligned}$$

Equations describing phenomenological logarithmic activation of innate immune cells

1 Dendritic Cell Maturation (DC)

$$\begin{aligned}\frac{dDC}{dt} &= \alpha_{DC} \left[k_V \cdot \log(V) + k_I \cdot \log(I) + k_D \log(DAT1 + DAT2) \right] \left[k_{DC(TNF)} \cdot \left(\frac{(TNF\alpha)}{K_{DC(TNF)} + (TNF\alpha)} \right) + \right. \\ &k_{DC(IFN\gamma)} \cdot \left(\frac{(IFN\gamma)}{K_{DC(IFN\gamma)} + (IFN\gamma)} \right) + k_{DC(GM-CSF)} \cdot \left(\frac{(GM-CSF)}{K_{DC(GM-CSF)} + (GM-CSF)} \right) \left. \right] \\ &\left[\frac{K_{DC(IL10)}}{K_{DC(IL10)} + (IL10)} \right] - \beta_{DC} \cdot (DC)\end{aligned}$$

Mature (DC) are formed upon recognition of viral particles (V), infected cells (I), and damaged ($AT1$) and ($AT2$) cells. (DC) maturation is further induced by ($TNF\alpha$), ($IFN\gamma$), ($GM-CSF$), and inhibited by ($IL10$). Mature (DC) undergo nonspecific clearance at a rate β_{DC} .

2 Macrophage Activation ($M1$)

$$\begin{aligned} \frac{dM1}{dt} = & \alpha_{M1} \left[k_V \cdot \log(V) + k_I \cdot \log(I) + k_D \log(DAT1 + DAT2) \right] \left[k_{M1(TNF)} \cdot \left(\frac{(TNF\alpha)}{K_{M1(TNF)} + (TNF\alpha)} \right) + \right. \\ & \left. k_{M1(IFN\gamma)} \cdot \left(\frac{(IFN\gamma)}{K_{DC(IFN\gamma)} + (IFN\gamma)} \right) + k_{M1(GM-CSF)} \cdot \left(\frac{(GM-CSF)}{K_{M1(GM-CSF)} + (GM-CSF)} \right) \right] \\ & \left[\frac{K_{M1(IL10)}}{K_{M1(IL10)} + (IL10)} \right] - \beta_{M1} \cdot (M1) \end{aligned}$$

Activated ($M1$) are formed upon recognition of viral particles (V), infected cells (I), and damaged ($AT1$) and ($AT2$) cells. ($M1$) activation is induced by ($TNF\alpha$), ($IFN\gamma$), ($GM-CSF$), and inhibited by ($IL10$). Activated ($M1$) undergo nonspecific clearance at a rate β_{M1} .

3 Neutrophil Activation (N)

$$\begin{aligned} \frac{dN}{dt} = & \alpha_N \left[k_V \cdot \log(V) + k_I \cdot \log(I) + k_D \log(DAT1 + DAT2) \right] \left[k_{M1(TNF)} \cdot \left(\frac{(TNF\alpha)}{K_{M1(TNF)} + (TNF\alpha)} \right) + \right. \\ & \left. k_{M1(IFN\gamma)} \cdot \left(\frac{(IFN\gamma)}{K_{DC(IFN\gamma)} + (IFN\gamma)} \right) + k_{M1(GM-CSF)} \cdot \left(\frac{(GM-CSF)}{K_{M1(GM-CSF)} + (GM-CSF)} \right) \right] \\ & k_{trN} \cdot N_c \cdot \left(\frac{(IL17)}{k_{mN_IL17} + (IL17)} \right) - \beta_N \cdot N - k_{tr(N)} \cdot N \end{aligned}$$

Activated (N) are formed upon recognition of viral particles (V), infected cells (I), and damaged ($AT1$) and ($AT2$) cells. (N) activation is induced by ($TNF\alpha$), ($IFN\gamma$), ($GM-CSF$), and inhibited by ($IL10$). Activated (N) undergo nonspecific clearance at a rate β_N . Activated (N) migration is also induced by ($IL17$).