Characterization of the significant decline in humoral immune response six months post-SARS-CoV-2 mRNA vaccination: A systematic review

Kin Israel Notarte^{1*}, Israel Guerrero-Arguero^{2*}, Jacqueline Veronica Velasco¹, Abbygail Therese Ver¹, Maria Helena Santos de Oliveira³, Jesus Alfonso Catahay¹, Md. Siddiqur Rahman Khan², Adriel Pastrana¹, Grzegorz Juszczak⁴, Jordi B. Torrelles², Giuseppe Lippi⁵, Luis Martinez-Sobrido², Brandon Michael Henry²

¹*Faculty of Medicine and Surgery, University of Santo Tomas, Manila, Philippines*

²*Disease Intervention & Prevention and Population Health Programs, Texas Biomedical Research Institute, San Antonio, Texas, USA*

³*Department of Biostatistics, State University of Maringa, Brazil*

⁴*Department of Public Health, Medical University of Warsaw, Warsaw, Poland*

⁵*Section of Clinical Biochemistry, University of Verona, Verona, Italy*

**Equal contributors*

Corresponding Author:

Brandon Michael Henry, M.D.

Disease Intervention & Prevention and Population Health Programs

Texas Biomedical Research Institute

San Antonio, Texas, USA

Tel/Fax: +1 (210) 258-9400

Email: bhenry@txbiomed.org

Abstract

Accumulating evidence shows a progressive decline in the efficacy of coronavirus disease 2019 (COVID-19) mRNA vaccines such as Pfizer-BioNTech (mRNA BNT161b2) and Moderna (mRNA-1273) in preventing breakthrough infections due to diminishing humoral immunity over time. Thus, this review characterizes the kinetics of anti-SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) antibodies after the second dose of a primary cycle of COVID-19 mRNA vaccination. A systematic search of literature was performed and a total of 18 studies ($N=15,980$) were identified and reviewed. The percent difference of means of reported antibody titers were then calculated to determine the decline in humoral response after the peak levels post-vaccination. Findings revealed that the peak humoral response was reached at 21-28 days after the second dose, after which serum levels progressively diminished at 4-6 months post-vaccination. Additionally, results showed that regardless of age, sex, serostatus and presence of comorbidities, longitudinal data reporting antibody measurement exhibited a decline of both anti-receptor binding domain (RBD) IgG and anti-spike IgG, ranging from 94-95% at 90-180 days and 55-85% at 140-160 days, respectively, after the peak antibody response. This suggests that the rate of antibody decline may be independent of patient-related factors and peak antibody titers but mainly a function of time and antibody class/molecular target. Hence, this study highlights the necessity of more efficient vaccination strategies to provide booster administration in attenuating the effects of waning immunity, especially in the appearance of new variants of concerns (VoCs).

Keywords: COVID-19, Pfizer-BioNTech, mRNA BNT162b2, Moderna, mRNA 1273, IgG, immunoglobulins, vaccines

Introduction

The coronavirus disease 2019 (COVID-19) pandemic claimed millions of lives worldwide and remains an unprecedented challenge to global public health.^{1,2} Despite the increasing availability of therapeutics against Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) infection, mass vaccination is the mainstay for limiting new infections, reinfections, breakthrough infections, and unwanted sequelae from COVID-19.³⁻⁶ Although nationwide immunization programs have already been implemented in most countries, the mass production, allocation, and accessibility to COVID-19 vaccines remain a hurdle, especially in low-income countries where administration of primary and adjunctive doses could be challenging.⁷

Among different types of COVID-19 vaccines, the utilization of mRNA-based vaccines is unprecedented.⁸ Despite the efficacy of mRNA vaccines in reducing the risk of developing severe COVID-19 illness, ample evidence showed the progressive decline in their efficacy for preventing SARS-CoV-2 infections.⁹ Such vulnerability is correlated to the significant decline in anti-SARS-CoV-2 antibodies (Abs) over time.¹⁰⁻¹² This waning humoral response could lead to breakthrough infections, thus highlighting the potential role of serologic testing to assess vaccine immunogenicity and protection efficacy. While the role of T cells and B memory cells from vaccine-induced immunity has yet to be fully elucidated, multiple studies showed that Ab levels directly correlate with risk of vaccine reinfection and breakthrough, with low levels of anti-SARS-CoV-2 IgG in patients with infections correlating with high viral load and duration of viral shedding.^{3,4} In addition, the emergence of the new SARS-CoV-2 Omicron variant (B.1.1.529), with potential for high immune escape, highlights the necessity for vaccine boosters and high Ab titers for an improved immunoprotection.¹³ Thus, to better understand the rate of decay of anti-SARS-CoV-2 Abs through time, this rapid systematic review was designed to analyze humoral

response within a 6-month period among recipients of the Pfizer-BioNTech or Moderna COVID-19 mRNA-based vaccines to provide further evidence on the necessity of boosting vulnerable populations.

Methods

Search strategy and eligibility criteria

A systematic literature search was conducted to identify studies reporting the decline of humoral response of individuals post-mRNA vaccination. As shown in **Figure 1**, a comprehensive search was carried out in PubMed, Cochrane CENTRAL, Google Scholar, Science Direct, medRxiv, and Research Square for articles published from January 2021 to December 2021. The search keywords used included “SARS-CoV-2”, “COVID-19”, “declining humoral response”, “post-vaccination”, “mRNA vaccine”, “Pfizer-BioNTech”, “Moderna”, “mRNA-1273”, and “mRNA BNT162b2” which resulted in 51 journal articles.

Figure 1. Screening and appraisal of journal articles for inclusion in the systematic review.

For the inclusion criteria, articles reporting the following data were considered: (1) all participants who received a primary vaccination cycle with two complete doses of Pfizer-BioNTech (mRNA BNT162b2) or Moderna (mRNA-1273), (2) with at least a record of 10 patients and above (3) individual IgG or IgA, total Ig, or neutralizing anti-SARS-CoV-2 antibody titers, (4) quantitative or semi-quantitative antibody tests, (6) articles reporting humoral response 4-8–months post-vaccination, (7) articles available in the English language, and (8) randomized controlled, cohort, preprint or published papers and (9) providing complete and extractable data given the limited papers available for this novel disease and the mRNA vaccine.

The exclusion criteria were: (1) participants who received a single dose of mRNA vaccine only, (2) participants who received other types of vaccines such as Sinovac, AstraZeneca, Johnson

& Johnson, Novavax, and Sanofi-GSK, (3) articles reporting humoral response <4 months post-vaccination, and (4) anti-SARS-CoV-2 IgM response and cell-mediated immunity; the former because it is now increasingly clear that IgM plays a minor role against COVID-19, as it has lower sensitivity (64%), specificity (99%), and accuracy (94%) compared to IgG (93%, 100%, and 98%, respectively).¹⁴ IgM antibodies were also found to decline early (*i.e.*, at day 20 post-vaccination) and have lower neutralizing potential.¹⁵ The latter because we only focused on humoral immunity due to limited data and lack of standardization of available cell-mediated immunity assays, hence imposing difficulty in data analysis.

Data extraction

Data were extracted independently from each article by two authors into a spreadsheet. A third author checked the extracted data for completeness and accuracy. Any disagreements were resolved by consensus among the authors.

Descriptive and outcome data were extracted from the included studies, including type of vaccine, country of origin, sample size, age range or median age of the population, type of serologic test employed and its manufacturing company, analyzer used, immunoglobulin measured, and molecular target of immunoglobulin. In addition, serial or non-serial serologic measurement determination was done, to differentiate which study performed better patient follow-up, thus reducing methodologic bias due to greater accountability for inter-individual variability despite general lower sample size for serial serologic sampling. Additional data were requested from the original study authors when necessary.

Data analysis

All data reported in this study were in reference to the peak humoral response after a primary vaccination cycle with 2 doses. However, due to significant heterogeneity in the assays used to probe antibody titers, the antibody measurements reported in the articles were standardized using the percent difference of means. This shows the absolute value of the ratio of the difference between two groups, (groups A and B which pertain to antibody titer measurements on the peak and after the peak of humoral response) and their average, expressed as a percentage to enable standard comparison of these data regardless of their units of measurements and the diagnostic tools used for their quantification. It is computed using the formula below:

$$\text{Percent (\%) Difference of Means} = \left| \frac{(\text{Group A Mean} - \text{Group B Mean})}{\text{Group A Mean}} \right|$$

Group A - Antibody titer on the peak humoral response; **Group B** - Antibody titer after the peak humoral response

Therefore, values are reported as percentual decrease from the peak. Time points were also standardized to represent number of days after the second vaccine dose.

Scope and limitations

The demographic parameters used in this study were limited to age, sex, serostatus, and comorbidities such as hemodialysis or chronic kidney disease, immunologic disorders, metabolic derangements including heart disease, hypertension, and diabetes mellitus. It also includes a general section wherein no specific parameter was identified in determining the percent decrease in humoral response. These factors were analyzed independently of each other.

Moreover, the authors utilized both preprint and published journal articles written from January 2021 up to the present which included vaccination time points from days 0 to 201 of either

Pfizer-BioNTech (mRNA BNT162b2) or Moderna (mRNA-1273) in demonstrating the decline of humoral response. These vaccines were only discussed in this systematic review as they were the leading vaccine utilized worldwide and due to the wide range of available resources at the time of writing.

A quantitative meta-analysis was not done because of the heterogeneity determined between studies, as well as between different immunoassays used in each study, their units, sensitivities/specificities, and antigenic targets, not appropriate to perform a reliable and sensible pooled analysis. Thus, percent (%) differences in titers between groups were calculated, as an attempt to standardize the antibody data and enable a combined evaluation of different studies.

Lastly, we did not evaluate cell-mediated immunity and the kinetics of T cells after vaccination. The measurement of T-cell responses against SARS-CoV-2 is complex and lacks standardization to allow for comparisons in the literature, but is a priority topic for future research. Nonetheless, Gilbert et al. demonstrated that the majority (68.5%) of vaccine efficacy is mediated by neutralizing antibody responses, with the remaining 31.5% of vaccine efficacy may be attributed to cell-mediated responses, other functional antibodies, and innate immunity.¹⁶ These findings are in agreement with studies demonstrating the relationship between neutralizing antibody titers and breakthrough infections.^{3,4}

Results

A total of 18 articles ($N=15,980$) were included in this systematic review in which 14 studies focused on Pfizer-BioNTech ($n=13,364$), 2 on Moderna ($n=234$), and 2 included both Pfizer-BioNTech and Moderna ($n=2,382$). These were further divided into four categories, humoral immunity post-second dose vaccination influenced by age, sex, baseline serostatus, and

presence of comorbidities. Thirteen articles were included in the general section, four under the age category, four under sex, four under serostatus, and three articles under comorbidities. For the sex category, a total of 3,128 males and 6,237 females were included in the analysis. Meanwhile, the comorbidities cohort were further grouped according to the reported condition. Three studies reported on hemodialysis or end stage renal disease, one study reported on heart disease and hypertension, one for diabetes mellitus, and one for immunologic disorders. It is also worth noting that articles under these categories were non-exclusive, as most studies included more than one factor in their analysis. With respect to the geographical distribution, seven studies were from Italy, three studies from the United States, three from Israel, two from Belgium, and one each from Japan, Estonia and Austria. Of the reviewed articles that conducted serial study sampling, the largest sample size was from Italy ($n=787$).

Tables 1-5 provide a summary of the following trends observed in this study. In general, there is a substantial decrease of Ab titers against SARS-CoV-2 following 6 months post-second dose mRNA vaccination (**Figure 2**); however, the amount and rate of decline of the Ab titers varied according to the factors analyzed. Participants exhibited up to 95% decline in Ab levels at least 120 days after the peak rise of the IgG titers post-mRNA vaccination. In studies reporting longitudinal data on Ab measures, an observed decline from peak levels of anti-RBD IgG ranged from 94 to 95% at 90 to 180 days post-peak, while the anti-spike IgG showed a decline that ranged from 55 to 85% at 140 to 160 days after the peak (**Figure 2**). With respect to the vaccine brand, Pfizer-BioNTech displayed a 90 to 94% decline of anti-RBD IgG at 150 to 161 days following the peak, and a decline of IgG anti-spike protein ranging from 55 to 95% at 140 to 180 days after the peak. Meanwhile, Moderna exhibited a 69 to 96% decrease in anti-RBD IgG at 150-174 days from the peak, and a 45% decline in anti-spike IgG 90 days post-titer peak.

A consistent decrease in humoral immunity post-vaccination was found across all age groups ranging from 53 to 96% (**Figure 3**). Similarly, a consistent decline in Ab titers regardless of sex was also noted. Females reported a 91 to 93% decrease in their anti-RBD IgG levels, similar to males (ranging from 89 to 92%) (**Figure 4**). Anti-spike IgG levels in females decreased by 83%, also close to the decline seen in males (84%) (**Figure 4**). Regardless of serostatus, a diminished humoral immunity post-mRNA vaccination was evident. However, the Ab titers post-second dose of mRNA vaccine displayed varying trends (**Figure 5**). One study showed a 90% decrease of titers in the seronegative group, which is higher than the decline observed in the seropositive group (79%) (**Figure 5**). This is in contrast with two studies, one showing a 52% and the other a 60% decline in their seronegative participants, which is lower than the observed decline in their seropositive participants (68% and 74%, respectively) (**Figure 5**). For comorbidities cohorts, there was a consistent decrease in the total antibody and IgG levels among those individuals with kidney disease and undergoing hemodialysis (93%), heart disease and hypertension (91%), diabetes mellitus (90%), and immunologic disorders (85%).

Although most of the studies focused on IgG titers, one study measured the levels of anti-spike IgA with a decline up to 72% at 150 days after its peak, consistent with the rate of decay of other immunoglobulins. Conversely, a study from Italy reported the decline in total Ig reaching only up to 56% at 150 days post-peak, which is lower than the rate of decline measured in other studies.

Discussion

Declining Ab levels post-mRNA vaccination

In the midst of the COVID-19 pandemic, the effectiveness and lasting protective effects of anti-SARS-CoV-2 vaccines are topics of interest for the general population. Here, we highlight the

trends of Ab titers post-mRNA vaccination (**Table 1** and **Figure 2**). Most studies reported Ab peak titers within 1-4 weeks post-second dose vaccination, followed by a steady decline up to 90% in anti-SARS-CoV-2 Abs by 4-6 months post-vaccination. Consequently, as weeks post-mRNA vaccination increased, some studies reported an increase in the proportion of vaccine recipients whose Ab titers were below estimated protective thresholds.^{17,18} Some studies noted that at 6 months post-vaccination, 16.1% of the study population had non-protective levels of anti-RBD IgG.¹⁷ Similarly, other studies reported that at 6 months post-vaccination, 95% and 27% of the study population had anti-RBD IgG and anti-spike trimeric IgG levels below the protective threshold, respectively.¹⁸ Although results are slightly dependent on the immunoassay used, the kinetics of Ab titer decline display similar properties.¹⁸

Across all studies, the rate of decline was noted to be greater during the initial weeks after reaching peak levels compared to 4-6 months post-mRNA vaccination (**Figure 2**). While the Ab level needed for protection is not established yet, there is a correlation between seroimmunity and protection from infection as well as seroimmunity and the likelihood of transmission.¹⁹ Therefore, this declining trend, with data demonstrating up to 95% decline in Ab levels depending on the antigenic target, supports the need for boosters and serologic monitoring moving forward to identify low responders and timely need for additional booster administration.

Table 1. Declining humoral response post-second dose administration of SARS-CoV-2 mRNA vaccine.

Author	Vaccine	Country of Origin	Sample Size	Mean / Median Age	Immunoglobulin Measurement	Serial / non-serial Serologic Measurement	Reported Values	% Difference from Peak
Campo et al. ¹⁰	Pfizer-BioNTech	Italy	Healthcare workers: 274	Median (range)	Diasorin, LIAISON, IgG, S1/S2	Serial	Geometric means and 95% confidence Intervals	D28 There is 9% decrease 21 days after the peak rise of IgG titer on day 7 after full vaccination
			Male: 99 Female: 175	46.1 years (23-69 years)				D61 There is 41% decrease 54 days after the peak rise of IgG titer on day 7 after full vaccination
								D147 There is 55% decrease 140 days after the peak rise of IgG titer on day 7 after full vaccination
Salvagno et al. ¹²	Pfizer-BioNTech	Italy	Healthcare workers: 787		Roche, Elecsys, total Ig, RBD	Serial	Medians and interquartile ranges	D90 There is 38% decrease 60 days after the peak rise of IgG titer on day 30 after full vaccination
			<65 years: 754	Mean age 44±12 years				D180 There is 55% decrease 150 days after the peak rise of IgG titer on day 30 after full vaccination
			≥65 years: 33 Male: 268 Female: 519					
Ponticelli et al. ²⁰	Pfizer-BioNTech	Italy	Healthcare workers: 162 Male: 68 Female: 94	Mean age 42.5±11.9 years	SNIBE, MAGLUMI, IgG, RBD	Serial	Medians and interquartile ranges	D180 There is 90% decrease 150 days after the peak rise of IgG titer on day 30 after full vaccination
Salvagno et al. ²¹	Pfizer-BioNTech	Italy	Baseline SARS-CoV-2 seronegative	Median age: 42 years	Euroimmun, IgA, spike S1 subunit	Serial	Medians and interquartile ranges	D180 There is 71% decrease 150 days after the peak rise of IgA titer on day 30 after full vaccination

			healthcare workers: 97	IQR: 31-52 years				
Kato et al. ²²	Pfizer-BioNTech	Japan	Healthcare workers: 98 Male: 24 Female: 74	Median age: 43 years IQR: 38-49 years	Tosoh, AIA-CL, IgG, nucleocapsid/spike	Serial	Geometric means and 95% confidence intervals	D201 There is 93% decrease 180 days after the peak rise of IgG titer on day 21 after full vaccination
								D42 There is 48% decrease 35 days after the peak rise of IgG titer on day 7 after full vaccination
Naaber et al. ²³	Pfizer-BioNTech	Estonia	Laboratory employees: 122 Male: 21 Female: 101	Median age: 34 years Range: 21-69 years	Abbott, IgG, RBD	Serial	Medians and interquartile ranges	D84 There is 79% decrease 87 days after the peak rise of IgG titer on day 7 after full vaccination D180 There is 94% decrease 161 days after the peak rise of IgG titer on day 7 after full vaccination
Laing et al. ²⁴	Pfizer-BioNTech	USA	Healthcare workers: 187 Male: 56 Female: 131	Mean age: 42.8 years IQR: 21-69 years	LakePharma, Multiplex microsphere-based immunoassay, IgG, nucleocapsid/spike	Serial	Geometric means and 95% confidence intervals	D180 There is 77% decrease 150 days after the peak rise of IgG titer on day 30 after full vaccination
							Medians and interquartile ranges	D30-59 There is 48% decrease 30 days after the peak rise of total antibody titer on day 0-29 after full vaccination
Israel et al. ¹⁷	Pfizer-BioNTech	Israel	2653	Mean age: 56.45±15.87 years	Abbott, Alinity i, IgG, RBD	Non-serial		D60-89 There is 78% decrease 60 days after the peak rise of total

								antibody titer on day 0-29 after full vaccination
								<p>D90-119</p> <p>There is 87% decrease 90 days after the peak rise of total antibody titer on day 0-29 after full vaccination</p> <p>D120-149</p> <p>There is 89% decrease 120 days after the peak rise of total antibody titer on day 0-29 after full vaccination</p> <p>D150-179</p> <p>There is 92% decrease 150 days after the peak rise of total antibody titer on day 0-29 after full vaccination</p> <p>D180+</p> <p>There is 95% decrease 180 days after the peak rise of total antibody titer on day 0-29 after full vaccination</p>
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	8395	No data	Abbott, Quant II, IgG, RBD	Non-serial	Means and 95% confidence intervals	<p>D30-59</p> <p>There is 42% decrease 30 days after the peak rise of total antibody titer on day 7-29 after full vaccination</p> <p>D60-89</p> <p>There is 78% decrease 60 days after the peak rise of total antibody titer on day 7-29 after full vaccination</p>

								<p>D90-119 There is 87% decrease 90 days after the peak rise of total antibody titer on day 7-29 after full vaccination</p> <p>D120-149 There is 89% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination</p> <p>D150-179 There is 92% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination</p> <p>D 180+ There is 95% decrease 180 days after the peak rise of total antibody titer on day 7-29 after full vaccination</p>
Salvagno et al. ¹⁸	Pfizer-BioNTech	Italy	86	<p>Median age: 45 years</p> <p>IQR: 31-53 years</p>	<p>DiaSorin, LIAISON, IgG, spike trimeric</p> <p>New Industries Biomedical Engineering Co., LTD, MAGLUMI, IgG, RBD</p>	Serial	Medians and interquartile ranges	<p>D180 There is 83% and 92% decrease 150 days after the peak rise of anti-spike trimeric IgG and RBD IgG titers, respectively, on day 30 after full vaccination</p>
Khoury et al. ²⁶	Pfizer-BioNTech	Israel	<p>Healthcare workers: 100</p> <p>Male: 55</p> <p>Female: 45</p>	<p>Mean age: 40.7±13.7 years</p>	<p>Abbot's © quant assay, IgG, RBD</p>	Serial	Medians and ranges	<p>D60 There is 56% decrease 30 days after the peak rise of total antibody titer on day 30 after full vaccination</p>

								D90 There is 88% decrease 60 days after the peak rise of total antibody titer on day 30 after full vaccination
								D120 There is 94% decrease 90 days after the peak rise of total antibody titer on day 30 after full vaccination
Ciabattini et al. ²⁷	Pfizer-BioNTech	Italy	145	Mean Age: 48.8±14.3 years	In-house ELISA, IgG, S1/S2	Non-serial	Geometric means and 95% confidence Intervals	D97 There is 65% decrease 90 days after the peak rise of anti-spike IgG titer on day 7 after full vaccination D167 There is 85% decrease 160 days after the peak rise of anti-spike IgG titer on day 7 after full vaccination
Brisotto et al. ²⁸	Pfizer-BioNTech & Moderna	Italy	516 Male: 145 Female: 371	Median age: 46.5 years IQR: 36-55 years	SNIBE, MAGLUMI, IgG, RBD	Serial	Medians and interquartile ranges	D20 There is 83% decrease 90 days after the peak rise of IgG titer on day 30 after full vaccination

A

B

C

Anti-SARS-CoV-2 Anti-RBD and Anti-Spike Antibody Decline Levels

Figure 2. Kinetics of antibodies in studies reporting longitudinal data showing percentage decline from peak titer after vaccination for anti-spike (A) and anti-RBD (B) IgG antibodies. Data summary of antibody percentage decline levels at ~6 months post-vaccination (C).

This trend was also observed in both Pfizer-BioNTech and Moderna mRNA vaccines, and regardless of the vaccine brand used, the absolute Ab titers of those with older age, seronegative status, or presence of major comorbidities were consistently lower at each time point, findings consistent with previous studies.²⁹ Finally, with respect to the percentual decline in Ab titers 6 months post-second dose mRNA vaccination, there is a consistent rate of declining humoral response regardless of age, sex, serostatus, and comorbidities. Possible effects of the aforementioned factors on declining humoral response post-second dose mRNA vaccination are summarized in **Tables 2-5** and discussed in subsequent sections.

Contributing factors to declining humoral immunity following complete mRNA vaccination

Age

Both humoral and cellular immune responses are impaired as individuals age, ultimately leading to poorer vaccine response.³⁰ Studies looking into the relationship of age and humoral response after undergoing Pfizer-BioNTech vaccination noted an inverse relationship on the antibody titers mounted.^{12,31} The studies included in **Table 2** highlighted a similar trend. Regardless of the age group, there was a decline of titers starting one month after the second dose of vaccination (**Figure 3**). The mounted antibody titers are initially higher in the age group <60 years, but the rate of decline was consistent across all age groups. These all suggest the need for a booster vaccine dose for all age groups, but in low resource settings, prioritization should be given to older age groups who mount weaker humoral responses and overall lower antibody titers

Table 2. Effect of age in declining humoral response post-second dose administration of SARS-CoV-2 mRNA vaccine.

Author	Vaccine	Country of Origin	Sample Size	Mean / Median Age	Immunoglobulin Measurement	Serial / Non-serial Serologic Measurement	% Difference from Peak
Salvagno et al. ¹²	Pfizer-BioNTech	Italy	Healthcare workers: 787	Mean age: 44±12 years	Roche Elecsys, total Ig, RBD	Serial	D180
			<65 years: 754				<65 years Male There is 52% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
			≥65 years: 33				Female There is 56% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
			Male: 268 Female: 519				≥65 years Male There is 56% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
Salvagno et al. ¹⁸	Pfizer-BioNTech	Italy	86	Median age: 45 years	Diasorin, LIAISON, IgG, spike trimeric	Serial	D180
			Male: 41 Female: 45	IQR: 31-53 years	New Industries Biomedical Engineering Co., LTD, MAGLUMI, IgG, RBD		<60 years There is 83% and 81% decrease 150 days after the peak rise of anti-spike trimeric IgG and RBD IgG titers, respectively, on day 30 after full vaccination
							≥60 years

						There is 88% and 93% decrease 150 days after the peak rise of anti-spike trimeric IgG and RBD IgG titers, respectively, on day 30 after full vaccination
						D180
						18-55 years There is 83% decrease 167 days after the peak rise of IgG titer on day 14 after full vaccination
Doria-Rose et al. ³²	Moderna	USA	Healthy adult participants: 33	No data	Anti-SARS-CoV-2, IgG, RBD	Serial
						56-70 years There is 96% decrease 174 days after the peak rise of IgG titer on day 7 after full vaccination
						≥ 71 years There is 93% decrease 174 days after the peak rise of IgG titer on day 7 after full vaccination
						D120-150
						<60 years old There is 89% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	8395	No data	Abbott, Quant II, IgG, RBD	Non-serial
			Age in years < 18: 92 18-44: 2199 45-59: 3016 60-74: 2515 75+: 573			
						D150+ <60 years old There is 92% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination

D120-150**≥60 years old**

There is 87% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination

D150+**≥60 years old**

There is 86% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination

Figure 3. Comparison by age of the percentage decline of anti-SARS-CoV-2 antibodies from peak to last measured timepoint (~ 6 months) in each study by immunoglobulin class and molecular target.

Sex

Ab titers of females were higher even after humoral decline in all studies (**Table 3**); however, the percentage decline from the peak was similar for both sexes (**Figure 4**).^{12,25} While it is important to note that overall Ab titers may be influenced by the larger female population cohort (6,238 females vs. 3,128 males) involved in the included studies, this might be attributed to the sex-based difference in humoral immunity which affects vaccine responses. Previous studies indicated that females have a greater humoral response attributed to hormonal differences and X chromosomes leading to biallelic expression of certain genes.³³ Several studies support this given the higher Ab titers mounted by females after vaccination against SARS-CoV-2.^{31,34} Further, studies on the morbidity/mortality relationship due to SARS-CoV-2 infection between sexes found

that males have three times higher probability of requiring intensive care unit admission and higher odds of mortality.³⁵ While these studies point to the increased risk and mortality among males, the overall percentage decline of Ab titers identified post-second dose vaccination indicates the need for boosters regardless of sex.

Table 3. Effect of sex in declining humoral response post-second dose administration of SARS-CoV-2 mRNA vaccine.

Author	Vaccine	Country of Origin	Sample Size	Mean / Median Age	Immunoglobulin Measurement	Serial / Non-serial Serologic Measurement	% Difference from Peak
							D180
							<65 years
							Male
							There is 52% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							Female
							There is 56% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							≥65 years
							Male
							There is 56% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							Female
							There is 52% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							D180
							Males
							There is 72% decrease 150 days after the peak rise of IgA titer on day 30 after full vaccination

							Females There is 71% decrease 150 days after the peak rise of IgA titer on day 30 after full vaccination
							D180 Male There is 84% and 92% decrease 150 days after the peak rise of anti-spike trimeric IgG and RBD IgG titers, respectively, on day 30 after full vaccination
Salvagno et al. ¹⁸	Pfizer-BioNTech	Italy	86 Male: 41 Female: 45	Median age: 45 years IQR: 31-53 years	Diasorin, LIAISON, IgG, spike trimeric New Industries Biomedical Engineering Co., LTD, MAGLUMI, IgG, RBD	Serial	Female There is 83% and 93% decrease 150 days after the peak rise of anti-spike trimeric IgG and RBD IgG titers, respectively, on day 30 after full vaccination
							D120-150 Males There is 86% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	8395 Males: 2774 Females: 5621	No data	Abbott, Quant II, IgG, RBD	Non-serial	Females There is 88% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination

D150+**Males**

There is 89% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination

Females

There is 91% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination

Figure 4. Comparison by sex of the percentage decline of anti-SARS-CoV-2 antibodies from peak to last measured timepoint (~ 6 months) in each study by immunoglobulin class and molecular target.

Serostatus

Infection with SARS-CoV-2 elicits a humoral immune response *via* production of immunoglobulins targeting the spike, nucleocapsid, as well as other viral proteins.³⁶ This translates in detectable Ab titers prior to SARS-CoV-2 mRNA vaccine administration among previously infected individuals, and significantly higher Ab levels after the first dose compared to those with no history of prior infection.³⁶ However, Ab titers measured after the second vaccine dose report varying trends (**Figure 5**). Some studies reported that seropositive groups continue to have increased Ab levels compared to seronegative groups, while others reported that there is no significant difference between the two groups.^{37–42} These disparities in trend are conveyed in the rate of decline of Ab titers at 5-6 months after full immunization compared to peak levels. Among the four journal articles included in this study, two reported that seropositive groups had a larger

percent decline in Ab titers from peak compared with seronegative groups.^{10,12} This is possibly due to higher peak values making Ab catabolism more pronounced in seropositive groups.¹² Another possible explanation is due to timing, as seropositive groups reach peak Ab titers right after the first dose. Studies have observed that a second vaccine dose in seropositive groups does not significantly boost antibody titers higher, therefore these groups start their decline from peak much earlier than seronegative groups, who reach peak levels only after the second dose.^{36,38} Conversely, two studies reported that seronegative groups had a larger percent decline.^{43,44} This led to a recommendation to prioritize seronegative groups for booster administration over seropositive in cases of limited booster supply.⁴⁴ However, given the heterogeneity of these findings, as well as a poor understanding of the immunity arising from the combination of native infection supplemented by vaccinated, booster administration should likely be recommended to everyone regardless of serostatus.

Table 4. Effect of serostatus in declining humoral response post-second dose administration of SARS-CoV-2 mRNA vaccine.

Author	Vaccine	Country of Origin	Sample Size	Mean / Median Age	Immunoglobulin Measurement	Serial / Non-serial Serologic Measurement	% Difference from Peak
							D147
Campo et al. ¹⁰	Pfizer-BioNTech	Italy	Healthcare workers: 274 Male: 99 Female: 175	Median (range) 46.1 (23-69) years	Diasorin, LIAISON, IgG, S1/S2	Serial	Seronegative There is 60% decrease in IgG titer 140 days after the peak rise of IgG titer on day 7 after full vaccination Seropositive There is 68% decrease in IgG titer 140 days after the peak rise of IgG titer on day 7
							D180
Salvagno et al. ¹²	Pfizer-BioNTech	Italy	Healthcare workers: 787 Baseline seronegative: 624 Baseline seropositive: 163	Mean age 44±12 years	Roche Elecsys, total Ig, RBD	Serial	Seronegative There is 52% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination Seropositive There is 74% decrease 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							D159
Bayart et al. ⁴³	Pfizer-BioNTech	Belgium	Laboratory employees: 217 Male: 61 Female: 170	Male Mean age: 43 years Range: 23-64 years Female Mean age: 43 years Range: 23-66 years	Abbott Architect, IgG, RBD	Serial	Seronegative There is 90% decrease in IgG titer 152 days after the peak rise of IgG titer on day 7 after full vaccination Seropositive There is 79% decrease in IgG titer 138 days after the peak rise of IgG titer on day 21 after full vaccination

							D159
Tré-Hardy et al. ⁴⁴	Moderna	Belgium	Healthcare workers: 201	Median age:	LIAISON®, IgG, S1 and S2 subunit	Serial	Seronegative There is 45% decrease 90 days after the peak rise of IgG titer on day 69 after full vaccination
				50.1 (46.9-52.4) years			Seropositive IgG titer remains the same even 90 days after the peak rise on day 69 after full vaccination

Figure 5. Comparison of antibody levels of seronegative and seropositive kinetics in studies reporting longitudinal data showing percentage decline from peak titer after vaccination. Red lines indicate seronegative data and green lines indicate seropositive data.

Comorbidities

Several studies established that elderly patients (age >60 years) with stage-specific comorbidities along with other factors such as sex and serostatus elicit a comparably diminished humoral response post-vaccination.^{29,45} Moreover, the robustness of initial Ab response post-vaccination is reported to predict rapidity of Ab titer decline.¹⁹ Those with higher maximum Ab response within 2 months from full vaccination were less likely to become seronegative or develop SARS-CoV-2 infection.¹⁹ As such, it is important to follow the course of Ab titers several weeks after full vaccination among those with stage-specific comorbidities in order to predict better when best to give additional booster vaccine doses and ensure appropriate protection.

In this systematic review, all three studies involving populations with comorbidities reported a consistent decrease in Ab titers from peak values at 4 months or later post-second dose SARS-CoV-2 mRNA vaccination.^{19,25,46} Interestingly, studies reported that the percent decline of Ab titer for those with chronic kidney disease, heart disease, diabetes mellitus, and immunosuppressive disorders was greater at 120 days after peak levels (28-50 days after first dose) compared to at 150 days after.²⁵ These findings, however, may be attributed to the study design, as it was a non-serial study where the sample size at day 120 was less than at day 150, which could have skewed the results. This particular study was also limited by the lack of control groups, preventing us from directly comparing the percent decline of those with comorbidities *versus* healthy populations, although the absolute values of Ab levels were significantly lower in those with comorbidities compared to the total sample population.²⁵ Regardless of this limitation, when taken together with other studies in this systematic review, those with comorbidities exhibited a similar trend in percent decline with the general population and other cohorts.

Table 5. Effect of comorbidities in declining humoral response post-second dose administration of SARS-CoV-2 mRNA vaccine.

Author	Vaccine	Country of Origin	Sample Size	Mean / Median Age	Immunoglobulin Measurement	Serial / Non-serial Serologic Measurement	% Difference from Peak
<i>Hemodialysis and kidney disease</i>							
Hsu et al. ¹⁹	Pfizer-BioNTech & Moderna	USA	Dialysis patients: 1866 Male: 1084 Female: 782	Mean age $\pm$ SD 63.9 $\pm$ 13.7 years	ADVIA Centaur®, IgG, RBD	Non-serial	Pfizer-BioNTech
							D120 There is 86% decrease in IgG titer 90 days after the peak rise of total antibody titer on day 30 after full vaccination
							D180 There is 93.5% decrease in IgG titer 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							Moderna
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	Without chronic kidney disease (CKD): 7715	No data	Abbott, Quant II, IgG, RBD	Non-serial	D120 There is no decrease in IgG titer 90 days after the peak rise of total antibody titer on day 30 after full vaccination
							D180 There is 69% decrease in IgG titer 150 days after the peak rise of total antibody titer on day 30 after full vaccination
							With CKD D120-150

			With chronic kidney disease (CKD): 680					There is 87% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
								D150+ There is 72% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination
								All
			Hemodialysis Patients: 41					D180 There is 92% decrease 136 days after the peak rise in IgG titers at day 42 after full vaccination
Davidovic et al. ⁴⁶	Pfizer-BioNTech	Austria	Positive seroconversion six months after vaccination: 27	Mean±SD 67.3±15.5 years	Diasorin, LIAISON, IgG, spike protein trimer	Serial		Positive seroconversion D180 There is 91% decrease 136 days after the peak rise in IgG titers at day 42 after full vaccination
			No seroconversion six months after vaccination: 14					No seroconversion D180 There is 86% decrease 136 days after the peak rise in IgG titers at day 42 after full vaccination
Heart disease and hypertension								
								With heart disease
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	Without heart disease: 7790	No data	Abbott, Quant II, IgG, RBD	Non-serial		D120-150 There is 91% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
			With heart disease: 605					D150+ There is 79% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination

							With hypertension
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	Without hypertension: 5960	No data	Abbott, Quant II, IgG, RBD	Non-serial	D120-150 There is 86% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
			With hypertension: 2435				D150+ There is 86% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination
Diabetes mellitus							
							With diabetes mellitus
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	Without diabetes: 7342	No data	Abbott, Quant II, IgG, RBD	Non-serial	D120-150 There is 90% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
			With diabetes: 1053				D150+ There is 84% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination
Immunologic disorders							
							With immunosuppressive disorder
Kertes et al. ²⁵	Pfizer-BioNTech	Israel	Without immuno suppressive disorder: 7411	No data	Abbott, Quant II, IgG, RBD	Non-serial	D120-150 There is 85% decrease 120 days after the peak rise of total antibody titer on day 7-29 after full vaccination
			With immuno suppressive disorder: 984				D150+ There is 73% decrease 150 days after the peak rise of total antibody titer on day 7-29 after full vaccination

Conclusion

In this systematic review, we characterized the substantial decline in Abs at ~6 months after COVID-19 mRNA vaccinations. To date, several studies identified a significant correlation between time-dependent waning of Ab levels and an increased risk for breakthrough infections.^{47,48} In addition to the rise of SARS-CoV-2 VoC, like the Delta and more recently the Omicron variant, studies also suggest a correlation between cold weather and an increase in COVID-19 cases.⁴⁹ This is consistent with the established association between the circulation of respiratory viruses and climatic factors, displaying a peak incidence in the winter months.⁴⁹ As such, the waning immunity, new VoC such as Omicron, uncontrolled community transmission of the Delta variant, and winter seasons could mark the culmination of major waves of COVID-19, which could in part be abated by booster doses. Indeed, boosters are likely to provide cross-reactive protection against SARS-CoV-2 infection, including VoC.⁵⁰

Individuals in the older age group and with pre-existing comorbidities have lower levels of Abs compared to other individuals.²⁹ Hence, even with the same rate of decline in humoral immunity post-second dose mRNA vaccination as observed in our study, their titers are consistently lower, and may in many cases be below adequate thresholds for protection. Thus, it is prudent to prioritize the elderly and those with comorbidities for booster administration in the coming months.

As evidence of waning Ab levels accumulate, several countries have expanded their vaccine coverage to include giving off booster doses to all adults after completion of primary vaccination.^{51,52} As such, the rate of confirmed infection and the rate of severe illness were lower by a factor of 11.3 and 19.5 respectively, in the group that received the booster vaccine.⁵³ A secondary analysis had also shown that the rate of confirmed infection at least 12 days after booster

administration was lower than the rate after 4 to 6 days by a factor of 5.4.⁵³ Further research is needed to understand the kinetics of antibody responses after vaccine boosters and to determine if a similar rate of decline is seen following adjunctive doses as observed with initial vaccination regimen.

In conclusion, our findings show that the levels of protective Abs significantly and consistently decline 6-8 months after the second dose of anti-SARS-CoV-2 mRNA vaccines regardless of age, sex, baseline serostatus and comorbidities. This supports that the rate of Ab decline is mostly independent from patient-related factors and peak Ab titer achieved after the second mRNA dose, but instead mainly a function of time and Ab class/molecular target, with some variability arising from the sensitivity of the immunoassay.⁵⁴ Given this information, it may then be possible to predict the decline in Ab levels over time to an insufficient titer, based only on the patient's peak Ab titer (~4 weeks after second dose for seronegative patients), and thus identify in advance the optimal time when a booster vaccine dose should be indicated without the need for serial serologic monitoring. Such a strategy should be investigated for those who are likely to be low responders or at high risk of breakthrough infection.⁵⁵ A similar strategy could be used following booster vaccine doses upon evaluation of Ab kinetics. Alternatively, research should explore the use of age, sex, serostatus, and comorbidity to gauge the most suitable time to measure the antibody titer in order to assess the need for a booster dose based on the data presented in our report. Given the significant decline of Abs observed in our systematic review, as well as studies showing correlation between Ab titers and breakthrough infections, it is of utmost importance to provide equal opportunities in obtaining the vaccines, create more efficient vaccination strategies, and to emphasize the necessity of providing booster shots in attenuating the effects of waning immunity, especially in the appearance of new VoC, in addition to the forthcoming winter season.

Credit Authorship Contribution Statement

Kin Israel Notarte: Conceptualization, Visualization, Methodology, Validation, Formal analysis, Data curation, Writing – original draft, Writing – review & editing. **Israel Guerrero-Arguero:** Conceptualization, Formal analysis, Data curation, Writing – review & editing. **Jacqueline Veronica Velasco:** Methodology, Validation, Formal analysis, Data curation, Writing – original draft, Writing – review & editing. **Abbygail Therese Ver:** Methodology, Validation, Formal analysis, Data curation, Writing – original draft, Writing – review & editing. **Maria Helena Santos de Oliveira:** Formal analysis, Data curation, Writing – review & editing. **Jesus Alfonso Catahay:** Methodology, Validation, Formal analysis, Data curation, Writing – original draft, Writing – review & editing. **Md. Siddiquir Rahman Khan:** Writing – review & editing. **Adriel Pastrana:** Methodology, Validation, Formal analysis, Data curation, Writing – original draft, Writing – review & editing. **Grzegorz Juszczuk:** Writing – review & editing. **Jordi B. Torrelles:** Writing – review & editing. **Giuseppe Lippi:** Writing – review & editing. **Luis Martinez-Sobrido:** Writing – review & editing. **Brandon Michael Henry:** Conceptualization, Visualization, Validation, Formal analysis, Writing – review & editing.

Declaration of Competing Interests

The authors report no conflict of interests.

Ethical Approval

Not required.

REFERENCES

1. Albano PM, Notarte KI, Macaranas I, Maralit B. Cross-contamination in molecular diagnostic laboratories in low- and middle-income countries: A challenge to COVID-19 testing. *Philippine Journal of Pathology*. Published online 2020:7-11. doi:10.21141/PJP.2020.09
2. Ciotti M, Ciccozzi M, Terrinoni A, Jiang WC, Wang C bin, Bernardini S. The COVID-19 pandemic. *Critical Reviews in Clinical Laboratory Sciences*. 2020;57(6):365-388. doi:10.1080/10408363.2020.1783198
3. Bergwerk M, Gonen T, Lustig Y, et al. Covid-19 breakthrough Infections in vaccinated healthcare workers. *New England Journal of Medicine*. 2021;385(16):1474-1484. doi:10.1056/nejmoa2109072
4. Lippi G, Henry BM, Plebani M. Anti-sars-cov-2 antibodies testing in recipients of covid-19 vaccination: Why, when, and how? *Diagnostics*. 2021;11(6):1-10. doi:10.3390/diagnostics11060941
5. Lippi G, Henry BM, Plebani M. Optimizing effectiveness of COVID-19 vaccination: Will laboratory stewardship play a role? *Clinical Chemistry and Laboratory Medicine*. 2021;59(12):1885-1888. doi:10.1515/ccm-2021-0972
6. Noor R. A Review on the effectivity of the current COVID-19 drugs and vaccines: Are they really working against the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) variants? *Current Clinical Microbiology Reports*. 2021;8(3):186-193. doi:10.1007/s40588-021-00172-w
7. Asundi A, O'Leary C, Bhadelia N. Global COVID-19 vaccine inequity: The scope, the impact, and the challenges. *Cell Host and Microbe*. 2021;29(7):1036-1039. doi:10.1016/j.chom.2021.06.007
8. Huang Q, Zeng J, Yan J. COVID-19 mRNA vaccines. *Journal of Genetics and Genomics*. 2021;48(2):107-114. doi:10.1016/j.jgg.2021.02.006
9. Levin EG, Lustig Y, Cohen C, et al. Waning immune humoral response to BNT162b2 COVID-19 vaccine over 6 months. *New England Journal of Medicine*. Published online 2021:1-11. doi:10.1056/nejmoa2114583
10. Campo F, Venuti A, Pimpinelli F, et al. Antibody persistence 6 months post-vaccination with bnt162b2 among health care workers. *Vaccines*. 2021;9(10):1-9. doi:10.3390/vaccines9101125
11. Klompas M. Understanding breakthrough infections following mRNA SARS-CoV-2 vaccination. *JAMA*. Published online 2021:1-3. doi:10.1001/jama.2021.19063
12. Salvagno GL, Lippi G, Henry BM, Pighi L, de Nitto S. Total anti-SARS-CoV-2 antibodies measured 6 months after Pfizer-BioNTech COVID-19 vaccination in healthcare workers. *Journal of Medical Biochemistry*. 2021;(October):0-5. doi:10.5937/jomb0-33999
13. Pulliam JRC, van Schalkwyk C, Govender N, et al. Increased risk of SARS-CoV-2 reinfection associated with emergence of the Omicron variant in South Africa. *medRxiv*. Published online 2021. doi:10.1101/2021.11.11.21266068
14. Hoffman T, Nissen K, Krambrich J, et al. Evaluation of a COVID-19 IgM and IgG rapid test; an efficient tool for assessment of past exposure to SARS-CoV-2. *Infection Ecology and Epidemiology*. 2020;10(1). doi:10.1080/20008686.2020.1754538

15. Liu X, Wang J, Xu X, Liao G, Chen Y, Hu CH. Patterns of IgG and IgM antibody response in COVID-19 patients. *Emerging Microbes and Infections*. 2020;9(1):1269-1274. doi:10.1080/22221751.2020.1773324
16. Gilbert PB, Montefiori DC, McDermott A, et al. Immune Correlates Analysis of the mRNA-1273 COVID-19 Vaccine Efficacy Trial. *Science*. Published online November 23, 2021. doi:10.1126/science.abm3425. *Epub ahead of print*
17. Israel A, Shenhar Y, Green I, et al. Large-scale study of antibody titer decay following BNT162b2 mRNA vaccine or SARS-CoV-2 infection. *medRxiv*. Published online August 21, 2021. doi:10.1101/2021.08.19.21262111
18. Salvagno GL, Henry BM, Pighi L, de Nitto S, Lippi G, Gianfilippi G. The pronounced decline of anti-SARS-CoV-2 spike trimeric IgG and RBD IgG in baseline seronegative individuals 6 months after BNT162b2 vaccination is consistent with the need for vaccine boosters. Published online 2021. doi:10.21203/rs.3.rs-1063499/v1
19. Hsu CM, Weiner DE, Manley HJ, et al. Seroresponse to SARS-CoV-2 vaccines among maintenance dialysis patients over six months. *medRxiv*. Published online 2021. doi:10.1101/2021.09.13.21263535
20. Ponticelli D, Antonazzo I, Caci G, et al. Dynamics of antibody response to BNT162b2 mRNA COVID-19 vaccine after 6 months. *Journal of Travel Medicine*. Published online October 26, 2021. doi:10.1093/jtm/taab173
21. Salvagno GL, Henry BM, Pighi L, de Nitto S, Lippi G. Six-month decline of serum anti-spike S1 subunit IgA in SARS-CoV-2 in seronegative healthcare workers after mRNA-based COVID-19 vaccination. Published online 2021. doi:10.21203/rs.3.rs-1019658/v1
22. Kato H, Miyakawa K, Ohtake N, et al. Vaccine-induced humoral and cellular immunity against SARS-CoV-2 at 6 months post BNT162b2 vaccination. doi:10.1101/2021.10.30.21265693
23. Naaber P, Tserel L, Kangro K, et al. Dynamics of antibody response to BNT162b2 vaccine after six months: a longitudinal prospective study. *The Lancet Regional Health - Europe*. 2021;10:100208. doi:10.1016/j.lanepe.2021.100208
24. Laing ED, Weiss CD, Samuels EC, et al. Durability of antibody responses and frequency of clinical and subclinical SARS-CoV-2 Running title: Antibody durability post-BNT162b2 Mitre. doi:10.1101/2021.10.16.21265087
25. Kertes J, Baruch Gez S, Saciuk Y, et al. Effectiveness of the mRNA BNT162b2 vaccine six months after vaccination: findings from a large Israeli HMO. *medRxiv*. Published online 2021. doi:10.1101/2021.09.01.21262957
26. Khoury J, Najjar-Debbiny R, Hanna A, et al. COVID-19 vaccine – Long term immune decline and breakthrough infections. *Vaccine*. 2021;39(48):6984-6989. doi:10.1016/j.vaccine.2021.10.038
27. Ciabattini A, Pastore G, Fiorino F, et al. Evidence of SARS-CoV-2-Specific Memory B Cells Six Months After Vaccination With the BNT162b2 mRNA Vaccine. *Frontiers in Immunology*. 2021;12. doi:10.3389/fimmu.2021.740708
28. Brisotto G, Muraro E, Montico M, et al. IgG antibodies against SARS-CoV-2 decay but persist 4 months after vaccination in a cohort of healthcare workers. *Clinica Chimica Acta*. 2021;523:476-482. doi:10.1016/j.cca.2021.10.035
29. Notarte KI, Ver AT, Velasco JV, et al. Effects of Age, Sex, Serostatus and Underlying Comorbidities on Humoral Response Post-SARS-CoV-2 Pfizer-BioNTech Vaccination –

- A Systematic Review. *medRxiv*. Published online 2021.
doi:10.1101/2021.10.10.21264825
30. Frasca D, Diaz A, Romero M, Landin AM, Blomberg BB. Age effects on B cells and humoral immunity in humans. *Ageing Research Reviews*. 2011;10(3):330-335.
doi:10.1016/j.arr.2010.08.004
 31. Jabal KA, Ben-Amram H, Beiruti K, et al. Impact of age, ethnicity, sex and prior infection status on immunogenicity following a single dose of the BNT162b2 mRNA COVID-19 vaccine: Real-world evidence from healthcare workers, Israel, December 2020 to January 2021. *Eurosurveillance*. 2021;26(6). doi:10.2807/1560-7917.ES.2021.26.6.2100096
 32. Doria-Rose N, Suthar MS, Makowski M, et al. Antibody Persistence through 6 Months after the Second Dose of mRNA-1273 Vaccine for Covid-19. *New England Journal of Medicine*. 2021;384(23):2259-2261. doi:10.1056/nejmc2103916
 33. Fink AL, Klein SL. The evolution of greater humoral immunity in females than males: implications for vaccine efficacy. *Current Opinion in Physiology*. 2018;6:16-20.
doi:10.1016/j.cophys.2018.03.010
 34. Pellini R, Venuti A, Pimpinelli F, et al. Initial observations on age, gender, BMI and hypertension in antibody responses to SARS-CoV-2 BNT162b2 vaccine. *EClinicalMedicine*. Published online 2021. doi:10.1016/j.eclinm.2021.100928
 35. Peckham H, de Grujter NM, Raine C, et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ICU admission. *Nature Communications*. 2020;11(1). doi:10.1038/s41467-020-19741-6
 36. Fraley E, LeMaster C, Geanes E, et al. Humoral immune responses during SARS-CoV-2 mRNA vaccine administration in seropositive and seronegative individuals. *BMC Medicine*. 2021;19(1). doi:10.1186/s12916-021-02055-9
 37. Callegaro A, Borleri D, Farina C, et al. Antibody response to SARS-CoV-2 vaccination is extremely vivacious in subjects with previous SARS-CoV-2 infection. *Journal of Medical Virology*. 2021;93(7):4612-4615. doi:10.1002/jmv.26982
 38. Ebinger JE, Fert-Bober J, Printsev I, et al. Antibody responses to the BNT162b2 mRNA vaccine in individuals previously infected with SARS-CoV-2. *Nature Medicine*. 2021;27(6):981-984. doi:10.1038/s41591-021-01325-6
 39. Kelsen SG, Braverman AS, Aksoy MO, et al. A Longitudinal Study of BNT162b2 Vaccine-Induced Humoral Response and Reactogenicity in Health Care Workers with Prior COVID-19 Disease. doi:10.1101/2021.03.18.21253845
 40. Modenese A, Paduano S, Bargellini A, et al. Neutralizing anti-SARS-CoV-2 antibody titer and reported adverse effects, in a sample of italian nursing home personnel after two doses of the BNT162b2 vaccine administered four weeks apart. *Vaccines*. 2021;9(6).
doi:10.3390/vaccines9060652
 41. Padoan A, Dall'Olmo L, Rocca F della, et al. Antibody response to first and second dose of BNT162b2 in a cohort of characterized healthcare workers. *Clinica Chimica Acta*. 2021;519:60-63. doi:10.1016/j.cca.2021.04.006
 42. Salvagno GL, Henry BM, di Piazza G, et al. Anti-sars-cov-2 receptor-binding domain total antibodies response in seropositive and seronegative healthcare workers undergoing covid-19 mrna bnt162b2 vaccination. *Diagnostics*. 2021;11(5).
doi:10.3390/diagnostics11050832

43. Bayart JL, Douchfils J, Gillot C, et al. Waning of igt, total and neutralizing antibodies 6 months post-vaccination with bnt162b2 in healthcare workers. *Vaccines*. 2021;9(10). doi:10.3390/vaccines9101092
44. Tré-Hardy M, Cupaiolo R, Wilmet A, et al. Six-month interim analysis of ongoing immunogenicity surveillance of the mRNA-1273 vaccine in healthcare workers: A third dose is expected. *Journal of Infection*. Published online November 1, 2021. doi:10.1016/j.jinf.2021.08.031
45. Salmerón Ríos S, Mas Romero M, Cortés Zamora EB, et al. Immunogenicity of the BNT162b2 vaccine in frail or disabled nursing home residents: COVID-A study. *Journal of the American Geriatrics Society*. 2021;69(6):1441-1447. doi:10.1111/jgs.17153
46. Davidovic T, Schimpf J, Abbassi-Nik A, et al. Waning humoral response 6 months after SARS-CoV-2 vaccination with the mRNA-BNT162b2 vaccine in hemodialysis patients: time for a boost. *Kidney International*. 2021;100(6):1334-1335. doi:10.1016/j.kint.2021.10.006
47. Mizrahi B, Lotan R, Kalkstein N, et al. Correlation of SARS-CoV-2 Breakthrough Infections to Time-from-vaccine; Preliminary Study. *medRxiv*. Published online 2021. doi:10.1101/2021.07.29.21261317
48. Shrotri M, Navaratnam AMD, Nguyen V, et al. Spike-antibody waning after second dose of BNT162b2 or ChAdOx1. *The Lancet*. 2021;398(10298):385-387. doi:10.1016/S0140-6736(21)01642-1
49. Bolaño-Ortiz TR, Camargo-Cacedo Y, Puliafito SE, et al. Spread of SARS-CoV-2 through Latin America and the Caribbean region: A look from its economic conditions, climate and air pollution indicators. *Environmental Research*. 2020;191. doi:10.1016/j.envres.2020.109938
50. Corbett KS, Gagne M, Wagner DA, et al. Protection against SARS-CoV-2 beta variant in mRNA-1273 vaccine–boosted nonhuman primates. *Science*. Published online October 21, 2021. doi:10.1126/SCIENCE.ABL8912/SUPPL_FILE/SCIENCE.ABL8912_SM.PDF
51. Mahase E. Covid-19 booster vaccines: What we know and who’s doing what. *BMJ (Clinical research ed)*. 2021;374:n2082. doi:10.1136/bmj.n2082
52. Mbaeyi S, Oliver SE, Collins JP, et al. The Advisory Committee on Immunization Practices’ Interim Recommendations for Additional Primary and Booster Doses of COVID-19 Vaccines — United States, 2021. *MMWR Morbidity and Mortality Weekly Report*. 2021;70(44):1545-1552. doi:10.15585/mmwr.mm7044e2
53. Bar-On YM, Goldberg Y, Mandel M, et al. Protection of BNT162b2 Vaccine Booster against Covid-19 in Israel. *New England Journal of Medicine*. 2021;385(15):1393-1400. doi:10.1056/nejmoa2114255
54. Danese E, Montagnana M, Salvagno GL, et al. Comparison of five commercial anti-SARS-CoV-2 total antibodies and IgG immunoassays after vaccination with BNT162b2 mRNA. *Journal of medical biochemistry*. 2021;40(4):335-340. doi:10.5937/JOMB0-31475
55. Dickerman BA, Gerlovin H, Madenci AL, et al. Comparative Effectiveness of BNT162b2 and mRNA-1273 Vaccines in U.S. Veterans. *New England Journal of Medicine*. Published online December 1, 2021. doi:10.1056/NEJMoa2115463