The Omega-3 Index is Inversely Associated with the Neutrophil-Lymphocyte Ratio in Adults

Michael I. McBurney¹⁻³, Nathan L. Tintle^{1,4}, William S. Harris^{1,5}

¹Fatty Acid Research Institute, Sioux Falls, SD 57106, USA (MIM, NLT, WSH); ²Department of Human Health and Nutritional Sciences, University of Guelph, Guelph, ON N1G 2W1, Canada (MIM), ³Division of Biochemical and Molecular Biology, Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA 02111, USA (MIM); ⁴Department of Population Health Nursing Science, College of Nursing, University of Illinois – Chicago, Chicago, IL 60612, USA(NLT); ⁵Sanford School of Medicine, University of South Dakota, Sioux Falls, SD 57105, USA (WSH).

Authors last names: McBurney, Tintle, Harris

Corresponding author: M.I. McBurney, Fatty Acid Research Institute, Sioux Falls, SD 57106, USA. E-mail: mim@faresinst.com

Reprint requests: W.S. Harris, Fatty Acid Research Institute, Sioux Falls, SD 57106, USA. E-mail: wsh@faresinst.com

Sources of Support: This work was supported by the Fatty Acid Research Institute (FARI). FARI is a non-profit foundation bringing together nutrition scientists and biostatistical experts to accelerate discovery of the relationships between fatty acids, especially omega-3 fatty acids, and health. Pending application and approval, data described in the manuscript, code book, and analytic code will be made available upon request to the Fatty Acid Research Institute (<https://www.faresinst.org/>).

25

26 **Summary**

27 The neutrophil-lymphocyte ratio (NLR) is a biomarker of systemic inflammation and measures
 28 innate-adaptive immune system balance. The omega-3-index (O3I) measures the amount of
 29 EPA+DHA in blood. Both a low O3I and an elevated NLR are associated with increased risk for
 30 chronic disease and mortality, including cardiovascular diseases and cancer. Hypothesizing that
 31 low O3I may partly contribute to systemic chronic inflammation, we asked if a relationship
 32 existed between O3I and NLR in healthy adults (≥ 18 y, $n=28,871$, 51% female) without
 33 inflammation [C-reactive protein (CRP) $< 3\text{mg/mL}$] who underwent a routine clinical
 34 assessment. NLR was inversely associated with O3I before ($p < 0.0001$) and after adjusting for
 35 age, sex, BMI, and CRP ($p < 0.0001$). Pearson correlations of other variables with NLR were
 36 $r=0.06$ (CRP), $r=0.14$ (age), and $r=0.01$ (BMI). In this healthy population, an O3I $< 6.6\%$ was
 37 associated with increasing NLR whereas NLR remained relatively constant (low) when O3I
 38 $> 6.6\%$, suggestive of a quiescent, balanced immune system.

39

40 **Keywords:** neutrophils, lymphocytes, neutrophil-lymphocyte ratio, omega-3 index,
41 eicosapentaenoic acid, docosahexaenoic acid
42
43 **Abbreviations:** BMI, body mass index; CRP, high-sensitivity C-reactive protein; DHA,
44 docosahexaenoic acid; EPA, eicosapentaenoic acid; NLR, neutrophil-lymphocyte ratio; O3I,
45 omega-3 index; SPM, specialized pro-resolving lipid mediators.
46

1. Introduction

Cells from both the innate and adaptive immune system circulate in blood. Cells of the former (i.e., neutrophils, phagocytes, dendritic cells, macrophages, eosinophils, basophils, mast cells, and natural killer cells) work together to neutralize pathogens, in part by releasing pro-inflammatory mediators that modulate the latter. Neutrophils are the most abundant leukocytes in circulation. They migrate to sites of tissue injury or damage and are responsible for a non-specific inflammatory response. The adaptive system (acquired immunity) responds to specific pathogens and cytokines by producing antibodies and maintains immunological memory via lymphocytes, i.e., T and B cells. The hallmark of the adaptive immune system is clonal expansion of lymphocytes. Inflammation is generally self-limiting, with its resolution being an active rather than passive process [1]. Chronic or unresolved inflammation can damage host tissues and increase risk for the development of a range of non-communicable diseases, e.g. cardiovascular, cancer, chronic respiratory diseases, diabetes, and autoimmune disorders, e.g. rheumatoid arthritis [2–4].

The neutrophil-lymphocyte ratio (NLR) is an emerging biomarker of systemic inflammation [5–8] and innate-adaptive immune system balance [6,9] that is readily accessible from a complete blood count. It has the benefit of not being influenced by physiological conditions such as dehydration and exercise [6]. NLR has been used to monitor astronaut immune function during long-duration missions [10]. Elevated NLR is also a biomarker of systemic inflammation [5,11–14], cardiovascular disease and events [6,15–18], the severity of covid-19 [19–22], mood disorders [23], cognitive impairment [24], cancer [6,7,25], dry eye [26], periodontitis [27],

rheumatoid arthritis [28], cancer [12,25,29], diabetic kidney disease [30] and total mortality [6,9,14,17,31,32].

Nutrient deficiencies or inadequacies can impair immune system function and weaken the immune response [3,33–36]. The proportion of EPA and DHA in immune cell membranes can modulate inflammation via the synthesis of prostaglandins, leukotrienes, lipoxins and specialized pro-resolving lipid mediators (SPMs, e.g., resolvins, protectins, and maresins) [1,37]. The EPA and DHA-derived metabolites are less inflammatory than those derived from the omega-6 precursor, arachidonic acid [1,33,37–42]. SPMs derived from EPA and DHA reduce neutrophil infiltration and production of reactive oxygen species, regulate the cytokine-chemokine axis, and temper the inflammatory response without immunosuppression [37,43–45]. The omega-3 index (O3I) [46] is a stable biomarker of long term EPA and DHA intake [47,48]. A low O3I [49] and an elevated NLR [14,18,50] are both associated with risk of cardiovascular events and mortality (**Supplemental Table 1**). Hypothesizing that low blood EPA+DHA levels may partly contribute to systemic chronic inflammation, we asked if there was a relationship between the O3I and the NLR in healthy individuals without inflammation.

2. Patients and Methods

This is a cross-sectional analysis of data from blood samples submitted for testing to Health Diagnostic Laboratory, Inc (HDL, Inc., Richmond VA; now defunct) as part of routine clinical assessment between 2011-2012. Subjects were adults (≥ 18 y) with data on O3I, NLR, body mass index (BMI), age, sex, and high-sensitivity C-reactive protein (CRP) (n=44,925). Individuals with extreme O3I values (i.e., highest and lowest 0.5%) were excluded. Because CRP > 3 mg/L is indicative of inflammation arising from infection, trauma or chronic disease [51–54],

individuals with CRP >3 mg/L were also excluded. Thus, the final study population consisted of 28,871 individuals (**Figure 1**).

2.1 Laboratory Methods

Blood samples were drawn after an overnight fast and shipped with cold packs to HDL, Inc. for testing. Samples were prepared at each clinical site according to standardized instructions as previously described [55]. Absolute concentrations of blood cells (neutrophils, lymphocytes, NLR) were determined with a Beckman-Coulter DxH 800 analyzer (Brea, CA, USA) and biomarker data (CRP, O3I) were extracted without any linked patient identifiers except age, sex, and BMI. For fatty acid analysis, RBCs were separated from plasma by centrifugation and analyzed using gas chromatography as previously described [56]. The University of South Dakota Institutional Review Board reviewed and approved the use of such de-identified clinical data for research purposes (IRB-21-147).

2.2 Statistical methods

Sample characteristics were summarized using standard statistical methods (e.g., means, SDs, correlations) with t-tests and adjusted linear models used to compare characteristics of male and female participants above and below O3I values that were ultimately identified as primary cut points in the NLR-O3I curves (see below). Splines were fit using 3rd degree polynomials with knots at each decile in R (version 3.6.2; splines package). Unadjusted models used a linear model to predict NLR values by splines of O3I. Adjusted models accounted for sex, age, BMI and CRP values in the linear models accounting for potential non-linear relationships using splines. To identify significant changes in the NLR-O3I relationship, we used a “sliding O3I

window” approach. The width of each window was three O3I percentage points (e.g., 4% to 7%). By moving the window up by 0.1% increments and repeatedly testing for significant differences between the mean NLR in the lower vs. the upper half of the window, we sought to discover O3I cut points where the NLR-O3I relationship appeared to flatten. These would be O3I values above which the “effect” of an increase in O3I on NLR had little impact. We began with a window midpoint of O3I =2.6%, and we used adjusted linear models in R to test for upper vs lower half differences. We identified the first window which did not have a statistically significant difference in upper and lower mean NLR values. The midpoint of this window was chosen as the O3I cut point to be used in further analysis.

Statistical interactions with O3I values above and below the subsequently identified inflection points were tested by placing the interaction term in separate models predicting O3I, NLR, age, sex, BMI, and CRP. Pearson correlations were used to assess strength and direction of linear association between covariates and NLR. Statistical significance was set to 0.05 for all analyses and 95% confidence bands are provided where appropriate.

3. Results

The final dataset consisted of 28,871 healthy adults (**Figure 1**). The average age was 55.1±15.1 years, the average BMI was 27.6±5.4 kg/m², and there were slightly more females (51%) than males. Females were significantly younger and had slightly but significantly higher O3I and CRP levels and lower NLR, neutrophil count, and BMI than males (**Table 1**). When the model was also adjusted for age, BMI and CRP, small but significant sex differences remained with the exception of BMI (**Table 1**).

NLR was significantly ($p < 0.0001$) and inversely associated with O3I (**Figure 2A**). Pearson correlations (r) of other variables with NLR were $r = 0.06$ (CRP), $r = 0.14$ (age), $r = 0.01$ (BMI), $r = 0.65$ (neutrophil count), and $r = -0.24$ (lymphocyte count). Adjustment for age, sex, BMI, and CRP strengthened the relationship (**Figure 2B**). The O3I value where the relationship between O3I and NLR began to flatten (i.e., the cut point determined by the sliding window analysis described in Methods) was 6.6% although NLR continued to decline until O3I $> 8.5\%$ (**Figure 2B**). Below 6.6%, the curve was clearly steeper than it was above 6.6%. The O3I cut point of 6.6% was explored in more detail. An O3I $< 6.6\%$ was associated with significantly higher NLR, neutrophil number, lymphocyte number, CRP, BMI, and lower age (versus O3I $\geq 6.6\%$) (**Table 2**). These O3I-related differences persisted, except for lymphocyte number, when the model was adjusted for age, sex, BMI and CRP (**Table 2**).

When adults with inflammation (CRP $> 3\text{mg/L}$) were included ($n = 43,851$), NLR was significantly and inversely related with O3I before ($p < 0.0001$) and after adjusting for age, sex, BMI and CRP ($p < 0.0001$) (**Supplemental Figure 1**). As noted above, we chose to restrict our analysis to a healthy population (i.e., without evident inflammation) to establish EPA+DHA standards that could be used to develop DRIs.

4. Discussion and Conclusions

We report a significant inverse relationship between O3I and NLR. By excluding individuals with CRP $> 3\text{ mg/L}$, we could examine the O3I-NLR relationship in individuals without any evidence of tissue injury, infection, or systemic acute inflammation [57,58], i.e. a cohort suitable for deriving O3I standards on which Dietary Reference Intakes (DRIs) could be developed for planning and assessing EPA+DHA intakes of healthy people. NLR values measured

in this cohort (**Table 1**) were similar to those measured in healthy adults globally [8,24,59–63]. NLR is known to increase with age [9,32,60], obesity [13,60], CRP [64,65], and to sometimes differ by sex [32,59]. These same relationships were mostly observed here as well (data not reported), suggesting that the findings from this clinical laboratory cohort are generally representative.

Inflammation is a normal physiological response to infection and injury, but unrestrained inflammation can cause tissue damage leading to disease. As noted earlier, EPA and DHA serve as precursors for the synthesis of prostaglandins, leukotrienes, lipoxins and SPMs that can mediate immune responses [1,29,33,37,39–41]. SPMs derived from EPA and DHA protect tissues by limiting acute inflammatory responses and helping achieve homeostasis without immunosuppression [37,44,45]. In a randomized, double-blind, crossover trial, supplementation (10wk) with high doses of EPA or DHA (~8g/d) in individuals with abdominal obesity and low-grade inflammation (baseline CRP 3.3 ± 2.4 mg/L) separated by a 9-wk washout significantly increased blood EPA and DHA concentrations (vs corn oil control) [66]. DHA, but not EPA, supplementation decreased CRP concentrations (vs control) and the authors concluded EPA and DHA had more consistent effects on anti-inflammatory gene expression than pro-inflammatory gene expression [66]. Using pairwise and network meta-analyses of randomized controlled trials, DHA and EPA seem to have similar effects on CRP, IL-6, TNF- α , and adiponectin [67]. Although EPA and DHA trigger anti-inflammatory responses in vitro [68,69], the effects of long-chain omega-3 fatty acids on plasma CRP concentrations are mixed [70–73]. A threshold blood EPA+DHA concentration may be required to fuel sufficient SPM biosynthesis to achieve meaningful resolution of inflammation [37].

Omega-3 supplementation increases the EPA+DHA content of neutrophil membranes in vivo and inhibits the 5-lipoxygenase pathway and leukotriene B₄-mediated functions of neutrophils *in vitro* [38]. Omega-3 supplementation significantly increased plasma EPA+DHA levels (50-100%) and decreased concentrations of IL-6 (10-12% vs 36% increase in placebo group) and TNFα (0.2 to -2.3% vs 12% increase in placebo group) in healthy, middle-aged and older individuals [74]. Meta-analyses have confirmed this effect [72]. In a study of 8,237 participants living in Japan without a history of cardiovascular disease, eating fish >4 days per week was associated with significant lower NLR [75]. High omega-3 dietary intake was associated with a trend towards lower CRP (p=0.09) and NLR (p=0.17) in men with acute coronary syndrome [76]. A systematic review and meta-analysis of 18 randomized controlled trials found marine-derived omega-3 fatty acids lowered pro-inflammatory eicosanoid concentrations, e.g., in neutrophil leukotriene B₄ [77]. In the long-term, chronically lower levels of inflammation may have health benefits. For example, a pooled analysis of data from 17 prospective cohort studies involving 15,720 deaths among 43,466 individuals over a median of 16 years of follow-up found blood EPA+DHA levels were inversely associated with risk for death from all causes, CVD, cancer and other causes [49].

Despite the strength of the NLR-O3I relationship, the most obvious limitation of this study is that confounding metabolic events, dietary behaviors, smoking habits and/or physical activity levels associated with O3I may also affect innate-adaptive immune system balance. However, a nationally representative study of 9,427 individuals living in the US did not find significant differences in NLR with respect to sex, education, insurance status, or drinking habits [59]. Moreover, this cross-sectional population is similar to 2/3 of the US population who have CRP

199 <3mg/L [78] and the ~95% of adults living in the USA with circulating O3I percentages below 5%
200 [79,80].

201 In this cross-sectional study of healthy individuals, 81% of participants had O3I <6.6%.
202 Below this threshold, an increase in NLR, reflecting an increasingly imbalanced innate-adaptive
203 immune system, was observed. NLR continued to decrease with increasing O3I until O3I >8.5%,
204 similar to the target proposed for cardiovascular health, i.e. O3I >8% [46,81]. Clearly additional
205 evidence, including intervention studies, is needed to determine whether the NLR-O3I
206 relationship is causal or coincidental.

207 In conclusion, red blood cell EPA+DHA levels are significantly and inversely associated with
208 NLR, a biomarker of inflammation and innate-adaptive immune system balance. Based on our
209 observations in this healthy cohort, an O3I >6.6%, and possibly as high as 8.5%, is associated
210 with lower NLR values that are indicative of a quiescent, balanced innate-adaptive immune
211 system.

212

Acknowledgements: The authors wish to thank Steven Varvel, PhD for his help in acquiring and collating the dataset used in this study.

CRediT authorship contribution statement: **M.I. McBurney:** Conceptualization & analytical design. Writing -original draft, review & editing. **W.S. Harris:** Conceptualization & analytical design. Funding acquisition. Writing- review & editing. **N.L. Tintle:** Formal analysis. Writing- review & editing.

Disclosures: M.I. McBurney serves on the Board of Directors of the American Society for Nutrition and has or has held consulting agreements in the past 3 years with Council for Responsible Nutrition; Church & Dwight; DSM Nutritional Products; International Life Sciences Institute, North America; McCormick; OmegaQuant Analytics; PepsiCo; and VitaMe Technologies. W.S. Harris holds an interest in OmegaQuant Analytics, a lab that offers omega-3 blood testing; and is a member of the RB Schiff Science and Innovation Advisory Board. N.L. Tintle has no conflicts to disclose.

References

- [1] C.N. Serhan, Pro-resolving lipid mediators are leads for resolution physiology, *Nature*. 510 (2014) 92–101. <https://doi.org/10.1038/nature13479>.
- [2] P.C. Calder, Marine omega-3 fatty acids and inflammatory processes: Effects, mechanisms and clinical relevance, *Bioch. Biophys. Acta: Mol Cell Biol Lipids*. 1851 (2015) 469–484. <https://doi.org/10.1016/j.bbalip.2014.08.010>.
- [3] Childs, Calder, Miles, Diet and Immune Function, *Nutrients*. 11 (2019) 1933. <https://doi.org/10.3390/nu11081933>.
- [4] D. Furman, J. Campisi, E. Verdin, P. Carrera-Bastos, S. Targ, C. Franceschi, L. Ferrucci, D.W. Gilroy, A. Fasano, G.W. Miller, A.H. Miller, A. Mantovani, C.M. Weyand, N. Barzilai, J.J. Goronzy, T.A. Rando, R.B. Effros, A. Lucia, N. Kleinstreuer, G.M. Slavich, Chronic inflammation in the etiology of disease across the life span, *Nat Med*. 25 (2019) 1822–1832. <https://doi.org/10.1038/s41591-019-0675-0>.
- [5] F. Imtiaz, K. Shafique, S.S. Mirza, Z. Ayoob, P. Vart, S. Rao, Neutrophil lymphocyte ratio as a measure of systemic inflammation in prevalent chronic diseases in Asian population, *International Archives of Medicine*. 5 (2012) 2. <https://doi.org/10.1186/1755-7682-5-2>.
- [6] T. Bhat, S. Teli, J. Rijal, H. Bhat, M. Raza, G. Khoueiry, M. Meghani, M. Akhtar, Neutrophil to lymphocyte ratio and cardiovascular diseases: a review, *Expert Rev. Cardiovasc. Ther*. (2013) 6.
- [7] S. Socorro Faria, P.C. Fernandes Jr, M.J. Barbosa Silva, V.C. Lima, W. Fontes, R. Freitas-Junior, A.K. Eterovic, P. Forget, The neutrophil-to-lymphocyte ratio: a narrative review, *Ecancer*. 10 (2016). <https://doi.org/10.3332/ecancer.2016.702>.

- 249 [8] J. Fest, R. Ruiter, M.A. Ikram, T. Voortman, C.H.J. van Eijck, B.H. Stricker, Reference values
250 for white blood-cell-based inflammatory markers in the Rotterdam Study: a population-
251 based prospective cohort study, *Sci Rep.* 8 (2018) 10566. [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-018-28646-w)
252 018-28646-w.
- 253 [9] M. Song, B.I. Graubard, C.S. Rabkin, E.A. Engels, Neutrophil-to-lymphocyte ratio and
254 mortality in the United States general population, *Nature Res.* 11 (2021) 464.
255 <https://doi.org/10.1038/s41598-020-79431-7>.
- 256 [10] A.M. Paul, S.D. Mhatre, E. Cekanaviciute, A.-S. Schreurs, C.G.T. Tahimic, R.K. Globus, S.
257 Anand, B.E. Crucian, S. Bhattacharya, Neutrophil-to-Lymphocyte Ratio: A Biomarker to
258 Monitor the Immune Status of Astronauts, *Front. Immunol.* 11 (2020) 564950.
259 <https://doi.org/10.3389/fimmu.2020.564950>.
- 260 [11] R. Zahorec, Ratio of neutrophil to lymphocyte counts - rapid and simple parameter of
261 systemic inflammation and stress in critically ill, *Bratisl. Lek. Listy.* 102 (2001) 5–14.
- 262 [12] L. Venkatraghavan, T.P. Tan, J. Mehta, A. Arekapudi, A. Govindarajulu, E. Siu, Neutrophil
263 Lymphocyte Ratio as a predictor of systemic inflammation - A cross-sectional study in a
264 pre-admission setting., *F1000Res.* 4 (2015) 123.
265 <https://doi.org/10.12688/f1000research.6474.1>.
- 266 [13] E. Rodríguez-Rodríguez, A.M. López-Sobaler, R.M. Ortega, M.L. Delgado-Losada, A.M.
267 López-Parra, A. Aparicio, Association between Neutrophil-to-Lymphocyte Ratio with
268 Abdominal Obesity and Healthy Eating Index in a Representative Older Spanish Population,
269 *Nutrients.* 12 (2020) 855. <https://doi.org/10.3390/nu12030855>.

- [14] H. Wada, T. Dohi, K. Miyauchi, R. Nishio, M. Takeuchi, N. Takahashi, H. Endo, M. Ogita, H. Iwata, T. Kasai, S. Okazaki, K. Isoda, S. Suwa, H. Daida, Neutrophil to Lymphocyte Ratio and Long-Term Cardiovascular Outcomes in Coronary Artery Disease Patients with Low High-Sensitivity C-Reactive Protein Level, *Int. Heart J.* 61 (2020) 447–453. <https://doi.org/10.1536/ihj.19-543>.
- [15] B. Suh, D.W. Shin, H.-M. Kwon, J.M. Yun, H.-K. Yang, E. Ahn, H. Lee, J.H. Park, B. Cho, Elevated neutrophil to lymphocyte ratio and ischemic stroke risk in generally healthy adults, *PLoS ONE*. 12 (2017) e0183706. <https://doi.org/10.1371/journal.pone.0183706>.
- [16] T. Angkananard, T. Anothaisintawee, M. McEvoy, J. Attia, A. Thakkinstian, Neutrophil Lymphocyte Ratio and Cardiovascular Disease Risk: A Systematic Review and Meta-Analysis, *BioMed Research International*. 2018 (2018) 1–11. <https://doi.org/10.1155/2018/2703518>.
- [17] N.H. Adamstein, J.G. MacFadyen, L.M. Rose, R.J. Glynn, A.K. Dey, P. Libby, I.A. Tabas, N.N. Mehta, P.M. Ridker, The neutrophil–lymphocyte ratio and incident atherosclerotic events: analyses from five contemporary randomized trials, *European Heart Journal*. 42 (2021) 896–903. <https://doi.org/10.1093/eurheartj/ehaa1034>.
- [18] L. Zhan, Y. Liu, Y. Cheng, W. Guo, J. Yang, Predictive Value of Neutrophil/Lymphocyte Ratio (NLR) on Cardiovascular Events in Patients with COVID-19, *IJGM*. Volume 14 (2021) 3899–3907. <https://doi.org/10.2147/IJGM.S317380>.
- [19] S. Qun, Y. Wang, J. Chen, X. Huang, H. Guo, Z. Lu, J. Wang, C. Zheng, Y. Ma, Y. Zhu, D. Xia, Y. Wang, H. He, Y. Wang, M. Fei, Y. Yin, M. Zheng, Y. Xu, W. Ge, F. Hu, J. Zhou, Neutrophil-

to-Lymphocyte Ratios Are Closely Associated With the Severity and Course of Non-mild
COVID-19, *Front. Immunol.* 11 (2020) 2160. <https://doi.org/10.3389/fimmu.2020.02160>.

[20] S. Sun, X. Cai, H. Wang, G. He, Y. Lin, B. Lu, C. Chen, Y. Pan, X. Hu, Abnormalities of
peripheral blood system in patients with COVID-19 in Wenzhou, China, *Clinica Chimica
Acta.* 507 (2020) 174–180. <https://doi.org/10.1016/j.cca.2020.04.024>.

[21] M. Seyit, E. Avci, R. Nar, H. Senol, A. Yilmaz, M. Ozen, A. Oskay, H. Aybek, Neutrophil to
lymphocyte ratio, lymphocyte to monocyte ratio and platelet to lymphocyte ratio to
predict the severity of COVID-19, *The American Journal of Emergency Medicine.* 40 (2021)
110–114. <https://doi.org/10.1016/j.ajem.2020.11.058>.

[22] E. Vafadar Moradi, A. Teimouri, R. Rezaee, N. Morovatdar, M. Foroughian, P. Layegh, B.
Rezvani Kakhki, S.R. Ahmadi Koupaei, V. Ghorani, Increased age, neutrophil-to-lymphocyte
ratio (NLR) and white blood cells count are associated with higher COVID-19 mortality, *The
American Journal of Emergency Medicine.* 40 (2021) 11–14.
<https://doi.org/10.1016/j.ajem.2020.12.003>.

[23] M.G. Mazza, S. Lucchi, A.G.M. Tringali, A. Rossetti, E.R. Botti, M. Clerici,
Neutrophil/lymphocyte ratio and platelet/lymphocyte ratio in mood disorders: A meta-
analysis, *Progress in Neuro-Psychopharmacology and Biological Psychiatry.* 84 (2018) 229–
236. <https://doi.org/10.1016/j.pnpbp.2018.03.012>.

[24] J.-H. Liu, Y.-J. Zhang, Q.-H. Ma, H.-P. Sun, Y. Xu, C.-W. Pan, Elevated blood neutrophil to
lymphocyte ratio in older adults with cognitive impairment, *Archives of Gerontology and
Geriatrics.* 88 (2020) 104041. <https://doi.org/10.1016/j.archger.2020.104041>.

- [25] A.J. Templeton, M.G. McNamara, B. Šeruga, F.E. Vera-Badillo, P. Aneja, A. Ocaña, R. Leibowitz-Amit, G. Sonpavde, J.J. Knox, B. Tran, I.F. Tannock, E. Amir, Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Solid Tumors: A Systematic Review and Meta-Analysis, *JNCI: Journal of the National Cancer Institute*. 106 (2014). <https://doi.org/10.1093/jnci/dju124>.
- [26] Y.-F. Meng, Q. Pu, Q. Ma, W. Zhu, X.-Y. Li, Neutrophil/lymphocyte ratio as an inflammatory predictor of dry eye: A case-control study, *Therapeutics and Clinical Risk Management*. 17 (2021) 259–266.
- [27] R. Lu, W. Li, X. Wang, D. Shi, H. Meng, Elevated neutrophil-to-lymphocyte ratio but not platelet-to-lymphocyte ratio is associated with generalized aggressive periodontitis in a Chinese population, *J Periodontol*. 92 (2021) 507–513. <https://doi.org/10.1002/JPER.20-0282>.
- [28] Z. Jin, G. Cai, P. Zhang, X. Li, S. Yao, L. Zhuang, M. Ren, Q. Wang, X. Yu, The value of the neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio as complementary diagnostic tools in the diagnosis of rheumatoid arthritis: A multicenter retrospective study, *J Clin Lab Anal*. 35 (2021). <https://doi.org/10.1002/jcla.23569>.
- [29] L. Xia, W. Li, J. Zhai, C. Yan, J. Chen, H. Yang, Significance of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio and prognostic nutritional index for predicting clinical outcomes in T1–2 rectal cancer, *BMC Cancer*. 20 (2020) 208. <https://doi.org/10.1186/s12885-020-6698-6>.

- [30] L. Winter, L.A. Wong, G. Jerums, J. Seah, M. Clarke, S.M. Tan, M.T. Coughlan, R.J. MacIsaac, E.I. Ekinici, Use of Readily Accessible Inflammatory Markers to Predict Diabetic Kidney Disease, *Front. Endocrinol.* 9 (2018) 225. <https://doi.org/10.3389/fendo.2018.00225>.
- [31] S. Balta, S. Demirkol, M. Unlu, Z. Arslan, T. Celik, Neutrophil to lymphocyte ratio may be predict of mortality in all conditions, *Br J Cancer.* 109 (2013) 3125–3126. <https://doi.org/10.1038/bjc.2013.598>.
- [32] J. Fest, T.R. Ruiter, B. Groot Koerkamp, D. Rizopoulos, M.A. Ikram, C.H.J. van Eijck, B.H. Stricker, The neutrophil-to-lymphocyte ratio is associated with mortality in the general population: The Rotterdam Study, *Eur J Epidemiol.* 34 (2019) 463–470. <https://doi.org/10.1007/s10654-018-0472-y>.
- [33] D. Wu, E.D. Lewis, M. Pae, S.N. Meydani, Nutritional Modulation of Immune Function: Analysis of Evidence, Mechanisms, and Clinical Relevance, *Front. Immunol.* 9 (2019) 3160. <https://doi.org/10.3389/fimmu.2018.03160>.
- [34] P.C. Calder, A.C. Carr, A.F. Gombart, M. Eggersdorfer, Optimal Nutritional Status for a Well-Functioning Immune System Is an Important Factor to Protect against Viral Infections, *Nutrients.* 12 (2020) 1181. <https://doi.org/10.3390/nu12041181>.
- [35] A.F. Gombart, A. Pierre, S. Maggini, A Review of Micronutrients and the Immune System—Working in Harmony to Reduce the Risk of Infection, *Nutrients.* 12 (2020) 236. <https://doi.org/10.3390/nu12010236>.
- [36] C.A. Reider, R.-Y. Chung, P.P. Devarshi, R.W. Grant, S. Hazels Mitmesser, Inadequacy of Immune Health Nutrients: Intakes in US Adults, the 2005–2016 NHANES, *Nutrients.* 12 (2020) 1735. <https://doi.org/10.3390/nu12061735>.

- [37] P.C. Calder, Eicosapentaenoic and docosahexaenoic acid derived specialised pro-resolving mediators: Concentrations in humans and the effects of age, sex, disease and increased omega-3 fatty acid intake, *Biochimie*. 178 (2020) 105–123.
<https://doi.org/10.1016/j.biochi.2020.08.015>.
- [38] T.H. Lee, R.L. Hoover, J.D. Williams, R.I. Sperling, J. Ravalese III, B.W. Spur, D.R. Robinson, E.J. Corey, R.A. Lewis, K.F. Austen, Effect of dietary enrichment with eicosapentaenoic acid and docosahexaenoic acids on in vitro neutrophil and monocyte leukotriene generation and neutrophil function, *New England Journal of Medicine*. 312 (1985) 217–224.
- [39] P. Yaqoob, P.C. Calder, Fatty acids and immune function: new insights into mechanisms, *Br J Nutr*. 98 (2007) S41–S45. <https://doi.org/10.1017/S0007114507832995>.
- [40] E.A. Dennis, P.C. Norris, Eicosanoid storm in infection and inflammation, *Nat Rev Immunol*. 15 (2015) 511–523. <https://doi.org/10.1038/nri3859>.
- [41] U. Radzikowska, A.O. Rinaldi, Z. Çelebi Sözen, D. Karaguzel, M. Wojcik, K. Cypryk, M. Akdis, C.A. Akdis, M. Sokolowska, The Influence of Dietary Fatty Acids on Immune Responses, *Nutrients*. 11 (2019) 2990. <https://doi.org/10.3390/nu11122990>.
- [42] J. McDaniel, Regulation of Neutrophil Function by Marine n-3 Fatty Acids—A Mini Review, *Cell Biochem Biophys*. (2021). <https://doi.org/10.1007/s12013-021-01001-z>.
- [43] C.N. Serhan, N. Chiang, T.E. Van Dyke, Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators, *Nat Rev Immunol*. 8 (2008) 349–361.
<https://doi.org/10.1038/nri2294>.

- [44] J. Dalli, J.W. Winkler, R.A. Colas, H. Arnardottir, C.-Y.C. Cheng, N. Chiang, N.A. Petasis, C.N. Serhan, Resolvin D3 and Aspirin-Triggered Resolvin D3 Are Potent Immunoresolvents, *Chemistry & Biology*. 20 (2013) 188–201. <https://doi.org/10.1016/j.chembiol.2012.11.010>.
- [45] J.K. Colby, R.-E.E. Abdunour, H.P. Sham, J. Dalli, R.A. Colas, J.W. Winkler, J. Hellmann, B. Wong, Y. Cui, S. El-Chemaly, N.A. Petasis, M. Spite, C.N. Serhan, B.D. Levy, Resolvin D3 and Aspirin-Triggered Resolvin D3 Are Protective for Injured Epithelia, *The American Journal of Pathology*. 186 (2016) 1801–1813. <https://doi.org/10.1016/j.ajpath.2016.03.011>.
- [46] W.S. Harris, C. von Schacky, The Omega-3 Index: a new risk factor for death from coronary heart disease?, *Prev. Med.* 39 (2004) 212–220. <https://doi.org/10.1016/j.ypmed.2004.02.030>.
- [47] M.R. Flock, A.C. Skulas-Ray, W.S. Harris, T.D. Etherton, J.A. Fleming, P.M. Kris-Etherton, Determinants of erythrocyte omega-3 fatty acid content in response to fish oil supplementation: A dose–response randomized controlled trial, *J. Am. Heart Assoc.* 2 (2013) e000513. <https://doi.org/10.1161/JAHA.113.000513>.
- [48] R.E. Walker, K.H. Jackson, N.L. Tintle, G.C. Shearer, A. Bernasconi, S. Masson, R. Latini, B. Heydari, R.Y. Kwong, M. Flock, P.M. Kris-Etherton, A. Hedengran, R.M. Carney, A. Skulas-Ray, S.S. Gidding, A. Dewell, C.D. Gardner, S.M. Grenon, B. Sarter, J.W. Newman, T.L. Pedersen, M.K. Larson, W.S. Harris, Predicting the effects of supplemental EPA and DHA on the omega-3 index, *Am. J. Clin. Nutr.* 110 (2019) 1034–1040. <https://doi.org/10.1093/ajcn/nqz161>.
- [49] W.S. Harris, N.L. Tintle, F. Imamura, F. Qian, A.V. Ardisson Korat, M. Marklund, L. Djoussé, J.K. Bassett, P.-H. Carmichael, Y.-Y. Chen, Y. Hirakawa, L.K. Kupers, M. Lankinen, R.A.

Murphy, C. Samieri, M.K. Senn, P. Shi, J.K. Virtanen, I.A. Brouwer, K.-L. Chien, G.
 Eiriksdottir, N.G. Forouhi, J.M. Geleijnse, G.G. Giles, V. Gudnason, C. Helmer, A. Hodge, R.
 Jackson, K.-T. Khaw, M. Laakso, H. Lai, D. Laurin, J. Lindsay, R. Micha, J. Mursu, T.
 Ninomiya, W. Post, B.M. Psaty, U. Riserus, J.G. Robinson, A.H. Shadyab, L. Snetselaar, A.
 Sala-Vila, Y. Sun, L.M. Steffen, M.Y. Tsai, N.J. Wareham, A.C. Wood, J.H.Y. Wu, F. Hu, Q.
 Sun, D.S. Siscovick, R.N. Lemaitre, D. Mozaffarian, F. The Fatty Acids and Outcomes
 Research Consortium, Blood n-3 fatty acid levels and total and cause-specific mortality
 from 17 prospective studies, *Nature Commun.* 12 (2021) 1–9.
<https://doi.org/10.1038/s41467-021-22370-2>.
 [50] S. Kim, M. Eliot, D.C. Koestler, W.-C. Wu, K.T. Kelsey, Association of Neutrophil-to-
 Lymphocyte Ratio With Mortality and Cardiovascular Disease in the Jackson Heart Study
 and Modification by the Duffy Antigen Variant, (2018) 8.
 [51] M.A. Zimmerman, Diagnostic Implications of C-Reactive Protein, *Arch Surg.* 138 (2003)
 220. <https://doi.org/10.1001/archsurg.138.2.220>.
 [52] A. Villaseñor, S.W. Flatt, C. Marinac, L. Natarajan, J.P. Pierce, R.E. Patterson, Postdiagnosis
 C-Reactive Protein and Breast Cancer Survivorship: Findings from the WHEL Study, *Cancer*
Epidemiol Biomarkers Prev. 23 (2014) 189–199. [https://doi.org/10.1158/1055-9965.EPI-](https://doi.org/10.1158/1055-9965.EPI-13-0852)
 13-0852.
 [53] M.L. Knight, The Application of High-Sensitivity C-Reactive Protein in Clinical Practice: A
 2015 Update, *US Pharmacist.* 40 (2015) 50–53.
 [54] S. Jafarnejad, V. Boccardi, B. Hosseini, M. Taghizadeh, Z. Hamedifard, A Meta-analysis of
 Randomized Control Trials: The Impact of Vitamin C Supplementation on Serum CRP and

Serum hs-CRP Concentrations, (2018).

<https://doi.org/10.2174/1381612824666181017101810>.

[55] T.D. Dayspring, S.A. Varvel, L. Ghaedi, D.L. Thiselton, J. Bruton, J.P. McConnell, Biomarkers of cholesterol homeostasis in a clinical laboratory database sample comprising 667,718 patients, *Journal of Clinical Lipidology*. 9 (2015) 807–816.

<https://doi.org/10.1016/j.jacl.2015.08.003>.

[56] W.S. Harris, J.V. Pottala, R.S. Vasan, M.G. Larson, S.J. Robins, Changes in erythrocyte membrane trans and marine fatty acids between 1999 and 2006 in older Americans, *J. Nutr.* 142 (2012) 1297–1303. <https://doi.org/10.3945/jn.112.158295>.

[57] WHO, C-reactive protein concentrations as a marker of inflammation or infection for interpreting biomarkers of micronutrient status., World Health Organization, Geneva, Switzerland, 2014.

http://apps.who.int/iris/bitstream/10665/133708/1/WHO_NMH_NHD_EPG_14.7_eng.pdf?ua=1.

[58] M. Antonelli, I. Kushner, It's time to redefine inflammation, *FASEB J.* 31 (2017) 1787–1791. <https://doi.org/10.1096/fj.201601326R>.

[59] B. Azab, M. Camacho-Rivera, E. Taioli, Average Values and Racial Differences of Neutrophil Lymphocyte Ratio among a Nationally Representative Sample of United States Subjects, *PLoS ONE*. 9 (2014) e112361. <https://doi.org/10.1371/journal.pone.0112361>.

[60] J. Li, Q. Chen, X. Luo, J. Hong, K. Pan, X. Lin, X. Liu, L. Zhou, H. Wang, Y. Xu, H. Li, C. Duan, Neutrophil-to-Lymphocyte Ratio Positively Correlates to Age in Healthy Population: NLR

Correlates to Age, *J. Clin. Lab. Anal.* 29 (2015) 437–443.

<https://doi.org/10.1002/jcla.21791>.

[61] Y. Chen, Y. Zhang, G. Zhao, C. Chen, P. Yang, S. Ye, X. Tan, Difference in Leukocyte

Composition between Women before and after Menopausal Age, and Distinct Sexual

Dimorphism, *PLoS ONE*. 11 (2016) e0162953.

<https://doi.org/10.1371/journal.pone.0162953>.

[62] P. Forget, C. Khalifa, J.-P. Defour, D. Latinne, M.-C. Van Pel, M. De Kock, What is the

normal value of the neutrophil-to-lymphocyte ratio?, *BMC Res Notes*. 10 (2017) 12.

<https://doi.org/10.1186/s13104-016-2335-5>.

[63] J.S. Lee, N.Y. Kim, S.H. Na, Y.H. Youn, C.S. Shin, Reference values of neutrophil-lymphocyte

ratio, lymphocyte-monocyte ratio, platelet-lymphocyte ratio, and mean platelet volume in

healthy adults in South Korea, *Medicine*. 97 (2018) e11138.

<https://doi.org/10.1097/MD.00000000000011138>.

[64] C.P. de Jager, P.T. van Wijk, R.B. Mathoera, J. de Jongh-Leuvenink, T. van der Poll, P.C.

Wever, Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia

better than conventional infection markers in an emergency care unit, *Crit Care*. 14 (2010)

R192. <https://doi.org/10.1186/cc9309>.

[65] J.C. Yombi, P.E. Schwab, E. Thienpont, Neutrophil-to-lymphocyte ratio (NLR) distribution

shows a better kinetic pattern than C-reactive protein distribution for the follow-up of

early inflammation after total knee arthroplasty, *Knee Surg Sports Traumatol Arthrosc*. 24

(2016) 3287–3292. <https://doi.org/10.1007/s00167-015-3921-0>.

- [66] C. Vors, J. Allaire, J. Marin, M.-C. Lépine, A. Charest, A. Tchernof, P. Couture, B. Lamarche, Inflammatory gene expression in whole blood cells after EPA vs. DHA supplementation: Results from the ComparED study, *Atherosclerosis*. 257 (2017) 116–122. <https://doi.org/10.1016/j.atherosclerosis.2017.01.025>.
- [67] C. Vors, J. Allaire, S.B. Mejia, T.A. Khan, J.L. Sievenpiper, B. Lamarche, Comparing the Effects of Docosahexaenoic and Eicosapentaenoic Acids on Inflammation Markers Using Pairwise and Network Meta-Analyses of Randomized Controlled Trials, *Advances in Nutrition*. 12 (2021) 128–140. <https://doi.org/10.1093/advances/nmaa086>.
- [68] S. Weldon, A. Mullen, C. Loscher, L. Hurley, H. Roche, Docosahexaenoic acid induces an anti-inflammatory profile in lipopolysaccharide-stimulated human THP-1 macrophages more effectively than eicosapentaenoic acid, *The Journal of Nutritional Biochemistry*. 18 (2007) 250–258. <https://doi.org/10.1016/j.jnutbio.2006.04.003>.
- [69] R. De Caterina, J.K. Liao, P. Libby, Fatty acid modulation of endothelial activation, *The American Journal of Clinical Nutrition*. 71 (2000) 213S–223S. <https://doi.org/10.1093/ajcn/71.1.213S>.
- [70] E.M. Balk, A.H. Lichtenstein, M. Chung, B. Kupelnick, P. Chew, J. Lau, Effects of omega-3 fatty acids on serum markers of cardiovascular disease risk: A systematic review, *Atherosclerosis*. 189 (2006) 19–30. <https://doi.org/10.1016/j.atherosclerosis.2006.02.012>.
- [71] W. Xin, W. Wei, X. Li, Effects of fish oil supplementation on inflammatory markers in chronic heart failure: a meta-analysis of randomized controlled trials, *BMC Cardiovascular Disord*. 12 (2012) 77. <https://doi.org/10.1186/1471-2261-12-77>.

- [72] K. Li, T. Huang, J. Zheng, K. Wu, D. Li, Effect of Marine-Derived n-3 Polyunsaturated Fatty Acids on C-Reactive Protein, Interleukin 6 and Tumor Necrosis Factor α : A Meta-Analysis, PLoS ONE. 9 (2014) e88103. <https://doi.org/10.1371/journal.pone.0088103>.
- [73] S. AbuMweis, S. Jew, R. Tayyem, L. Agraib, Eicosapentaenoic acid and docosahexaenoic acid containing supplements modulate risk factors for cardiovascular disease: a meta-analysis of randomised placebo-control human clinical trials, Journal of Human Nutrition and Dietetics. (2017). <https://doi.org/10.1111/jhn.12493>.
- [74] J.K. Kiecolt-Glaser, M.A. Belury, R. Andridge, W.B. Malarkey, B.S. Hwang, R. Glaser, Omega-3 supplementation lowers inflammation in healthy middle-aged and older adults: A randomized controlled trial | Elsevier Enhanced Reader, Brain, Behav. Immun. 26 (2012) 988–995. <https://doi.org/10.1016/j.bbi.2012.05.011>.
- [75] S. Tani, R. Matsuo, W. Atsumi, K. Kawauchi, T. Ashida, T. Yagi, K. Imatake, Y. Suzuki, A. Takahashi, N. Matsumoto, Y. Okumura, Higher Frequency of Fish Intake May Be Associated with a Lower Neutrophil/Lymphocyte Ratio: Anti-Atherosclerotic Effects of Fish Consumption, Ann Nutr Metab. (2021) 1–8. <https://doi.org/10.1159/000515915>.
- [76] A. Sut, K. Chiżyński, M. Różalski, J. Golański, Dietary intake of omega fatty acids and polyphenols and its relationship with the levels of inflammatory markers in men with chronic coronary syndrome after percutaneous coronary intervention, Kardiologia Polska. 78 (2020) 117–123. <https://doi.org/10.33963/KP.15078>.
- [77] J. Jiang, K. Li, F. Wang, B. Yang, Y. Fu, J. Zheng, D. Li, Effect of Marine-Derived n-3 Polyunsaturated Fatty Acids on Major Eicosanoids: A Systematic Review and Meta-Analysis

from 18 Randomized Controlled Trials, PLoS One. 11 (2016).

<https://doi.org/10.1371/journal.pone.0147351>.

[78] I. Kushner, D. Rzewnicki, D. Samols, What Does Minor Elevation of C-Reactive Protein Signify?, The American Journal of Medicine. 119 (2006) 166.e17-166.e28.

<https://doi.org/10.1016/j.amjmed.2005.06.057>.

[79] R. Murphy, E. Yu, E. Ciappio, S. Mehta, M. McBurney, Suboptimal Plasma Long Chain n-3 Concentrations are Common among Adults in the United States, NHANES 2003–2004, Nutrients. 7 (2015) 10282–10289. <https://doi.org/10.3390/nu7125534>.

[80] R.A. Murphy, P.P. Devarshi, S. Ekimura, K. Marshall, S. Hazels Mitmesser, Long-chain omega-3 fatty acid serum concentrations across life stages in the USA: an analysis of NHANES 2011–2012, BMJ Open. 11 (2021) e043301. <https://doi.org/10.1136/bmjopen-2020-043301>.

[81] W.S. Harris, The Omega-3 Index: Clinical Utility for Therapeutic Intervention, Curr Cardiol Rep. 12 (2010) 503–508. <https://doi.org/10.1007/s11886-010-0141-6>.

Table 1. Characteristics of the study population. Mean±SD				
Variable ¹	All (n=28,871)	Males (n=14,211)	Females (n=14,660)	Male vs. Female Adjusted ² p-value
O3I (%)	4.97±1.84	4.9±1.82	5.05±1.85	0.016
NLR	2.09±1.10	2.13±1.13	2.05±1.07	<0.0001
Neutrophil (10 ³ /μL)	3.41 ±1.4	3.46±1.4	3.37±1.39	0.0002
Lymphocyte (10 ³ /μL)	1.81±1.47	1.82±1.77	1.80±1.10	<0.0001
CRP (mg/L)	1.23±0.77	1.21±0.75	1.25±0.79	0.0001
BMI (kg/m ²)	27.6±5.4	28.8±4.9	26.6±5.6	0.54
Age (years)	55.1±15.1	55.3±14.7	54.9±15.5	<0.001
¹ O3I, Omega-3 index; NLR, neutrophil-lymphocyte ratio; CRP, high-sensitivity C-reactive protein; BMI, body mass index. ² Models were adjusted for age, BMI, and CRP (except when the model was predicting these variables).				

518

519

520

Table 2. Stratification by omega-3 index (O3I) classification (n=28,871). Mean±SD			
Variable ¹	O3I <6.6% (n=23,510)	O3I ≥6.6% (n=5,361)	Adjusted ² p-value
O3I (%)	4.28±1.11	8.03±1.16	<0.0001
NLR	2.12±1.11	1.96±1.07	<0.0001
Neutrophil (10 ³ /μL)	3.48±1.40	3.11±1.35	<0.0001
Lymphocyte (10 ³ /μL)	1.82±1.56	1.76±0.95	0.591
CRP (mg/L)	1.27±0.78	1.07±0.73	<0.0001
BMI (kg/m ²)	28.0±5.4	26.3±4.9	<0.0001
Age (years)	54.1±15.4	59.4±13.1	<0.0001
¹ O3I, Omega-3 index; NLR, neutrophil-lymphocyte ratio; CRP, high-sensitivity C-reactive protein; BMI, body mass index.			
² Models were adjusted for sex, age, BMI, and CRP (except when the model was predicting these variables).			

521

522

523

524 Figure 1. Analytical sample flow. NLR, neutrophil-lymphocyte ratio; O3I, omega-3 index; CRP,
525 high-sensitivity C-reactive protein; BMI, body mass index.

526

527 Figure 2. The unadjusted (A) and adjusted for sex, age, BMI, and CRP (B) relationship between
528 the neutrophil-lymphocyte ration (NLR) and the omega-3 index (O3I) in 28,871 healthy adults.
529 (Means and 95% confidence bands).