

Association of Urinary Sodium Excretion with the Incidence of End-Stage Kidney Disease and Kidney-Related Death: Results from the UK Biobank

Ying Shan, Yong Bai, Jingwen Zhang, Yueqi Lu, Sike Yu, Congying Song, Juehan Liu, Min Jian, Junjie Xu, Zuying Xiong, and Xiaoyan Huang

Supplemental Table of Contents

Figure S1. Histograms of 24-h urinary potassium excretion calculated by the INTERSALT (a) and Kawasaki (b) equations.

Vertical dashed lines indicate the upper limit of daily sodium intake (2 g/d) recommended by the World Health Organization.

Table S1. Incidence rates and adjusted hazard ratios for primary and secondary outcomes according to 24-h urinary sodium excretion with subdistribution hazards models.

Table S2. Adjusted hazard ratios for primary outcomes according to 24-h urinary sodium excretion under 3 sensitivity analyses.




Figure S1. Histograms of 24-h urinary potassium excretion calculated by the INTERSALT (a) and Kawasaki (b) equations. Vertical dashed lines indicate the upper limit of daily sodium intake (2 g/d) recommended by the World Health Organization.

Table S1. Incidence rates and adjusted hazard ratios for primary and secondary outcomes according to 24-h urinary sodium excretion with subdistribution hazards models.

	Crude		Model 1		Model 2		Model 3	
	HRs (95% CIs)	P	HRs (95% CIs)	P	HRs (95% CIs)	P	HRs (95% CIs)	P
<i>Composite of end-stage kidney disease and kidney-related death</i>								
Continuous, per 1 g/d increment	1.67 (1.55, 1.79)	<0.001	1.43 (1.26, 1.62)	<0.001	1.03 (0.91, 1.16)	0.66	0.96 (0.84, 1.10)	0.60
Binary								
Below 2 g/d	Reference		Reference		Reference		Reference	
Above 2 g/d	0.78 (0.55, 1.10)	0.15	0.79 (0.53, 1.18)	0.25	0.84 (0.56, 1.27)	0.40	0.76 (0.49, 1.18)	0.23
Multicategory								
Quartile 1	Reference		Reference		Reference		Reference	
Quartile 2	0.97 (0.80, 1.18)	0.79	1.31 (1.02, 1.66)	0.03	1.03 (0.80, 1.31)	0.84	1.08 (0.83, 1.42)	0.56
Quartile 3	1.51 (1.27, 1.80)	<0.001	1.38 (1.01, 1.88)	0.04	1.04 (0.80, 1.36)	0.76	1.06 (0.79, 1.43)	0.71
Quartile 4	2.60 (2.22, 3.06)	<0.001	1.87 (1.36, 2.58)	<0.001	1.09 (0.82, 1.46)	0.56	1.01 (0.72, 1.41)	0.96
P for linear trend	<0.001		<0.001		0.58		>0.99	
P for quadratic trend	<0.001		0.79		0.89		0.42	
<i>End-stage kidney disease</i>								
Continuous, per 1 g/d increment	1.57 (1.40, 1.75)	<0.001	1.31 (1.08, 1.58)	0.006	1.07 (0.88, 1.30)	0.48	0.93 (0.75, 1.16)	0.54
Binary								
Below 2 g/d	Reference		Reference		Reference		Reference	
Above 2 g/d	1.63 (0.73, 3.65)	0.23	1.83 (0.67, 4.98)	0.24	2.06 (0.74, 5.75)	0.17	1.68 (0.60, 4.71)	0.32
Multicategory								
Quartile 1	Reference		Reference		Reference		Reference	
Quartile 2	0.92 (0.67, 1.27)	0.63	1.12 (0.77, 1.64)	0.55	0.92 (0.62, 1.38)	0.69	1.02 (0.67, 1.55)	0.93
Quartile 3	1.63 (1.23, 2.16)	<0.001	1.25 (0.82, 1.89)	0.30	1.10 (0.73, 1.66)	0.63	1.03 (0.65, 1.62)	0.91
Quartile 4	2.25 (1.74, 2.94)	<0.001	1.38 (0.90, 2.12)	0.14	1.08 (0.68, 1.72)	0.73	0.88 (0.53, 1.47)	0.63

P for linear trend	<0.001		0.15		0.59		0.68	
P for quadratic trend	0.04		0.95		0.80		0.50	
<i>Kidney-related death</i>								
Continuous, per 1 g/d increment	1.74 (1.60, 1.88)	<0.001	1.52 (1.31, 1.76)	<0.001	1.00 (0.87, 1.15)	>0.99	0.96 (0.82, 1.12)	0.57
Binary								
Below 2 g/d	Reference		Reference		Reference		Reference	
Above 2 g/d	0.67 (0.47, 0.97)	0.03	0.67 (0.44, 1.02)	0.06	0.67 (0.44, 1.04)	0.08	0.63 (0.39, 1.00)	0.05
Multicategory								
Quartile 1	Reference		Reference		Reference		Reference	
Quartile 2	0.97 (0.77, 1.22)	0.80	1.37 (1.03, 1.83)	0.03	1.03 (0.77, 1.38)	0.84	1.12 (0.82, 1.55)	0.47
Quartile 3	1.55 (1.26, 1.90)	<0.001	1.57 (1.06, 2.32)	0.02	1.09 (0.79, 1.50)	0.59	1.17 (0.81, 1.68)	0.40
Quartile 4	2.79 (2.32, 3.36)	<0.001	2.19 (1.46, 3.29)	<0.001	1.07 (0.75, 1.52)	0.71	1.05 (0.71, 1.57)	0.79
P for linear trend	<0.001		<0.001		0.67		0.78	
P for quadratic trend	<0.001		0.90		0.78		0.25	

Model 1 was adjusted for age, sex, Townsend deprivation index, education, ethnicity, smoking status, alcohol consumption, metabolic equivalents, and 24h urinary potassium excretion.

Model 2 was adjusted for age, sex, Townsend deprivation index, education, ethnicity, smoking status, alcohol consumption, metabolic equivalents, 24-h urinary potassium excretion, waist circumference, hypertension, diabetes, cardiovascular disease, stroke, diuretics, angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers, and estimated glomerular filtration rate.

Model 3 was adjusted for age, sex, Townsend deprivation index, education, ethnicity, smoking status, alcohol consumption, metabolic equivalents, 24-h urinary potassium excretion, waist circumference, hypertension, diabetes, cardiovascular disease, stroke, diuretics, angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers, estimated glomerular filtration rate, systolic blood pressure, diastolic blood pressure, and urine albumin-to-creatinine ratio.

Abbreviations: CIs, confidence intervals; HRs, hazard ratios.

Table S2. Adjusted hazard ratios for primary outcomes according to 24-h urinary sodium excretion under 3 sensitivity analyses.

Scenarios	Sensitivity analysis 1		Sensitivity analysis 2		Sensitivity analysis 3	
	HRs (95% CIs)	P	HRs (95% CIs)	P	HRs (95% CIs)	P
Excluded participants who took diuretics, angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers, or glucocorticoids at baseline	360,549		440,003		444,328	
Excluded participants who had prevalent congestive heart failure or CKD stage 4-5 at baseline, or developed ESKD or died within 2 years of follow-up					Used the exposure of 24-h urinary sodium excretion calculated by the Kawasaki equations	
Participants included, n						
Continuous, per 1 g/d increment	0.92 (0.75, 1.14)	0.44	0.97 (0.84, 1.11)	0.62	1.03 (0.98, 1.10)	0.25
Binary						
Below 2 g/d	Reference		Reference		Reference	
Above 2 g/d	0.71 (0.40, 1.25)	0.23	0.72 (0.47, 1.10)	0.13	1.22 (0.89, 1.67)	0.23
Multicategory						
Quartile 1	Reference		Reference		Reference	
Quartile 2	0.90 (0.60, 1.36)	0.62	0.99 (0.74, 1.32)	0.93	0.89 (0.73, 1.09)	0.25
Quartile 3	1.03 (0.65, 1.61)	0.91	0.97 (0.71, 1.33)	0.87	1.01 (0.83, 1.23)	0.93
Quartile 4	0.85 (0.51, 1.42)	0.54	0.97 (0.69, 1.38)	0.88	1.08 (0.88, 1.31)	0.47
P for linear trend	0.69		0.88		0.29	
P for quadratic trend	0.75		0.94		0.20	

Models for sensitivity analysis 1 were adjusted for age, sex, Townsend deprivation index, education, ethnicity, smoking status, alcohol consumption, metabolic equivalents, 24-h urinary potassium excretion, waist circumference, hypertension, diabetes, cardiovascular disease, stroke, and estimated glomerular filtration rate.

Models for sensitivity analysis 2 and 3 were adjusted for age, sex, Townsend deprivation index, education, ethnicity, smoking status, alcohol consumption, metabolic equivalents, 24-h urinary potassium excretion, waist circumference, hypertension, diabetes, cardiovascular disease, stroke, diuretics, angiotensin-converting enzyme inhibitors or angiotensin II receptor blockers, and estimated glomerular filtration rate.

Note that more participants were included in sensitivity analysis 3 than in the main analysis due to fewer participants with incomplete 24-h urinary sodium excretion data calculated by the Kawasaki equations.

Abbreviations: CIs, confidence intervals; CKD, chronic kidney disease; ESKD, end-stage kidney disease; HRs, hazard ratios.