

Supplementary material

Chronic kidney disease

To demonstrate the public health burden (rather than likely causal mechanisms), within patients with chronic kidney disease (CKD), Table S1 shows both crude and adjusted rates of positive SARS-CoV-2 swab test, COVID-19 related hospital admissions, COVID-19 related critical care admissions and COVID-19 related death, broken down by CKD stage. Incidence rates for each outcome were estimated by dividing the count by person-years, with 95% confidence intervals. Crude and adjusted rates were also calculated using cox regression and show that even after adjustment, rates remain very high. These results highlight that fully vaccinated individuals with kidney disease are at far higher risk of serious outcomes when infected with COVID-19 than individuals without kidney disease.

Table S1: Number of fully vaccinated (2 doses + 2 weeks) patients with chronic kidney disease by stage, in OpenSAFELY-TPP, and associated crude and adjusted rates of positive SARS-CoV-2 swab test, COVID-19 related hospital admissions, COVID-19 related critical care admissions and COVID-19 related death, broken down by CKD stage.

Population/Outcome		Kidney disease					
		No CKD	Stage 3a	Stage 3b	Stage 4	Stage 5	
Fully vaccinated	Count*	14622275	527255	205755	51035	5505	
	Follow-up time, medium days (IQR)	147 (103-177)	187 (173-201)	192 (178-205)	191 (174-204)	184 (164-200)	
	Tests conducted (%)	0	59	61	60	55	37
		1	17	14	13	12	12
		2	7	6	6	7	8
3+		17	18	21	26	44	
Positive SARS-CoV-2 test	Positivity rate, %	2.00	1.80	1.77	1.77	1.42	
	Events* (PYs)	557715 (5474016)	12135 (267915)	4440 (106744)	1275 (25816)	215 (2607)	
	Incidence rate per 10,000 person-years (95% CI)	101.88 (101.75-102.02)	45.29 (44.88-45.7)	41.59 (40.97-42.21)	49.35 (47.99-50.75)	82.86 (77.41-88.69)	
	Hazard ratio	1.00	0.41 (0.40-0.41)	0.37 (0.36-0.38)	0.44 (0.42-0.47)	0.76 (0.66-0.86)	
	Age-adjusted hazard ratio	1.00	0.79 (0.77-0.80)	0.80 (0.78-0.83)	0.95 (0.90-1.00)	1.29 (1.13-1.48)	
COVID-19 related hospital admission	Events* (PYs)	11865 (5513255)	2040 (267918)	1270 (106641)	465 (25788)	80 (2615)	
	Incidence rate per 10,000 person-years (95% CI)	2.15 (2.13-2.17)	7.61 (7.44-7.78)	11.89 (11.56-12.23)	18.11 (17.29-18.97)	30.97 (27.72-34.61)	

	Hazard ratio	1.00	3.00 (2.86-3.14)	4.61 (4.35-4.88)	7.08 (6.45-7.77)	12.47 (10.02-15.52)
	Age-adjusted hazard ratio	1.00	1.84 (1.75-1.93)	2.56 (2.41-2.72)	3.88 (3.53-4.27)	8.01 (6.43-9.97)
COVID-19 related critical care admission	Events* (PYs)	815 (5514965)	105 (268206)	55 (106814)	25 (25846)	10 (2625)
	Incidence rate per 10,000 person-years (95% CI)	0.15 (0.14-0.15)	0.4 (0.36-0.44)	0.51 (0.45-0.59)	1.04 (0.86-1.27)	3.43 (2.46-4.78)
	Hazard ratio	1.00	2.45 (2.00-3.00)	3.21 (2.44-4.23)	6.57 (4.47-9.64)	21.81 (11.30-42.07)
	Age-adjusted hazard ratio	1.00	1.92 (1.55-2.38)	2.40 (1.55-2.38)	4.88 (3.29-7.23)	17.52 (9.05-33.90)
COVID-19 related death	Events* (PYs)	2370 (5554595)	690 (269616)	500 (107352)	215 (25976)	30 (2639)
	Incidence rate per 10,000 person-years (95% CI)	0.43 (0.42-0.44)	2.55 (2.46-2.65)	4.67 (4.46-4.88)	8.28 (7.73-8.86)	11.37 (9.47-13.65)
	Hazard ratio	1.00	4.23 (3.88-4.61)	7.36 (6.68-8.12)	13.27 (11.53-15.28)	19.53 (13.62-28.00)
	Age-adjusted hazard ratio	1.00	1.68 (1.54-1.84)	2.33 (2.10-2.58)	3.98 (3.45-4.60)	8.08 (5.63-11.60)