Decreased retinal vascular complexity is an early biomarker of MI supported by a shared genetic control.

Ana Villaplana-Velasco¹², Justin Engelmann², Konrad Rawlik¹, Oriol Canela-Xandri³, Claire Tochel², Frida Lona-Durazo⁴, Muthu Rama Krishnan Mookiah⁵, Alex Doney⁵, Esteban J. Parra⁴, Emanuele Trucco⁵, Tom MacGillivray⁶, Kristiina Rannikmae², Albert Tenesa¹²³, Erola Pairo-Castineira^{1*}, and Miguel O. Bernabeu^{2,7*}.

¹The Roslin Institute, Royal (Dick) School of Veterinary Studies, The University of Edinburgh

²Centre for Medical Informatics, Usher Institute, The University of Edinburgh

³MRC Human Genetics Unit, IGC, The University of Edinburgh

⁴University of Toronto at Mississauga, Mississauga, Ontario, Canada

⁵VAMPIRE project, Computing, School of Science and Engineering, University of Dundee

⁶VAMPIRE project, Centre for Clinical Brain Sciences, The University of Edinburgh

⁷The Bayes Centre, The University of Edinburgh

*Equally contributing authors.

Abstract

There is increasing evidence that the complexity of the retinal vasculature (measured as fractal dimension, D_f) might offer earlier insights into the progression of coronary artery disease (CAD) before traditional biomarkers can be detected. This association could be partly explained by a common genetic basis; however, the genetic component of D_f is poorly understood. We present here a genome-wide association study (GWAS) aimed to elucidate the genetic component of D_f and to analyse its relationship with CAD. To this end, we obtained D_f from retinal fundus images and genotyping information from ~38,000 white-British participants in the UK Biobank. We discovered 9 loci associated with D_f , previously reported in pigmentation, retinal width and tortuosity, hypertension, and CAD studies. Significant negative genetic correlation estimates endorse the inverse relationship between D_f and CAD, and between D_f and myocardial infarction (MI), one of CAD fatal outcomes. This strong association motivated us to developing a MI predictive model combining clinical information, D_f , a CAD polygenic risk score and using a random forest algorithm. Internal cross validation evidenced a considerable improvement in the area under the curve (AUC) of our predictive model (AUC=0.770) when comparing with an established risk model, SCORE, (AUC=0.719). Our findings shed new light on the genetic basis of D_f , unveiling a common control with CAD, and highlights the benefits of its application in individualised MI risk prediction.

36 Introduction

37 Coronary artery disease (CAD) remains the leading cause of death and disability
 38 worldwide¹. Early-diagnosis and preventive therapies are essential strategies to control
 39 CAD morbidity and the mortality associated with its outcomes, such as myocardial
 40 infarction (MI). There is increasing evidence that morphological changes in the retinal
 41 vasculature, for instance in vessel width and vascular complexity, might offer insights
 42 into CAD before traditional risk factors (such as systolic blood pressure and cholesterol
 43 levels)^{2,3}. Recent studies reported that a reduced degree of vascular complexity,
 44 quantified through estimates of the fractal dimension (D_f), is found in individuals who had
 45 a higher CAD risk, independent of their age². This suggests that D_f could be a promising
 46 non-invasive and highly accessible biomarker. Nevertheless, this finding has not
 47 translated so far to a substantial increase in prediction accuracy for major adverse
 48 cardiac events (MACE) risk, compared to models based on patient demographics and
 49 lifestyle risk factors^{4,5}.

50 Evidence points towards coronary and retinal vessels experiencing similar
 51 pathophysiological changes at even early CAD stages⁶⁻⁸, plausibly influenced by a
 52 shared genetic basis^{7,9-14}. Population-based studies have demonstrated that both
 53 tortuosity and width of arteries and veins have a moderate genetic basis^{12,13}. Veluchamy
 54 *et al.* described two novel loci near the *COL4A2* and *ACTN4* genes associated with
 55 retinal tortuosity which were previously reported in genetic atrial fibrillation and CAD¹¹
 56 studies. However, the genetic component of retinal vascular complexity remains poorly
 57 understood.

58 We report here a genome-wide association study (GWAS) on the topic, using ~38,000
 59 white British participants from the UK Biobank. The aim is twofold: to comprehensively
 60 study the genetic control of D_f and to assess the extent of its relationship with CAD. We
 61 discovered 9 loci associated with D_f . Two of these loci (*SLC12A9* and *RDH5* genes) were
 62 previously associated with cardiovascular risk factors and diseases¹⁵. Genetic
 63 correlation estimates and enrichment analysis support the shared genetic control
 64 between D_f and CAD, suggesting that decreasing D_f might be influenced by clinical CAD
 65 manifestations and, in part, by common genetic effects. Given this strong connection, we
 66 developed a model to predict incident MI cases in the UK Biobank over the 10 years
 67 succeeding the ophthalmic examination at baseline, including D_f and a CAD polygenic
 68 risk score (PRS_{CAD}). Internal 10-fold cross validation shows a considerable performance
 69 improvement compared with the SCORE model¹⁶, an established CAD risk prediction
 70 score based on epidemiological variables. This enhancement can be partly explained

by the additional predictive power of retinal and genetic determinants, as these respectively capture early vascular morphological abnormalities and personalised MI risk. Our findings shed new light on the genetic component of D_f , suggesting a common genetic basis with CAD aetiology, and demonstrate its potential for individual MI risk prediction.

Results

Automated Quality Control and fractal dimension calculation in UK Biobank fundus images reveals interocular asymmetry in vascular complexity at an individual level.

For this study, we first developed a semi-automated pipeline to segment the vasculature and select good-quality segmentations in 175,611 fundus images available in the UK Biobank (Fig. 1a) using VAMPIRE software (version 3.1, Universities of Edinburgh and Dundee)^{17,18} and a previously published fundus image classifier¹⁹. An image quality score (IQS) was computed as part of the classification process (see Section “Methods”). D_f was subsequently calculated from binary vessel maps produced automatically by VAMPIRE for ~98,600 good quality images.

We completed the, to our knowledge, largest within individual interocular D_f comparison ($n=39,656$ participants) reported so far. The population median (1.492 ± 0.043) and D_f distributions appear identical between left and right eyes (Fig. 1b). However, their moderate correlation ($r^2=0.61$, $P\text{-value}=2 \cdot 10^{-16}$, Fig. 1d) and the significant difference between left and right D_f (paired T-test $P\text{-value}=1.59 \cdot 10^{-75}$) highlight an individual interocular asymmetry (Fig. 1c), where 50% of the individuals have a right D_f 1 SD unit larger than their respective left D_f . As shown in Fig. 1d, differences occur in both directions and are more pronounced when any of the D_f is lower than the median. To control for this individual asymmetrical effect (Fig. 1e), we performed further analysis in both eyes separately.

Figure 1. Pipeline and D_f characteristics. **a** Study design's diagram describing the stepwise development of this project. **b** Left and right D_f histogram. **c** Individual variation distribution between left and right D_f . **d** Overlapping left and right D_f histograms including the regression line. **e** Example of individual interocular asymmetry in UKBB fundus images.

We next fitted univariate linear models using D_f as dependent variable and estimated the Pearson correlation between D_f and 779 UKBB binary and quantitative traits (see methods below) and IQS. Amongst these 780 variables, IQS has the strongest effect ($\beta_{\text{right}} = 0.033$, $P\text{-value} < 10^{-300}$; $\beta_{\text{left}} = 0.024$, $P\text{-value} < 10^{-300}$; $r^2_{\text{right}} = 0.39$, $P\text{-value} < 10^{-300}$; $r^2_{\text{left}} = 0.36$, $P\text{-value} < 10^{-300}$). Supplementary Figure 1 illustrates this association and that a larger interocular IQS difference moderately affects D_f variation ($\beta = 0.014$, $P\text{-value} < 10^{-300}$). Hence, we account for IQS influence in our following analysis.

Besides IQS, 75 quantitative and 161 binary traits were significantly associated with D_f after Bonferroni correction²⁰ ($P\text{-value} < 0.05/780 = 6.41 \times 10^{-5}$). Age, sex, height, retinal disorders, smoking, hypertension, and CAD have the greatest significant effect on D_f in both eyes amongst all measurements (Supplementary Table 1).

GWAS reveals nine fractal dimension loci and their association with cardiovascular risk factors.

Here we present a GWAS on D_f . This was completed with 38,811 and 38,017 unrelated white-British UK Biobank participants that had a right and left D_f measure, respectively. After QC (see Section "Methods"), there were 9,275,849 imputed SNPs with $HWE > 10^{-6}$, $MAF > 5 \times 10^{-3}$, a call rate > 0.9 , and an imputation score > 0.9 . The GWAS model included

hair and skin colour to control for spurious associations given the influence of eye and skin colour on fundus colour^{21,22}. Hair colour replaced eye colour because the latter is not recorded during UKBB assessments. In addition, we completed a supplementary GWAS including an eye colour PRS based on the study by Lona-Durazo *et al.*²³, which indicated no eye colour effect in our GWAS results (see Section “Methods”). The quantile-quantile plot of both GWASs indicated an adequate control of the genomic inflation in our analysis ($\lambda_{GC} = 1.065$ and $\lambda_{GC} = 1.067$ in the right and left eye, respectively, see Supplementary Fig. 2). Fig. 2c illustrates the SNPs effects comparison between eyes GWAS studies, highlighting analogous results.

The strongest associations are located around *OCA2* (rs916977, P-value= $7.41 \cdot 10^{-48}$) and *HERC2* (rs12913832, P-value= $2.16 \cdot 10^{-96}$) genes in chromosome 15 (Fig. 2a, Fig. 2b and Supplementary Table 2). We observed another significant association near the *IRF4* gene (rs12203592, P-value= $6.59 \cdot 10^{-24}$). Phenome-wide association studies (PheWAS), using GeneATLAS²⁴ and GWASCatalog²⁵, have commonly reported these SNPs in skin, hair, and eye colour analyses. Recent ocular studies demonstrated their implication in lens disorders, cataract, glaucoma, visual acuity, and retinal venular and arteriolar width and tortuosity. Additionally, Zekavat *et al.*²⁶ recently published a paper completing a D_f GWAS with UKBB which reported similar results to our GWAS. However, we couldn't make a complete comparison between studies as they didn't make the summary statistics available yet. The comparison between reported variants of ²⁶ is in Supplementary Table 3.

In addition to these 3 genetic variants, we found 6 SNPs with a P-value $<10^{-05}$ which did not reach genomic-wide significance (Fig. 2d). The majority of these SNPs are located around genes previously reported in pigmentation analyses: *SLC45A2*, *SLC12A9*, *RDH5/ORMDL2* and *MAP1LC3A*, whereas those near *EIF2B5* and *AGPAT3* genes were described in blood content and inflammation GWASs. Those SNPs located close to the *SLC12A9*, *RDH5/ORMLD2* and *AGPAT3* genes have also a strong effect in multiple ocular traits and diseases (such as macular thickness and retinal detachment), hypertension, and arterial disorders.

The SNP heritability (h^2_{SNP}) of the left and right D_f estimate are respectively 0.09 (0.015) and 0.10 (0.014). These h^2_{SNP} magnitude is in line with previous results from retinal vascular tortuosity^{11,14}, retinal width¹², and the recently published D_f ²⁶ GWAS.

We completed additional D_f GWAS using independent UKBB participants with white European ($n_{left}=4340$ and $n_{right}=4288$), Asian ($n_{left}=562$ and $n_{right}=568$), and African ($n_{left}=498$ and $n_{right}=509$) ancestry. Only the GWAS including white European ancestry

replicated the effect of the strongest associations (P -value $<0.05/11=0.0045$), which can be explained by the considerably larger number of participants in this analysis when compared with Asian and African ancestries. Little heterogeneity and forest plots of D_f loci indicate that multiple significant genetic variants (rs16891982, rs12203592, rs12913832 and rs31381412) have similar effect across Asian, African, white-European, and white-British ancestries (Supplementary Fig 3).

Figure 2. GWAS of both eyes' D_f . Manhattan plot of **a** right and **b** left D_f . Points are truncated at $-\log_{10}(P)=50$ for clarity. **c** Comparison of the genetic variant effects between left and right D_f results. Colour depth indicates the significance of each variant (navy, violet, and purple for non-significant, close to genome-wide significance and significant, respectively). Genetic variants included are truncated at a minimum $-\log_{10}(p) = 3$ for clarity. **d** Summary statistics of D_f -associated SNPs and its nearest located gene.

Genetic correlation estimates and functional analysis indicate shared genetic control between fractal dimension and coronary artery disease.

To assess the link between D_f and CAD risk factors and outcomes, we calculated their genome-wide genetic correlation using LD score regression (LDSC)²⁷. Genetic correlation estimates (r_g^2) suggest a negative correlation between D_f and hypertension ($r_g^2=-0.30$, P-value= $4.52 \cdot 10^{-06}$), acute MI ($r_g^2=-0.16$, P-value=0.03), and CAD ($r_g^2=-0.18$, P-value=0.025) (Table 2). Atherosclerosis and D_f also have a negative r_g^2 , but significance is above the commonly accepted significance threshold (P-value<0.05). All these estimates agree in direction with phenotypic correlations (see Supplementary Table 1) and published studies, which reported that retinal D_f decreases as people develop these conditions^{2,3,6,28}. Therefore, our results suggest that these correlations of phenotypes are partly explained by its shared genetic basis.

Table 2. Genetic correlation estimates and significance (P-value) between D_f and associated cardiovascular events.

Correlated trait	LEFT EYE			RIGHT EYE		
	r_g^2	SE	P value	r_g^2	SE	P value
Hypertension	-0.2229	0.0534	3.0261e-05	-0.3020	0.0659	4.52e-06
Acute myocardial infarction	-0.1717	0.0809	0.0801	-0.1585	0.1075	0.0308
Self reported acute myocardial infarction	-0.2071	0.0754	0.006	-0.2663	0.0982	0.0067
Coronary artery disease	-0.2214	0.0591	1.7854e-04	-0.1776	0.0795	0.025
Atherosclerosis	-0.3585	0.1819	0.084	-0.2668	0.2100	0.051
Right Fractal dimension	0.9468	0.0962	1.75E-26	-		-

Moreover, we estimated the r_g^2 between pigmentation traits and D_f to examine the similarities among their genetic basis (Supplementary Fig 4). Although the estimates are non-significant ($r_g^2=-0.0751$, P-value=0.64), genomic regions around common genetic variants in these traits might have a significant value. We did not further pursue this hypothesis experimentally.

We investigated possible causal relationships between CAD, hypertension, MI and D_f . We found evidence of horizontal pleiotropy on the loci of interest (P=0.0056), suggesting that the link between D_f and such cardiovascular events may not be causal (Supplementary table 4).

FUMA functional analysis showed that D_f loci have a significant enrichment in melanin synthesis and pigmentation diseases as well as inflammation pathways. In addition, plasma activator-inhibitor levels type I, age-related macular degeneration, heart rate response and red blood cell count share significant overlapping gene sets with D_f (Supplementary Fig. 5).

Fractal dimension improves prediction of incident myocardial infarction in UK Biobank cases.

Given our findings, we hypothesized that D_f and PRS_{CAD} can provide additional information for MI risk estimation at an individual patient level. We thus developed a model to predict incident cases of MI over the 10 years succeeding the ophthalmic examination at baseline (Fig. 3a). Briefly, the model includes PRS_{CAD} derived from meta-analysis completed by the CARDIoGRAMplusC4D Consortium¹⁵, clinical variables from an established CAD risk assessment strategy named SCORE¹⁶ (age, sex, SBP and BMI), and the D_f of both eyes.

a MYOCARDIAL INFARCTION MODEL

Figure 3. Development and performance of MI predictive models. **a** Diagram illustrating the development of our MI model. **b** ROC curve of MI predictive models. **c** Kaplan-Meier curve of incident MI cases separated

203 by predicted MI probability.* D_f : fractal dimension; PRS_{CAD}: CAD polygenic risk score; BMI: Body-mass index; SBP:
204 Systolic blood pressure.

Model	MI MODEL		
	Precision $\pm$ SE	Recall $\pm$ SE	AUC $\pm$ SE
SCORE model ¹⁶	0.716 $\pm$ 0.0025	0.725 $\pm$ 0.0038	0.719 $\pm$ 0.0021
Random Forest including D_f	0.756 $\pm$ 0.0012	0.778 $\pm$ 0.0019	0.763 $\pm$ 0.0011*
Random Forest including PRS _{CAD}	0.735 $\pm$ 0.001	0.756 $\pm$ 0.0021	0.741 $\pm$ 0.0012*
Random Forest including D_f + PRS _{CAD}	0.763 $\pm$ 0.0016	0.788 $\pm$ 0.0014	0.770 $\pm$ 0.0013*

205 **Table 3.** Internal 10-fold cross validation of MI models evaluated with precision, recall and AUC.* AUC
206 estimates significantly different (Wilcoxon signed-rank test P-value<0.005) from the ones obtained with the SCORE
207 model. The obtained Wilcoxon signed-rank P-value for each model comparison is included in Supplementary Table 6.

208 The MI model was trained with the 526 individuals who experienced an MI event after
209 their UKBB ophthalmic examination. We created an age-matched control group with an
210 equal number of individuals and no underlying MI and CAD (Supplementary Table 5).
211 The mean age and its SD in case and control group is respectively 57.31 $\pm$ 6.47 and
212 54.21 $\pm$ 7.84 years. We chose the random forest classifier (RFC) as this method allows
213 one to model non-linear associations with the outcome and interactions between the
214 predictor variables, which might boost the prediction while being interpretable^{29,30}. We
215 also considered model versions excluding either PRS_{CAD} or D_f . As a baseline for
216 comparison, we retrained the original SCORE model¹⁶ on the same dataset. Internal 10-
217 fold cross validation (FCV) indicates that our models dominate the ROC curve of the
218 SCORE model, achieving a greater precision, recall, and AUC (Fig 3b and Table 3).
219 Amongst our considered models, only the one including PRS_{CAD} (AUC=0.741 $\pm$ 0.001)
220 yielded an AUC significantly different from the model introducing D_f (AUC=0.763 $\pm$ 0.001),
221 and the one combining both D_f and PRS_{CAD} (AUC=0.770 $\pm$ 0.001) (Table 3 and
222 Supplementary Table 6).

223 Next, we investigated potential survival rate differences between low and high MI risk
224 groups. These groups were defined by the predictions obtained with our top-performing
225 MI model and by subsequently separating these with a probability threshold of 0.5 (high
226 MI-risk>0.5 and low MI-risk= $\leq$ 0.5). The Kaplan-Meier curve (Fig. 3c) illustrates a
227 significant divergence between these groups (Log-rank test P=3.52 $\cdot$ 10⁻³⁰), which can be

explained by the pronounced decrease in survival during the first 4 years in the high MI risk group.

Finally, we performed an ablation study to understand the origin of the performance improvement in our new model. Briefly, we evaluated the performance of all possible variations between SCORE and the top-performing model (see Section “Methods”). This assessment revealed three key contributors to the reported improvement: the use of quantitative variables, the introduction of PRS_{CAD} and D_f, and the use of a random forest classifier. An extended discussion can be found in the Supplementary Material and Supplementary Table 7. The added predictive value of D_f is supported by the RFC development analysis which reveals that age, BMI and D_f are the most important features in its architecture (Supplementary Fig.6). PRS_{CAD} is also determinant to the model’s development as its RFC importance is equivalent to SBP and smoking taken together, which in line with recently published results⁸. Additional assessments suggest that the replacement of D_f measures with D_f adjusted by IQS or just the introduction of one-eye D_f measurements in our MI model yield a performance comparable with the aforementioned ones (Supplementary Table 8).

Discussion

This work provides a comprehensive examination of the genetic basis of D_f and unveiled a shared genetic basis with cardiovascular risk factors and CAD outcomes. Given the strong D_f and MI connection, we presented a predictive model for MI based on a random forest algorithm which includes D_f and a CAD polygenic risk score. This novel model improves MI individual risk prediction compared to state-of-the-art approaches, demonstrating the additional predictive power of these complementary traits to early identify high risk groups.

We identified an individual interocular D_f asymmetry in UKBB that led us to perform most of the analyses in both eyes separately. This finding is in line with published studies which reported lateral asymmetry in D_f, tortuosity, and retinal width³¹. We observed that this asymmetry is more pronounced when one of the two eyes has a D_f below the population median. Interestingly, the regression coefficients and the Pearson’s correlation estimates between D_f and UKBB traits, and the genetic findings are equivalent in both eyes independently, suggesting that the asymmetrical effect has a negligible influence at a population level. A quantitative assessment of the asymmetry of retinal vascular measurements between eyes seems crucial for studies on retinal vascular biomarkers, often conducted on a single eye, and require further work.

We found that age, sex, smoking, and developing ocular and cardiovascular diseases have a significant effect on D_f , agreeing with studies reporting that D_f decreases with age or by developing these conditions^{2,6,9,13}. Interestingly, IQS has the strongest effect on this trait. To overcome quality imaging differences, various studies insist on the importance of assessing quantitatively image quality, especially in large cohorts analysed automatically³². In our case, IQS is computed from the binary vessel map and encapsulates the vessels segmentation's sharpness and connectivity, which are key features frequently used^{32,33} to compute vascular branching complexity.

We discovered 9 loci associated with D_f with similar effect across European, Asian, and African UKBB participants. Most of these genetic variants are relevant to multiple traits and diseases; for instance, the one located near *HERC2* has been previously associated with hair³⁴, skin³⁵, and eye colour³⁶; but recent studies also suggest a strong effect in AMD³⁷, glaucoma³⁸, intraocular pressure³⁹, visual acuity⁴⁰, retinal arterial width⁴¹ and arterial and venular retinal tortuosity^{11,14}. Another interesting associated SNP is the one near the *SLC12A9* gene as it has been reported in pigmentation^{34–36}, mean arterial pressure⁴², and resting heart rate⁴³ GWAS. We found a significant negative r^2_g estimates between D_f and hypertension, CAD, and MI. The direction of these estimates agrees with their phenotypical correlations and published papers⁶, suggesting a partial explanation by their common genetic basis. This finding agrees with three aforementioned studies^{11,14,41} which identified novel retinal width and tortuosity loci associated with CAD but did not estimate a genetic correlation between retinal phenotypes and CAD. In summary, our analysis characterises the genetic architecture of D_f and highlights a common genetic basis with cardiovascular diseases, which partly explains the inverse association between D_f and coronary vessels dysfunction. Future analyses are needed to characterise the genetic expression and regulation of retinal tissue to better understand this shared genetic basis with MI and CAD.

The potential of the retinal vasculature for stratifying risk of Major Adverse Cardiac Events (MACE) has already been assessed in diabetic⁵ and non-diabetic^{4,44} individuals. Several predictive models have included retinal traits, either in a semantic⁵ or a non-semantic construction⁴, but reported very modest improvements in terms of AUC compared to the established risk estimation strategies based on epidemiological variables (e.g., 0.73 vs 0.72 in ⁴). This discrepancy with our results might be attributed to the different clinical definition of MACE, comprising normally a heterogeneous group of cardiovascular events where some of which might be not well captured in secondary care data. This situation reduces the model's statistical power as there might be an overlap in case-controls groups. In the case of diabetic population studies, both cases

and controls also have comorbidities directly affecting the architecture of the retinal vasculature that might reduce predictive power for MACE risk. In this work, we focused on MI events and considered available ICD10 guidelines and UKBB validation reports of MI data to characterize cases, achieving the maximum possible statistical power.

Recent papers have addressed the additional predictive value of a CAD PRS in MACE and CAD risk stratification^{5,8,45–47}. These approaches, although mainly developed in European populations, achieve a better identification of high-risk MI individuals than those strategies based only on epidemiological variables^{8,45–47}. Given this promising finding and the observed shared genetic basis between D_f and MI, we examined the effect of both retinal and genetic determinants on MI event risk stratification. We found that adequate clinical phenotyping is key to our models' performance, but, as shown by our ablation study, the choice of the random forest algorithm, the use of continuous variables and the introduction of D_f and PRS_{CAD} in the model all independently improve traditional individual MI risk predictions. Additionally, the model including these three modifications achieves the greatest performance. We therefore speculate that D_f provides an early indication of coronary abnormalities not captured in clinical variables and that PRS accounts for the individual protective/risk-conferring effect on the genetic architecture of the disease. Hence, the proposed model has potential, as illustrated in the Kaplan-Meier analysis, to stratify patients by MI risk. This could allow for targeted preventive efforts, like the administration of cholesterol-lowering treatments.

Our work has multiple limitations. Firstly, there are only 526 MI cases with a good-quality fundus image taken in UKBB. Higher numbers of such participants would allow us to train and evaluate our models more robustly. Secondly, PRS_{CAD} is obtained from a meta-analysis of primarily European ancestry. This restricts the application of the predictive strategy to European samples. It is of utmost importance to complete GWAS in non-European populations to provide input for PRS estimations and to be included in its applications. Thirdly, the stability of numerical estimates of the fractal dimension is the object of a continuing debate in the retinal image analysis community^{48–50}. Finally, there is little information about the genetic expression profiles and the regulation mechanisms of retinal and ocular tissues in public databases. This might be influenced by the minority of studies across these tissues and the complicated protocols to extract and characterise them.

In conclusion, our study contributes to a growing body of evidence showing associations between abnormal morphologic characteristics in coronary vessels and retinal vascular remodelling. In particular, we characterised retinal fractal dimension at a population level

and uncovered the association of 9 novel loci and a shared genetic basis with CAD. Remarkably, our MI model improved the stratification of the high-risk population. This is of great interest as it discloses a promising holistic strategy that can prevent MI incidence and triage those with an elevated MI hazard. This study ultimately sheds new light on the value of easily accessible vascular imaging phenotypes and their promising application in personalised medicine.

Materials and methods

UK Biobank

UK Biobank (<https://www.ukbiobank.ac.uk/>) is a large multi-site cohort study that consist of 502,655 individuals aged between 40 and 69 years at baseline, recruited from 22 centres across the UK during 2006-2010. The study was approved by the National Research Ethics Committee, reference 11/NW/0382, and informed consent was obtained from all participants as part of the recruitment and assessment process. From these, a baseline questionnaire, physical measurements, and biological samples were undertaken for each participant. Ophthalmic examination was not included in the original baseline assessment and was introduced as an enhancement in 6 UKBB centres across the UK. This examination consisted of capturing paired retinal fundus with a 45° primary field of view and optical coherence tomography images obtained with Topcon 3D OCT-1000 MKII (Topcon Corporation). This project was completed using fundus images collected in the first and the repeated ophthalmic examination which took place in 2012 and 2013. It includes 175,709 fundus images (87,552 left and 88,157 from the right) from 67,725 participants.

Image classification

Image quality was not reported in the cohort, and was found wanting for the purpose of automatic analysis in the first study of this kind⁵¹. A previous study defined an automated classifier for this dataset using three imaging features following vessels segmentation: white pixel ratio (WPR), largest connected component ratio (LCCR) and number of connected components (NCC) on a support vector machine (SVM) classifier¹⁹. We reproduced this classifier using a data subset of 448 random fundus images and VAMPIRE 3.1 software running in MATLAB 2018a^{17,18}. The software performs automatic detection of the retinal vasculature, creating a binary vessel map for each image. A.V.V. manually classified the quality of these images based on the connectivity and the sharpness of the binary vessel map, and the lack of imaging artefacts. Manual classification was repeated 2 times using the same random subset of 100 images and the intra-classifier agreement coefficient was 0.897. This dataset was subsequently split

in a training (n=278) and validation (n=170) sets. Both data subsets included an even number of manually classified good and bad quality images. We obtained a precision of 0.95, and a recall of 0.87, agreeing with the original study.

The classifier found 98,603 images with good quality from a total of 175,709 fundus images, of which 49,903 were from the right eye and 48,700 from the left eye. These images derived from ~45,000 participants with different ancestries and included individuals with both or one eye examined at least one time. In the case of those participants that had two good quality images from one eye, following analyses are completed using the images obtained at the first examination.

Beside classification, the classifier returns an imaging quality score (IQS) based on the distance of an image from the classification boundary computed at the training phase of the SVM. We retrieved IQS using the score parameter in the prediction function running in MATLAB 2018a. We thus quantify individually the reliability of each image being classified as bad and good image.

Calculating fractal dimension

Retinal fractal dimension, D_f , was computed from the binarized good-quality images using VAMPIRE software based on the multifractal analysis method⁵². This process was parallelised using 12 cores and 10GB per core.

Statistical analysis

To compare left and right D_f values we used participants who had both eyes scanned at the same UKBB examination and whose images were classified as good quality. 39,659 participants met these criteria. Both D_f distributions were compared using a paired T-Test and by estimating the Pearson correlation with the SciPy package in python 3. We also fitted a linear regression using respectively left and right D_f as dependent and independent variables.

We estimated the Pearson correlation and the effect of 779 UKBB traits on D_f by fitting univariate linear regressions with each variable and using D_f as dependent variable. This included 121 quantitative variables (such as age, height, and BMI) and 658 binary variables (such as sex, diagnosed with myopia, and diagnosed with hypertension) which were extracted as reported elsewhere in ²⁴. The effect of IQS was also analysed following this approach. In addition, we evaluated the IQS difference effect on D_f variability by fitting univariate linear regression using participants who had a good-quality image of both eyes scanned at the same UKBB examination. These analyses was completed

using SciPy in python 3. Allied graphs were created using matplotlib and seaborn graphical packages in python 3.

Genome-wide association studies

We included 38,811 and 38,017 individuals in the right and left GWAS, respectively, with a self-reported and genotyped confirmed unrelated white-British ancestry⁵³. Variants included were autosomal SNPs present in the genotyping arrays employed by UKBB and from the UKBB imputation panel with $HWE > 10^{-6}$, $MAF > 5 \cdot 10^{-3}$, call rate > 0.9 in unrelated white British individuals and imputation score > 0.9 in the latest ones. The number of total SNPs analysed after quality control was 9,275,849.

Following genotype-level QC, a linear regression model was used to analyse the association of each SNP genotype with D_f using PLINK v2.0. We assumed an additive genetic model, adjusting for age at examination, sex, IQS, assessment centre, the first 10 genomic principal components and genotyping batch. Additionally, we included hair and skin colour as covariates to control for the influence of skin and eye colour on the fundus image colour, which can affect image segmentation and D_f calculation. Hair colour replaced eye colour as the latter was not recorded during UKBB assessments, and it has a similar genetic control to eye pigmentation. Besides, we performed an additional GWAS including a polygenic risk score (PRS) for eye colour to assess its influence on our GWAS results. This PRS derives from an eye colour GWA study that defines it quantitatively (i.e. 1 = blue or grey, 2 = green, 3 = hazel, and 4 = brown) completed by Lona-Durazo et al. using the CanPath cohort, which includes ~5000 participants with European ancestry²³. We estimated this PRS for each participant by extracting those genetic variants with a $P\text{-value} < 5 \cdot 10^{-8}$ from the summary statistics and applying linear regression to the effects of these SNPs and the genotypes of our UKBB participants. We then included this PRS as covariate in an additional GWAS. Supplementary Fig. 7 demonstrates that the results of these GWAS are analogous to those of GWAS including both skin and hair colour.

QQ plots were generated using the R package qqman and ggplot2, and Manhattan plots and GWAS comparisons plots were generated using Matplotlib and seaborn libraries in python 3.

We completed a PheWAS to assess whether D_f loci have a significant effect in other traits. To this end, we searched D_f associated SNPs in GWASCatalog²⁵ and GeneATLAS²⁴. GWASCatalog contains hundreds of GWAS performed in different traits and populations and it constantly updates new GWAS to its database. GeneATLAS contains GWAS summary statistics for 778 UKBB traits and diseases using white

European and white British participants from UKBB. These genetic variants have a P-value smaller than $5 \cdot 10^{-8}$ on the trait in order to assume a strong association common to D_f .

GWAS and meta-analysis of D_f loci across UKBB ancestries

We performed additional GWAS including UKBB participants with white European non-British ($n_{\text{left}}=4340$ and $n_{\text{right}}=4288$), Asian ($n_{\text{left}}=562$ and $n_{\text{right}}=568$) and African ancestries ($n_{\text{left}}=498$ and $n_{\text{right}}=509$) following the aforementioned model and procedure.

The meta-analysis was completed with those significant and independent SNPs from the D_f GWAS including white British participants. We extracted the summary statistics of these SNPs from the Asian, African, and white-European GWAS and performed a fixed-effect meta-analysis across UKBB ancestries. The meta-analysis and the forest plots were carried out with Meta package in R 4.0 software.

Genetic correlation and heritability estimation

To investigate the shared genetic control between D_f and associated traits, we estimated their genome-wide genetic correlation. For this purpose, we obtained the GWAS summary statistics of traits of interest to our study from GeneATLAS and the eye colour study²³. These calculations were computed with LD Score, a toolbox that estimates genetic correlation using GWAS summary statistics considering possible inflation caused by SNPs in linkage disequilibrium (LD). To ascertain the LD blocks within each variant, the software uses the 1000 Genomes panel as reference. Heatmaps were created with the genetic correlation estimate using the seaborn library in python 3.

The LD Score was also used to calculate the SNP heritability of both eyes' D_f . In this case, the software uses the reference map and the GWAS summary statistics to estimate the fraction of D_f variance explained by the SNPs' additive effect.

Mendelian Randomization

In order to infer the causality between the shared genetic basis of CAD, MI, hypertension and D_f , we performed Mendelian Randomization analysis. For this procedure, we extracted the summary statistics of MI, hypertension, and CAD from GeneATLAS. We next selected for each cardiovascular condition separately those SNPs with a P-value $< 5 \cdot 10^{-8}$, $r^2 < 0.001$, $MAF > 0.01$ which weren't palindromic. The effect and the significance of these variants was also extracted from D_f GWAS summary statistics. We then estimated the causal effect of these genetic variants through different methods (inverse variance weighted regression, Egger's regression, and Maximum likelihood) to analyse whether using different scenarios could better characterise the causality. This

process was completed with TwoSamplesMR package in R 4.0. This package applies a quality control and a sensitivity analysis to evaluate the presence of palindromic SNPs, pleiotropy and heterogeneity which might influence the results of the study.

Enrichment analysis

Functional and enrichment analysis were completed with FUMA to further analyse those metabolic pathways underlying significant D_f loci and its associated traits with a similar expression pattern. FUMA software not only characterise significant genetic variants from a GWAS study, but it also uses available GTEX data and available studies who implemented it to establish GO terms and traits with a similar expression profile^{54,55}. We set the threshold of significant variants in a $P\text{-value} < 5 \cdot 10^{-06}$ and a reference genomic panel of UKBB white Europeans.

Development of MI predictive model

We used a subset of the UKBB data for MI model training and evaluation. We extracted white British UKBB participants who had good-quality images and a MI event after UKBB recruitment. MI events were defined in UKBB as a participant self-reporting MI at first repeated assessment visit [code 1075 from UKBB data field 20002] and MI hospitalizations identified using ICD10 codes [codes I21.1, I21.2, I21.3, I21.4, I21.9, I22, I22.0, I22.1, I22.8, I22.9, I23, I23.0, I23.1, I23.2, I23.3, I23.4, I23.5, I23.6, I23.8, I24.1 and I25.2 from UKBB data field 41204 and 41202]. The UKBB team previously validated this MI extraction algorithm and reported a minimum precision of 75%⁵⁶. To define incident cases occurring after UKBB recruitment we used the date of the MI event [UKBB algorithmically defined MI event date from data field 42000] and the approximate time period when participants underwent to the ophthalmic examination, resulting in 526 incident cases. We randomly selected an equal number of age-matched participants (40-70 years) with good-quality images of both eyes no cardiovascular event within the CAD spectrum and no known risk factor (e.g., hypertension, and family history of heart disease).

Our MI predictive model uses age, sex, systolic blood pressure, smoking, BMI, a polygenic risk score for CAD and D_f of both eyes separately as features in a classification algorithm. We chose a random forest classifier algorithm allowing both non-linear associations between outcome and variables as well as inter-variable interaction in the model. Permutation-based feature importance scores⁵⁷ were extracted in the modelling phase to assess the effect of each variable in the random forest construction using the *feature_importances_* function from the scikit-learn package. Given the influence of IQS on D_f , we trained an additional model replacing D_f to D_f adjusted by IQS with to assess

the existence of major differences in the model's performance. We also tested whether introducing just one eye D_i in the model implied major differences in its performance.

To evaluate the performance of the predictive model, we reproduced SCORE with this MI dataset. SCORE uses age, sex, systolic blood pressure, and BMI as input variables for a logistic regression, with quantitative variables being discretized using healthcare guidelines⁴. We then assessed each model's performance by using internal 10-fold cross validation and computing its AUC, precision, and recall. We used the same data partitions across SCORE and our MI models. A Wilcoxon signed-rank test was completed across all the trained models to evaluate the significance of the AUC differences.

We used Kaplan-Meier curves to assess the difference in survival rate difference between patients with high and low predicted MI probability, dichotomised at a probability of 0.5. This probability was obtained with our top-performing MI model. A Log-rank test was completed to evaluate the difference between these groups' curves.

We investigated the sources of improvement of our MI model compared to the SCORE model through an ablation study. The model differs from SCORE in four key aspects: 1) introducing D_i , 2) the use of not-discretized quantitative variables, 3) using Random Forest instead of logistic regression, and 4) introducing PRS_{CAD} . This ablation study consisted of assessing the performance of a modified version of SCORE through its AUC, recall and precision. These modifications included all the possible independent combinations across these alterations.

The code for this part of the study was written in Python 3.5.7 using the scikit-learn, NumPy and Pandas packages and is available at https://github.com/Anavillaplana/MI_risk_prediction. ROC curves were plotted using the predicted MI probability from each model using the ROCurve plot package in R 4.0. Both Kaplan-Meier curves and the Log-rank test were completed with the lifelines Python package.

Estimating CAD polygenic risk score.

PRS_{CAD} derives from the CARDIoGRAMplusC4D Consortium¹⁵ which is one of the largest completed CAD meta-analysis. This study doesn't include UKBB data, but it's developed with multiple CAD databases with different ancestries to better characterise the genetic control of this outcome. We estimated PRS_{CAD} for each participant in the MI dataset by using PRSice-2 software⁵⁸, the summary statistics of the meta-analysis, and

the genotypes of this MI dataset. We then included this PRS as variable in our MI predictive model.

Acknowledgements

This research has been conducted using the UK Biobank Resource under project 788. This work is supported by the Medical Research Council grant (MR/N013166/1) to A.V.V; the Roslin Institute Strategic Programme Grant from the BBSRC (BBS/E/D/30002275 and BBS/E/D/30002276) to A.T; Health Data Research UK grants (references HDR-9004 and HDR-9003) to A.T; the Engineering and Physical Sciences Research Council (EPSRC) grant (EP/R029598/1, EP/T008806/1) to M.O.B; Fondation Leducq grant (17 CVD 03) to M.O.B, the European Union's Horizon 2020 research and innovation programme under grant agreement No 801423 to M.O.B; Diabetes UK grant (20/0006221) to M.O.B; Fight for Sight grant (5137/5138) to M.O.B; and British Heart Foundation and The Alan Turing Institute (which receives core funding under the EPSRC grant EP/N510129/1) as part of the Cardiovascular Data Science Awards Round 2 (SP/19/9/34812) to M.O.B. Support from NHS Lothian R&D and the Edinburgh Clinical Research Facility is acknowledged.

References

1. Roth, G. A. *et al.* Global Burden of Cardiovascular Diseases and Risk Factors, 1990–2019. *J. Am. Coll. Cardiol.* **76**, 2982–3021 (2020).
2. Wong, T. Y. *et al.* Retinal Vascular Caliber, Cardiovascular Risk Factors, and Inflammation: The Multi-Ethnic Study of Atherosclerosis (MESA). *Invest. Ophthalmol. Vis. Sci.* **47**, 2341–2350 (2006).
3. MacGillivray, T. J. *et al.* Retinal imaging as a source of biomarkers for diagnosis, characterization and prognosis of chronic illness or long-term conditions. *Br. J. Radiol.* **87**, (2014).
4. Poplin, R. Prediction of cardiovascular risk factors from retinal fundus photographs via deep learning. *Nat. Biomed. Eng.* **2**, 9 (2018).
5. Fetit, A. E. *et al.* A multimodal approach to cardiovascular risk stratification in patients with type 2 diabetes incorporating retinal, genomic and clinical features. *Sci. Rep.* **9**, 3591 (2019).
6. Liew, G. *et al.* Fractal analysis of retinal microvasculature and coronary heart disease mortality. *Eur. Heart J.* **32**, 422–429 (2011).
7. Fedele, F., Pucci, M. & Severino, P. *Genetic Polymorphisms and Ischemic Heart Disease. Genetic Polymorphisms* (IntechOpen, 2017). doi:10.5772/intechopen.69621.
8. Riveros-Mckay, F. *et al.* An integrated polygenic and clinical risk tool enhances coronary artery disease prediction. 2020.06.01.20119297

- 575 <https://www.medrxiv.org/content/10.1101/2020.06.01.20119297v1> (2020)
- 576 doi:10.1101/2020.06.01.20119297.
- 577 9. Agrawal, H., Choy, H. K., Liu, J., Auyoung, M. & Albert, M. A. Coronary Artery
- 578 Disease. *Arterioscler. Thromb. Vasc. Biol.* **40**, (2020).
- 579 10. Elosua, R. & Sayols-Baixeras, S. The Genetics of Ischemic Heart Disease:
- 580 From Current Knowledge to Clinical Implications. *Rev. Esp. Cardiol. Engl. Ed.* **70**, 754–
- 581 762 (2017).
- 582 11. Veluchamy, A. *et al.* Novel Genetic Locus Influencing Retinal Venular Tortuosity
- 583 Is Also Associated With Risk of Coronary Artery Disease. *Arterioscler. Thromb. Vasc.*
- 584 *Biol.* (2019) doi:10.1161/ATVBAHA.119.312552.
- 585 12. Ikram, M. K. *et al.* Four Novel Loci (19q13, 6q24, 12q24, and 5q14) Influence
- 586 the Microcirculation In Vivo. *PLOS Genet.* **6**, e1001184 (2010).
- 587 13. Fahy, S. J. *et al.* The Relationship between Retinal Arteriolar and Venular
- 588 Calibers Is Genetically Mediated, and Each Is Associated with Risk of Cardiovascular
- 589 Disease. *Invest. Ophthalmol. Vis. Sci.* **52**, 975–981 (2011).
- 590 14. Tomasoni, M. *et al.* Genome-Wide Association Studies of retinal vessel
- 591 tortuosity identify 173 novel loci, capturing genes and pathways associated with
- 592 disease and vascular tissue pathomechanics. *medRxiv* 2020.06.25.20139725 (2021)
- 593 doi:10.1101/2020.06.25.20139725.
- 594 15. Nikpay, M. *et al.* A comprehensive 1000 Genomes–based genome-wide
- 595 association meta-analysis of coronary artery disease. *Nat. Genet.* **47**, 1121–1130
- 596 (2015).
- 597 16. Conroy, R. M. *et al.* Estimation of ten-year risk of fatal cardiovascular disease in
- 598 Europe: the SCORE project. *Eur. Heart J.* **24**, 987–1003 (2003).
- 599 17. Mookiah, M. R. K., Hogg, S., MacGillivray, T. & Trucco, E. On the quantitative
- 600 effects of compression of retinal fundus images on morphometric vascular
- 601 measurements in VAMPIRE. *Comput. Methods Programs Biomed.* **202**, 105969
- 602 (2021).
- 603 18. Trucco, E. *et al.* Novel VAMPIRE algorithms for quantitative analysis of the
- 604 retinal vasculature. *2013 ISSNIP Biosignals Biorobotics Conf. Biosignals Robot. Better*
- 605 *Safer Living BRC*.
- 606 19. Welikala, R. A. *et al.* Automated retinal image quality assessment on the UK
- 607 Biobank dataset for epidemiological studies. *Comput. Biol. Med.* **71**, 67–76 (2016).
- 608 20. Bonferroni, C. E. *Il calcolo delle assicurazioni su gruppi di teste*. (Tipografia del
- 609 Senato, 1935).
- 610 21. Rochtchina, E., Wang, J. J., Taylor, B., Wong, T. Y. & Mitchell, P. Ethnic
- 611 Variability in Retinal Vessel Caliber: A Potential Source of Measurement Error from
- 612 Ocular Pigmentation?—The Sydney Childhood Eye Study. *Invest. Ophthalmol. Vis. Sci.*
- 613 **49**, 1362–1366 (2008).
- 614 22. Li, X. *et al.* Racial Differences in Retinal Vessel Geometric Characteristics: A
- 615 Multiethnic Study in Healthy Asians. *Invest. Ophthalmol. Vis. Sci.* **54**, 3650–3656
- 616 (2013).

23. Lona-Durazo, F. *et al.* Investigating the genetic architecture of eye colour in a Canadian cohort. 2021.09.29.462299
<https://www.biorxiv.org/content/10.1101/2021.09.29.462299v1> (2021)
doi:10.1101/2021.09.29.462299.
24. Canela-Xandri, O., Rawlik, K. & Tenesa, A. An atlas of genetic associations in UK Biobank. *Nat. Genet.* **50**, 1593–1599 (2018).
25. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* **47**, D1005–D1012 (2019).
26. Zekavat, S. M. *et al.* Deep Learning of the Retina Enables Phenome- and Genome-wide Analyses of the Microvasculature. *Circulation* **0**,.
27. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* **47**, 1236–1241 (2015).
28. Lemmens, S. *et al.* Systematic Review on Fractal Dimension of the Retinal Vasculature in Neurodegeneration and Stroke: Assessment of a Potential Biomarker. *Front. Neurosci.* **14**, (2020).
29. Lee, J., Cai, J., Li, F. & Vesoulis, Z. A. Predicting mortality risk for preterm infants using random forest. *Sci. Rep.* **11**, 7308 (2021).
30. Chowdhury, A. R., Chatterjee, T. & Banerjee, S. A Random Forest classifier-based approach in the detection of abnormalities in the retina. *Med. Biol. Eng. Comput.* **57**, 193–203 (2019).
31. Cameron, J. R. *et al.* Lateral thinking – Interocular symmetry and asymmetry in neurovascular patterning, in health and disease. *Prog. Retin. Eye Res.* **59**, 131–157 (2017).
32. Sevik, U., Kose, C., Berber, T. & Erdol, H. Identification of suitable fundus images using automated quality assessment methods. *J. Biomed. Opt.* **19**, 046006 (2014).
33. Chalakkal, R. J., Abdulla, W. H. & Thulaseedharan, S. S. Quality and content analysis of fundus images using deep learning. *Comput. Biol. Med.* **108**, 317–331 (2019).
34. Morgan, M. D. *et al.* Genome-wide study of hair colour in UK Biobank explains most of the SNP heritability. *Nat. Commun.* **9**, 5271 (2018).
35. Liu, F. *et al.* Genetics of skin color variation in Europeans: genome-wide association studies with functional follow-up. *Hum. Genet.* **134**, 823–835 (2015).
36. Adhikari, K. *et al.* A GWAS in Latin Americans highlights the convergent evolution of lighter skin pigmentation in Eurasia. *Nat. Commun.* **10**, 358 (2019).
37. Guenther, F. *et al.* Chances and challenges of machine learning-based disease classification in genetic association studies illustrated on age-related macular degeneration. *Genet. Epidemiol.* **44**, 759–777 (2020).
38. Craig, J. E. *et al.* Multitrait analysis of glaucoma identifies new risk loci and enables polygenic prediction of disease susceptibility and progression. *Nat. Genet.* **52**, 160–166 (2020).

39. Zhou, W. *et al.* Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nat. Genet.* **50**, 1335–1341 (2018).
40. Currant, H. *et al.* Genetic variation affects morphological retinal phenotypes extracted from UK Biobank Optical Coherence Tomography images. *medRxiv* 2020.07.20.20157180 (2020) doi:10.1101/2020.07.20.20157180.
41. Jensen, R. A. *et al.* Novel Genetic Loci Associated With Retinal Microvascular Diameter. *Circ. Cardiovasc. Genet.* **9**, 45–54 (2016).
42. Liu, C. *et al.* Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat. Genet.* **48**, 1162–1170 (2016).
43. Eijgelsheim, M. *et al.* Genome-wide association analysis identifies multiple loci related to resting heart rate. *Hum. Mol. Genet.* **19**, 3885–3894 (2010).
44. Theuerle, J. D. *et al.* Impaired Retinal Microvascular Function Predicts Long-Term Adverse Events in Patients with Cardiovascular Disease. *Cardiovasc. Res.* (2020) doi:10.1093/cvr/cvaa245.
45. Fahed, A. C. *et al.* Polygenic background modifies penetrance of monogenic variants for tier 1 genomic conditions. *Nat. Commun.* **11**, 3635 (2020).
46. Yanes, T., McInerney-Leo, A. M., Law, M. H. & Cummings, S. The emerging field of polygenic risk scores and perspective for use in clinical care. *Hum. Mol. Genet.* **29**, R165–R176 (2020).
47. Mars, N. *et al.* Polygenic and clinical risk scores and their impact on age at onset and prediction of cardiometabolic diseases and common cancers. *Nat. Med.* **26**, 549–557 (2020).
48. Huang, F. *et al.* Reliability of Using Retinal Vascular Fractal Dimension as a Biomarker in the Diabetic Retinopathy Detection. *J. Ophthalmol.* **2016**, e6259047 (2016).
49. Cosatto, V. F. *et al.* Retinal vascular fractal dimension measurement and its influence from imaging variation: results of two segmentation methods. *Curr. Eye Res.* **35**, 850–856 (2010).
50. McGrory, S. *et al.* Towards Standardization of Quantitative Retinal Vascular Parameters: Comparison of SIVA and VAMPIRE Measurements in the Lothian Birth Cohort 1936. *Transl. Vis. Sci. Technol.* **7**, 12 (2018).
51. MacGillivray, T. J. *et al.* Suitability of UK Biobank Retinal Images for Automatic Analysis of Morphometric Properties of the Vasculature. *PLOS ONE* **10**, e0127914 (2015).
52. T, S. & Bd, S. Multifractal analysis of human retinal vessels. *IEEE Trans. Med. Imaging* **25**, (2006).
53. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

699 54. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional
700 mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826
701 (2017).

702 55. GTEx Consortium *et al.* Genetic effects on gene expression across human
703 tissues. *Nature* **550**, 204–213 (2017).

704 56. SCHNIER, C., Bush, K., Nolan, J. & Sudlow, C. *Definitions of Acute Myocardial*
705 *Infarction and Main Myocardial Infarction Pathological Types UK Biobank Phase 1*
706 *Outcomes Adjudic.*
707 https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/alg_outcome_mi.pdf (2017).

708 57. Breiman, L. Random Forests. *Mach. Learn.* **45**, 5–32 (2001).

709 58. Choi, S. W., Mak, T. S.-H. & O'Reilly, P. F. Tutorial: a guide to performing
710 polygenic risk score analyses. *Nat. Protoc.* **15**, 2759–2772 (2020).

711