Antibody Responses to 3rd Dose mRNA Vaccines in Nursing Home and Assisted Living Residents

Authors:

Ali Zhang, MSc^{1,2,4}; Jessica A. Breznik, PhD¹⁻³; Rumi Clare³, BSc; Ishac Nazy, PhD^{3,5}; Matthew S. Miller, PhD^{1,2,4}; Dawn M. E. Bowdish, PhD^{1-3,6}; Andrew P. Costa PhD^{3,7-8}

¹McMaster Immunology Research Centre, McMaster University, Hamilton, Ontario, Canada

²Michael G. DeGroot Institute for Infectious Disease Research, McMaster University, Hamilton, Ontario, Canada

³Department of Medicine, Michael G. DeGroot School of Medicine, McMaster University, Hamilton, Ontario, Canada

⁴Department of Biochemistry & Biomedical Sciences, McMaster University, Hamilton, Ontario, Canada

⁵McMaster Centre for Transfusion Research, McMaster University, Hamilton, Ontario, Canada

⁶Firestone Institute of Respiratory Health, St Joseph's Healthcare, Hamilton, Ontario, Canada

⁷Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Canada

⁸Centre for Integrated Care, St. Joseph's Health System, Hamilton, Ontario, Canada

Corresponding authors:

Dawn Bowdish, PhD

McMaster University, MDCL 4020

1280 Main Street West, Hamilton, L8S 4L8 ON, Canada.

Email: bowdish@mcmaster.ca

Andrew P. Costa, PhD

McMaster University, MIP 309A

1280 Main Street West, Hamilton, L8S 4L8 ON, Canada.

Email: acosta@mcmaster.ca

Funding Sources: This work was funded by a grant from Canadian COVID-19 Immunity Task Force and Public Health Agency of Canada awarded to APC and DMEB. DMEB is the Canada Research Chair in Aging & Immunity. APC is the Schlegel Chair in Clinical Epidemiology and Aging. Funding support for this work was provided by grants from the Ontario Research Foundation, COVID-19 Rapid Research Fund, and by the Canadian COVID-19 Immunity Task Force awarded to IN. MSM is supported, in part, by an Ontario Early Researcher Award.

Additional Information: Data in this study were collected by the COVID-in-LTC Study Group.

Members of the COVID-in-LTC Study Group include Jonathan L. Bramson, PhD; Eric D. Brown, PhD; Kevin Brown, PhD; David C. Bulir, MD, PhD; Judah A. Denburg, MD; George A. Heckman, MD, MSc; Michael P. Hillmer, PhD; John P. Hirdes, PhD; Aaron Jones, PhD; Mark Loeb, MD, MSc; Janet E. McElhaney, MD; Ishac Nazy, PhD; Nathan M. Stall, MD; Parminder Raina, PhD; Marek Smieja, MD, PhD; Kevin J. Stinson, PhD; Ahmad Von Schlegell; Arthur Sweetman, PhD; Chris Verschoor, PhD; Gerry Wright, PhD. The authors declare no conflicts of interest.

Acknowledgements: We acknowledge administrative and technical assistance from Tara Kajaks, PhD; Ahmad Rahim, MSc; Komal Aryal, MSc; Megan Hagerman; Braeden Cowbrough MSc; Lucas Bilaver; Sheneice Joseph; Angela Huynh, PhD; and Leslie Tan who were compensated for their contributions by a grant funded by the Canadian COVID-19 Immunity Task Force at McMaster University.

Conflict of Interest Disclosures: None reported. The authors have no financial relationship with any organization that may have an interest, and have no other relationships or activities that could appear to have influenced the submitted work.

Abstract

A comparison of SARS-CoV-2 wild-type and the beta variant virus neutralization capacity between 2 and 3 mRNA vaccine series in nursing home residents, and between nursing home and assisted living residents strongly supports 3rd dose vaccine recommendations, and equivalent policies for nursing homes and assisted living settings. Findings suggest that residents mount a robust humoral response to a 3rd mRNA vaccination, and have greater neutralization capacity compared to a 2 dose series.

Introduction

A 3rd dose COVID-19 vaccination was recommended in U.S. long-term care settings on September 24, 2021¹, though only half of eligible nursing home residents have received a 3rd dose as of Dec 5 2021². On August 17, 2021, the province of Ontario, Canada, was relatively early to recommend a 3rd dose mRNA vaccination for residents in nursing home and assisted living settings in response to waning protection³. Early real-world vaccine effectiveness data suggests substantially greater protection from Omicron (B.1.1.529; hereafter Omicron) infection for persons who have received a 3rd dose mRNA vaccine, compared to a 2-dose vaccination series⁴. It is unclear if residents in long-term care settings mount a robust response to a 3rd mRNA vaccination that may confer protection.

Methods

In this serial cross-sectional analysis, we examined SARS-CoV-2 neutralizing antibody titers after both 2nd and 3rd vaccination in nursing home residents, and after 3rd vaccination in assisted

living (also known as retirement home) residents from Ontario, Canada. Residents were recruited from seventeen nursing homes between March and November 2021, and eight assisted living facilities between August and November 2021. Nursing home residents were administered Moderna Spikevax 100 mcg (mRNA-1273; hereafter Moderna) or Pfizer-BioNTech Comirnaty 30 mcg (BNT163b2; hereafter Pfizer) as per recommended 2 dose schedules. Nursing home and assisted living participants were administered either Moderna or Pfizer 3rd dose vaccine no earlier than 6 months post 2nd dose. All protocols were approved by an ethics board, and informed consent was obtained.

Neutralization capacity of antibodies was assessed by cell culture assays with live SARS-CoV-2 virus, with data reported as geometric microneutralization titers at 50% (MNT₅₀) which ranged from below detection (MNT₅₀ = 10) to MNT₅₀ = 1280^{3,5}. Antibody neutralization was measured against the wild-type strain of SARS-CoV-2 and the beta variant of concern (B.1.351; hereafter beta variant). The beta variant, the most immunologically similar to omicron to date, was obtained through BEI Resources, NIAID, NIH: SARS-Related Coronavirus 2, Isolate hCoV-19/South Africa/KRISP-K005325/2020, NR-54009, contributed by Alex Sigal and Tulio de Oliveira.

Median neutralization levels by days since vaccination for the SARS-CoV-2 wild-type strain and the beta variant were plotted for comparisons. Median neutralization levels were compared by Kruskal-Wallis test for overlapping time points, and proportion below detection by chi-square, using SAS 9.4 (SAS Institute Inc.).

Results

A total of 418 nursing home residents were recruited 12-240 days post 2nd dose, with an average (standard deviation; SD) age of 82.3 (11.5) years and 63.5% (263) being female sex. The Moderna-Moderna series was administered to 52.9% (221) residents, and the Pfizer-Pfizer series to 46.4% (194) residents. A total of 103 nursing home residents were recruited 12-77 days post 3rd dose, with an average (SD) age of 83.8 (10.3) years and 62.8% (64) being female sex; whereas 95 assisted living residents were also recruited 12-77days post 3rd dose, with an average (SD) age of 85.0 (7.0) years and 65.3% (62) being female sex ($p>0.050$ for comparisons). Nursing home residents received Moderna-Moderna-Moderna (66.0%), Pfizer-Pfizer-Moderna (17.5%), and Pfizer-Pfizer-Pfizer (16.5%); whereas assisted living residents received Pfizer-Pfizer-Pfizer (61.1%), Moderna-Moderna-Moderna (22.1%), Pfizer-Pfizer-Moderna (14.7%), and Moderna-Moderna-Pfizer (2.1%).

For the equivalent 12-77 days post vaccine period, residents post 3rd dose had substantially higher neutralization titers for both the wild type and beta variant ($p<0.001$). For beta variant neutralization, 20.8% (11) of post 2nd dose residents were below the level of detection compared to none of the post 3rd dose residents ($p<0.001$). Neutralization titers were substantially lower for beta variant neutralization ($p<0.001$). There were no differences in median neutralization titers between nursing home residents and assisted living residents 12-77 days post 3rd dose, though 5.3% (5) of assisted living residents were below detection ($p<0.05$).

Discussion

Our data strongly support 3rd dose vaccine recommendations, and equivalent policies for nursing homes and assisted living settings. Consistent with non-frail populations⁵, our findings suggest that residents mount a robust humoral response to a 3rd mRNA vaccination, with greater neutralization capacity compared to a 2nd dose series. Continued monitoring of neutralization titers over time will determine the rate of degradation. Neutralization performance for the beta variant may not mimic performance for the omicron variant, which at time of writing remains a concern.

References

1. Centers for Disease Control and Prevention. Press Release - CDC Statement on ACIP Booster Recommendations. Available at: <https://www.cdc.gov/media/releases/2021/p0924-booster-recommendations-.html>. Accessed December 16, 2021.
2. Centers for Disease Control and Prevention. Nursing Home COVID-19 Vaccination Data Dashboard. Available at: <https://www.cdc.gov/nhsn/covid19/ltc-vaccination-dashboard.html#4> Accessed December 16, 2021.
3. Breznik JA, Zhang A, Huynh A, et al. Antibody Responses 3-5 Months Post-Vaccination with mRNA-1273 or BNT163b2 in Nursing Home Residents. J Am Med Dir Assoc 2021; 22(12):2512-2514.
4. UK Health Security Agency. SARS-CoV-2 variants of concern and variants under investigation in England. Technical briefing 31. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1040076/Technical_Briefing_31.pdf. Accessed December 16, 2021.
5. Huynh A, Arnold DM, Smith JW, et al. Characteristics of Anti-SARS-CoV-2 Antibodies in Recovered COVID-19 Subjects. Viruses. 2021;13(4):697.
6. Falsey AR, Fenck RW, Jr., Walsh EE, et al. SARS-CoV-2 Neutralization with BNT162b2 Vaccine Dose 3. N Engl J Med. 2021;385(17):1627-1629.

Figure. Median Neutralization (MNT₅₀) Plots for SARS-Cov-2 Wild-Type Strain (top) and Beta Variant of Concern (B.1.351) (bottom). Nursing Home Residents post 2nd compared to 3rd mRNA dose (left), and post 3rd mRNA dose for Nursing Home Residents compared to Assisted Living Residents (right).