1

Benefits of integrated screening and vaccination for infection control 2

Marie Jeanne Rabil^{1*}¶, Sait Tunc^{1¶}, Douglas R. Bish^{2&}, Ebru K. Bish^{2&} 3

¹ Grado Department of Industrial and Systems Engineering, Virginia Polytechnic Institute and State University, Blacksburg, VA, 24061, United States of America 4
5
6
7

² Department of Information Systems, Statistics, and Management Science, Culverhouse College of Business, The University of Alabama, Tuscaloosa, AL, 35487, United States of America 8
9
10

* Corresponding author 11
E-mail: mariejeanne@vt.edu (MJR) 12
13
14

¶These authors contributed equally to this work. 15

&These authors also contributed equally to this work. 16

Abstract

Importance Screening and vaccination are essential in the fight against infectious diseases, but need to be integrated and customized based on community and disease characteristics.

Objective To develop effective screening and vaccination strategies, customized for a college campus, to reduce COVID-19 infections, hospitalizations, deaths, and peak hospitalizations.

Design, Setting, and Participants We construct a compartmental model of disease spread for vaccination and routine screening, and study the efficacy of four mitigation strategies (routine screening only, vaccination only, vaccination with partial routine screening, vaccination with full routine screening), and a no-intervention strategy. The study setting is a hypothetical college campus of 5,000 students and 455 faculty members, with 11 undetected, asymptomatic SARS-CoV-2 infections at the start of an 80-day semester. For sensitivity analysis, we vary the screening frequency, daily vaccination rate, initial vaccination coverage, and screening and vaccination compliance; and consider three scenarios that represent low/medium/high transmission rates and test efficacy. Model parameters come from publicly available or published sources.

Results With low initial vaccination coverage, even aggressive vaccination and screening result in a high number of infections: 1,024/2,040 (1,532/1,773) with routine daily (every other day) screening of the unvaccinated; 275/895 with daily screening extended to the newly vaccinated in base- and worst-case scenarios, with reproduction numbers 4.75 and 6.75, respectively, representative of COVID-19 Delta variant. With the emergence of the Omicron variant, the reproduction number may increase and/or effective vaccine coverage may decrease if a booster shot is needed to maximize vaccine efficacy.

Conclusion Integrated vaccination and routine screening can allow for a safe opening of a college when initial vaccination coverage is sufficiently high. The interventions need to be customized considering the initial vaccination coverage, estimated compliance, screening and vaccination capacity, disease transmission and adverse outcome rates, and the number of infections/peak hospitalizations the college is willing to tolerate.

Introduction

The current COVID-19 pandemic has illustrated how unprepared many countries were to combat such a highly contagious disease with potentially severe health outcomes. On the science side, screening kits, then vaccines, were developed at record speed, and their availability increased over time. On the implementation side, however, screening and vaccination efforts were initially administered on a mostly *ad hoc* basis, and more importantly, these efforts were initially uncoordinated.

First came the screening kits (around January 2020 [10]), making COVID-19 screening possible. Early on, the number of screening kits was limited [37, 50], and screening focused mainly on symptomatic individuals (with isolation of those detected with the infection). As availability increased, more routine screening, especially in residential facilities where people lived in close proximity to each other (e.g., college campuses, nursing homes, prisons, military bases), was deployed. Starting in December 2020 [17], vaccines started receiving emergency approval by the Food and Drug Administration (FDA) in the United States (U.S.), and by respective regulatory agencies in other parts of the world. As vaccine supplies increased, vaccines were progressively made available to the public based on pre-established priority groups. Initially these policies lacked coordination among screening and vaccination efforts, necessitating research to develop decision-making tools as well insights on effective, coordinated interventions. By developing such a tool and insights on effective vaccination and routine screening practices for college campuses, this paper contributes to this stream of research.

The literature on screening and vaccination for infectious diseases is vast, with many studies on the benefits of, and effective strategies for, lab-based screening, e.g. [1, 14, 34, 43, 56], and vaccination, e.g. [3, 22, 31, 35, 40]; and we refer the interested reader to the many references in these works. The study by [19] pairs antibody testing with vaccination, but does not consider routine screening. The studies by [29, 33] discuss routine screening on a college campus, but without on-campus vaccination efforts, that is, only considering that some portion of campus residents are already vaccinated prior to the start of the semester (“initial vaccine coverage”); in this setting, these studies investigate the impact of different levels of vaccine efficacy and initial vaccine coverage on the number of infections. The study by [61] discusses the combination of initial vaccine coverage, on-campus screening, and on-campus population size (e.g., 100% on-campus population size implies

that all students take classes in person) to control outbreaks on a college campus. 72

We contribute to this stream of literature by building a detailed compartmental model to construct 73
integrated on-campus vaccination and routine screening strategies for a college campus, and discuss 74
how the effectiveness of the different strategies depend on community and disease characteristics, 75
including initial vaccine coverage, estimated compliance for on-campus screening and vaccination, 76
available screening and vaccination supplies/capacity, disease transmission and adverse outcome 77
rates, as well as the number of infections and peak hospitalizations the college is willing to tolerate. 78
In particular, our model is more detailed than those in the literature, and is built specifically to 79
model strategies that not only include initial vaccine coverage, but also on-campus vaccination, 80
as well as different variants of routine on-campus screening, to allow for the targeted screening 81
population to include only the unvaccinated, or the unvaccinated and the newly vaccinated. We 82
perform a detailed comparative study of the different interventions, characterized by the level 83
of on-campus vaccination and screening, for various community and disease characteristics and 84
disease-related outcome measures. 85

Various protective and/or preventative interventions, such as screening, vaccination, and isola- 86
tion of those infected, can be employed to combat a pandemic. An optimal strategy may involve 87
a combination of the different interventions, implemented with different frequencies, and needs to 88
be revised over time and customized for each region, as the available interventions (e.g., screening 89
kits, vaccine) and their supplies, and the local situation of the pandemic change. In this paper, we 90
focus on integrated strategies of routine screening and vaccination, which we always pair with the 91
isolation of infected individuals detected through screening. We show, through a case study of a 92
college campus, that vaccination and screening work synergistically, and lower the total number of 93
infections and hospitalizations, and the number of infections and hospitalizations at the peak day, 94
thus flattening the infection curve, over what either intervention can achieve alone given an initial 95
vaccine coverage; however, the effectiveness of such on-campus interventions also depend heavily 96
on the initial vaccine coverage. In addition to the timely detection and quarantine of infected indi- 97
viduals, routine screening further allows for an efficient allocation of the limited vaccine supplies: 98
It does so by removing those individuals who are infected and asymptomatic, or those with a prior 99
infection, from the vaccination pool. This redirects on-campus vaccination efforts from those who 100
would not receive much benefit, to those who would. On the other hand, vaccination shrinks the 101

susceptible population. We also provide insight on strategies that are effective in various phases of a pandemic response, characterized by different availability of screening and vaccination supplies, current vaccine coverage, and the severity of infection spread (transmission rate) in the community, as well as adverse outcome rates of the infection.

Methods

Study Design and Parameters

We develop an extended compartmental epidemic model to study the spread of an infectious disease in a population composed of two groups (students and faculty/staff, hereafter students and faculty) under protective and/or preventative interventions, including screening, isolation, and vaccination; and utilize our model to study the spread of COVID-19 on a college campus. Our model provides an extension to an existing compartmental framework [34], to model integrated screening and vaccination strategies under various compliance levels, while capturing different disease and transmission dynamics of the faculty (indexed by “f”) and students (indexed by “s”). The details of the new compartmental model can be found in the S1 Appendix.

Strategy Description: We consider routine screening that excludes symptomatic individuals who go into isolation immediately upon the start of their symptoms (following a separate testing process, as needed). Regarding routine screening, the subjects to be screened each day are selected according to the given screening strategy (i.e., screening population and frequency). *Partial screening* applies to those subjects that are either susceptible, or “thought” to be susceptible; while *full screening* extends routine screening to “vaccinated-but-not-yet-immune” subjects. All screening uses the polymerase chain reaction (PCR) test, which can detect, with high accuracy, the presence of the virus. [26, 52] It takes 8 hours to receive the test result, and a positive test result leads to the immediate isolation of the affected subject. For vaccination, we consider that some proportion of each group starts the semester as fully vaccinated (i.e., with initial immunity). Then, subjects to be vaccinated each day on campus are randomly selected from those subjects that are either susceptible, or “thought” to be susceptible (thus excluding those initially immune), and is restricted by the given vaccination rate per day. We assume that the vaccines administered on campus correspond to either the two-dose Pfizer vaccine (60%) or the two-dose Moderna vaccine (40%)

[46]. We do not consider one-dose vaccines, because they represent less than 4% of the total number of vaccines administered in the U.S. [46]; and we do not consider waning immunity for those vaccinated (either at the outset or while on campus) due to the relatively short, 80-day period of the semester. Overall, we consider four strategies that represent various combinations of vaccination and screening efforts, and the no-intervention strategy, as summarized in Table 1. We assume perfect compliance for all isolation orders, and model imperfect compliance for screening and vaccination.

Table 1. Strategy description ($V/\bar{V}$ denotes vaccination/no vaccination; $S_P/S_F/\bar{S}$ denotes partial/full/no screening)

Strategy	Definition
Strategy $\bar{V} - \bar{S}$ (no intervention)	No vaccination and no screening
Strategy $\bar{V} - S_F$	No vaccination and full screening
Strategy $V - \bar{S}$	Vaccination with no screening
Strategy $V - S_P$	Vaccination with partial screening
Strategy $V - S_F$	Vaccination with full screening
Sensitivity Analysis	
Daily vaccination rate	30/day, 60/day, 120/day
Screening frequency	1 day, 2 days, 3 days, 7 days, 14 days
Initial vaccine coverage for students (L_s)	30%, 60%, 90%
Initial vaccine coverage for faculty (L_f)	30%, 60%, 90%
Vaccination compliance rate (α)	50%, 75%, 95%
Screening compliance rate (η)	50%, 75%, 95%
Severity scenarios	worst-case, base-case, best-case

Sensitivity Analysis: We conduct various sensitivity analyses through three scenarios, which represent different levels of pandemic severity and test efficacy, and through varying the values for the daily vaccination rate, screening frequency, compliance (for each of screening and vaccination), and initial vaccine coverage (for each of faculty and students). Each scenario is characterized by the level of infection spread (modeled by group transmission rates, $\beta(f)$ and $\beta(s)$, reported in terms of an overall reproduction number, R_T , computed as a weighted average of each group's reproduction number, see S1 Appendix); exogenous conditions (X weekly imported infections, i.e., X new infections introduced to campus residents each week from external contacts, such as family

or community members); and test efficacy, in terms of test specificity (*spec*) and sensitivity (*sens*).
We label these scenarios as the *worst-case* (high infection spread, low test sensitivity and specificity, high imported infections), *best-case* (low infection spread, high test specificity, medium test sensitivity, low imported infections), and *base-case* (medium infection spread, low test specificity, medium test sensitivity, medium imported infections), Table 2. For each scenario, we vary the screening frequency (every 1,2,3,7, and 14 days) for all strategies that involve screening; and the daily vaccination rate (30/day, 60/day, 120/day) for all strategies that involve vaccination, Table 1. These strategies represent different situations where screening frequency and daily vaccination rate may be restricted due to limited testing capacity, testing kits, or vaccine supply; the proportion of students who are exempt from a vaccination mandate or are unwilling to be vaccinated; or other logistical issues involved with test or vaccine administration. Moreover, we consider different compliance rates for screening (η) and vaccination (α), Table D in S1 Appendix, and different levels of initial vaccine coverage for students (L_s), Table B in S1 Appendix and faculty (L_f), Table C in S1 Appendix, for the different strategies.

Table 2. Summary of key parameters

Model Parameter	Value(s)	Comments/References
Disease dynamics		
Mean incubation time	3 days	Paltiel et al. [34], 2020
Time to recovery	14 days	CDC [8], 2021
Probability of symptoms if infected	30%	Assumption (similar to Paltiel et al. [34], 2020), also based on Poletti et al. [36], 2021
Fatality rate (students)	0.05%	See S1 Appendix for calculations
Fatality rate (faculty)	2%	Statista [45, 47], 2021. See S1 Appendix for calculations
Hospitalization rate (students)	1.4%	CDC [6], 2021 and COVID-Net [13], 2021. See S1 Appendix for calculations
Hospitalization rate (faculty)	8.4%	CDC [6], 2021 and COVID-Net [13], 2021. See S1 Appendix for calculations
Population reproduction number (R_T)	4.75, 5.75, 6.75	See S1 Appendix for calculations
Exogenous infections (X)	5, 10, 25 per week	Assumption (similar to Paltiel et al. [34], 2020)
Screening test characteristics		
Test sensitivity (<i>sens</i>)	70% 80%	Based on Kortela et al. [23], 2021 Based on Stites & Wilen [48], 2020 and Woloshin et al. [58], 2020 and Yohe [59], 2020
Test specificity (<i>spec</i>)	98% , 99.7%	Yohe [59], 2020, Stites & Wilen [48], 2020
Time to false-positive return	1 day	Assumption (similar to Paltiel et al. [34], 2020)
Vaccine characteristics		
Vaccine efficacy (ϵ)	94.64%	See S1 Appendix for calculations
Time to reach full immunity	5.4 weeks	See S1 Appendix for calculations
Severity scenarios		
Scenario	Parameter Values	Comments/References
Base-case scenario	$R_T = 5.75$, $X = 10/\text{week}$, $sens = 80\%$, $spec = 98\%$	See comments/references above
Best-case scenario	$R_T = 4.75$, $X = 5/\text{week}$, $sens = 80\%$, $spec = 99.7\%$	See comments/references above
Worst-case scenario	$R_T = 6.75$, $X = 25/\text{week}$, $sens = 70\%$, $spec = 98\%$	See comments/references above

Setting and Parameters: We consider a hypothetical college campus of 5,000 students and 455 faculty members, with 10 students and one faculty member having undetected, asymptomatic SARS-CoV-2 infections at the outset (representing 0.2% of each group), and with some individuals arriving to campus as fully vaccinated (i.e., with initial immunity), over an 80-day academic semester. We consider the COVID-19 PCR test, which is the primary test used to detect SARS-CoV-2 [11], with sensitivity 70% to 80%, and specificity 98% to 99.7%, based on published works [23, 48, 58, 59]. Vaccination rates of 60/day or higher lead to the ability to vaccinate all the eligible subjects within the 80-day period (i.e., by the end of the semester) under all scenarios, assuming initial vaccine coverage of at least 30% for both faculty and students. Table 2 reports all key parameters, along with references, see S1 Appendix for details.

Outcome Measures of Interest: Total number of infections, hospitalizations, and deaths over the 80-day semester for each of the student and faculty groups; the peak number and peak day of new infections, the peak number of hospitalizations.

Compartmental Model

We have expanded the SEIR-based (Susceptible, Exposed, Infectious, Removed) compartmental model of a published study [34], to study the integrated screening and vaccination strategies in Table 1 under various compliance levels, for a campus population consisting of two groups (faculty and students), with group-specific disease and transmission dynamics and initial vaccination coverage. In particular, our model contains additional compartments to capture the details of screening and vaccination, when applied, as well as the characteristics of the faculty and student groups, and has the following distinct features:

- We model that some individuals start the semester as fully vaccinated (i.e., with initial immunity). For the given daily vaccination rate, the subjects to be vaccinated are randomly selected from the unvaccinated and susceptible population (i.e., the “uninfected,” “exposed,” “asymptomatic,” and “recovered & unknown” compartments), and excludes those unvaccinated individuals whose infection is detected through routine screening (i.e., the “knowingly immune” compartment, as explained below) for all strategies that involve routine screening. However, due to imperfect compliance, not all available vaccines will be used on a given day,

leading to waste.

- We consider a two-dose vaccine with 94.64% efficacy upon the full dose. The time to reach full immunity is stochastic with a mean of 5.4 weeks from the day of the first dose (see S1 Appendix for calculations), before which the subject has no immunity. Hence, vaccinated subjects can be infected before reaching full immunity. This modeling necessitates additional compartments for the vaccinated-but-not-yet-immune subjects, namely the “vaccinated-uninfected,” “vaccinated-exposed,” and “vaccinated-asymptomatic” compartments.
- Each day, routine screening is offered to everyone eligible for screening on that day according to the given screening frequency and the screening strategy utilized (i.e., partial or full screening). Due to imperfect compliance, however, not all selected subjects will undergo screening. The test has imperfect sensitivity and specificity (Table 2); and our model contains additional compartments to account for false-positives.
- A subject with a positive test result during routine screening can be either a “false-positive,” or “asymptomatic & infected.” It takes 8 hours to receive the test result, and any subject with a positive test result moves to isolation immediately upon receiving the test result, where no transmission can occur (similar to a published study [34]). All false-positives are corrected after being in isolation for one day (thus leave isolation), while both the “asymptomatic & infected” and “symptomatic & infected” subjects have an isolation time with a mean of 14 days, after which they move to the “knowingly immune” compartment, unless the symptomatic subjects are hospitalized before. We assume sufficient isolation capacity and perfect compliance for isolation.
- We differentiate between subjects that are knowingly vs unknowingly immune, via two new compartments: The “knowingly immune” compartment is for those subjects who gained immunity through either vaccination (at the outset or while on campus), a prior asymptomatic infection that was verified by a positive routine test result (leading to isolation), or a prior symptomatic infection (leading to isolation); and the “recovered & unknown” compartment is for those subjects who gained immunity through a prior asymptomatic infection that was not tested (hence not detected), i.e., these individuals became immune without their knowledge.

As a result, the subjects in the “recovered & unknown” compartment cannot be differentiated from “uninfected” individuals, and hence they might get vaccinated or tested.

Statistical Analysis

We implement the compartmental model in C++. The results are extracted to Microsoft Excel, which is used to plot different curves for the purpose of visualizing the results. No statistical tests are run, hence we do not report any statistical significance levels.

Results

Integrating Screening with Vaccination

In our hypothetical college campus of 5,000 students and 455 faculty members, 10 students and one faculty member have undetected, asymptomatic SARS-CoV-2 infections (0.2% of each group) at the start of the semester.

We first discuss our results for the base-case scenario, considering 60% initial vaccination coverage for both students and faculty, and 75% compliance for both on-campus vaccination and screening. In this setting, vaccination alone (strategy $V - \bar{S}$), at a rate of 60/day, results in 1,200 infections (1,185 students, 15 faculty) and a daily peak of 5 hospitalizations, while full routine screening alone (strategy $\bar{V} - S_F$), with an 80% sensitive and 98% specific test, results in 1,408 infections (1,392 students, 16 faculty) if conducted weekly, and 294 infections (292 students, 2 faculty) if conducted daily, compared to the no-intervention strategy (strategy $\bar{V} - \bar{S}$) that results in 1,834 infections (1,802 students, 32 faculty) and a daily peak of 8 hospitalizations over the 80-day semester in the base-case scenario. Augmenting the 60/day vaccination with routine screening of the unvaccinated population (strategy $V - S_P$) every 1, 2, 3, 7, and 14 days results in 337, 513, 633, 880, and 1,024 infections, respectively, and 2 to 4 peak hospitalizations (Table 3). Including the vaccinated-but-not-yet-immune population in routine screening (strategy $V - S_F$) further diminishes the number of infections (e.g., from 880 to 711 for weekly screening). When the vaccination rate increases (reduces) to 120/day (30/day), vaccination alone results in 448 (1,561) infections, and a peak of 1 (7) hospitalization(s) in the base-case. Augmenting the 30/day vaccination with routine screening of the unvaccinated population (strategy $V - S_P$) every 1, 2, 3, 7, and 14 days results in 484 to 1,399 infections, and 3 to 6 peak hospitalizations (Table 3). All strategies with 120/day

vaccination, and most strategies with 60/day vaccination lead to zero fatality, but strategies with 244
30/day vaccination or no vaccination lead to 1-2 deaths (per our campus of 5,455), highlighting the 245
importance of vaccination in preventing deaths. 246

Table 3. Peak number, peak day, and number (students, faculty, overall) of infections, number of deaths, and peak and total number of hospitalizations (students, faculty, overall) over the 80-day semester for all strategies, implemented at various screening frequencies and vaccination rates, under the base-case scenario, considering 75% compliance ($\eta = \alpha = 75\%$) and 60% initial campus-wide immunity ($L_s = L_f = 60\%$).

		Infections					Deaths	Hospitalizations			
Strategy	Test Frequency	Peak number	Peak day	Student number	Faculty number	Total number	Total number	Peak number	Student number	Faculty number	Total number
120 vaccines/day											
$V - S_F$	every 14d	9	25	344	3	347	0	1	1	1	2
	every 7d	7	25	278	3	281	0	1	1	1	2
	every 3 d	4	23	169	2	171	0	1	0	1	1
	every 2d	3	16	124	1	125	0	1	0	1	1
	every 1d	2	16	65	0	65	0	1	0	1	1
$V - S_P$	every 14d	11	25	377	4	381	0	1	1	1	2
	every 7d	10	25	333	3	336	0	1	1	1	2
	every 3d	8	25	247	3	250	0	1	1	1	2
	every 2d	7	25	205	2	207	0	1	0	1	1
	every 1d	5	25	142	1	143	0	1	0	1	1
$V - \bar{S}$	N/A	12	25	444	4	448	0	1	1	1	2
60 vaccines/day											
$V - S_F$	every 14d	11	30	914	10	924	1	4	5	3	8
	every 7d	8	24	704	7	711	0	3	4	3	7
	every 3 d	5	24	388	4	392	0	2	3	3	6
	every 2d	3	23	267	2	269	0	2	2	3	5
	every 1d	2	16	133	2	135	0	2	1	3	4
$V - S_P$	every 14d	12	30	1,012	12	1,024	1	4	6	3	9
	every 7d	10	31	871	9	880	0	4	5	3	8
	every 3d	7	31	627	6	633	0	3	3	3	6
	every 2d	6	37	508	5	513	0	2	3	3	6
	every 1d	5	46	334	3	337	0	2	2	3	5
$V - \bar{S}$	N/A	15	26	1,185	15	1,200	1	5	7	3	10
30 vaccines/day											
$V - S_F$	every 14d	13	31	1,307	16	1,323	2	6	16	7	23
	every 7d	10	31	1,067	11	1,078	2	5	13	7	20
	every 3 d	5	24	630	5	635	2	3	8	7	15
	every 2d	3	23	440	3	443	2	3	6	7	13
	every 1d	2	16	222	2	224	1	2	4	7	11
$V - S_P$	every 14d	14	31	1,381	18	1,399	2	6	17	7	24
	every 7d	11	31	1,224	13	1,237	2	5	14	7	21
	every 3d	6	31	906	8	914	2	4	10	7	17
	every 2d	5	37	729	6	735	1	4	8	7	15
	every 1d	3	44	480	4	484	1	3	5	7	12
$V - \bar{S}$	N/A	17	30	1,538	23	1,561	2	7	19	8	27
No vaccination											
$\bar{V} - \bar{S}$	N/A	20	31	1,802	32	1,834	2	8	23	9	32
$\bar{V} - S_F$	every 14d	15	31	1,622	22	1,644	2	7	20	8	28
	every 7d	11	31	1,392	16	1,408	2	6	16	8	24
	every 3d	6	31	857	8	865	2	4	10	7	17
	every 2d	4	30	594	5	599	2	3	7	7	14
	every 1d	2	23	292	2	294	1	2	4	7	11

Regarding compliance, at only 50% compliance for both vaccination and screening, 60 vac- 248

cines/day & weekly screening of the unvaccinated population results in 1,226 infections and a peak
of 6 hospitalizations (see Table D in S1 Appendix). Increasing the screening (vaccination) com-
pliance to 95%, while keeping vaccination (screening) compliance at 50%, results in 1,051 (765)
infections and 5 (3) peak hospitalizations. Thus, increasing the vaccination compliance reduces the
number of infections and the peak number of hospitalizations more substantially than increasing
screening compliance, that is, compliance for vaccination and screening do not have symmetric
effects.

Fig 1. Peak number of hospitalizations versus total number of infections for an initial campus-wide vaccination coverage of 60% ($L_s = L_f = 60\%$) under 75% compliance ($\eta = \alpha = 75\%$) for the base-case with full screening

Impact of Initial Campus-wide Immunity on Strategy Effectiveness

The effectiveness of any strategy is impacted by initial vaccination coverage. For example, at 60% initial vaccine coverage, 120 vaccines/day & no screening lowers the number of infections compared to 60 vaccines/day & weekly screening (Fig 1), however, the opposite is true for 30% initial vaccination coverage (Table C, Table B, Fig B and Fig C in S1 Appendix).

To understand how initial vaccine coverage affects the number of infections, consider, for example, 60 vaccines/day & full weekly screening (strategy $V - S_F$), which results in 2,749, 711, and 13 infections, and 12, 3, and 0 peak hospitalizations in the base-case, for initial vaccination coverage of 30%, 60%, 90%, respectively (Table 4). For low initial vaccination coverage of 30%, augmenting the 60 vaccines/day with aggressive (daily and full) screening still leads to 487 infections over the semester, representing 8.9% of the college's population, compared to the no-intervention case with 3,512 infections. At 30% initial vaccination coverage, even more aggressive on-campus vaccination (120 vaccines/day) and screening efforts continue to result in a high number of infections: 1,024 to 2,040 for daily partial screening; 275 to 895 with daily full screening; and 1,532 to 1,773 infections with 2-day partial screening in the base- and worst-case scenarios, respectively considering reproduction numbers of 4.75 to 6.75.

Table 4. Total number of infections and peak number of hospitalizations for the 80-day semester for some strategies, implemented at various screening frequencies and a vaccination rate of 60 vaccines/day, under the base-case scenario, considering 75% compliance ($\eta = \alpha = 75\%$) and initial campus-wide immunity ($L_s = L_f$) of 30%, 60%, 90%.

		$L_s : 30\% , L_f : 30\%$		$L_s : 60\% , L_f : 60\%$		$L_s : 90\% , L_f : 90\%$	
Strategy	Test Frequency	Total number of infections	Peak number of hospitalizations	Total number of infections	Peak number of hospitalizations	Total number of infections	Peak number of hospitalizations
60 vaccines/day							
$V - S_F$	every 14d	3,014	13	924	4	15	0
	every 7d	2,749	12	711	3	13	0
	every 3 d	1,822	8	392	2	10	0
	every 2d	1,185	5	269	2	8	0
	every 1d	487	3	135	2	5	0
$V - S_P$	every 14d	3,075	14	1,024	4	15	0
	every 7d	2,938	13	880	4	14	0
	every 3d	2,536	10	633	3	12	0
	every 2d	2,219	9	513	2	10	0
	every 1d	1,603	6	337	2	7	0
$V - \overline{S}$	N/A	3,184	15	1200	5	17	0

Discussion

Integrating routine screening with vaccination during various phases of a pandemic can play a key role in managing the spread of the infection and mitigating the impact of the pandemic. Results from the extended compartmental model in this study suggest that routine screening and vaccination work synergistically to contain a SARS-CoV-2 outbreak and achieve a level of success that neither strategy can achieve alone. However, to maximize the effectiveness, the strategy needs to be customized for each setting, considering community and disease characteristics.

Vaccination is an effective and sustainable tool for fighting disease outbreaks [60], and reduces infection transmission [15, 57] and the consequent hospitalization and death [18, 28, 31]. Although it is of critical importance to deliver and administer the available vaccines in a swift manner, vaccination efforts might be limited by resource scarcity [4, 41, 44], especially during the earlier stages of a pandemic, as well as vaccine hesitancy due, for example, to lack of trust and confidence in vaccines or health systems, or fear of side effects [9, 38, 42, 55]. Currently, 60% of the U.S. population is fully vaccinated for COVID-19, with 71% of the population receiving at least one dose of a two-dose vaccine [54], and only 17% receiving a booster shot [32], despite that vaccines have been available for every adult in the U.S. since April 2021 [2], and for 12 and over since May 2021 [27]. Because the proportion of the fully vaccinated population is still below the 70% – 85% threshold estimated to contain an infection [21, 25], and because vaccines do not provide immediate protection against the infection [7] and vaccine-induced immunity wanes over time [20], it is critical to understand how vaccination, screening, and isolation can be utilized in the containment of a SARS-CoV-2 outbreak for a college campus, and whether such on-campus efforts can substitute for low vaccination rates at the start of the semester.

Our results suggest that the effectiveness of any intervention strategy, relatedly the college's ability to offer a viable in-person teaching experience, depends heavily on initial vaccination coverage. For the study population of 5,455 over an 80-day semester, with low initial vaccination coverage (30% in our study), even aggressive on-campus vaccination and screening efforts with moderate compliance (75%) result in a high number of infections: 1,024 to 2,040 for routine daily screening of the unvaccinated population; 275 to 895 with extended screening to also cover the newly vaccinated population; and 1,532 to 1,773 infections for every 2-day screening of the unvacci-

nated population, in the base- and worst-case scenarios (see Table F in S1 Appendix), respectively
 considering reproduction numbers of 4.75 and 6.75, representative of the COVID-19 Delta variant
 [24]. These results suggest that aggressive on-campus interventions throughout the semester cannot
 typically compensate for low initial vaccination rates, that is, when starting the semester with a
 highly susceptible population, especially when the disease reproduction number is relatively high.
 Further, with the emergence of the COVID-19 Omicron variant, it is likely that the reproduction
 number will increase and/or the effective vaccine coverage will decrease if a booster shot is needed
 to maximize vaccine efficacy. These findings underscore the importance of heavily promoting, even
 mandating, full vaccination at the start of the semester.

Our study also highlights the need to customize on-campus vaccination and screening efforts
 based on community and disease characteristics. For example, at a moderate initial vaccine coverage
 of 60%, an integrated vaccination and screening strategy of 60 vaccines per day and 2-day screening
 dominates both an aggressive vaccination-only (at 120 vaccines per day) strategy and an aggressive
 screening-only (daily screening) strategy in terms of the number of infections. Of course, the
 dominating strategy depends on initial vaccine coverage, and needs to be carefully chosen. Equally
 important is the finding that integrating routine screening (both partial and full, at every 14 days or
 less) with daily vaccination is beneficial at all vaccination rates and initial vaccine coverage levels,
 and is associated with a lower number of infections and improved peak number of hospitalization
 over an 80-day semester. This result suggests that even a mild regime of routine screening, followed
 by immediate isolation of the positive cases, can complement vaccination and help mitigate the
 spread of the disease and alleviate the burden on the hospital systems, especially in institutions
 like college campuses, nursing homes, prisons, or military bases in which higher contact rates are
 expected.

Another important policy question regarding the addition of routine screening to vaccination
 is whether the vaccinated population should continue to be screened until they build immunity
 against the infection. The findings in this study suggest that including the vaccinated population
 in the screening pool can potentially decrease the total number of infections, although the degree of
 these benefits varies with screening frequency as well as the daily number of vaccines administered.
 Another policy decision related to an integrated intervention effort is the frequency of administering
 each intervention. Although the daily vaccination rate is expected to be constrained by the number

of available doses as well as the public’s willingness to be vaccinated, especially in the early stages
of vaccination efforts, screening resources are generally abundant, and their frequency may be set
considering the health return and feasibility of each strategy. The results of this study show that the
incremental benefit of more frequent screening diminishes as the daily vaccination rate increases.

Most colleges have established plans and guidelines for containing a campus-wide outbreak
of SARS-CoV-2. While many colleges mandate either vaccination, or medium to high frequency
screening if unvaccinated [30, 49], others require neither, even in regions with low vaccination rates.
These differences underscore the importance of an outcome-based analysis of different potential
interventions, such as the study conducted in this paper, to guide the decision-maker (e.g., college)
in the most appropriate strategy given their unique characteristics. Our findings imply that multiple
factors and constraints need to be included in assessing the best strategy and measures for fighting
disease outbreaks. These factors may include the daily or total number of infections a college is
willing to tolerate, the peak level of hospitalization that local hospitals can cope with, screening and
vaccination capacity that the campus is able to achieve, initial vaccine coverage on campus, and
the transmission severity scenario. It should also be considered that a partially-vaccinated campus
environment (i.e., without a vaccine mandate) could provide a false sense of security for the overall
student population, including those who are unvaccinated, and, as a result, infections may rise
quite rapidly during the academic semester. Our results also suggest that increasing vaccination
compliance has a more substantial effect in reducing the number of infections than increasing
routine screening compliance. This implies that compliance for vaccination and screening do not
have symmetric effects.

Finally, early studies indicate that existing COVID-19 vaccines, without a booster shot, may
have low efficacy for the emerging Omicron variant [12, 51]. Our findings indicate that a sufficiently
high level of initial immunity, through full vaccination (which may imply a booster shot), before
arriving on campus, is key to reducing the spread of a disease on a college campus. For low initial
vaccine coverage (e.g., 30% in our study), even aggressive on-campus vaccination and screening
efforts may not be sufficient to keep the spread under control.

Recently, several U.S. colleges have added the booster shot to their current mandate of full
vaccination for COVID-19, thus requiring all students and faculty/staff to take the booster shot
prior to the start of the Spring 2021/2022 semester, e.g., [5, 16], whereas others have delayed in-

person teaching until a safe re-opening is feasible, e.g., [53]. Given the uncertain and dynamic nature of the pandemic, our analysis indeed confirms that academic institutions need to remain flexible, and adjust their plans as needed prior to the start of each academic semester, and take immediate action based on the most recent guidelines and the current state of the pandemic. If done right, an integrated routine screening and vaccination strategy can contain any potential disease outbreak and maintain a safe environment for the local community.

Limitations

We use a compartmental model, which, by definition, is based on assumptions such as homogeneous mixing, within and between, student and faculty compartments, and uniform contact, transmission, hospitalization, and fatality rates for all individuals within a compartment. We also assume that the vaccine-induced immunity follows a step function: zero immunity in the earlier stages after vaccination, followed by full immunity after two doses (if the vaccine is effective for the individual), which takes a stochastic amount of time. We do not consider waning immunity or booster shots. While we model noncompliance with on-campus screening and vaccination guidelines, we assume that the population with symptoms or positive test results fully comply with isolation recommendations. We do not consider other mitigation tools, such as mask wearing and contact tracing.

While many parameters in our model are assumed deterministic, sensitivity analysis has been conducted on key parameters, which shows that effective strategies are robust to a wide range of parameter values. Therefore, we expect the qualitative results to continue to hold under stochasticity in parameter values. Still it would be interesting to study our strategies by explicitly modeling the stochasticity.

Finally, we consider only two groups (students and faculty), as this captures the main heterogeneity that is present on a college campus. However, more groups (compartments) can be added to the model, but this comes at the expense of increased complexity and data requirements. It would be interesting to develop a more detailed model through, for example, an agent-based simulation, which is particularly useful for modeling this type of complexity [39].

Conclusions

Integrated routine screening and vaccination is effective for containing a SARS-CoV-2 outbreak during an academic semester on a college campus, as long as initial vaccination coverage is sufficiently high, depending on the reproduction rate of the disease. Vaccination and routine screening work synergistically to achieve a level of success that neither strategy can achieve alone. However, the incremental benefit of integrated screening decreases as vaccination rate and/or initial vaccination coverage increases. Further, to maximize efficacy, the intervention strategy needs to be customized for the setting, considering community and disease characteristics, including the vaccine coverage at the start of the semester, estimated screening and vaccination compliance rates, as well as the threshold number of infections and the peak number of hospitalization a campus is willing to tolerate. The model developed in this paper can be used to establish effective strategies for containing disease outbreaks in high-density settings, including college campuses, nursing homes, and military bases.

References

- [1] 8-2-8 Urgent Care. Advantages of onsite laboratory investigations and screening services, Accessed on June 2021. <https://www.828urgentcare.com/blog/advantages-of-onsite-laboratory-investigations-screening-services>.
- [2] Allassan, Fadel. All U.S. adults now eligible for COVID-19 vaccine, Accessed on July 2021. <https://www.axios.com/covid-vaccine-eligibility-all-states-2d842548-892a-40c1-91a6-07ae9a3b0ca1.html>.
- [3] R. M. Anderson. *The impact of vaccination on the epidemiology of infectious diseases*. Elsevier, 2016.
- [4] A. D. Barrett. Yellow fever in Angola and beyond—the problem of vaccine supply and demand. *New England Journal of Medicine*, 375(4):301–303, 2016.
- [5] Best Colleges. These are the colleges requiring vaccine boosters now, Accessed on December

2021. <https://www.bestcolleges.com/news/2021/12/14/what-colleges-require-covid-vaccine-booster-omicron/>.
423
424
- [6] CDC. Demographic trends of COVID-19 cases and deaths in the US reported to CDC, Accessed
425
on November 2021. <https://covid.cdc.gov/covid-data-tracker/#demographics>.
426
- [7] Centers for Disease Control. Key things to know about COVID-19 vaccines, Accessed on July
427
2021. <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/keythingstoknow.html>.
428
- [8] Centers for Disease Control. When to quarantine, Accessed on May 2021. <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/quarantine.html>.
429
430
- [9] W.-Y. S. Chou and A. Budenz. Considering emotion in COVID-19 vaccine communica-
431
tion: Addressing vaccine hesitancy and fostering vaccine confidence. *Health Communication*,
432
35(14):1718–1722, 2020.
433
- [10] CIDRAP News. China releases genetic data on new coronavirus, now deadly, Accessed on
434
July 2021. <https://www.cidrap.umn.edu/news-perspective/2020/01/china-releases-genetic-data-new-coronavirus-now-deadly>.
435
436
- [11] Cleveland Clinic. COVID-19 and PCR testing, Accessed on June 2021. <https://my.clevelandclinic.org/health/diagnostics/21462-covid-19-and-pcr-testing>.
437
438
- [12] CNN. Cornell University reports more than 900 COVID-19 cases this week. many are Omicron
439
variant cases in fully vaccinated students, Accessed on December 2021. <https://www.cnn.com/2021/12/14/us/cornell-university-covid-cases/index.html>.
440
441
- [13] COVID-Net. Laboratory-confirmed COVID-19-associated hospitalizations, Accessed on
442
November 2021. https://gis.cdc.gov/grasp/COVIDNet/COVID19_3.html.
443
- [14] A. Deckert, T. Bärnighausen, and N. N. Kyei. Simulation of pooled-sample analysis strategies
444
for COVID-19 mass testing. *Bulletin of the World Health Organization*, 98(9):590, 2020.
445
- [15] J. Ehreth. The global value of vaccination. *Vaccine*, 21(7-8):596–600, 2003.
446
- [16] Emory University. COVID-19 vaccine, Accessed on December 2021. <https://www.emory.edu/forward/covid-19/vaccine.html>.
447
448

- [17] FDA. FDA takes key action in fight against COVID-19 by issuing emergency use author-
ization for first COVID-19 vaccine, Accessed on July 2021. [https://www.fda.gov/news-
events/press-announcements/fda-takes-key-action-fight-against-covid-19-issuing-emergency-
use-authorization-first-covid-19](https://www.fda.gov/news-events/press-announcements/fda-takes-key-action-fight-against-covid-19-issuing-emergency-use-authorization-first-covid-19).
[18] J. C. Fitzgerald. Updating the epidemiology of severe bacterial infections reveals old truths:
Vaccination saves life and limb. *Pediatric Critical Care Medicine*, 21(6):595–596, 2020.
[19] A. B. Fujimoto, I. Yildirim, and P. Keskinocak. Significance of SARS-CoV-2 specific antibody
testing during COVID-19 vaccine allocation. *Vaccine*, 2021. doi: [https://doi.org/10.1016/j.
vaccine.2021.06.067](https://doi.org/10.1016/j.vaccine.2021.06.067).
[20] Y. Goldberg, M. Mandel, Y. M. Bar-On, O. Bodenheimer, L. Freedman, E. J. Haas, R. Milo,
S. Alroy-Preis, N. Ash, and A. Huppert. Waning immunity after the BNT162b2 vaccine in
Israel. *New England Journal of Medicine*, 2021.
[21] Holcombe, Madeline. More than 91 million live in U.S. counties with high COVID-19 infections.
it’s time to reset and put masks back on, expert says, Accessed on July 2021. [https://edition.
cnn.com/2021/07/21/health/us-coronavirus-wednesday/index.html](https://edition.cnn.com/2021/07/21/health/us-coronavirus-wednesday/index.html).
[22] B. Kahn, L. Brown, W. Foege, H. Gayle, E. National Academies of Sciences, Medicine, et al.
A framework for equitable allocation of COVID-19 vaccine. *National Academies Press (U.S.)*,
2020.
[23] E. Kortela, V. Kirjavainen, M. J. Ahava, S. T. Jokiranta, A. But, A. Lindahl, A. E.
Jääskeläinen, A. J. Jääskeläinen, A. Järvinen, P. Jokela, et al. Real-life clinical sensitivity
of SARS-CoV-2 RT-PCR test in symptomatic patients. *PloS one*, 16(5):e0251661, 2021.
[24] Y. Liu and J. Rocklöv. The reproductive number of the delta variant of SARS-CoV-2 is far
higher compared to the ancestral SARS-CoV-2 virus. *Journal of travel medicine*, 2021.
[25] Mayo Clinic. Herd immunity and COVID-19 (coronavirus): What you need to know, Accessed
on July 2021. [https://www.mayoclinic.org/diseases-conditions/coronavirus/in-depth/herd-
immunity-and-coronavirus/art-20486808](https://www.mayoclinic.org/diseases-conditions/coronavirus/in-depth/herd-immunity-and-coronavirus/art-20486808).

- [26] Medical Device Network. Different paths to the same destination: Screening for COVID-19, Accessed on June 2021. <https://www.medicaldevice-network.com/features/types-of-covid-19-test-antibody-pcr-antigen/>.
- [27] Mervosh, Sarah. ‘I just got my vaccine’: U.S. rollout expands to children ages 12 to 15, Accessed on July 2021. <https://www.nytimes.com/2021/05/13/us/coronavirus-vaccine-children-ages-12-15.html>.
- [28] J. Meyer, M. Sibanda, and R. Burnett. Vaccination against influenza saves lives. *SA Pharmaceutical Journal*, 87(2):26–30, 2020.
- [29] F. C. Motta, K. A. McGoff, A. Deckard, C. R. Wolfe, M. Bonsignori, M. A. Moody, K. Cavanaugh, T. N. Denny, J. Harer, and S. B. Haase. Assessment of simulated surveillance testing and quarantine in a SARS-CoV-2-vaccinated population of students on a university campus. In *JAMA Health Forum*, volume 2, pages e213035–e213035. American Medical Association, 2021.
- [30] New York University. Vaccine requirement, Accessed on July 2021. <https://www.nyu.edu/life/safety-health-wellness/coronavirus-information/safety-and-health/covid-19-vaccine/vaccine-questions-and-support/vaccine-requirement.html>.
- [31] W. A. Orenstein and R. Ahmed. Simply put: Vaccination saves lives. *National Acad Sciences*, 2017.
- [32] Our World in Data. COVID-19 vaccine booster doses administered per 100 people, Accessed on December 2021. https://ourworldindata.org/grapher/covid-vaccine-booster-doses-per-capita?country=BRA~CHL~ISR~RUS~USA~URY~OWID_WRL.
- [33] A. D. Paltiel and J. L. Schwartz. Assessing COVID-19 prevention strategies to permit the safe opening of residential colleges in Fall 2021. *Annals of internal medicine*, 2021.
- [34] A. D. Paltiel, A. Zheng, and R. P. Walensky. Assessment of SARS-CoV-2 screening strategies to permit the safe reopening of college campuses in the United States. *JAMA Network Open*, 3(7):e2016818–e2016818, 2020.

- [35] T. Philipson. Economic epidemiology and infectious diseases. *Handbook of Health Economics*, 1:1761–1799, 2000.
- [36] P. Poletti, M. Tirani, D. Cereda, F. Trentini, G. Guzzetta, G. Sabatino, V. Marziano, A. Castrofino, F. Grosso, G. Del Castillo, et al. Association of age with likelihood of developing symptoms and critical disease among close contacts exposed to patients with confirmed SARS-CoV-2 infection in Italy. *JAMA Network Open*, 4(3):e211085–e211085, 2021.
- [37] QUARTS. Here are the coronavirus testing materials that are in short supply in the U.S., Accessed on July 2021. <https://qz.com/1822596/all-the-coronavirus-test-materials-in-short-supply-in-the-us/>.
- [38] R. Rabi, B. Maraqa, Z. Nazzal, and T. Zink. Factors affecting nurses’ intention to accept the COVID-19 vaccine: A cross-sectional study. *Public Health Nursing*, 2021.
- [39] H. Rahmandad and J. Sterman. Heterogeneity and network structure in the dynamics of diffusion: Comparing agent-based and differential equation models. *Management Science*, 54(5):998–1014, 2008.
- [40] R. Rappuoli, M. Pizza, G. Del Giudice, and E. De Gregorio. Vaccines, new opportunities for a new society. *Proceedings of the National Academy of Sciences*, 111(34):12288–12293, 2014.
- [41] A. Ribas, R. Sengupta, T. Locke, S. K. Zaidi, K. M. Campbell, J. M. Carethers, E. M. Jaffee, E. J. Wherry, J.-C. Soria, and G. D’Souza. Priority COVID-19 vaccination for patients with cancer while vaccine supply is limited. *Cancer Discovery*, 11(2):233–236, 2021.
- [42] B. Roy, V. Kumar, and A. Venkatesh. Health care workers’ reluctance to take the COVID-19 vaccine: A consumer-marketing approach to identifying and overcoming hesitancy. *NEJM Catalyst Innovations in Care Delivery*, 1(6), 2020.
- [43] S. Shahangian. Laboratory-based health screening: Perception of effectiveness, biases, utility, and informed/shared decision making. *Laboratory Medicine*, 37(4):210–216, 2006.
- [44] E. Shim. Optimal allocation of the limited COVID-19 vaccine supply in South Korea. *Journal of Clinical Medicine*, 10(4):591, 2021.

- [45] Statista. Number of coronavirus disease 2019 (COVID-19) deaths in the U.S. as of November 24, 2021, by age*, Accessed on November 2021. <https://www.statista.com/statistics/1191568/reported-deaths-from-covid-by-age-us/>.
- [46] Statista. Number of COVID-19 vaccine doses administered in the United States as of November 17, 2021, by vaccine manufacturer, Accessed on November 2021. <https://www.statista.com/statistics/1198516/covid-19-vaccinations-administered-us-by-company/>.
- [47] Statista. Total number of cases of COVID-19 in the United States as of November 24, 2021, by age group, Accessed on November 2021. <https://www.statista.com/statistics/1254271/us-total-number-of-covid-cases-by-age-group/>.
- [48] E. C. Stites and C. B. Wilen. The interpretation of SARS-CoV-2 diagnostic tests. *Med*, 2020.
- [49] Syracuse University. COVID-19 update: Campus access — required testing — masks, Accessed on July 2021. <https://news.syr.edu/blog/2021/06/01/covid-19-update-campus-access-required-testing-masks/>.
- [50] The New Yorker. Why widespread coronavirus testing isn't coming anytime soon, Accessed on July 2021. <https://www.newyorker.com/news/news-desk/why-widespread-coronavirus-testing-isnt-coming-anytime-soon>.
- [51] UK Health Security Agency. SARS-CoV-2 variants of concern and variants under investigation in England, Accessed on December 2021. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1040076/Technical_Briefing_31.pdf.
- [52] University of Michigan Health. Human Immunodeficiency Virus (HIV) test, Accessed on June 2021. <https://www.uofmhealth.org/health-library/hw4961>.
- [53] University of Toronto. COVID-19 planning updates, Accessed on December 2021. <https://www.utoronto.ca/utogether/covid-19-planning-update>.
- [54] U.S.A. Facts. U.S. coronavirus vaccine tracker, Accessed on July 2021. <https://usafacts.org/visualizations/covid-vaccine-tracker-states/>.

- [55] P. Verger and E. Dubé. Restoring confidence in vaccines in the COVID-19 era, 2020. 553
- [56] S. Wacharapluesadee, T. Kaewpom, W. Ampoot, S. Ghai, W. Khamhang, K. Worachotsuep- 554
trakun, P. Wanthong, C. Nopvichai, T. Supharatpariyakorn, O. Putcharoen, et al. Evaluating 555
the efficiency of specimen pooling for PCR-based detection of COVID-19. *Journal of Medical* 556
Virology, 92(10):2193–2199, 2020. 557
- [57] A. M. Wendelboe, C. Avery, B. Andrade, J. Baumbach, and M. G. Landen. Importance of 558
employee vaccination against influenza in preventing cases in long-term care facilities. *Infection* 559
Control & Hospital Epidemiology, 32(10):990–997, 2011. 560
- [58] S. Woloshin, N. Patel, and A. S. Kesselheim. False negative tests for SARS-CoV-2 infec- 561
tion—challenges and implications. *New England Journal of Medicine*, 383(6):e38, 2020. 562
- [59] S. Yohe, C. of American Pathologists, et al. How good are COVID-19 (SARS-CoV-2) diagnostic 563
PCR tests. *College of American Pathologists*, 2020. 564
- [60] W. T. Zha, F. R. Pang, N. Zhou, B. Wu, Y. Liu, Y. B. Du, X. Q. Hong, and Y. Lv. Research 565
about the optimal strategies for prevention and control of varicella outbreak in a school in a 566
central city of China: Based on an SEIR dynamic model. *Epidemiology & Infection*, 148, 2020. 567
- [61] X. Zhao, H. Tatapudi, G. Corey, and C. Gopalappa. Threshold analyses on combinations of 568
testing, population size, and vaccine coverage for COVID-19 control in a university setting. 569
Plos one, 16(8):e0255864, 2021. 570

Supporting Information 571

S1 Appendix. Supplementary Online Content 572