

Supplemental Material

Supplemental Table 1a. SARS-CoV-2 PCR positivity by SARS-CoV-2 serostatus and CD4+ cell count in PLWH.

	Omicron detected	Omicron not detected	Total	% Omicron detected
Group 1 (SARS-CoV-2 seronegative)				
500 and above	10	48	58	17%
200 - 500	16	19	35	46%
less than 200	1	6	7	14%
Total	27	73	100	27%
Group 2 (SARS-CoV-2 seropositive)				
500 and above	12	31	43	28%
200 - 500	3	12	15	20%
less than 200	1	1	2	50%
Total	16	44	60	27%
Total	43	117	160	27%

Supplemental Table 1b: Summary of cycle threshold (CT) values by Fisher Taqpath for those samples with S gene drop out (n=56).

Target	N	Mean CT	SD CT	Min CT	Median CT	Max CT
N gene	56	25.8	5.3	14.5	25.8	35.7
ORF1ab	56	25.2	5.5	14.4	25.7	34.1
Average N and ORF1a	56	25.4	5.4	14.4	25.8	34.9


Note CT values of 20 and 25 correspond to log₁₀ viral loads of approximately 6.5 and 5, respectively.

Supplemental Table 2. Prevalence of asymptomatic SARS-CoV-2 infection by PCR in Sisonke sub study from enrollment through 6 months.

	Predominant variant(s)*	No. of participants	SARS-CoV-2 PCR+	Prevalence
Enrollment	Beta, Delta	1,604	39	2.4%
Week 3	Delta	1,447	51	3.5%
Week 6	Delta	1,260	25	2.0%
Month 6	Omicron	577	91	16%
<i>HIV positive</i>	<i>Omicron</i>	<i>169</i>	<i>27</i>	<i>16%</i>
<i>HIV negative</i>	<i>Omicron</i>	<i>405</i>	<i>62</i>	<i>15.3%</i>

*Source: <https://covariants.org/per-country>)

Month 6 participants sub divided by HIV status are in italics.



Supplemental Figure 1. Summaries of cycle threshold values. CoVPN 3008 baseline positive swabs (x axis) with S gene target failure (n=56). Fisher Taqpath CTs arranged in order of increasing Abbott cycle number. Dotted line denotes CT of 25. Note CT values of 20 and 25 correspond to log₁₀ viral loads of approximately 6.5 and 5, respectively.