Trends and country-level variation in age at first sex in sub-Saharan Africa among birth cohorts entering adulthood between 1985 and 2020

Van Knh Nguyen PhD^{1,*}, Jeffrey W Eaton PhD¹

¹MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, United Kingdom

* Corresponding author:

Van Knh Nguyen
MRC Centre for Global Infectious Disease Analysis, School of Public Health
Imperial College London
St. Mary's Hospital Campus
Norfolk Place
London
W2 1PG
k.nguyen@imperial.ac.uk

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background: Debuting sexual intercourse is a life course event marking exposure to pregnancy or fatherhood, and sexually transmitted infections (STIs), including HIV. We systematically analysed the timing, distribution, and trends in age at first sex (AFS) in 42 sub-Saharan Africa (SSA) countries.

Methods: We collated individual-level AFS data from nationally representative household surveys across SSA. We used a log-skew-logistic distribution to model the distribution of AFS in a Bayesian spatio-temporal hierarchical random-effect model to estimate national trends of AFS over time and space, adjusting for age at report biases.

Results: Small changes in AFS are observed between the birth cohorts entering adulthood between 1985 and 2020, ranging 0,79 years [-0,01–1,51] and 0,48 [-0,03–1,92] for female and male, respectively. Northern SSA countries show appreciable increase in AFS but its gender gap remains the widest compared with minimal gender gap in the southern countries. The gender gap shows little evidence of change over time in most of the countries. Female's AFS approach to a similar age across SSA while male's AFS varies between regions. Proportion ever had sex under fifteen and eighteen are as high 34% and 83%, respectively. AFS distribution is typically asymmetric with most of sexual debut events occur in a span of 3,9 [3,4–5,0] years. Female teen often reports higher AFS compared to their late twenties while male teen report lower AFS; both sexes tend to recall a higher AFS in older ages compared to their thirties.

Conclusions: Women debut sexually earlier and in a shorter span of age than men. Northern and southern SSA gender gap are distinctively different. Since the ratifying of HIV/AIDS intervention programs in SSA, a stagnant trend in AFS had remained in the countries most affected by the epidemic.

Introduction

Debuting sexual intercourse is an important life course event for myriad developmental, social, and health reasons. Initiation of sexual activity marks the time when an individual becomes at risk for pregnancy or fatherhood and of acquiring and transmitting sexually transmitted infections (STIs) (1), including HIV infection. Early sexual debut has been associated with a higher rate of STIs (2–5) and an increased risk of invasive cervical carcinoma (6). Early AFS could also heighten the likelihood of engaging risky sexual behaviours that facilitate STIs (7). At the population level, the median age of first sex (AFS) varies between populations according to cultural factors, religious practices, education, socio-economic status (8–11), and has changed over time along with these determinants and in response to public health efforts, particularly for HIV/AIDS. Delayed sexual debut has been associated with decline in HIV incidence (4,12) and contributes to curtailing HIV epidemics in sub-Saharan Africa (SSA) (13). Consequently, population-level data about AFS has become a key sexual risk indicator for determining and monitoring appropriate HIV and STI prevention activities (14,15), and predicting the vulnerability to and modelling the transmission of STIs within populations, especially those with generalized HIV epidemics in sub-Saharan Africa.

Data on AFS are routinely collected from nationally-representative household surveys using face-to-face interviews in a retrospective manner (16,17). Surveys are typically conducted roughly every five years to monitor key health and development indicators. These data are susceptible to systematic response biases related to difficulty recalling dates of life events that happened long ago and desirable standards of society, culture, and health or political campaigns (18–21). These types of bias are not evident from checking within-questionnaire consistency and cannot be corrected during the data cleaning process, necessitating the need to incorporate biases into the modelling process.

Assessing the reliability in reporting sexual behaviour in longitudinal studies showed that the percentage of inconsistent AFS reports ranged from 30% to 56% between survey rounds

(22,23), with men less reliable than women (17,22,23), and a tendency to report older AFS over time (23–25). In the same birth cohort, the absolute difference in the reported AFS between survey rounds were in both directions, mostly in a magnitude of a few years but can be as high as 10 years, however median age difference between surveys were consistently close to zero in both sexes (17,23); as the cohort aged from teenager to adulthood, men reported higher AFS while women reported lower AFS (24,26,27). Including inconsistency report samples in survey analyses, however, was found not affecting the estimated median AFS (23,28) or multivariate model (17). Covariates including sex, ethnicity, education, residence, reporting age, and HIV serostatus had been included AFS analyses to accounting for the differences in AFS reports (7,24,29); in some cases, the correction resulted in a loss of significant in cohort trends (24).

While many studies have documented evidence of reporting biases in AFS, few have attempted to correct for them to reconstruct how trends in AFS have changed within and between populations. We applied a model for the distribution of age of sexual debut by birth cohort that also uses data from multiple surveys within the same population to explicitly model reporting bias as a function of age at the time of interview. We used the model to estimate the national-level distribution of AFS and trends of sexual debut over the last four decades for countries across SSA.

Material and Methods

We used a skew logistic distribution to model the distribution of the log of reported age of first sex. To characterise temporal trends and geographic variation in the AFS distribution, we allowed the skew logistic distribution parameters to vary by country and birth cohort. Since members of each birth cohort reported data about their AFS distribution in multiple household surveys spaced, typically, approximately five years apart, we used the differences in the AFS reported among the same cohort in successive surveys to quantify and adjust for a systematic reporting bias in the AFS according to the age of the cohort at the time of the survey. This adjustment is described in more detail elsewhere (30). Changes in the AFS

over time for successive birth cohorts were represented by a first-order random walk (RW1) for males and females in each country. The AFS reporting bias as function of age at report was smoothed with a RW1 term for each sex and country. To accommodate countries with only one survey available (making identification of country-specific bias from successive surveys impossible), the effect was estimated borrowing from other countries with a spatially structured model.

Household survey data

Nationally representative surveys which collected AFS were collated, including Demographic and Health Surveys (DHS) (31), Malaria Indicator Survey (MIS) (31), Multiple Indicators Cluster Surveys (MICS) (32), AIDS Indicators Survey (AIS) (31), Performance Monitoring and Accountability (PMA) (33), Population and Health Survey (PHS) (34), Population-based HIV Impact Assessment (PHIA) (35), Sexual Behavior Survey (SBS) (36), South Africa HIV Prevalence, HIV Incidence, Behaviour and Communication Survey (SABSSM) (37). A complete list of surveys can be found in S1 Table. We used only individual records datasets. Respondents' age was calculated from the century-month-code records of data of birth and date of interview; if those variables were not presented, we calculated the century-month-code from the month and year of birth; if only the year was available, we set month of birth to June; otherwise, the age of respondent variable in the questionnaires were used.

Model for age at first sex

AFS was reported as integer ages. We modelled the integer AFS as an interval censored event in which the exact event occurring between the reported AFS and AFS + 1. Individuals who reported not yet sexually debuting were right censored at their age at the interview. The likelihood for the reported AFS by each survey respondent was modelled using a time to event, interval censored model. A weighted pseudo-likelihood approach was used to account for unequal survey weights, which were normalized to the total sample size of each survey (38). The three-parameter skew log-logistic distribution was used to describe the AFS

distribution. This distribution permits a non-monotonic and asymmetric functional form of the sexual debut hazard by age which other commonly used distributions, e.g., log-normal, gamma, were not able to capture. It was also shown to have a better predictive value than the simpler two-parameter log-logistic distribution, which has a symmetric hazard function (39). Illustrations of the distribution and its parameters are in S1 Text.

The model was fitted separately by sex but simultaneously for all countries, with independent shape and skewness parameters for each country. The linear predictor was modelled on the distribution's scale parameter as an additive effect of the spatial correlation, birth cohort, country-birth cohort interaction, age at report effect, and country-age interaction. Both birth cohort and age at report main effects were structured as first-order random walk (RW1) models and an intrinsic conditional auto-regressive (ICAR) model (40) was used to model the correlation between neighbouring countries. We assumed interactions that allowed the temporal trend and age effect to vary between countries but are correlated between neighbouring countries. Model parameters were estimated with empirical Bayes using Template Model Builder (41). Further technical specifications of the model constraints, priors, and implementations are in S1 Text. We computed posterior medians and 95% credible intervals of the outcomes of interests, including median AFS and proportion ever had sex by specified age thresholds, using posterior simulations of 3000 parameters samples.

Results

Survey data were collated for 42 SSA countries. Eight countries were excluded from our analysis because they did not have household surveys with AFS data (Cape Verde, Djibouti, Equatorial Guinea, Guinea Bissau, Mauritania, Mauritius, Seychelles, and Somalia). Among 290 surveys included (consisting of 288 female and 168 male datasets), 23 datasets had more than 10% missing the AFS variables (AFS or ever had sex indicator variable), mainly from PMA, HSRC, and MICS3 surveys; 34 datasets had more than 10% improper records by the survey's definition (e.g., AFS coded in the data as don't know, not applicable, refused,

or inconsistent depending on a specific survey, ever had sex indicator coded other than zero or one). These observations were removed from the analyses. In addition, 101 datasets had reports of AFS under seven years, ranging 0,004%–5% of all observations. We considered those observations irregularities and removed from the analyses. We included only respondents aged 15–49 years at the time of interview and born after 1950 (94,2% of all observations). The final dataset contained 2,968,614 observations for 55 birth cohorts from 1950 to 2005 (entering adulthood at age 15 between 1965 and 2020). The sample size per dataset ranged from 774 (male, Comoros DHS 1996) to 40,759 (female, Nigeria DHS 2018). Further details are in S1 Table.

Spatial patterns of median AFS and proportion sexually active by age 18

For the birth cohort entering adulthood (turning age 15) in 2015, the median AFS ranged from 15,8 [IQR: 14,5–17,2] in Central Africa Republic to 24,3 [IQR: 21,3–27,9] in Niger (Fig 1). Females had lower median AFS than males in most countries, but across countries the AFS between the two sexes were correlated such that countries where females had low median AFS also males had low median AFS in male and vice versa. This was substantiated by high correlation between male and female's estimates of the skew-logistic distribution parameters (S1 Fig). The median AFS for men was less than women only in Lesotho and South Africa. Only females in Gambia, Burundi, Comoros, and Eritrea had median AFS above twenty.

Fig 1. Distribution of the age at first sex by region, country, and sex. The median of each distribution is shown with the vertical lines. The estimates are for the birth cohort turn fifteen in 2015 and reporter aged 23. Within each region, countries sequenced by increase median female AFS.

The spatial variation in sexual debut is further evident in the percentage who ever had sex before age 18 years (Fig 2). A very low percentage of men in the Sahel region of Africa had debuted sex by age 18, with the lowest 6,3% [4,5–8,7%] percent in Niger. Among women,

the percentage sexually active by age 18 was lowest, below 45%, in the southern countries, and as low as 31,8% [23,3–41,3%] in Botswana. In both sexes, Botswana, Burundi, and Rwanda all had older AFS than the surrounding region whereas Mozambique had notably younger AFS in the southeast Africa. At the other extreme, women in Central Africa Republic and men in Angola were most likely to debut sex under eighteen with 84,2% [79,2–88,0%] and 75,7% [70,9–79,9%], respectively. AFS were spatially homogenous in the remained locations. Similar patterns were observed for the percentage ever had sex under fifteen (S2 Fig) where the largest estimate was 35,4% [30,2–40,7%] in Angola male following by Central Africa female at 32,9% [25,7–40,8]. In half of cases (41/82) less than 10% had initiated sexual activity by age 15. (S2 Fig, S2 Table).

Fig 2. Percentage ever had sex by age 18 among the birth cohort turning 15 in the year 2015. The two countries Comoros and São Tomé & Príncipe are represented in the bottom left. Countries without data are coloured the same as the background.

Trends in AFS by cohort

The median AFS increased between birth cohorts in 1985 and 2017 in 58 of 82 time-series estimated by country and sex (**Error! Reference source not found.**), and in 34 increased more than one year. During this period, the average change in median AFS across all countries was 0,79 years [IQR: -0,01-1,51] for females and 0,48 [IQR: -0,03 -1,92] for males. The changes, however, have not been monotonic but fluctuated over time (such as in Gambia, Guinea Bissau, Uganda, and Rwanda) and varied between countries. The estimated trends were similar for male and female in most countries with few exceptions of Comoros, South Africa, and Lesotho in the most recent birth cohorts. Increases in AFS were especially large for men in some West African countries (e.g., Senegal, Niger, Mali, Chad, Nigeria, Burkina Faso, Benin). The largest estimated decreases were also among men (e.g., Tanzania, Sierra Leon, Lesotho, Mozambique). Eastern and Southern Africa exhibited

minimal changes in AFS during the time frame with the average increase of 0,31 and 0,12 years, respectively.

Fig 3. Trend in median age at sexual debut for men and women in Sub Saharan-Africa countries by year turned age 15 (x-axis). For countries where the most recent survey is before 2015, trends are extrapolated for recent birth cohorts based on the overall trend across the region and its neighbours. Countries without AFS data are excluded in this figure.

In the same birth cohort, the gap in median AFS between women and men was under two years in most countries and remained unchanged in the region. The gap was larger in the West Africa countries, especially those in the Sahel and Nigeria, and in the Horn of Africa (Ethiopia, Eritrea). Median AFS increased most in Western Africa (Fig 4). Middle Africa had the lowest median AFS on average for both men and women. In Southern Africa, which is dominated by the large population of South Africa, there was no sex difference in median AFS and median AFS decreased by approximately two years since the 1985 adult cohort, in contrast to the increasing trend in other regions.

Fig 4. Aggregated trend of median age at first sex by region for men and women. Sudan was included in Eastern Africa. The regional trend is a weighted average of the countries included in the analyses by their respective population size in 2020 sourced from the UN World Population Prospects.

Distribution of age at sexual debut by sex

The distribution of AFS was asymmetric in most countries with various degree of skewness (Fig 1). The inter-quartile range (IQR) of the AFS distribution represent the age range with the highest sexual debut activity and its magnitude reflects how fast the rate of most of the population becomes sexually active. In most countries the IQR was under five years with an average of 3,9 [3,4–5,0] years. It took as short as 2,25 years (Liberia - female) or up to 8,8 years (Sudan - male) for 50% of a birth cohort entering sexually active life. Between sexes, in most countries (37/40) the IQR was shorter for females than for males; among these, 17

countries had a difference of more than a year. The distribution of AFS spanned a longer range for countries of the Sahel and Horn of Africa.

Biases in reported AFS by age at interview

The estimated pattern for the effect of age at the time of interview on the reported AFS for each country are shown in Fig 5. Data in different countries yielded similar information on the effect of age at report bias. Among females, there was a tendency to report higher AFS in their teens compared to when they reached their late twenties, indicating under-reporting of sexual activity among the youngest survey respondents. The opposite pattern was observed among men—young men tended to report a younger AFS distribution. For both sexes, the reported distribution of AFS increased as the cohort aged. The direction of both bias's directions varied across countries. The coefficient estimates of the biases, however, only show the relative difference between ages, but do not identify at which age people would be both truthful and accurately recall their AFS. To arrive at a marginal estimate for the AFS of each birth cohort, we chose age 23 as the reference age in both female and male and report estimates using this reference age throughout the results presented.

Fig 5. Direction of the effect of age at report bias on the median age at first sex estimate relative to other ages. A positive coefficient in the figure corresponds to an increase in median AFS estimate reported by the respective age compared to a median AFS reported by the age with that coefficient equals zero.

Discussion

In this study, we systematically analysed trends and patterns in debuting sexual activity across sub-Saharan Africa among cohorts entering adulthood from 1985 to 2020, a period during which the continent combatted the HIV/AIDS pandemic and experienced rapid development and large efforts to increase access to education (42). We applied a statistical model to account for reporting biases that have previously hampered consistent

interpretation of AFS recorded in household surveys and used a parsimonious but flexible model to characterise the distribution of AFS over ages.

Gender gaps in AFS differed greatly between the southern and northern countries with little evidence of changes in most of the countries. The gap in AFS remained large and widened over time in the north while remained small and narrowed over time in the south. In the Western Africa region, the gap of more than two years in median AFS between male and female in many countries suggested more disparity in sexual partner mixing compared to countries in the south. Large gender gaps suggested adolescent girls could remain vulnerable to poor sexual health outcomes (43).

The median AFS has increased in many parts of sub-Saharan Africa over the past three decades, especially in the northern-most countries, but the timing and magnitude of changes varied. In countries heavily affected by HIV/AIDS epidemic in southern and eastern Africa in which there has been a multitude of intervention programmes focused on reducing sexual risk, median AFS only slightly increased or even decreased. Other analyses have reported that trends towards earlier sexual debut are not pronounced (44) and median AFS in SSA increased over time (45). Previous findings from individual countries also found similar results as we have reported such as a small increase in men's AFS in Zambia (25) and both sexes in Uganda (46). In contrast, AFS increased more steeply in western Africa. These changes could be related to large gains in female education (42) leading to delaying age at marriage in female while modern life's economic uncertainty pressuring male to postponing age at marriage (47). These combined with stricter religious norms of abstaining sex until marriage in the region may have contributed to larger increases in AFS compared to other regions.

Reproductive health and HIV monitoring indicators have tended to focus on median AFS in a population or the proportion initiating sexual activity by specified ages. Our analysis calculated the full distribution of debuting sexual activity, reflecting the range of ages at which adolescents and young adults may need to become engaged in sexual health, family

planning, and HIV/STI prevention services. The rate at which populations become sexually active varied across countries, but typically was slightly more peaked for young women. For young women, the percentage sexually experienced increased from 25% to 75% in around four years compared to five years for men. This varied across countries, however. In some, sexual debut of a population occurred in a very short span of age while in others in a gradual manner (Fig 1). Two populations with the same median AFS, but different shapes parameters may have different demand for reproductive health and STIs services. Programming prevention services efforts should be directed accordingly to deliver services to the crucial period of a sexually active life.

Previous research using longitudinal data or sexual activity biomarkers has identified reporting biases in AFS that can obfuscate trends assessed from cross-sectional surveys (23,43,45,48). This study addressed this bias by comparing reports among the same cohort in successive surveys. Young men tended to report earlier AFS than when asked again in later surveys, while young women tended to report older AFS. Above age 30, the reported AFS tended to increase as the cohort aged, especially for men, indicative of recall biases for exact age for events that occurred long ago. These results are consistent with previous findings documenting biased self-reporting of sexual activity by norms that boys are expected to engage in sexual activity and girls are expected to abstain until marriage (20)(24,26,27). Other country-specific analyses have also documented that respondents were less likely to report a young AFS in later surveys (25). While adjusting for these biases to reconstruct consistent time trends is a strength of our model, the variation in magnitude and direction of biases across countries implies caution about confidently distinguishing actual trends from changing reporting biases among the youngest cohorts in new surveys. This will be especially challenging in settings with fewer surveys from which to triangulate across different time points in life of a birth cohort (20).

This study has several limitations. First, while we were able to adjust for relative reporting differences by age at report, there was no 'gold standard' measure to identify at which age

reporting was most accurate. The chosen reference age of twenty-three is at most ten years difference from the AFS and above the average age of marriage of women in SSA, thus it is expected that reports at this age are less influenced by social desirability compared to a younger age and recall more accurate compared to older reporting age. In addition, as digit preference for age reporting ending in zero or five is a common issue (49), using ages around 25 or 30 might introduce biases to the reported estimates. Second, country- and sex-specific differences of potential confounders to median AFS such as religion (50) were not modelled. Third, our model allowed the median AFS to change over time, and for the shape of the distribution and reporting bias to vary by country but did not allow the shape or reporting bias to change over time within each country. Fourth, survey series varied slightly in the eligibility of respondents asked about AFS, the way AFS was recorded, and data processing procedures. For example, respondents older than 24 were not asked AFS in some in MICS or FHS. In DHS it standard practice to recode internally inconsistent data such as reported AFS that is older than the age at the time of survey, after the conception of the first child, or after the first marriage (45), but this is not done in all surveys, as evidenced in larger proportions of excluded observations (S1 Table). Fifth, we accounted for survey design by incorporating survey weights into the model pseudo likelihood (38), but did not model stratified clustered sampling designs. Finally, results in this paper are limited to SSA at national level. Similar methods to ours could be applied for subnational estimates and globally with available age at first sex data.

In conclusion, AFS has increased slightly in most, but not all, SSA countries. However, in most cases changes over-time within a country were small compared to large and relatively persistent variation between countries and genders, reflecting deeply rooted cultural norms around sexual debut. These information on patterns of initiating sexual activity are essential for planning sexual health, sexual education, family planning, STI, and HIV prevention services. Incorporating our estimates of age- and sex-specific sexual debut rates into epidemic models of HIV and STIs may also help to shed light on national and regional

340 variation in epidemic burdens and transmission dynamics, for example the role of the larger
341 age gaps in west and middle African countries compared eastern and southern regions.

Declarations

Ethics approval and consent to participate

The study involved secondary analysis of anonymised publicly available data following project approval from the Demographic and Health Surveys Program, the Performance Monitoring for Action project, the Population-based HIV Impact Assessment project, and South African National HIV Prevalence, Incidence, Behaviour and Communication Survey. Primary survey protocols were reviewed and approved by the relevant national ethics committee in each country.

Consent for publication

Not applicable

Availability of data and materials

The datasets analysed during the current study are available in the Demographic and Health Surveys Program (<https://dhsprogram.com/Data/>), the Performance Monitoring for Action project (<https://www.pmadata.org/data>), the Population-based HIV Impact Assessment project (<https://phia-data.icap.columbia.edu>), and South African National HIV Prevalence, Incidence, Behaviour and Communication Survey (<http://www.hsrb.ac.za/>).

Competing interests

The Authors declare that there is no conflict of interest.

Funding

This research was supported the Bill and Melinda Gates Foundation (OPP1190661), National Institute of Allergy and Infectious Disease of the National Institutes of Health under award numbers R01AI136664 and R01AI152721, and the MRC Centre for Global Infectious Disease Analysis (reference MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and the UK Foreign, Commonwealth & Development Office (FCDO), under the MRC/FCDO Concordat agreement and is also part of the EDCTP2 programme

supported by the European Union. The funders have no roles in the study design or manuscript preparation.

Authors' contributions

VKN and JWE conceived the work. VKN and JWE designed the work. VKN implemented the models. Both authors critically reviewed model results throughout the model development process. VKN wrote the first draft of the manuscript. Both authors critically edited the manuscript for intellectual content.

Acknowledgements

This research was supported the Bill and Melinda Gates Foundation (OPP1190661). We acknowledge joint MRC Centre funding from the UK Medical Research Council and Department for International Development via MRC MR/R015600/1.

Supplemental Materials

S1 Text. Skew log-logistic distribution & specifications of the survival model

S1 Fig: Parameter distribution correlation

S2 Fig: Spatial distribution of the percentage ever had sex under fifteen.

S3 Fig: Proportion ever had sex and fitted value by country and sex

S1 Table. List of surveys used in the analyses

S2 Table. Parameter estimates by country and sex

S3 Table. Proportion ever had sex under 15 & 18 by country and sex

References

1. WHO. Report on global sexually transmitted infection surveillance 2018. Who. 2018.
2. Pettifor AE, van der Straten A, Dunbar MS, Shiboski SC, Padian NS. Early age of first sex. AIDS. 2004;
3. Kaestle CE, Halpern CT, Miller WC, Ford CA. Young age at first sexual intercourse and sexually transmitted infections in adolescents and young adults. Am J Epidemiol. 2005 Apr;161(8):774–80.
4. Hallett TB, Gregson S, Lewis JJC, Lopman BA, Garnett GP. Behaviour change in generalised HIV epidemics: Impact of reducing cross-generational sex and delaying age at sexual debut. Sex Transm Infect. 2007 Aug 1;83(SUPPL. 1):i50-54.
5. Wand H, Ramjee G. The relationship between age of coital debut and HIV seroprevalence among women in Durban, South Africa: a cohort study. BMJ Open [Internet]. 2012 Jan 5 [cited 2020 Nov 20];2(1). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3253418/>
6. Louie KS, De Sanjose S, Diaz M, Castellsagué X, Herrero R, Meijer CJ, et al. Early age at first sexual intercourse and early pregnancy are risk factors for cervical cancer in developing countries. Br J Cancer. 2009 Apr 10;100(7):1191–7.
7. Kaplan DL, Jones EJ, Olson EC, Yunzal-Butler CB. Early Age of First Sex and Health Risk in an Urban Adolescent Population. J Sch Health. 2013;
8. Upchurch DM, Levy-Storms L, Sucoff CA, Aneshensel CS. Gender and ethnic differences in the timing of first sexual intercourse. Fam Plann Perspect. 1998 Jun;30(3):121–7.
9. Wróblewska W. Religiosity differences in the transition to first intercourse before age 18 among Polish students. Genus. 2020 Jul 8;76(1):14.
10. Kågesten A, Blum RWm. Characteristics of Youth Who Report Early Sexual Experiences in Sweden. Arch Sex Behav. 2015 Apr 1;44(3):679–94.
11. Zaba B, Pisani E, Slaymaker E, Boerma JT. Age at first sex: Understanding recent trends in African demographic surveys. Sex Transm Infect. 2004;80(SUPPL. 2):28–35.
12. Gregson S, Garnett GP, Nyamukapa CA, Hallett TB, Lewis JJC, Mason PR, et al. HIV decline associated with behavior change in eastern Zimbabwe. Science. 2006 Feb 3;311(5761):664–6.
13. Halperin DT, Steiner MJ, Cassell MM, Green EC, Hearst N, Kirby D, et al. The time has come for common ground on preventing sexual transmission of HIV. The Lancet. 2004 Nov;364(9449):1913–5.
14. UNAIDS. Monitoring the Declaration of Commitment on HIV/AIDS: guidelines on construction of core indicators.
15. Dehne, K. L., Gabriele, Riedner. Sexually transmitted infections among adolescents : the need for adequate health services.

- 423 16. Aisha NZD, Philipp U, Vladimíra K. Sexual Activity by Marital Status and Age: A
424 Comparative Perspective, Technical Paper No. 11. 2017;
- 425 17. Assche SB Van. Are we measuring what we want to measure? Individual consistency
426 in survey response in rural Malawi. *Demogr Res.* 2003 Sep 19;9(SUPPL. 1):77–108.
- 427 18. Catania JA. A framework for conceptualizing reporting bias and its antecedents in
428 interviews assessing human sexuality. Vol. 36, *Journal of Sex Research*. Taylor &
429 Francis Group; 1999. p. 25–38.
- 430 19. Wellings K, Collumbien M, Slaymaker E, Singh S, Hodges Z, Patel D, et al. Sexual
431 behaviour in context: a global perspective. *Lancet Lond Engl.* 2006 Nov
432 11;368(9548):1706–28.
- 433 20. Gersovitz M. HIV, ABC and DHS: age at first sex in Uganda. *Sex Transm Infect.* 2007
434 Apr 1;83(2):165–8.
- 435 21. United Nations. World population monitoring: adolescents and youth. U N. 2012;1.
- 436 22. Eggleston E, Leitch J, Jackson J. Consistency of Self-Reports of Sexual Activity among
437 Young Adolescents in Jamaica. *Int Fam Plan Perspect.* 2000;26(3):79–83.
- 438 23. Wringe A, Cremin I, Todd J, McGrath N, Kasamba I, Herbst K, et al. Comparative
439 assessment of the quality of age-at-event reporting in three HIV cohort studies in sub-
440 Saharan Africa. *Sex Transm Infect.* 2009;85(SUPPL. 1).
- 441 24. Zaba B, Pisani E, Slaymaker E, Boerma JT. Age at first sex: Understanding recent
442 trends in African demographic surveys. *Sex Transm Infect.* 2004;80(SUPPL. 2):28–35.
- 443 25. Slaymaker E. Monitoring trends in sexual behaviour in Zambia, 1996-2003. *Sex*
444 *Transm Infect.* 2004 Dec 1;80(suppl_2):ii85–90.
- 445 26. Mensch BS, Hewett PC, Erulkar AS. The reporting of sensitive behavior by
446 adolescents: A methodological experiment in Kenya. *Demography.* 2003
447 May;40(2):247–68.
- 448 27. Luke N, Clark S, Zulu EM. The relationship history calendar: improving the scope and
449 quality of data on youth sexual behavior. *Demography.* 2011 Aug;48(3):1151–76.
- 450 28. Cremin I, Mushati P, Hallett T, Mupambireyi Z, Nyamukapa C, Garnett GP, et al.
451 Measuring trends in age at first sex and age at marriage in Manicaland, Zimbabwe. *Sex*
452 *Transm Infect.* 2009 Apr 1;85(Suppl 1):i34–40.
- 453 29. Pettifor AE, van der Straten A, Dunbar MS, Shiboski SC, Padian NS. Early age of first
454 sex. *AIDS.* 2004;
- 455 30. Nguyen VK, Eaton JW. A model for reconstructing trends and distribution in age at first
456 sex from multiple household surveys with reporting biases [Internet]. 2021 Dec [cited
457 2021 Dec 15] p. 2021.12.14.21267770. Available from:
458 <https://www.medrxiv.org/content/10.1101/2021.12.14.21267770v1>
- 459 31. ICF. Demographic and Health Surveys (various). Funded by USAID. Rockville,
460 Maryland: ICF [Distributor]. 2020.
- 461 32. UNICEF. Multiple Indicator Cluster Surveys (MICS). 2020.

33. Elizabeth Heger Boyle, Devon Kristianse, Matthew Sobek. IPUMS PMA: Version 4.1 [dataset]. Minneapolis, MN: IPUMS. [Internet]. 2020. Available from: <https://doi.org/10.18128/D081.V4.1>
34. UNICEF. Eritrea Population and Health Survey. 2020.
35. PHIA Collaborating Institutions. Population-based HIV Impact Assessment (PHIA) [Internet]. 2020. Available from: <https://doi.org/10.18128/D081.V4.1>
36. MEASURE Evaluation Project. Zambia sexual behaviour surveys. [Internet]. 2020. Available from: <https://doi.org/10.18128/D081.V4.1>
37. The Human Sciences Research Council. South African National HIV Prevalence, Incidence and Behaviour Survey. [Internet]. 2020. Available from: <https://doi.org/10.18128/D081.V4.1>
38. Asparouhov T. General Multi-Level Modeling with Sampling Weights. *Commun Stat - Theory Methods*. 2006 Apr;35(3):439–60.
39. Van Kinh Nguyen, Jeffrey W Eaton. Model the age at first sex distribution and age at reporting bias in household survey data. *medRxiv*. 2021;
40. Besag J, Kooperberg C. On Conditional and Intrinsic Autoregression. *Biometrika*. 1995;82(4):733–46.
41. Kristensen K, Nielsen A, Berg CW, Skaug H, Bell BM. TMB: Automatic differentiation and laplace approximation. *J Stat Softw*. 2016;70.
42. Barro Robert, Jong-Wha Lee. A New Data Set of Educational Attainment in the World, 1950-2010. *J Dev Econ*. 2013;104:184–98.
43. Melesse DY, Mutua MK, Choudhury A, Wado YD, Faye CM, Neal S, et al. Adolescent sexual and reproductive health in sub-Saharan Africa: who is left behind? *BMJ Glob Health*. 2020 Jan 1;5(1):e002231.
44. Wellings K, Collumbien M, Slaymaker E, Singh S, Hodges Z, Patel D, et al. Sexual behaviour in context: a global perspective. *Lancet Lond Engl*. 2006 Nov;368(9548):1706–28.
45. Slaymaker E, Scott RH, Palmer MJ, Palla L, Marston M, Gonsalves L, et al. Trends in sexual activity and demand for and use of modern contraceptive methods in 74 countries: a retrospective analysis of nationally representative surveys. *Lancet Glob Health*. 2020 Mar 9;8(4):e567–79.
46. Slaymaker E, Bwanika JB, Kasamba I, Lutalo T, Maher D, Todd J. Trends in age at first sex in Uganda: evidence from Demographic and Health Survey data and longitudinal cohorts in Masaka and Rakai. *Sex Transm Infect*. 2009 Apr 1;85(Suppl 1):i12–9.
47. Smith DJ. Masculinity, Money, and the Postponement of Parenthood in Nigeria. *Popul Dev Rev*. 2020;46(1):101–20.
48. Plummer ML, Ross DA, Wight D, Chagalucha J, Mshana G, Wamoyi J, et al. 'A bit more truthful': The validity of adolescent sexual behaviour data collected in rural northern Tanzania using five methods. In: *Sexually Transmitted Infections*. 2004. p. ii49–56.

- 502 49. Amos M, Stones T. Trends in Demographic and Health Survey data quality: An
503 analysis of age heaping over time in 34 countries in Sub Saharan Africa between 1987
504 and 2015. BMC Res Notes. 2017 Dec 1;10.
- 505 50. Westoff, Charles F. K Bietsch. Religion and Reproductive Behavior in Sub Saharan
506 Africa. DHS Anal Stud No 48 [Internet]. 2015 [cited 2020 Nov 13]; Available from:
507 <https://dhsprogram.com/pubs/pdf/AS48/AS48.pdf>
- 508

Figures

Figure 1. Distribution of the age at first sex by region, country, and sex. The median of each distribution is shown with the vertical lines. The estimates are for the birth cohort turn fifteen in 2015 and reporter age 23. Within each region, countries sequenced by increase median female AFS.

Figure 2. Percentage ever had sex by age 18 among the birth cohort turning 15 in the year 2015. The two countries Comoros and São Tomé & Príncipe are represented in the bottom left. Countries without data are coloured the same as the background

Figure 3. Trend in median age at sexual debut for men and women in Sub Saharan-Africa countries by year turned age 15 (x-axis). For countries where the most recent survey is before 2015, trends are extrapolated for recent birth cohorts based on the overall trend across the region and its neighbours. Countries without AFS data are excluded in this figure.

Figure 4. Aggregated trend of median age at first sex by region for men and women. Sudan was included in Eastern Africa. The regional trend is a weighted average of the countries included in the analyses by their respective population size in 2020 sourced from the UN World Population Prospects.

Figure 5. Direction of the effect of age at report bias on the median age at first sex estimate relative to other ages. A positive coefficient in the figure corresponds to an increase in median AFS estimate reported by the respective age compared to a median AFS reported by the age with that coefficient equals zero.