A survey assessing antimicrobial prescribing at UNRWA primary health care centers in Jordan.

**Orubu ESF^{1,2*}, Albeik S³, Ching C², Hussein R², Mousa A³, Horino M³, Naqa R³, Elayyan M³,
Saadeh R³, Zaman MH²**

¹. Social Innovation on Drug Resistance, Boston University, Boston, United States of America

². Department of Biomedical Engineering, College of Engineering, Boston University, Boston,
United States of America

³. United Nations Relief and Works Agency for Palestinian Refugees in the Near East, UNRWA,
Jordan

* Corresponding author. Email: sforubu@bu.edu

Key words: antimicrobial resistance, stewardship, humanitarian, primary health care, UNRWA

Word count: Abstract 247 words; Text 3031

Number of tables: 5

Number of figures: 3

Supplemental materials: Appendix A – Questionnaire; Appendix B – Results, additional

Running Head: Antimicrobial resistance, use and stewardship in Jordan

Abstract

Antimicrobial resistance (AMR) is a public health emergency. There is insufficient information on AMR in the context of humanitarian settings. An understanding of behavioural and institutional level factors can strengthen antimicrobial stewardship. This study used a semi-structured questionnaire to assess both knowledge, attitudes and practices (KAP) on antimicrobial use, resistance and stewardship, and options to improving prescribing, among prescribers at the Primary Healthcare facilities of the UNRWA Jordan field office. Responses to the KAP questions were evaluated using the Capability, Opportunity, Motivation, Behaviour (COM-B) framework and Bloom's cutoffs. For each framework component, Blooms cutoffs and interpretations were: >80%, "good"; 60-79%, "moderate"; and <60% "poor". Fourteen options to improve prescribing were each assessed using 5-point Likert scales from very unhelpful to very helpful, aggregated by helpful and very helpful and ranked as: >90%, best/most acceptable; >80-90%, as acceptable; and, 70-80% as maybe acceptable/good. The questionnaire response rate was 59% (37/63) with a completion rate of 92% (34/37). Aggregate scores for real knowledge on AMR was 97%; opportunity to improve prescribing 88%; and motivation 16% - participants did not believe that there was a connection between their prescribing and AMR or that they had a key role in helping control AMR. Good options (74% aggregate score) to improving prescribing were the availability of guidelines and resistance data. There was good knowledge of AMR and good opportunities, but poor motivation for rational prescribing or behavioral change. There is a clinical need for antimicrobial resistance data to promote rational antibiotic prescribing.

Introduction

Antimicrobial resistance (AMR), an inherent or acquired ability of disease-causing microorganisms to survive therapeutic doses of antimicrobials is a pressing global health challenge^{1,2}. Anthropogenic factors including the non-evidence-based prescribing, dispensing and use of antimicrobials including for non-therapeutic indications in livestock, the use of substandard or falsified antimicrobials, poor adherence in patients and environmental (metal and) antibiotic run-offs accelerate AMR³. Low- and Middle-Income Countries (LMICs) are predicted to bear the greater burden of AMR⁴. Generally, communities living in cramped environments with limited resources are at a high risk of developing drug resistant infections³. This may be particularly true for refugees and the forcibly displaced, where geo-socio-economic determinants predisposing to disease combine with sub-optimal access and the excessive use of antibiotics⁵⁻⁷.

Jordan is one of the world's largest hosts for displaced populations. In 2020, it was the ranked the 10th largest host country for refugees and the internally displaced⁸. Jordan, historically, also has the highest number – over 2.3 million – of registered Palestine refugees, (PRs) under the United Nations Relief and Works Agency for Palestine Refugees in the Near East (UNRWA) program among the five fields of its operations (Gaza, the West bank, Syria, Lebanon, and Jordan). In Jordan, UNRWA provides free health care services (including outpatient services) to Palestine refugees through a network of 25 primary health care centers. There are scant studies on antibiotic utilization among PRs settlements in Jordan. One previous study from 2018 on the knowledge, attitude and behavior of PRs regarding antibiotics surveyed 250 adults at 4 different UNRWA health centers (HCs) in Jordan (Irbid, South Baqaa, Taybeh and Marka)⁹. The study revealed a poor understanding of antibiotics. Self-medication, over-the-counter (OTC) purchase as well as using left-over antibiotics were commons form of inappropriate antibiotics use. Further, there was a knowledge gap in the correct use, and of side effects from irrational use, of antibiotics among participants. Long waiting times drive the OTC purchase of antibiotics, even though there is high trust in doctors among patients in refugee camps⁹. This data suggests that the risk for AMR among refugees, arising from misuse of antibiotics, is high. The study, however, did not identify which specific antibiotics were most commonly used and did not consider the prescribing and dispensing practices of healthcare workers and pharmacists. Thus, there are gaps in our understanding.

This study, as part of a series assessing refugees' access to quality healthcare services and AMR, performs an evaluation within the context of a primary health care system. The aim was to assess and analyze antimicrobial prescribing in the Jordanian UNRWA sites. The objectives of the study were two-fold: (i) describe antibiotic prescribing patterns, as proxies of use (ii) assess knowledge, attitude, and practices on antimicrobial resistance and use, with a view to understanding barriers to antimicrobial stewardship.

Methods

Ethics: Ethics application was reviewed by the Institutional Review Board of Boston University and granted exemption (5681X). Ethical approval was also obtained from UNRWA's research review board before questionnaire administration.

Study design: Cross-sectional electronic survey design. This study adopted the Checklist for Reporting Results of Internet E-Surveys (CHEERIES) protocol for questionnaire development and reporting¹⁰.

Study setting: Primary Health Care (PHC) facilities run by the UNRWA in Jordan. A convenience sample of 16 PHCs were selected based on size and accessibility¹¹.

Participants' inclusion and exclusion criteria: Only medical officers, gynecologists, and dental surgeons working at the selected UNRWA PHCs who prescribe antimicrobials were included in the survey. All other healthcare workers at the selected PHCs and all healthcare workers outside these facilities were excluded.

Study instrument: The study instrument was a semi-structured questionnaire based primarily on a previously validated knowledge, attitude, and practice survey used in Europe¹². This was complemented by other previously used survey instruments^{13,14} further adapted to a humanitarian setting by including questions assessing prescribers' perceptions of AMR in the forcibly displaced¹⁵.

The questionnaire comprised 4 sections containing 43 questions assessing demographics, current prescribing behavior, and potential for behavioral change (Table 1 and Supplement 1). Section I had 8 questions on demographics. Section II included 16 questions to describe current prescribing practices and antibiotic prescribing patterns. Section III contained 16 questions assessing knowledge and attitudes. These questions were structured using the Capability, Opportunity and Motivation for behavioral change (COM-B) framework. Capability assessed prescribers' knowledge, awareness and perceptions or attitudes. In this study, we described capacity as consisting of decision making, knowledge, awareness/perceptions, attitudes, and infection prevention and control measures (hand hygiene). Knowledge was both real and assumed. Opportunity included perceptions about both individual- and facility-level factors that make for rational prescribing. Motivation was self-held beliefs about influence on rational prescribing and association between prescribing behavior and AMR. Section IV contained 1 question with 14 options to improve antimicrobial prescribing. There were also 1 introductory question on consent and 1 final question on additional comments.

Items in the questionnaires were randomized. Using adaptive questioning, certain questions were only required to be answered based on whether a preceding question was answered. To ensure completeness when filling the questionnaires, some questions were pre-specified and made mandatory, where these questions had to be filled before the questionnaire could be submitted. The questionnaires were made available in English and Arabic. The Arabic version

was provided by SH and verified by SA. Both versions incorporated an introductory information and consent section.

Piloting: The questionnaire was piloted in a PHC not among those selected to be sampled. Based on the results from this pilot, the instrument was modified to ensure validity in the study locations. Modifications included changes to the Arabic translation, a consent section with termination where consent was not granted, and the review step to ensure completeness.

Sample size: The sample size was all consenting physicians out of a list of 63 prescribers provided by Jordan field office at the 16 selected PHCs.

Questionnaire administration: Questionnaires were self-administered. Participants were contacted through a list of emails and/or WhatsApp numbers for all medical officers currently working in the selected 16 PHCs in Jordan. The questionnaires were shared (as links) with the participants. Consent was obtained from all participants by ensuring that only those who consented continued to fill the questionnaire. Questionnaires were administered from August 1 to September 30, 2021.

Data analysis: Data analysis was descriptive with responses expressed as proportions (%). Current prescribing practices were described. Aggregate scores across participants in the capability (real knowledge), opportunity and motivation components of behavior were computed and evaluated against Bloom's cutoffs. For real knowledge, opportunity and motivation a mean aggregate score >80% was assessed as "good", 60-79% as "moderate", and <60% as "poor"¹⁶. The perceived options to improve antibiotic prescribing were aggregated and ranked. On the 5-point Likert Scale, "very helpful" and "helpful" were aggregated into the new dummy variable "helpful" and neutral, unhelpful and very unhelpful aggregated into a new "unhelpful" dummy variable. The aggregated scores under the new "helpful" category for each of the 14 options to improve prescribing were then ranked with aggregate scores >90% assessed as the best, or most acceptable, option(s); >80-90%, as acceptable; and, 70-80% as maybe acceptable, or good.

Results

Characteristics: response rate and participants demographics

The questionnaire response, or participation, rate was 59% (37/63). However, the completion rate, or participants consenting to, and completing, the survey was 92% (34/37).

Participants were mostly general medicine specialists (75%, 25/34). The majority had more than 6 years' experience (35% with 6-10 years' experience; 38% with 11-15; and 9% with >15 years).

The complete demographic characteristics and detailed results are included in the supplement.

Antibiotic prescribing practices

All respondents had prescribed antibiotics in the 6 months preceding the study. All prescriptions were by generic names (100%, 34/34). Prescribers (91%, 31/34) reported the most frequently used antibiotic as amoxicillin (including amoxicillin-clavulanate), a penicillin (Fig. 1A). There was specificity, or a differentiation, to antibiotic classes according to disease category. The top antibiotic class for respiratory tract infections (RTIs), urinary tract infections (UTIs), gastrointestinal tract infections (GIT), and dental cases, were, penicillins, sulfonamides, macrolides and sulfonamides, and macrolides, respectively (Fig. 1B).

Physicians (65%, 22/34) report pressure to prescribe antibiotics. The reasons for this pressure, ranked from high to low, among participants who provided a reason, were the patients' condition, 78% (14/18); perceived patient demand, 60% (12/20); fear of complications, 50% (8/16); and need to provide rapid relief, 47% (7/15).

The main considerations for prescribing antibiotics were the patients' condition (94%, 32/34) and case clinical severity (62%, 21/34). The results from Antimicrobial Sensitivity Testing (AST) were a consideration only in about a third of respondents (29%, 10/34) – this being a Primary level facility without institutional AST.

Antibiotic choice was primarily determined by availability (87%, 26/30). Secondary considerations were inclusion in the Standard Treatment Guidelines (STG) or formulary (96%, 22/23); antibiotic spectrum (86%, 6/7); and patient symptoms (69%, 11/16). About one third (29%, 10/34) of respondents usually prescribe 2-3 antibiotics per antibiotic prescribing episode.

Generic prescribing, in line with existing protocols, was practiced by all prescribers (100%, 34/34). However, slightly over three quarters (76%, 26/34) also reported the practice of prescribing by brand names (likely outside the UNRWA system). The reasons for prescribing by brand was the perception that these are more efficacious (85%, 22/26) and/or of better quality (42%, 11/26) than generics.

The use of STG for prescribing was followed by 94% (32/34). There was limited AST capacity (Fig. 2). While about half (56%, n=34) responded that there was the capacity to perform AST, this was almost always (90%, n=34) off-site.

Capability, opportunity and motivation for behavior change

Knowledge: There was a high real knowledge of AMR among participants. The great majority of prescribers correctly assessed the 7 knowledge statements (Table 2). All (100%, 34/34) knew that antibiotics were ineffective against viruses, including those causing the common cold and flu. Similarly, all (100, 34/34) had knowledge that every person treated with antibiotics is at an increased risk for developing an antibiotic resistant infection. The correct responses to the remaining 5 questions ranged from 82-97% (Table 2). The greatest relative knowledge gap was

the fact that antibiotic resistance can spread from person to person (82% correct answer, n=34)¹⁷.

Awareness: There was good awareness of the WHO 5 moments of hand-hygiene. All but one (97%, 33/34) could list the 5 moments. In contrast, there was poor awareness of the WHO AWaRe classification of antibiotics. Among participants, 41% (14/34) had heard of this classification.

Attitude: There was good attitude towards Infection Prevention and Control (hand hygiene) measures. All (100%, 34/34) affirmed the need to perform hand hygiene as often as recommended after contact with a patient or biological material.

Perceptions: Antimicrobial resistance was perceived to be a problem at both the PHC and country-level (Fig. 3). Four out of every five (79%, n=34) participants thought AMR was a problem at their PHC. Almost all 91% (31/34), thought AMR to be a problem in Jordan.

Antimicrobial resistance was perceived as a high or very high public health challenge in Jordan. Most (88%, 30/34) rated the AMR challenge as either very high or high. Refugees were perceived by 85% of respondents (n=34) to have a higher risk of AMR.

Increasing awareness among patients was perceived by almost all (94%, 32/34) prescribers as a solution to the problem of AMR in Jordan. Other top suggestions were enforcing prescription laws (85%, 29/34 of prescribers), increased AST (47%, 16/34); and training (38%, 13/34).

Opportunity: The aggregate opportunity score was 88%, indicating good opportunity for rational prescribing (Table 3).

Motivation: There was poor motivation among participants (Table 4). Less than 1 in 5 either believed that there was a correlation between their antibiotic prescribing and AMR (18%, 6/34), or that they have a key role in the control of AMR (15%, 5/34).

Perceived solutions to improving prescribing

Among 14 options for improving prescribing, only 3 were “maybe acceptable” (Table 5). These were: availability of local guidelines (74%); availability of resistance data (74%); and educational sessions on prescribing (71%).

Discussion

Establishing needs through assessing behavioral, organizational and social barriers (including knowledge and awareness) and enablers provides evidence to design context-specific interventions aimed at curtailing the continuous spread of AMR. This study, conducted together with UNRWA, Jordan field of operation, provides a baseline evaluation to establish the need and parameters for intervention to designing, implementing and evaluation and monitoring of an effective AMS program at UNRWA.

In 2017, The Hashemite Republic of Jordan, in line with the global recommendation of the World Health Organization and the United Nations, formulated its National Action Plan to combat AMR for the period 2018-2022¹⁸. This plan, one of few that are fully costed, demonstrates the political commitment at a national level to addressing AMR in Jordan. The goals in this plan include a 30% reduction in antimicrobial use in animals, a 20% reduction in use in humans, and a 40% increase in public knowledge on AMR and awareness of appropriate use of antimicrobials. These goals are operationalized in two strategic objectives to improve knowledge on AMR and optimize use through AMS. These goals and objectives are shared by UNRWA.

Similarly, the UNRWA health department has a stated policy to maintain antibiotic prescription rates to less than 25%, reflecting earlier WHO recommendations on “optimum” antimicrobial medicines use/stewardship. In 2020, the Jordan field office reported a 16.6% antibiotic prescription rate, down from 21.3% in 2019, largely due to the impact of COVID-19 leading to a decrease in outpatient consultations and rational medicines prescription^{19,20}.

There was a high level of awareness of AMR among prescribers. By far a majority of participants rated AMR to be a challenge in Jordan as well as in their PHC. In addition, participants were of the opinion that refugees are at a higher risk of AMR than the general population. A few studies seem to suggest high carriage of AMR among the forcibly displaced probably as a result of certain socioeconomic factors^{21,22}.

Overall, measures for antibiotic prescribing align with the WHO/INRUD recommendations. Most prescribers prescribe only 1 antibiotic per patient encounter or consultation; there is generic prescribing; and the formulary or STG guides antibiotic selection. The UNRWA field office has an electronic system that uses generic names and allows prescribing based only on the included generic medicines as derived or selected from the WHO Essential Medicines List.

The antibiotic prescribing pattern at UNRWA’s PHC Jordan field seem to mirror the prescribing pattern in the Middle East/North Africa (MENA) geographical region, with amoxicillin (including amoxicillin/clavulanate) being the most commonly prescribed antibiotic in Yemen, as well as in other states¹⁵. In Jordan, this is an established trend. A 2007 study by Al-Niemat et al, found amoxicillin to be the most frequently prescribed²³. These results also agree broadly with a similar study conducted earlier in 2014 across five tertiary or referral outpatient public hospital pharmacies in Jordan which found amoxicillin to be the most prescribed antibiotic, followed by ciprofloxacin²⁴. Current national level antimicrobial consumption records also show amoxicillin and amoxicillin/clavulanate to be the most frequently consumed antibiotics in Jordan, accounting for 47% of total antibiotic consumption in Defined Daily Doses per 10, 000 population (DDD) in 2019 and 32% in 2020²⁵. The fall in 2020 was attributed to both COVID-19 and a simulated increase in other classes of antibiotics, compared to amoxicillin and the clavulanate.

The implications of this use pattern of amoxicillin on AMR, or clinical effectiveness, is not known. There is no information on the Global Antimicrobial Sensitivity Surveillance System (GLASS) for amoxicillin resistance. However, studies indicate very high levels of resistance (>80%) to this antibiotic in cases of Urinary Tract Infections, including those caused by Extended Spectrum Beta-Lactamase (ESBL) producing bacteria, in hospitalized children for the period 2012-2017²⁶. A more recent, smaller, study found *P. aeruginosa* to be 100% resistant to amoxicillin²⁷.

Antimicrobial prescribing, in practice, is a multi-layer exercise whereby the prescriber may have to balance certain considerations to resolve a case. These considerations are largely context specific. In this study, we found that physicians prioritized the condition of the patient in prescribing with the majority considering the condition of the patient in almost all cases, and or the clinical severity.

In line with common global practice, though, antibiotics were prescribed empirically, with limited considerations for AST. The apparent low prioritization of AST by prescribers is similar to another study in Yemen, though in this Jordan study, a higher proportion of prescribers rated as important the use of AST to guide rational prescribing than in Yemen, suggesting a higher level of awareness²⁸. The UNRWA PHC, it is to be noted, do not provide for AST services, being functionally out-patient level facilities. While there are no institutional AST services, some patients may go to private sector to do AST (as the Jordan field patients can access health services from both public and private sectors).

However, there may be a need to introduce AST or rapid diagnostic device services at the PHC to account for resistance rates. For example, results from GLASS, 2018, show that cotrimoxazole, the most commonly prescribed antibiotic for UTIs in this study would not be effective in 53% of cases. One study on antimicrobial use and resistance in a tertiary hospital in Jordan found an overall resistance prevalence of 26% across a panel of antibiotics including amoxicillin and the others commonly prescribed in this study in 33 patients receiving an antibiotic for a range of infections²⁹.

The use of a formulary, STG, or Essential Medicines List (EML) to guide antibiotic prescription is a AMS recommendation. This study suggests that prescribers follow this guideline in largely making decisions on choice of antibiotic to prescribe from the guidelines. Indeed, the UNRWA PHCs use an electronic prescribing and dispensing system that prevents prescribing outside this system. However, as the illustration with amoxicillin use and resistance shows, there is need for underlying dynamic resistance data to better guide antibiotic choice.

Conclusion

This study assessed the knowledge, attitude and practices of prescribers at the UNRWA primary healthcare facilities in Jordan. There was a high knowledge of AMR and good opportunities, but

poor motivation for rational prescribing or behavioral change. There is a clinical need for antimicrobial resistance data to promote rational antibiotic prescribing.

Acknowledgments

This work was made possible by support from the Boston University Social Innovation on Drug Resistance (SIDR) Postdoctoral Program to ESFO and the U.S. Pharmacopeial Convention (USP) Quality Institute to CC.

Financial Support

We confirm no financial support was received for this project.

Disclosures regarding real or perceived conflicts of interest.

None to declare.

Co-Author Contact Information.

1. Albeik S; Department of health, headquarters, Amman, Jordan; S.albaik@unrwa.org
2. Ching C; Department of Biomedical Engineering, College of Engineering, Boston University, Boston, United States of America; chingc@bu.edu
3. Hussein R; Department of Biomedical Engineering, College of Engineering, Boston University, Boston, United States of America; ranahu@bu.edu
4. Mousa A; Department of health, headquarters, Amman, Jordan; ay.mousa@unrwa.org
5. Horino M; Department of health, headquarters, Amman, Jordan; m.horino@unrwa.org
6. Naqa R; Department of health, Jordan Field office, Amman, Jordan; r.naqa@unrwa.org
7. Elayyan M; Department of health, Jordan Field office, Amman, Jordan; m.elayyan@unrwa.org
8. Saadeh R ; Department of health, headquarters, Amman, Jordan; r.saadeh@unrwa.org
9. Zaman MH; Department of Biomedical Engineering, College of Engineering, Boston University, Boston, United States of America; zaman@bu.edu

402
403
404
405
406
407

References:

1. United Nations., 2018. Antimicrobial resistance a “global health emergency,” UN, ahead of awareness week. Available at: <https://news.un.org/en/node/1025511/antimicrobial-resistance-a-global-health-emergency-un-ahead-of-awareness-week-2>. Accessed. November 2018
2. Toner E, Adalja A, Gronvall GK, Cicero A, Inglesby TV., 2015. Antimicrobial resistance is a global health emergency. *Health Secur* 13: 153–155
3. Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R., 2018. Anthropological and socioeconomic factors contributing to global antimicrobial resistance: a univariate and multivariable analysis. *The Lancet Planetary Health* 2: e398–e405
4. World Bank., 2017. *Drug-Resistant Infections: A Threat to Our Economic Future*. Washington, DC
5. Abbara A, Rawson TM, Karah N, El-Amin W, Hatcher J, Tajaldin B, Dar O, Dewachi O, Abu Sitta G, Uhlin BE, Sparrow A., 2018. A summary and appraisal of existing evidence of antimicrobial resistance in the Syrian conflict. *International Journal of Infectious Diseases* 75: 26–33
6. Kanapathipillai R, Malou N, Hopman J, Bowman C, Yousef N, Michel J, Hussein N, Herard P, Ousley J, Mills C, Seguin C, Saim M., 2019. Antibiotic resistance in conflict settings: lessons learned in the Middle East. *JAC-Antimicrobial Resistance* 1
7. Haraoui DL-P., 2019. Armed conflicts and antimicrobial resistance: A deadly convergence. *AMR Control*: 5
8. McAuliffe M, Khadria B, Bauloz C., 2019. *World migration report 2020*. Geneva: IOM
9. Al Baz M, Law MR, Saadeh R., 2018. Antibiotics use among Palestine refugees attending UNRWA primary health care centers in Jordan – A cross-sectional study. *Travel Medicine and Infectious Disease* 22: 25–29
10. Eysenbach G., 2004. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 6: e34
11. Turki Y, Saleh S, Albaik S, Barham Y, van de Vrie D, Shahin Y, Hababeh M, Armagan M, Seita A., 2020. Assessment of the knowledge, attitudes, and practices (KAP) among UNRWA* health staff in Jordan concerning mental health programme pre-implementation: a cross-sectional study. *International Journal of Mental Health Systems* 14: 54
12. European Centre for Disease Prevention and Control., 2019. *Survey of healthcare workers’ knowledge, attitudes and behaviours on antibiotics, antibiotic use and antibiotic resistance in the EU/EEA*. LU: Publications Office
13. Teixeira Rodrigues A, Roque F, Falcão A, Figueiras A, Herdeiro MT., 2013. Understanding physician antibiotic prescribing behaviour: a systematic review of qualitative studies. *International Journal of Antimicrobial Agents* 41: 203–212
14. Spervovasilis N, Ierodiakonou D, Milioni A, Markaki L, Kofteridis DP, Tsioutis C., 2020. Assessing the knowledge, attitudes and perceptions of junior doctors on antimicrobial use and antimicrobial resistance in Greece. *Journal of Global Antimicrobial Resistance* 21: 296–302
15. Orubu ESF, Al-Dheeb N, Ching C, Bu Jawdeh S, Anderson J, Sheikh R, Hariri F, Basaleem H, Zaman MH., 2021. Assessing Antimicrobial Resistance, Utilization, and Stewardship in Yemen: An Exploratory Mixed-Methods Study. *Am J Trop Med Hyg*: tpmd210101
16. Seid MA, Hussen MS., 2018. Knowledge and attitude towards antimicrobial resistance among final year undergraduate paramedical students at University of Gondar, Ethiopia. *BMC Infect Dis* 18: 312

17. Ashiru-Oredope D, Hopkins S, Vasandani S, Umoh E, Oloyede O, Nilsson A, Kinsman J, Elsert L, Monnet DL, Group the #ECDCAntibioticSurvey PA., 2021. Healthcare workers' knowledge, attitudes and behaviours with respect to antibiotics, antibiotic use and antibiotic resistance across 30 EU/EEA countries in 2019. *Eurosurveillance* 26: 1900633
18. Ministry of Health., 2017. National Action Plan on Antimicrobial Resistance Containment Jordan
19. United Nations Relief and Works Agency., 2021. UNRWA Health Department Annual Report 2020. Available at:
https://www.unrwa.org/sites/default/files/content/resources/unrwa_health_department_annual_report_2020_-_final-compressed.pdf. Accessed. 2021
20. United Nations Relief and Works Agency., 2019. UNRWA Health Department Annual Report. Available at:
https://www.unrwa.org/sites/default/files/content/resources/health_department_annual_report_2019_-_final.pdf. Accessed. 2019
21. Kassem DF, Hoffmann Y, Shahar N, Ocampo S, Salomon L, Zonis Z, Glikman D., 2017. Multidrug-Resistant Pathogens in Hospitalized Syrian Children. *Emerg Infect Dis* 23: 166–168
22. Maltezou HC, Theodoridou M, Daikos GL., 2017. Antimicrobial resistance and the current refugee crisis. *J Glob Antimicrob Resist* 10: 75–79
23. Al-Niemat SI, Bloukh DT, Al-Harasis MD, Al-Fanek AF, Salah RK., 2008. Drug use evaluation of antibiotics prescribed in a Jordanian hospital outpatient and emergency clinics using WHO prescribing indicators. *Saudi Med J* 29: 743–748
24. Al-Azayzih A, Al-Azzam SI, Alzoubi KH, Shawaqfeh M, Masadeh MM., 2017. Evaluation of drug-prescribing patterns based on the WHO prescribing indicators at outpatient clinics of five hospitals in Jordan: a cross-sectional study. *CP* 55: 425–432
25. Al-Azzam S, Mhaidat NM, Banat HA, Alfaour M, Ahmad DS, Muller A, Al-Nuseirat A, Lattyak EA, Conway BR, Aldeyab MA., 2021. An Assessment of the Impact of Coronavirus Disease (COVID-19) Pandemic on National Antimicrobial Consumption in Jordan. *Antibiotics* 10: 690
26. Albaramki JH, Abdelghani T, Dalaeen A, Khdaif Ahmad F, Allassaf A, Odeh R, Akl K., 2019. Urinary tract infection caused by extended-spectrum β -lactamase-producing bacteria: Risk factors and antibiotic resistance. *Pediatrics International* 61: 1127–1132
27. Tarazi YH, Abu-Basha E, Ismail ZB, Al-Jawasreh SI., 2021. Antimicrobial susceptibility of multidrug-resistant *Pseudomonas aeruginosa* isolated from drinking water and hospitalized patients in Jordan. *Acta Tropica* 217: 105859
28. Orubu ESF, Najwa A, Ching C, Jawdeh SB, Anderson J, Sheikh R, Hariri F, Basaleem H, Zaman MH., 2021. Assessing antimicrobial resistance, utilization and stewardship in Yemen: An exploratory mixed-methods study. *medRxiv*: 2021.01.27.21250628
29. Abu Hammour K, AL-Heyari E, Allan A, Versporten A, Goossens H, Abu Hammour G, Manaseer Q., 2020. Antimicrobial Consumption and Resistance in a Tertiary Care Hospital in Jordan: Results of an Internet-Based Global Point Prevalence Survey. *Antibiotics* 9: 598

Table 1 Interview guide for the knowledge, attitudes, and practices survey of prescribers at UNRWA PHC, Jordan showing sections, questions, and response types. Section I collected the characteristics of clientele and prescribers. Section II describes current prescribing behavior and practices. Section III assessed knowledge and attitudes on rational prescribing mapped onto the COM-B framework.

Section/Domain	Questions	Response
Section I. Demographic characteristics	<ol style="list-style-type: none"> 1) Have you prescribed an antibiotic within the past 6 months? 2) How many patients do you see on a daily basis? 3) Among the patients you see, approximately what percentage are children under 12 years old? 4) Among the patients you see, approximately what percentage are refugees (all nationalities)? 5) How many prescriptions for antibiotics have you written in the last 7 days? 6) How many patients, on average, do you see in a month? 7) How many years of practice experience do you have? 8) What is your specialty? 	Multiple choice questions – Yes/No; others
Section II. A. Prescribing behavior/practices	<ol style="list-style-type: none"> 1) Are antibiotic susceptibility tests performed before antibiotic selection? 2) If not, why? (Please, check all that apply) [Availability, Time pressure, Other ____] 3) If yes, how often? 4) If yes, in what cases are antibiotic susceptibility tests usually used? (List up to three) 5) What is your main consideration for prescribing antibiotics? (Please, specify all that apply) 6) Do you feel pressured to prescribe antibiotics? 7) If yes, are any of these reasons that you feel pressured to prescribed antibiotics: <ul style="list-style-type: none"> • Patient demand • Condition of patient • Fear of complications • Need to provide rapid relief 8) Please specify other reasons that you feel pressured to prescribe antibiotics _____ 9) Do you usually prescribe antibiotics by generic or brand name? 10) If you prescribe by brand, what are your reasons for doing so? (Please, specify all that apply) [Clinical efficacy [<i>high bacterial resistance to generics/better quality/others</i>]] 11) What are the top three things you considered for antibiotic CHOICE when you prescribe? <ul style="list-style-type: none"> • Availability of antibiotic in the facility • Antibiotic susceptibility testing information • Broad-spectrum nature of antibiotic • Symptoms of patient • Other: (Please list) _____ • Other: (Please list) _____ • Other: (Please list) _____ 12) What is your main consideration for prescribing antibiotics? (Please, specify all that apply) <ul style="list-style-type: none"> ○ Antibiotic susceptibility testing information ○ Symptoms of patient ○ Clinical severity of case ○ Other: Please, specify _____ 13) Is it possible to perform laboratory antibiotic susceptibility testing on-site or off-site? 	Multiple choice questions – Yes/No; others

Section	Questions	Response																																										
Section II. B. Antibiotic prescribing pattern	<div>1) Please select the top three antibiotics that you usually prescribe?<ul style="list-style-type: none">• Amoxicillin (including amoxicillin-clavulanate)• Azithromycin• Cefuroxime• Ciprofloxacin• Co-trimoxazole• Metronidazole• Other: (Please list) _____• Other: (Please list) _____• Other: (Please list) _____</div> <div>2) When you prescribe antibiotics for a patient, do you usually prescribe? [Combination products count as 1]</div> <div>3) Please indicate which GROUPs are MOST FREQUENTLY prescribed for the following infections?</div> <table><thead><tr><th></th><th>Respiratory Tract Infections</th><th>Urinary Tract Infections</th><th>Gastro-Intestinal Diseases</th><th>Dental Infections</th><th>Skin & Soft Tissue infections</th></tr></thead><tbody><tr><td>Penicillins</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Cephalosporins</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Fluoroquinolones</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Macrolides</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Trimethoprim/sulfonamides</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Other (please specify)</td><td></td><td></td><td></td><td></td><td></td></tr></tbody></table>		Respiratory Tract Infections	Urinary Tract Infections	Gastro-Intestinal Diseases	Dental Infections	Skin & Soft Tissue infections	Penicillins						Cephalosporins						Fluoroquinolones						Macrolides						Trimethoprim/sulfonamides						Other (please specify)						MCQ
	Respiratory Tract Infections	Urinary Tract Infections	Gastro-Intestinal Diseases	Dental Infections	Skin & Soft Tissue infections																																							
Penicillins																																												
Cephalosporins																																												
Fluoroquinolones																																												
Macrolides																																												
Trimethoprim/sulfonamides																																												
Other (please specify)																																												
Section III. Capability, Opportunity, Motivation-Behaviour	<div>Capability – Knowledge – assumed (1,3); real (2) & Infection Prevention and Control (4)</div> <div>1) To what extent do you agree or disagree with each of the following statements:<ul style="list-style-type: none">○ I know what antibiotic resistance is○ I know what information to give to individuals about prudent use of antibiotics and antibiotic resistance○ I know sufficient knowledge about how to use antibiotics appropriately for my current practice</div> <div>2) Please answer whether you believe each of these statements to be True or False:<ul style="list-style-type: none">• Antibiotics are effective against viruses• Antibiotics are effective against cold and flu• Unnecessary use of antibiotics makes them become ineffective• Taking antibiotics has associated side effects such as diarrhea, colitis, or allergy• Every person treated with antibiotics is at an increased risk of antibiotic-resistant infection• Antibiotic resistant bacteria can spread from person to person• Healthy people can carry antibiotic resistant bacteria</div> <div>3) To what extent do you agree or disagree that the following environmental and animal health factors are important in contributing to antibiotic resistance in bacteria from humans?<ul style="list-style-type: none">○ Environmental factors such as waste water in the environment○ Excessive use of antibiotics in livestock and food production</div> <div>4) Do you need to perform hand hygiene (as often as recommended) if you have used gloves in contact with patients or biological material?</div>	<div>5-point Likert Scale (from left to right; Strongly disagree to Strongly agree)</div> <div>MCQ – True or False</div> <div>5-point Likert Scale (from left to right; Strongly disagree to Strongly agree)</div> <div>Yes/No</div>																																										

Section	Questions	Response
	<p><i>Capability –Attitudes or assumptions on AMR (1-5); awareness of WHO Guidelines (6-7)</i></p> <ol style="list-style-type: none"> Do you think antibiotic resistance is a problem in Jordan? Do you think Refugees have a higher risk of antibiotic resistance than the general population? Do you think antibiotic resistance is a problem in your primary health care facility? How do you suggest the problem of antibiotic resistance can be tackled in Jordan? (Choose ALL that apply) <ol style="list-style-type: none"> Increase testing for antibiotic susceptibility Increase awareness among patients Enforce prescription laws for the dispensing of antibiotics Improved training of doctors and pharmacists Other _____ How often do you see treatment failure within a month (no response to therapy, switch to new antibiotic, or addition of new antibiotic)? [<10%, 11-15%, 26-50%, >50%] <ol style="list-style-type: none"> To optimize the use of antibiotics, the WHO has classified antibiotics as Access, Watch, & Reserve. Have you heard about these categories? If yes, do they affect your prescribing? Can you list the WHO's five moments of hand hygiene? 	MCQ – Yes/No; others
	<p><i>Motivation</i></p> <ol style="list-style-type: none"> To what extent do you agree or disagree with the following statements: <ul style="list-style-type: none"> There is a connection between my prescribing of antibiotics and emergence and spread of antibiotic resistant bacteria I have a key role in helping control antibiotic resistance 	5-point Likert Scale (from left to right; Strongly disagree to Strongly agree)
	<p><i>Opportunity</i></p> <ol style="list-style-type: none"> To what extent do you agree or disagree with each of the following statements: <ol style="list-style-type: none"> I have easy access to guidelines I need on managing infections I have easy access to the materials I need to give advice on prudent antibiotic use and antibiotic resistance I have good opportunities to provide advice on prudent use to individuals (patients/public) I am confident making antibiotic prescribing decisions I have confidence in the antibiotic guidelines available to me I consider antibiotic resistance when treating a patient I feel supported to not prescribe antibiotics when they are not necessary In your facility, do you follow Standard Treatment Guidelines (or a guide specific to prescribing antibiotics)? Does your facility have a Drug/Pharmacy and Therapeutic Committee (DTC/PTC)? <ol style="list-style-type: none"> Is there a Healthcare Facility (or hospital) Formulary or an Essential Medicine List? If yes, which (Please, check all that apply)? 	<p>5-point Likert Scale (from left to right; Strongly disagree to Strongly agree)</p> <p>Yes/No</p>

Section	Questions	Response
IV. Perceived strategies to improve prescribing	<p>1) What measures do you think would be the most helpful in improving antibiotic prescribing?</p> <ul style="list-style-type: none"> • Educational sessions on prescribing • Availability of local / national guidelines / policies / protocols • Availability of local/national resistance data • Computer-aided prescribing • Presence of an antimicrobial management team • Readily accessible microbiological advice • Readily accessible advice from Infectious Disease physician • Readily accessible advice from a pharmacist • Readily accessible advice from infection control team • Advice from senior colleagues • Speaking to a pharmaceutical representative • Restriction of prescription of certain antibiotics • Restriction of prescription of all antibiotics • Regular audit and feedback on antibiotic prescribing in your PHC 	5-point Likert Scale (from left to right; Very helpful to Very unhelpful)

Notes:

1. Capability here is a combination of knowledge (actual or real and assumed or perceived), AND infection prevention and control (hand washing).

Table2 Real knowledge scores among prescribers at the UNRWA PHC, Jordan

		TRUE, n	FALSE, n	Correct answer	Correct (%)
a	Antibiotics are effective against viruses	0	34	FALSE	100%
b	Antibiotics are effective against cold and flu	0	34	FALSE	100%
c	Taking antibiotics has associated side effects such as diarrhea, colitis, or allergy	34	0	TRUE	100%
d	Unnecessary use of antibiotics makes them become ineffective	33	1	TRUE	97%
e	Antibiotic resistant bacteria can spread from person to person	30	4	TRUE	88%
f	Every person treated with antibiotics is at an increased risk of antibiotic resistant infection	34	0	TRUE	100%
g	Healthy people can carry antibiotic resistant bacteria	32	2	TRUE	94%
Mean/aggregate real knowledge score					97%

Table 3 Opportunity to improve prescribing behavior

		Strongly Disagree	Disagree	Neither disagree nor agree	Agree	Strongly agree	Total	Agree
a	I have easy access to guidelines I need on managing infections	2			21	11	34	94%
b	I am confident making antibiotic prescribing decisions		2	1	21	10	34	91%
c	I have good opportunities to provide advice on prudent use to individuals (patients/public)		2	2	19	11	34	88%
d	I have confidence in the antibiotic guidelines available to me		3	1	20	10	34	88%
e	I consider antibiotic resistance when treating a patient	2		3	17	12	34	85%
f	I have easy access to the materials I need to give advice on prudent antibiotic use and antibiotic resistance		4	1	18	11	34	85%
g	I feel supported to not prescribe antibiotics when they are not necessary	1	2	3	18	10	34	82%
	Aggregate opportunity score							88%

Note: *Agree* is the combined proportion of “agree” and “strongly agree” responses expressed as percentage of all respondents (n=34).

Table 4 Motivation for behavior change towards rational prescribing assessed by two statements and aggregate scores

		Strongly Disagree	Disagree	Neither disagree nor agree	Agree	Strongly agree	Agree
a	There is a connection between my prescribing of antibiotics and emergence and spread of antibiotic resistant bacteria	4	18	6	3	3	18%
b	I have a key role in helping control antibiotic resistance	8	20	1	3	2	15%
	Aggregate motivation						16%

Note: *Agree* is the combined proportion of “agree” and “strongly agree” responses expressed as percentage of all respondents (n=34).

Table 5 Options to improve antibiotic prescribing ranked by aggregate perceptions of “helpful” options among prescribers at UNRWA PHC, Jordan

No.	Options	Responses (no. of participants)				Helpful
		Very helpful	Helpful	Neutral	Unhelpful	Very unhelpful
1	Availability of local / national guidelines / policies / protocols	7	18	9		74%
2	Availability of local/national resistance data	4	21	9		74%
3	Educational sessions on prescribing	8	16	10		71%
4	Regular audit and feedback on antibiotic prescribing in your facility	7	16	10	1	68%
5	Computer-aided prescribing	5	18	11		68%
6	Readily accessible microbiological advice	4	18	11	1	65%
7	Restriction of prescription of all antibiotics	6	16	10	2	65%
8	Restriction of prescription of certain antibiotics	5	16	12	1	62%
9	Presence of an antimicrobial management team	4	16	13	1	59%
10	Readily accessible advice from a pharmacist	6	14	13	1	59%
11	Readily accessible advice from infection control team	5	13	16		53%
12	Speaking to a pharmaceutical representative	5	13	14	2	53%
13	Readily accessible advice from Infectious disease physician	5	12	16	1	50%
14	Advice from senior colleagues	5	12	16	1	50%

Note: *Helpful* is the proportion of “very helpful” and “helpful” responses expressed as percentage of all respondents (34).

A

Prescribers Selection of Antibiotics

B

Selection by Disease Category

i AST Possible (on-site or off-site)

■ 56% Yes
■ 44% No

ii AST on-site or off-site

■ 9.5% On-site
■ 90.5% Off-site

A **B**