The statistical analysis of daily data associated with different parameters of the New Coronavirus COVID-19 pandemic in Georgia and their monthly interval prediction from September 1, 2021 to December 31, 2021

Avtandil G. Amiranashvili¹, Ketevan R. Khazaradze^{2,3}, Nino D. Japaridze^{3,4}

1 Mikheil Nodia Institute of Geophysics of Ivane Javakhishvili Tbilisi State University, Tbilisi, Georgia

2 Georgian State Teaching University of Physical Education and Sport, Tbilisi, Georgia

3 Ministry of Internally Displaced Persons from Occupied Territories, Labour, Health and Social Affairs of Georgia, Tbilisi, Georgia

4 Tbilisi State Medical University, Tbilisi, Georgia

* Corresponding author

E-mail: avtandilamiranashvili@gmail.com

Source(s) of support: Nil.

Conflicting Interest – No.

ORCID: 0000-0001-6152-2214

Abstract

The lockdown introduced in Georgia on November 28, 2020 contributed to positive trends in the spread of COVID-19 until February - the first half of March 2021. Then, in April-May 2021, the epidemiological situation worsened significantly, and from June to the end of December COVID - situation in Georgia was very difficult.

In this work results of the next statistical analysis of the daily data associated with New Coronavirus COVID-19 infection of confirmed (C), recovered (R), deaths (D) and infection rate (I) cases of the population of Georgia in the period from September 01, 2021 to December 31, 2021 are presented. It also presents the results of the analysis of monthly forecasting of the values of C, D and I. As earlier, the information was regularly sent to the National Center for Disease Control & Public Health of Georgia and posted on the Facebook page <https://www.facebook.com/Avtandil1948/>.

The analysis of data is carried out with the use of the standard statistical analysis methods of random events and methods of mathematical statistics for the non-accidental time-series of observations. In particular, the following results were obtained.

Georgia's ranking in the world for Covid-19 monthly mean values of infection and deaths cases in investigation period (per 1 million population) was determined. Among 157 countries with population ≥ 1 million inhabitants in October 2021 Georgia was in the 4 place on new infection cases, and in September - in the 1 place on death. Georgia took the best place in terms of confirmed cases of diseases (thirteenth) in December, and in mortality (fifth) - in October.

A comparison between the daily mortality from Covid-19 in Georgia from September 01, 2021 to December 31, 2021 with the average daily mortality rate in 2015-2019 shows, that the largest share value of D from mean death in 2015-2019 was 76.8 % (September 03, 2021), the smallest 18.7 % (November 10, 2021).

As in previous work [9,10] the statistical analysis of the daily and decade data associated with coronavirus COVID-19 pandemic of confirmed, recovered, deaths cases and infection rate of the population of Georgia are carried out. Maximum daily values of investigation parameters are following: C = 6024 (November 3, 2021), R = 6017 (November 15, 2021), D = 86 (September 3, 2021), I = 12.04 % (November 24, 2021). Maximum mean decade values of investigation parameters are following: C = 4757 (1 Decade of November 2021), R = 4427 (3 Decade of November 2021), D = 76 (2 Decade of November 2021), I = 10.55 % (1 Decade of November 2021).

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

It was found that as in spring and summer 2021 [9,10], from September to December 2021 the regression equations for the time variability of the daily values of C, R, D and I have the form of a tenth order polynomial.

Mean values of speed of change of confirmed -V(C), recovered - V(R), deaths - V(D) and infection rate V(I) coronavirus-related cases in different decades of months for the indicated period of time were determined. Maximum mean decade values of investigation parameters are following: V(C) = +139 cases/day (1 Decade of October 2021), V(R) = +124 cases/day (3 Decade of October 2021), V(D) = +1.7 cases/day (3 Decade of October 2021), V(I) = + 0.20 %/ day (1 decades of October 2021).

Cross-correlations analysis between confirmed COVID-19 cases with recovered and deaths cases shows, that from September 1, 2021 to November 30, 2021 the maximum effect of recovery is observed on 12 and 14 days after infection ($CR=0.77$ and 0.78 respectively), and deaths - after 7, 9, 11, 13 and 14 days ($0.70 \leq CR \leq 0.72$); from October 1, 2021 to December 31, 2021 - the maximum effect of recovery is observed on 14 days after infection ($RC=0.71$), and deaths - after 9 days ($CR=0.43$). In Georgia from September 1, 2021 to November 30, 2021 the duration of the impact of the delta variant of the coronavirus on people (recovery, mortality) could be up to 28 and 35 days respectively; from October 1, 2021 to December 31, 2021 - up to 21 and 29 days respectively.

Comparison of daily real and calculated monthly predictions data of C, D and I in Georgia are carried out. It was found that in investigation period of time daily and mean monthly real values of C, D and I practically fall into the 67% - 99.99% confidence interval of these predicted values.

Traditionally, the comparison of data about C and D in Georgia (GEO) with similar data in Armenia (ARM), Azerbaijan (AZE), Russia (RUS), Turkey (TUR) and in the World (WRL) is also carried out.

Key words: New Coronavirus COVID-19, statistical analysis, short-term prediction.

1. Introduction

Two years have passed since the outbreak of the new coronavirus (COVID-19) in China, which was recognized on March 11, 2020 as a pandemic due to its rapid spread in the World [1]. During this period of time, despite the measures taken (including vaccination), several strains of this virus have appeared. The overall level of morbidity and mortality in many countries of the world is still quite high. Scientists and specialists of various disciplines from all over the world continue intensive research of this unprecedented phenomenon (including in Georgia [2-11]), rendering their all possible assistance to epidemiologists.

In particular, in our works [6-11], it was noted that specialists in the field of physical and mathematical sciences make an important contribution to research on the spread of the new coronavirus COVID-19. Works on statistical analysis [2-5, 12-15], forecasting [16-32], forecasting systematization [33,34], spatial-temporary modeling of the spread of the new coronavirus [35-38] etc. was actively continuing in 2021.

This work is a continuation of the researches [7-11].

In this work results of a statistical analysis of the daily data associated with New Coronavirus COVID-19 infection of confirmed (C), recovered (R), deaths (D) and infection rate (I) cases of the population of Georgia in the period from September 01, 2021 to December 31, 2021 are presented. It also presents the results of the analysis of monthly forecasting of the values of C, D and I. The information was regularly sent to the National Center for Disease Control & Public Health of Georgia and posted on the Facebook page <https://www.facebook.com/Avtandil1948/>.

The comparison of data about C and D in Georgia with similar data in Armenia, Azerbaijan, Russia, Turkey and in the world is also carried out.

We used standard methods of statistical analysis of random events and methods of mathematical statistics for non-random time series of observations [7-11, 39-41].

2. Study areas, material and methods

The study area: Georgia. Data of John Hopkins COVID-19 Time Series Historical Data (with US State and County data) [<https://www.southsawyer.com/john-hopkins-time-series-data-with-us-state-and-county-city-detail-historical/>; <https://data.humdata.org/dataset/total-covid-19-tests-performed-by-country>] and <https://stopcov.ge> about daily values of confirmed, recovered, deaths and infection rate coronavirus-related cases,

from September 01, 2021 to December 31, 2021 are used. The work also used data of National Statistics Office of Georgia (Geostat) on the average monthly total mortality in Georgia in January-December 2015-2019 [<https://www.geostat.ge/en/>]. In the proposed work the analysis of data is carried out with the use of the standard statistical analysis methods of random events and methods of mathematical statistics for the non-accidental time-series of observations [7-11, 39-41].

The following designations will be used below: Mean – average values; Min – minimal values; Max – maximal values; Range – Max-Min; St Dev - standard deviation; σ_m - standard error; $C_v = 100 \cdot \text{St Dev} / \text{Mean}$ – coefficient of variation, %; R^2 – coefficient of determination; r – coefficient of linear correlation; CR – coefficient of cross correlation; Lag = 1, 2...60 Day; K_{DW} – Durbin-Watson statistic; Calc – calculated data; Real - measured data; Dec1, Dec2, Dec3 – numbers of the month decades; α - the level of significance; C, R, D - daily values of confirmed, recovered and deaths coronavirus-related cases; V(C), V(R) and V(D) - daily values of speed of change of confirmed, recovered and deaths coronavirus-related cases (cases/day); I - daily values of infection rate (or positive rate) coronavirus-related cases ($100 \cdot C / \text{number of coronavirus tests performed}$); V(I) - daily values of speed of change of I; DC – deaths coefficient, % = $(100 \cdot D / C)$. Official data on number of coronavirus tests performed are published from December 05, 2020 [<https://stopcov.ge>].

The statistical programs Data Fit 7, Mesosaur and Excel 16 were used for calculations.

The curve of trend is equation of the regression of the connection of the investigated parameter with the time at the significant value of the determination coefficient and such values of K_{DW} , where the residual values are accidental. If the residual values are not accidental the connection of the investigated parameter with the time we will consider as simply regression.

The calculation of the interval prognostic values of C, D and I taking into account the periodicity in the time-series of observations was carried out using Excel 16 (the calculate methodology was description in [7]).

The duration of time series of observations to calculating monthly forecasts of C and I values was as follows: for September - 120 days, for October - 122 days, for November - 118 days, for December - 127 days, periodicity - 7 days. The duration of time series of observations to calculating monthly forecasts of D values was as follows: for September - 119 days, periodicity - 8 days; for October - 124 days, periodicity - 22 days; for November - 118 days, periodicity - 8 days; for December - 127 days, periodicity - 14 days.

67%...99.99%_Low - 67%...99.99% lower level of confidence interval of prediction values of C, D and I; 67%...99.99%_Upp - 67%...99.99% upper level of confidence interval of prediction values of C, D and I.

In the Table 1 [8] the scale of comparing real data with the predicted ones and assessing the stability of the time series of observations in the forecast period in relation to the pre-predicted one (period for prediction calculating) is presented.

3. Results and Discussion

The results in the Fig. 1- 24 and Table 1-9 are presented.

3.1 Comparison of time-series of Covid-19 confirmed and deaths cases in Georgia, its neighboring countries and World in summer 2021

The time-series curves and statistical characteristics of Covid-19 confirmed and deaths cases (to 1 million populations) in Georgia, its neighboring countries and World from September 01, 2021 to December 31, 2021 in Fig. 1 - 2 and in Table 1-2 are presented.

Variability of the values of C per 1 million populations is as follows (Fig. 1, Table 1):

- Georgia. Range of change: 178-1616, mean value - 843.
- Armenia. Range of change: 20-877, mean value - 284.
- Azerbaijan. Range of change: 0-716, mean value - 154.
- Russia. Range of change: 117-276, mean value - 197.
- Turkey. Range of change: 0-898, mean value - 298.
- World. Range of change: 39-248, mean value - 73.

Fig.1. Time-series of Covid-19 confirmed cases to 1 million populations in Georgia, neighboring countries (Armenia, Azerbaijan, Russia, Turkey) and World from September 1, 2021 to December 31, 2021.

Table 1. The statistical characteristics of Covid-19 confirmed cases to 1 million populations in Georgia, neighboring countries (Armenia, Azerbaijan, Russia, Turkey) and World from September 1, 2021 to December 31, 2021 ($r_{\min} = \pm 0.18$, $\alpha = 0.05$).

Parameter	GEO	ARM	AZE	RUS	TUR	WRL
Mean	843	284	154	197	298	73
Min	178	20	0	117	0	39
Max	1616	877	716	276	898	248
Range	1438	857	716	159	898	208
Median	806	226	137	198	296	65
St Dev	358	214	119	49	84	33
σ_m	32.6	19.4	10.8	4.4	7.6	3.0
C_v (%)	42.5	75.2	77.1	24.8	28.1	45.7
Correlation Matrix						
GEO	1	0.50	0.23	0.67	0.01	-0.14
ARM	0.50	1	0.23	0.53	0.23	-0.45
AZE	0.23	0.23	1	0.09	-0.03	-0.14
RUS	0.67	0.53	0.09	1	-0.03	-0.27
TUR	0.01	0.23	-0.03	-0.03	1	0.17
WRL	-0.14	-0.45	-0.14	-0.27	0.17	1

The largest variations in C values were observed in Azerbaijan ($C_v=77.1\%$), the smallest - in Russia ($C_v=24.8\%$).

Significant linear correlation ($r_{\min} = \pm 0.18$, $\alpha = 0.05$) between these countries on C value varies from 0.23 (pairs Georgia-Azerbaijan, Armenia-Azerbaijan, Armenia-Turkey) to 0.67 (pair Georgia-Russia). Linear correlation between World and these countries is significant only for two pairs: World-Armenia ($r = -0.45$) and World-Russia ($r = -0.27$). The degree of correlation [41] is as follows: moderate correlation ($0.5 \leq R < 0.7$) – for pairs Georgia-Armenia, Georgia-Russia and Armenia-Russia; low correlation ($0.3 \leq R < 0.5$) – for pair World-Armenia; negligible correlation ($0 \leq R < 0.3$) – for pairs Georgia-Azerbaijan, Armenia-Azerbaijan, Armenia-Turkey and World-Russia.

Fig.2. Time-series of deaths cases from Covid-19 to 1 million population in in Georgia, neighboring countries (Armenia, Azerbaijan, Russia, Turkey) and World from September 1, 2021 to December 31, 2021.

Table 2. The statistical of deaths cases from Covid-19 to 1 million populations in Georgia, neighboring countries (Armenia, Azerbaijan, Russia, Turkey) and World from September 1, 2021 to December 31, 2021. ($r_{\min} = \pm 0.18$, $\alpha = 0.05$).

Parameter	GEO	ARM	AZE	RUS	TUR	WRL
Mean	14.1	8.7	2.2	6.9	2.5	1.0
Min	6.2	0.3	0.0	4.9	0.0	0.4
Max	23.1	23.6	10.9	8.4	3.7	1.6
Range	16.9	23.2	10.9	3.5	3.7	1.2
Median	13.5	7.4	2.0	6.8	2.5	1.0
St Dev	4.42	5.10	1.64	1.10	0.48	0.22
σ_m	0.40	0.46	0.15	0.10	0.04	0.02
C_v (%)	31.5	58.9	75.3	16.0	19.6	23.4
Correlation Matrix						
GEO	1	0.11	0.14	0.34	-0.02	0.12
ARM	0.11	1	0.07	0.56	0.08	0.00
AZE	0.14	0.07	1	-0.08	0.19	0.20
RUS	0.34	0.56	-0.08	1	-0.33	-0.27
TUR	-0.02	0.08	0.19	-0.33	1	0.37
WRL	0.12	0.00	0.20	-0.27	0.37	1

Variability of the values of D per 1 million populations is as follows (Fig. 2, Table 2):

- Georgia. Range of change: 6.2-23.1, mean value – 14.1.
- Armenia. Range of change: 0.3-23.6, mean value – 8.7.
- Azerbaijan. Range of change: 0.0-10.9, mean value – 2.2.
- Russia. Range of change: 4.9-8.4, mean value – 6.9.
- Turkey. Range of change: 0.0-3.7, mean value -2.5.
- World. Range of change: 0.4-1.6, mean value – 1.0.

The largest variations in D values were observed in Azerbaijan ($C_v=75.3\%$), the smallest - in Russia ($C_v=16.0\%$).

Significant positive linear correlation between these countries on D value varies from 0.19 (pair Azerbaijan-Turkey) to 0.56 (pair Armenia-Russia); negative – between Russia and Turkey ($r = -0.33$). Linear

positive correlation between World and these countries is significant for two pairs: World-Azerbaijan ($r = 0.20$) and World-Turkey ($r = 0.37$); negative – for pair World-Russia ($r = -0.27$).

The degree of correlation [41] is as follows: moderate correlation – only for pair Armenia-Russia; low correlation – for pairs Georgia-Russia, Russia-Turkey and World-Turkey; negligible correlation – for pairs Azerbaijan-Turkey, World-Azerbaijan and World-Russia.

It should be noted that, in general, in the studied period of time, level of correlations shown in Tables 1 and 2 are worse than in the summer of 2021 [10].

In Table 3 the statistical characteristics of mean monthly values of C and D related to Covid-19 for 157 countries with population ≥ 1 million inhabitants from September to December 2021 (normed per 1 million population) is presented.

Table 3. The statistical characteristics of mean monthly values of confirmed cases and deaths cases related to Covid-19 for 157 countries with population ≥ 1 million inhabitant from September to December 2021 (normed on 1 mln pop.).

Month	September		October		November		December	
Parameter	C	D	C	D	C	D	C	D
Mean	113	1.6	121	1.6	166	2.0	190	1.7
Min	0	0.0	0	0.0	0	0.0	0	0.0
Max	904	13.7	1051	18.1	1656	21.5	1748	16.3
Range	904	13.7	1051	18.1	1656	21.5	1748	16.3
St Dev	163	2.4	211	2.8	310	4.0	324	3.3
σ_m	13.1	0.2	16.9	0.2	24.8	0.3	26.0	0.3
C_V (%)	145	148	174	180	187	198	170	191

As follows from this Table range of change of mean monthly values of C for 157 countries varied from 0 (all months) to 1748 (December). Average value of C for 157 countries varied from 113 (September) to 190 (December). Value of C_V changes from 145% (September) to 187% (November).

Range of change of mean monthly values of D for 157 countries varied from 0 (all months) to 21.5 (November). Average value of D for 157 countries varied from 1.6 (September and October) to 2.0 (November). Value of C_V changes from 148% (September) to 198% (November).

Fig. 3. Linear correlation and regression between mean monthly values of deaths and confirmed cases related to Covid-19 for 157 countries with population ≥ 1 million inhabitants (normed on 1 mln pop.) from September to December 2021.

Between mean monthly values of Deaths and Confirmed cases related to Covid-19 for 157 countries with population ≥ 1 million inhabitant linear correlation and regression are observed (Fig. 3). As follows from Fig. 3 the highest growth rate of the monthly average values of D depending on C was observed in October, the smallest - in December (the corresponding values of the linear regression coefficients).

In Table 4 data about Covid-19 mean monthly values of infection (C) and deaths (D) cases from September to December 2021 (per 1 million population) and ranking of Georgia, Armenia, Azerbaijan, Russia and Turkey by these parameters (in brackets) among 157 countries with population ≥ 1 million inhabitant are presented. The corresponding values of the deaths coefficient (DC) are also given here.

In particular, as follows from this Table, mean monthly values of C for 5 country changes from 69.7 (Armenia, December, 66 place between 157 country) to 1130.0 (Georgia, November, 6 place between 157 country).

Mean monthly values of D for 5 country changes from 1.58 (Azerbaijan, December, 39 place between 157 country) to 17.96 (Georgia, November, 2 place between 157 country).

Table 4. Covid-19 mean monthly values of infection (C) and deaths (D) cases from September to December 2021 (per 1 million population) and ranking of Georgia, Armenia, Azerbaijan, Russia, Turkey by these parameters (in brackets) among 157 countries with population ≥ 1 million inhabitant, and correspondent values of deaths coefficient (DC).

Location	Georgia	Armenia	Azerbaijan	Russia	Turkey	World
Parameter/Month	September					
C (Rating C)	553.5 (5)	219.7 (27)	192.4 (32)	132.6 (44)	300.1 (15)	67.0
D (Rating D)	13.75 (1)	5.33 (14)	2.90 (29)	5.38 (12)	2.88 (30)	1.12
DC, %	2.48	2.43	1.51	4.06	0.96	1.67
Parameter/Month	October					
C (Rating C)	919.1 (4)	506.8 (13)	149.8 (37)	216.0 (26)	333.4 (18)	53.1
D (Rating D)	9.51 (5)	10.97 (3)	1.73 (39)	6.77 (12)	2.49 (28)	0.89
DC, %	1.03	2.16	1.16	3.14	0.75	1.67
Parameter/Month	November					
C (Rating C)	1130.0 (6)	339.1 (25)	185.7 (38)	249.2 (31)	299.7 (27)	66.2
D (Rating D)	17.96 (2)	13.91 (6)	2.55 (26)	8.16 (15)	2.44 (27)	0.91
DC, %	1.59	4.10	1.37	3.27	0.81	1.38
Parameter/Month	December					
C (Rating C)	770.8 (13)	69.7 (66)	90.3 (58)	188.3 (41)	260.6 (35)	104.0
D (Rating D)	15.11 (3)	4.46 (19)	1.58 (39)	7.24 (14)	2.09 (31)	0.90
DC, %	1.96	6.39	1.75	3.85	0.80	0.86

On the whole (Table 4) in October 2021 Georgia was in the 4 place on new infection cases, and in September - in the 1 place on death. Georgia took the best place in terms of confirmed cases of diseases (thirteenth) in December, and in mortality (fifth) - in October.

Mean monthly values of DC for 5 country changes from 0.75% (Turkey, October) to 6.39 (Armenia, December).

Note that the mean values of DC (%) from September to December 2021 are: Georgia – 1.67, Armenia – 3.05, Azerbaijan – 1.42, Russia – 3.50, Turkey – 0.83, World – 1.31 (according to data from Table 1 and 2).

In summer 2021 the mean values of DC (%) were: Georgia – 1.28, Armenia – 2.08, Azerbaijan – 0.80, Russia – 3.36, Turkey – 0.80, World – 1.79 [10]. Thus, during the study period, compared to the summer 2021, the DC values in all five indicated countries increased (in Turkey, this growth is insignificant). In the World, on the contrary, the value of DC has decreased.

3.2 Comparison of daily death from Covid-19 in Georgia with daily mean death in 2015-2019 from September 1, 2021 to December 31, 2021.

In Fig. 4-7 data of the daily death from Covid-19 in Georgia from October 1, 2020 to December 31, 2021 in comparison with daily mean death in 2015-2019 are presented. The daily mean death in 2015-2019 in different months are: January - 155, February - 146, March - 141, April - 137, May - 131, June - 124, July - 117, August - 120, September - 112, October - 123, November - 135, December - 144 (Fig. 4, 6).

Fig. 4. Daily death from Covid-19 in Georgia from September 1, 2020 to December 31, 2021 in comparison with daily mean death in 2015-2019.

Fig. 5. Share value of D from mean death in 2015-2019.

Fig. 6. Mean daily mortality from Covid-19 from October 2020 to December 2021, % and mean daily mortality from January to December in 2015-2019.

The daily death from Covid-19 from September 1, 2021 to December 31, 2021 (Fig. 4) changes from 23 (October 11, 2021) to 86 (September 3, 2021). A comparison between the daily mortality from Covid-19 in Georgia from September 01, 2021 to December 31, 2021 with the average daily mortality rate in 2015-2019 shows, that the largest share value of D from mean death in 2015-2019 (Fig. 5) was 76.8 % (September 03, 2021), the smallest 18.7 % (November 10, 2021).

Fig. 7. Share of mean daily mortality from Covid-19 of mean daily mortality in 2015-2019 from October 2020 to December 2021.

The most share of mean daily mortality from Covid-19 of mean daily mortality in 2015-2019 from October 2020 to December 2021 in November was observed – 49.7% (Fig. 7).

3.3 The statistical analysis of the daily and decade data associated with New Coronavirus COVID-19 pandemic from September to December 2021

Results of the statistical analysis of the daily and decade data associated with New Coronavirus COVID-19 pandemic in Georgia from September 1, 2021 to December 31, 2021 in Tables 5-7 and Fig. 8 – 20 are presented.

The mean and extreme values of the studied parameters are as follows (Table 5): C - mean - 3145, range: 662 - 6024; R - mean - 3293, range: 795 - 6017; D - mean - 52, range: 23 - 86; I (%) - mean - 7.74, range: 3.56– 12.04. The range of variability for all studied parameters: 23.2% ($I \leq C_v \leq 42.5\%$ (C)).

Mean decade values of confirmed and recovered coronavirus-related cases varies within the following limits (Fig. 8): C – from 1709 (3 Decade of September 2021) to 4757 (1 Decade of November 2021); R – from 1545 (1 Decade of October 2021) to 4427 (3 Decade of November 2021).

Table 5. Statistical characteristics of the daily data associated with coronavirus COVID-19 pandemic of confirmed, recovered, deaths cases and infection rate of the population of Georgia from 01.09.2021 to 31.12.2021.

Parameter	Confirmed	Recovered	Deaths	Infection Rate, %
Mean	3145	3293	52	7.74
Min	662 (Sep 20)	795 (Oct 2)	23 (Oct 11)	3.56 (Oct 2)
Max	6024 (Nov 3)	6017 (Nov 15)	86 (Sep 3)	12.04 (Nov 24)
Range	5362	5222	63	8.48
St Dev	1336	1274	16	1.79
σ_m	121	116	1.5	0.16
C_v (%)	42.5	38.7	31.5	23.2

Fig.8. Mean values of confirmed and recovered coronavirus-related cases in different decades of months in Georgia from September 2021 to December 2021.

Mean decade values of deaths coronavirus-related cases (Fig. 9) varies from 30 (1 Decade of October) to 76 (2 Decade of November).

Mean decade values of infection rate coronavirus-related cases (Fig. 10) varies from 5.38 % (3 Decade of December) to 10.55 % (1 Decade of November).

Fig.9. Mean values of deaths coronavirus-related cases in different decades of months in Georgia from September 2021 to December 2021.

Fig. 10. Mean values of infection rate coronavirus-related cases in different decades of months in Georgia from September 2021 to December 2021.

Time changeability of the daily values of C, R, D and I are satisfactorily described by the tenth order polynomial (Table 6, Fig. 11-13). For clarity, the data in Fig. 12 are presented in relative units (%) in relation to their average values.

Table 6. Form of the equations of the regression of the time changeability of the daily values of C, R and D from September 01, 2021 to December 31, 2021 in Georgia. The level of significance of R^2 is not worse than 0.001.

Parameter	Confirmed	Recovered	Deaths	Infection Rate
Regression	Tenth Order Polynomial			
R^2	0.577	0.565	0.789	0.726
K_{DW}	1.97	2.01	1.57	1.61
Autocorrelation of residual components	Absent

Fig.11. Changeability of the real and calculated daily values of coronavirus confirmed and recovered cases from September 01, 2021 to December 31, 2021 in Georgia.

Fig.12. Changeability of the real and calculated daily values of coronavirus-related confirmed and deaths cases from September 01, 2021 to December 31, 2021 in Georgia (normed on mean values, %).

Fig. 13. Changeability of the real and calculated daily values of coronavirus Infection Rate from September 01, 2021 to December 31, 2021 in Georgia.

Note that from Fig. 11 and 12, as in [8-10], clearly show the shift of the time series values of R and D in relation to C.

In Fig. 14-16 data about mean values of speed of change of confirmed, recovered, deaths coronavirus-related cases and infection rate in different decades of months from September to December 2021 are presented.

Fig. 14. Mean values of speed of change of confirmed and recovered coronavirus-related cases in different decades of months in Georgia from September 2021 to December 2021.

Fig. 15. Mean values of speed of change of deaths coronavirus-related cases in different decades of months in Georgia from September 2021 to December 2021.

Fig. 16. Mean values of speed of change of coronavirus infection rate in different decades of months in Georgia from September 2021 to December 2021.

Maximum mean decade values of investigation parameters are following: $V(C) = +139$ cases/day (1 Decade of October), $V(R) = +124$ cases/day (3 Decade of October), $V(D) = +1.7$ cases/day (3 Decade of October), $V(I) = +0.20$ %/ day (1 decades of October). Min mean decade values of investigation parameters are following: $V(C) = -155$ cases/day (1 Decade of September), $V(R) = -114$ cases/day (2 Decade of September), $V(D) = -2.9$ cases/day (1 Decade of September), $V(I) = -0.16$ %/day (2 Decade of September).

Data about mean monthly values of C, R, D, I and its speed of change in from September to December 2021 in Table 7 are presented. As follows from this Table there was an increase in average monthly values of C and I from September to November, and further decrease to December. Values of R and D decrease from September to October, increase from October to November, and further decrease to December.

Table 7. Mean monthly values of C, R, D, I and its speed of change in Georgia from September 2021 to December 2021.

Parameter	C	R	D	I, %	V(C)	V(R)	V(D)	V(I)
Sep_2021	2064	3052	51.3	6.74	-67	-88	-1.6	-0.07
Oct_2021	3427	2434	35.5	8.14	90	76	0.7	0.13
Nov_2021	4213	4256	67.0	9.59	-23	-2	0.3	-0.04
Dec_2021	2874	3455	56.3	6.54	-48	-56	-0.5	-0.10

The values of $V(C)$, $V(R)$, $V(D)$ and $V(I)$ changes as follows: $V(C)$ – from -67 cases/day (September) to +90 cases/day (October); $V(R)$ - from -88 cases/day (September) to +76 cases/day (October); $V(D)$ - from -1.6 cases/day (September) to +0.7 cases/day (October) and $V(I)$ - from -0.10 %/day (December) to +0.13 %/day (October).

Fig. 17. Connection of 14-day moving average of deaths cases due to COVID-19 in Georgia with 14-day moving average of infection rate from December 18, 2020 until December 31, 2021.

In Fig. 17 data about connection of 14-day moving average of deaths cases due to COVID-19 in Georgia with 14-day moving average of infection rate from December 18, 2020 until December 31, 2021 are presented. As follows from Fig. 17, in general, with an increase of the infection rate is observed increase of deaths cases.

Fig. 18. Mean monthly values of COVID-19 infection rate and deaths cases in Georgia from December 2020 to December 2021.

Fig. 19. Linear correlation and regression between the mean monthly values of COVID-19 infection rate and deaths cases in Georgia from December 2020 to December 2021.

$$R=0.58 \text{ (moderate correlation [41]), } \alpha \approx 0.03$$

Using the data in Fig. 18, a linear regression graph between the monthly mean values of D and I is obtained (Fig. 19). As follows from Fig. 19, in general a moderate level of linear correlation between these parameters is observed. This Fig. also clearly demonstrates the anomaly high mortality from coronavirus in November 2021 with relatively low value of infection rate in comparison with December 2020, which reduces the level of correlation between D and I.

As noted in [9,10] and above (Fig. 11 and 12), there is some time-lag in the values of the time series R and D with respect to C. An estimate of the values of this time-lag for September-November and October-December 2021 is given below (Fig. 20).

Fig. 20. Coefficient of cross-correlations between confirmed COVID-19 cases (normed to tests number) with recovered and deaths cases in Georgia from 01.09.2021 to 30.11.2021 and from 01.10.2021 to 31.12.2021.

Cross-correlations analysis between confirmed COVID-19 cases with recovered and deaths cases shows, that in spring 2021 the maximum effect of recovery is observed 9 and 13 days after infection, and deaths - after 12-17 days [9]. In spring 2021 the cross-correlation of the C values with the R and D values weakens completely 38-45 days after infection. Maximum values of CR were 0.81 for R, and 0.79 for D.

In summer 2021 the maximum effect of recovery is observed 19 days after infection (CR = 0.95), and deaths - after 16 and 18 days (CR = 0.94). In contrast to the spring of 2021, in the summer of 2021, a high level of cross-correlation of C values with R and D values to 40 days after infection is observed (CR $\geq$ 0.80). In summer 2021 the cross-correlation of the C values with the R and D values weakens completely 56-58 days after infection. So, in Georgia in summer 2021, the duration of the impact of the delta variant of the coronavirus on people (recovery, mortality) could be up to two months [10].

From September 1, 2021 to November 30, 2021 the maximum effect of recovery is observed on 12 and 14 days after infection (CR = 0.77 and 0.78 respectively), and deaths - after 7, 9, 11, 13 and 14 days ($0.70 \leq \text{CR} \leq 0.72$); from October 1, 2021 to December 31, 2021 - the maximum effect of recovery is observed on 14 days after infection (CR = 0.71), and deaths - after 9 days (RC = 0.43). In Georgia from September 1, 2021 to November 30, 2021 the duration of the impact of the delta variant of the coronavirus on people (recovery, mortality) could be up to 28 and 35 days respectively; from October 1, 2021 to December 31, 2021 - up to 21 and 29 days respectively (Fig. 20).

3.4 Comparison of real and calculated prognostic daily and monthly data related to the New Coronavirus COVID-19 pandemic in Georgia from September 1, 2021 to December 31, 2021

With September 1, 2021, we started monthly forecasting values of C, D and I. In Fig. 21-24 and Table 8 examples of comparison of real and calculated prognostic daily and mean monthly data related to the COVID-19 coronavirus pandemic in Georgia from September 1, 2021 to December 31, 2021 are presented. Note that the results of the analysis of monthly forecasting of the values of C, D and I, information about which was regularly sent to the National Center for Disease Control & Public Health of Georgia and posted on the Facebook page <https://www.facebook.com/Avtandil1948/>.

Fig. 21. Example of Interval Prediction of COVID-19 Infection Cases in Georgia from 01.12.2021 to 31.12.2021.

Fig. 22. Example of Verification of Interval Prediction of COVID-19 Infection Cases in Georgia from 01.12.2021 to 31.12.2021.

Fig. 23. Example of Verification Interval Prediction of COVID-19 Death Cases in Georgia from 01.12.2021 to 31.12.2021.

Fig. 24. Example of Verification Interval Prediction of COVID-19 Infection Rate in Georgia from 01.12.2021 to 31.12.2021.

Comparison of real and calculated predictions data of C, D and I shown, that monthly daily and mean monthly real values of C, D and I practically falls into the 67% - 99.99% confidence interval of these predicted values for the specified time periods (Fig. 21-24, data from <https://www.facebook.com/Avtandil1948/>, Table 8).

Table 8. Verification of monthly interval prediction of COVID-19 confirmed infection cases, deaths cases and infection rate in Georgia from 01.09.2021 to 31.12.2021. and from 01.01.2022 to 16.01.2022.

Date	Low Level				Forecast Central Point	Real Data	Forecast Central Point	Upp Level			
	99.99%	99%	95%	67%				67%	95%	99%	99.99%
	Noticeable, Significant, Sharp Improvement of Pre-Forecast State				Pre-Predicted State Preservation		Pre-Predicted State Preservation		Noticeable, Significant, Sharp Deterioration of Pre-Forecast State		
	COVID-19 Infection Cases										
9/01/2021- 9/30/2021	1070	1965	2447	3246	4035	2064	4035	4824	5622	6121	7186
10/01/2021- 10/31/2020	40	90	220	1056	2114	3427	2114	3202	4302	4990	6457
11/01/2021- 11/30/2021	1439	2275	2674	3314	3946	4213	3946	4577	5217	5616	6469
12/01/2021- 12/31/2021	956	1921	2411	3248	4076	2874	4076	4903	5741	6264	7381
01/01/2022- 01/16/2022	4	27	46	86	2102	3809	2102	3850	5619	6724	9083
	COVID-19 Death Cases										
9/01/2021- 9/30/2021	34	53	61	75	88	51	88	102	115	124	142
10/01/2021- 10/31/2020	1	11	18	30	41	35	41	53	64	72	87
11/01/2021- 11/30/2021	1	5	12	29	46	67	46	62	79	89	112
12/01/2021- 12/31/2021	5	16	29	53	77	56	77	101	125	140	172
01/01/2022- 01/16/2022	2	13	22	37	51	41	51	65	79	88	107
	COVID-19 Infection Rate, %										
9/01/2021- 9/30/2021	3.2	5.6	6.7	8.4	10.2	6.7	10.2	12.0	13.7	14.9	17.2
10/01/2021- 10/31/2020	0.0	0.5	1.4	3.6	5.8	8.1	5.8	8.0	10.2	11.6	14.6
11/01/2021- 11/30/2021	0.7	1.8	3.2	6.1	8.9	9.6	8.9	11.7	14.6	16.3	20.2
12/01/2021- 12/31/2021	0.6	1.7	3.1	6.1	9.1	6.5	9.1	12.1	15.1	16.9	20.9
01/01/2022- 01/16/2022	0.7	2.2	3.1	4.5	5.8	9.0	5.8	7.2	8.6	9.5	11.3

Table 9. Change in the forecast state of C, D and I in relation to the pre-predicted one according to Table 1 scale [8].

Date	COVID-19 Infection Cases
9/01/2021- 9/30/2021	Significant improvement, stability of a time-series of observations
10/01/2021-10/31/2020	Noticeable deterioration, stability of a time-series of observations
11/01/2021-11/30/2021	Preservation, stability of a time-series of observations
12/01/2021-12/31/2021	Noticeable improvement, stability of a time-series of observations
01/01/2022-01/16/2022	Preservation, stability of a time-series of observations
COVID-19 Death Cases	
9/01/2021- 9/30/2021	Sharp improvement, stability of a time-series of observations
10/01/2021-10/31/2020	Preservation, stability of a time-series of observations
11/01/2021-11/30/2021	Noticeable deterioration, stability of a time-series of observations
12/01/2021-12/31/2021	Preservation, stability of a time-series of observations
01/01/2022-01/16/2022	Preservation, stability of a time-series of observations
COVID-19 Infection Rate	
9/01/2021- 9/30/2021	Noticeable improvement, stability of a time-series of observations
10/01/2021-10/31/2020	Noticeable deterioration, stability of a time-series of observations
11/01/2021-11/30/2021	Preservation, stability of a time-series of observations
12/01/2021-12/31/2021	Preservation, stability of a time-series of observations
01/01/2022-01/16/2022	Significant deterioration, stability of a time-series of observations

For all forecast periods (4 monthly forecasting cases and one 16-days forecasting case), the stability of real time series of observations of these parameters (period for calculating the forecast + forecast period) remained (Table 9).

Thus, the daily monthly and mean monthly forecasted values of C, D and I quite adequately describe the temporal changes in their real values.

Despite the rapid spread of the omicron COVID variant an interval forecast check of confirmed COVID-19 cases, deaths and infection rates in Georgia from 01.01.2022 to 16.01.2022 confirms the representativeness of the monthly forecast for the specified time period (Table 8 and 9) [<https://www.facebook.com/Avtandil1948/>]. Further monitoring will determine the representativeness of this monthly forecast for the end of January 2022 in the context of the spread of mixed strains of coronavirus (Delta and Omicron) in Georgia.

Conclusion

In the future, it is planned to continue regular similar studies for Georgia in comparison with neighboring and other countries.

References

1. World Health Organization. (2020). Coronavirus Disease 2019 (COVID-19). Situation report. 67.
2. Covid-19 in Georgia. (2020). Report of the National Center for Disease Control & Public Health. The Fourth Revision. 60 p. <https://test.ncdc.ge/Handlers/GetFile.ashx?ID=363920f6-716c-4be0-8b46-fddeb0d60231>
3. One Year with COVID-19. (2021). Report of the National Center for Disease Control & Public Health. The Fifth Revision. Georgia. 68 p. <https://ncdc.ge/api/api/File/GetFile/20ed3f9c-54bd-42e2-a58a-dffea6169fc6>.
4. Covid-19 in Georgia. (2021). Report of the National Center for Disease Control & Public Health. The Six review. 93 p. (in Georgian). <http://test.ncdc.ge/Handlers/GetFile.ashx?ID=07ad44ba-95c0-4a9b-9682-f19ded67d51a>
5. Covid-19 in Georgia. (2021). Report of the National Center for Disease Control & Public Health. The Seventh Revision. 56 p. <https://test.ncdc.ge/Handlers/GetFile.ashx?ID=a8ec06ae-21c6-43f1-9ece-11d5fcd83e1d>
6. Amiranashvili A.G, Khazaradze K.R, Japaridze N.D. (2020). Twenty weeks of the pandemic of coronavirus Covid-19 in Georgia and neighboring countries (Armenia, Azerbaijan, Turkey, Russia). Preliminary comparative statistical data analysis. Int. Sc. Conf. „Modern Problems of Ecology“, Proc., ISSN 1512-1976, v. 7, Tbilisi-Telavi, Georgia, 26-28 September, 2020, pp. 364-370.
7. Amiranashvili A.G., Khazaradze K.R., Japaridze N.D. (2020). Analysis of twenty-week time-series of confirmed cases of New Coronavirus COVID-19 and their simple short-term prediction for Georgia and neighboring countries (Armenia, Azerbaijan, Turkey, Russia) in amid of a global pandemic. medRxiv preprint doi: <https://doi.org/10.1101/2020.09.09.20191494>, 13 p. Europe PMC, <https://europepmc.org/article/ppr/ppr213467>
8. Amiranashvili A.G., Khazaradze K.R., Japaridze N.D. (2021). The Statistical Analysis of Daily Data Associated with Different Parameters of the New Coronavirus COVID-19 Pandemic in Georgia and their Short-Term Interval Prediction from September 2020 to February 2021. medRxiv preprint doi: <https://doi.org/10.1101/2021.04.01.21254448>, 18 p.
9. Amiranashvili A.G., Khazaradze K.R., Japaridze N.D. (2021). The Statistical Analysis of Daily Data Associated with Different Parameters of the New Coronavirus COVID-19 Pandemic in Georgia and

- their Short-Term Interval Prediction in Spring 2021. medRxiv preprint doi: <https://doi.org/10.1101/2021.06.16.21259038>
10. Amiranashvili A.G., Khazaradze K.R., Japaridze N.D. (2021). The Statistical Analysis of Daily Data Associated with Different Parameters of the New Coronavirus COVID-19 Pandemic in Georgia and their Two-Week Interval Prediction in Summer 2021. medRxiv preprint, 2021, doi: <https://doi.org/10.1101/2021.09.08.21263265>, 20 p.
 11. Amiranashvili A., Khazaradze K., Japaridze N., Revishvili A. (2021). Analysis of the Short-Term Forecast of Covid-19 Related Confirmed Cases, Deaths Cases and Infection Rates in Georgia from September 2020 to October 2021. International Scientific Conference „Natural Disasters in the 21st Century: Monitoring, Prevention, Mitigation“. Proceedings, ISBN 978-9941-491-52-8, Tbilisi, Georgia, December 20-22, 2021. Publish House of Iv. Javakhishvili Tbilisi State University, Tbilisi, 2021, pp. 167 - 171.
 12. Leffler Ch. T., Joseph D. Lykins J.D., Yang E. (2021). Preliminary Analysis of Excess Mortality in India During the Covid-19 Pandemic (Update August 4, 2021). medRxiv preprint doi: <https://doi.org/10.1101/2021.08.04.21261604>
 13. Rethnagar N. (2021). Statistical Inferences and Analysis based on the COVID-19 Data from the United States. medRxiv preprint doi: <https://doi.org/10.1101/2021.11.30.21267096>, 42 p.
 14. Fahad Mostafa, Riya Ganji, Julie St. John, Hafiz Khan. (2021). Coronavirus (COVID-19) Spike in Georgia: An Epidemiologic Study of Data, Modelling, and Policy Implications to Understand the Gender-and Race-Specific Variations. medRxiv preprint doi: <https://doi.org/10.1101/2021.12.09.21267571>, 37 p.
 15. Maldonado A., Reyes M. (2021). Distribution of Confirmed with COVID-19 by Age and Gender in Mexico. medRxiv preprint doi: <https://doi.org/10.1101/2021.11.21.21264092>. 13 p.
 16. Ball S. (2021). Data Analysis of Covid-19 Pandemic and Short-Term Cumulative Case Forecasting Using Machine Learning Time Series Methods. Chaos Solitons Fractals. 142, 110512. doi:10.1016/j.chaos.2020.110512
 17. Oshinubi K., Al-Awadhi F., Rachdi M., Demongeot J. (2021). Data Analysis and Forecasting of COVID-19 Pandemic in Kuwait. medRxiv preprint doi: <https://doi.org/10.1101/2021.07.24.21261059>
 18. Reingruber J., Papale A., Ruckly S., Timsit J-F., Holcman D. (2021). Monitoring and Forecasting the SARS-Covid-19 Pandemic in France. medRxiv preprint doi: <https://doi.org/10.1101/2021.07.28.21260870>
 19. Liu Z., Magal P., Webb G. (2021) Predicting the Number of Reported and Unreported Cases for the Covid-19 Epidemics in China, South Korea, Italy, France, Germany and United Kingdom. Journal of theoretical biology, vol. 509, p. 110501
 20. Keeling M. J., Hill E., Gorsich E., Penman B., Guyver-Fletcher G., Holmes A., Leng T., Mc Kimm H., Tamborrino M., Dyson L., Tildesley M. (2021). Predictions of Covid-19 Dynamics in the UK: Short-Term Forecasting and Analysis of Potential Exit Strategies. PLOS Computational Biology, vol. 17, pp. 1-20.
 21. Khalilpourazari S., Doulabi H.H., Çiftçioğlu A. Ö., Weber G-W. (2021). Gradient-Based Grey Wolf Optimizer with Gaussian Walk: Application in Modelling and Prediction of the COVID-19 Pandemic. Expert Systems with Applications, Vol. 177, 114920. <https://doi.org/10.1016/j.eswa.2021.114920>
 22. Katris C. (2021). A Time Series-Based Statistical Approach for Outbreak Spread Forecasting: Application of COVID-19 in Greece. Expert Syst. Appl., 166, Article 114077.
 23. Liu L., Moon H.R., Schorfheide F. (2021). Panel Forecasts of Country-Level COVID-19 Infections. J. Econometrics, 220, pp. 2-22.
 24. Efimov D., Ushirobir R. (2021). On an Interval Prediction of COVID-19 Development Based on a SEIR Epidemic Model. Annual Reviews in Control, 51, pp. 477–487. <https://doi.org/10.1016/j.arcontrol.2021.01.006>

25. Leonardo Sestrem de Oliveira, Sarah Beatriz Gruetzmacher, João Paulo Teixeira. (2021). COVID-19 Time Series Prediction. *Procedia Computer Science*, 181, pp. 973–980.
26. Faheem Aslam, Tahir Mumtaz Awan, Ruqia Khan, Maira Aslam, Yasir Tariq Mohmand. (2021). Prediction of COVID-19 Confirmed Cases in Indo-Pak Sub-Continent. *J Infect Dev Ctries*, 15(3), pp. 382-388. doi:10.3855/jidc.13419
27. Gecili E., Ziady A., Szczesniak R.D. (2021). Forecasting COVID-19 Confirmed Cases, Deaths and Recoveries: Revisiting Established Time Series Modeling Through Novel Applications for the USA and Italy. *PLoS ONE* 16(1): e0244173, 11 p., <https://doi.org/10.1371/journal.pone.0244173>
28. Barriá-Sandoval C., Ferreira G., Benz-Parra K., López-Flores P. (2021) Prediction of confirmed cases of and deaths caused by COVID-19 in Chile through time series techniques: A comparative study. *PLoS ONE* 16(4), e0245414, 16 p., <https://doi.org/10.1371/journal.pone.0245414>
29. Zhao H., Merchant N.N., McNulty A., Radcliff T.A., Cote M.J., Fischer R.S.B., et al. (2021). COVID-19: Short Term Prediction Model using Daily Incidence Data. *PLoS ONE* 16(4), e0250110, 14 p., <https://doi.org/10.1371/journal.pone.0250110>
30. Reddy T., Shkedy Z., van Rensburg Ch.J., Mwambi H., Debba P., Zuma Kh., Manda S. (2021). Short-Term Real-Time Prediction of Total Number of Reported COVID-19 Cases and Deaths in South Africa: A Data Driven Approach. *BMC Medical Research Methodology*, 21:15, 11 p., <https://doi.org/10.1186/s12874-020-01165-x>
31. Tingqing Ye, Xiangfeng Yang. (2021). Analysis and Prediction of Confirmed COVID-19 Cases in China with Uncertain Time Series. *Fuzzy Optimization and Decision Making*, 20, pp. 209–228, <https://doi.org/10.1007/s10700-020-09339-4>
32. Keeling M.J. et al. (2021). Short-Term Projections Based on Early Omicron Variant Dynamics in England. *medRxiv preprint* doi: <https://doi.org/10.1101/2021.12.30.21268307>, 30 p.
33. Sheikh Muzaffar Shakeel, Nithya Sathya Kumar, Pranita Pandurang Madalli ,Rashmi Srinivasaiah, Devappa Renuka Swamy. (2021). COVID-19 Prediction Models: a Systematic Literature Review. *Osong Public Health Res Perspect*, 12(4), pp. 215-229, <https://doi.org/10.24171/j.phrp.2021.0100>
34. Iman Rahimi, Fang Chen, Amir H. Gandomi. (2021). A Review on COVID-19 Forecasting Models. *Neural Computing and Applications*, 11 p., <https://doi.org/10.1007/s00521-020-05626-8>
35. Brozaky S.J., Panty B., Safdary S., Gumely A.B. (2021). Dynamics of COVID-19 Pandemic in India and Pakistan: A Metapopulation Modelling Approach. *medRxiv preprint* doi: <https://doi.org/10.1101/2021.08.02.21261459>
36. Basak A., Rahaman S., Guha A., Sanyal T. (2021). Dynamics of the Third Wave, Modelling COVID-19 Pandemic with an Outlook Towards India. *medRxiv preprint* doi: <https://doi.org/10.1101/2021.08.17.21262193>
37. Bosman M., Esteve A., Gabbanelli L., Jordan X., López-Gay A., Manera M., Martínez M., Masjuan P., Mir Ll. M., Paradells J., Pignatelli A., Riu I, Vitagliano V. (2021). Stochastic simulation of successive waves of COVID-19 in the province of Barcelona. *medRxiv preprint* doi: <https://doi.org/10.1101/2021.12.02.21266468>, 17 p.
38. Sabara Parshad Rajeshbhai, Subhra Sankar Dhar, Shalabh. (2021). Statistical Forecasting: Third Wave of COVID-19- With an Application to India. *medRxiv preprint* doi: <https://doi.org/10.1101/2021.12.20.21268150>. 14 p.
39. Förster E., Rönz B. (1979). Methoden der korrelations - und regressions analyse. – Ein Leitfaden für Ökonomen. Verlag Die Wirtschaft Berlin. 324.
40. Kendall MG. (1981). Time-series. Moscow, 200, (in Russian).
41. Hinkle D. E., Wiersma W., Jurs S. G. (2003). Applied Statistics for the Behavioral Sciences. Boston, MA, Houghton Mifflin Company.