PLWHA's Perspective on Community Stigma for Getting Social Support and Improving Life Quality in the Digital Era

Rio Ady Erwansyah¹, Moses Glorino Rumambo Pandin²

¹Student of Doctoral Program in Nursing, Faculty of Nursing, Airlangga University

²Faculty of Humanities, Airlangga University

Corresponding author email : rio.ady.erwansyah-2021@fkip.unair.ac.id;
moses.glorino@fib.unair.ac.id

ABSTRACT

Community stigma against HIV/AIDS is still a problem that must be faced by people living with HIV/AIDS (PLWHA). The existence of stigmatization and discrimination received by PLWHA causes the lack of social support they get. Social support is very important for PLWHA to improve their quality of life. In this digital era where the development of technology and social media is an alternative for PLWHA to get social support online through their community, by sharing information and experiences. This systematic review was conducted through Scopus, Science Direct, ProQuest, and SAGE. The selection of articles followed inclusion criteria, including articles published in the last five years and published in English and discussing perspectives on stigma. Articles published not in English and articles with unclear literature reviews were excluded from this study. The PRISMA flow chart and JBI assessment checklist were used to assess the risk of bias and article quality. 20 relevant articles will be reviewed. The results of the study found that in this digital era PLWHA can take advantage of technological developments and social media to improve their quality of life by getting social support through the online communities they follow.

Keywords: *Social Support, Stigma, PLWHA/ODHA, HIV, AIDS*

1. Background

Stigma against HIV/AIDS remains a significant barrier to public health worldwide and a major barrier to HIV/AIDS treatment, prevention, and support. Stigma against people living with HIV/AIDS (PLWHA) leads to reduced economic income and increased isolation from family and friends regarding their HIV status¹. PLWHA often experience physical, psychological, social, and spiritual suffering, which harms their quality of life²⁻³

PLWHA face several difficulties when they try to achieve a satisfactory quality of life, from disturbances in their life history, disturbances in interpersonal and work relationships that can cause social isolation, and worsen social relationships that can harm their mental and physical health⁴. In this digital era, it is still difficult for them to adapt to their current status changes regarding the diagnosis they receive. Stigma and community discrimination against HIV/AIDS are still considered a major problem in the survival of PLWHA, making it difficult for them to get social support from both the community and family. Whereas social support

from the community, especially families, can help PLWHA improve their quality of life and assist in the acceptance and disclosure of their serostatus.

2. Objective

The purpose of this systematic review is to explore the perspective of PLWHA on community stigma in obtaining social support and improving their quality of life in the digital age.

3. Methodology

This research is a review of articles conducted to find out the perspective of PLWHA in the digital era. The checklist and flow chart of reporting items for systematic review (PRISMA) was used to present the results of the systematic review. We entered electronic databases to find relevant articles such as Scopus, Science Direct, ProQuest, and SAGE. The literature search process was determined in a recent study of the last five years. There are no regulatory restrictions, but only articles published in English are reviewed. The feasibility of this study was assessed using the PICOT framework as below.

Table 1 Picot Framework

PICOT Framework	Inclusion Criteria
Population	PLWHA
Intervention	Explore experiences and perspectives
Comparator	No comparator
Output	PLWHA perspective
Time	2017-2020

Articles are identified with the keywords “perspective, PLWHA, HIV, AIDS”.

Bias Risk

The risk of bias can be reduced by using critical assessment tools. The review used a critical assessment tool from the Joanna Briggs Institute (JBI). The JBI assessment tool is a checklist questionnaire with a list of required items, things to do, and points to consider. Each research design has different questions. Researchers must assess the articles that have been selected. The scoring results come from low risk, medium risk, high risk, or unclear risk.

Inclusion and exclusion criteria: the articles used, filtered, and selected based on the inclusion criteria that have been set with the provisions of 5 years from 2017 to 2021, contain a discussion of perspectives on stigma. Articles that pass the inclusion criteria are then eliminated by the exclusion criteria. Exclusion criteria were referred to when the article was not written in English, the article was more than five years old, and the literature had an unclear review. The data that has been obtained is reviewed and then selected to be further grouped and discussed based on the points

4. Result

An initial keyword search of the literature yielded 395 articles (59 from Scopus, 87 from Science Direct, 218 from ProQuest, and 31 from SAGE). After a review, 125 articles were eliminated because the population did not match the research objectives, the articles did not use English, and were not open access. Then 169 articles were excluded based on the last five years and did not include a literature review, and 101 complete articles that were assessed for eligibility were eliminated 81 articles because the articles did not discuss perspectives and the literature had an unclear review so that 20 articles were obtained that were following the feasibility of the study.

Figure 1 The results of the literature selection are summarized in the PRISMA flow diagram

The risk of bias in the articles that we found can be seen in the following table:

Table 2 Risk assessment bias using the JBI critical assessment checklist

No	Author, Year	1	2	3	4	5	6	7	8	9	10	11	12	13	Total
1	X. Han et al., 2018	v	v	v	v	v	x	v	v						87,5%
2	N. O'Brien et al., 2017	v	v	v	v	v	x	v	v	v	v				90%
3	Zepeda KGM et al., 2019	v	v	v	v	v	x	v	v	v	v				90%
4	Giraud, J.S, et al., 2020	v	v	v	x	x	v	v	v	v	v	v			81,8%
5	Mehraeen et al., 2021	v	v	v	v	x	x	v	v						75%
6	Haruna T. et al., 2021	v	v	v	v	v	x	x	v	v	v				80%
7	S.M. Rein et al., 2021	v	v	v	x	x	v	v	v	v	v	v			81,8%
8	C. R. Fumaz et al., 2019	v	v	v	v	x	x	v	v						75%
9	Ahmed et al., 2019	v	v	v	v	v	x	v	v	v	v				90%
10	M. Abboah-Offei et al., 2019	v	v	v	v	v	x	x	v	v	v				80%
11	Ernawati et al., 2021	v	v	v	v	v	v	x	v	v	v				90%
12	Fauk et al., 2021	v	v	v	v	v	v	x	v	v	v				90%
13	K. Srikant h Reddy, Seema Sahay, 2018	v	v	v	v	v	v	x	v	v	v				90%
14	Coelho B, Meirelles BHS., 2019	v	v	v	v	v	x	x	v	v	v				80%
15	Jesus GJ, et al., 2017	v	v	v	v	v	x	v	v	v	v				90%
16	Xie, Fie et al., 2019	v	v	v	v	x	v	v	v						87,5%
17	Rutakumwa e	v	v	v	v	v	x	v	v	v	v				90%

t al., 2021												80%
18	Rice et al., 2018	v	v	v	v	v	x	x	v	v		
19	Cherry S. at al., 2018	v	v	v	v	v	x	v	v	v		90%
20	N. S. Perry et al., 2017	v	v	v	v	v	x	v	v	v		90%

The Joanna Briggs Institute (JBI) Critical Assessment was used to assess and analyze the methodological quality of the articles obtained (n = 20). As agreed by the researcher himself that the final score of the methodological quality assessment if it reaches a minimum of 75% meets the critical assessment criteria, the article will be included for further data synthesis. All articles (n = 20) at the last screening achieved a score higher than 75% so they were ready for data synthesis.

Based on table 2, 14 articles were assessed for risk of bias using the JBI Critical Assessment Checklist for Qualitative Studies and the results were: a score of 90% (n=10 articles) (N. O'Brien et al., 2017, Zepeda KGM et al. , 2019, Ahmed et al., 2019, Ernawati et al., 2021, Fauk et al., 2021, K. Srikanth Reddy, Seema Sahay, 2018, Jesus GJ, et al., 2017, Rutakumwa et al., 2021, Cherry S. at al., 2018, NS Perry et al., 2017), score 80% (n=4 articles) (Haruna T. et al., 2021, M. Abboah-Offei at al., 2019, Coelho B, Meirelles BHS., 2019, Rice et al., 2018)

4 articles were assessed for risk of bias using the JBI critical assessment checklist for the Cross-Sectional Study and the results were: a score of 87.5% (n=2 articles) (X. Han et al., 2018, Xie, Fie, et al., 2019), score 75% (n=2 articles) (CR Fumaz et al., 2019, Mehraeen et al., 2021).

2 articles were assessed for risk of bias using the JBI critical appraisal checklist for a cohort study and the results were: a score of 81.8% (n=2 articles) (Giraud, J.S, et al., 2020, S.M. Rein et al., 2021). Based on the identification in table 1, 20 articles were assessed for risk of bias using the JBI critical assessment checklist for the design of Cross-Sectional Study, Cohort Study, Prevalence Study, and QuasiExperimental Study. The results found were 20 articles had scores above 75%.

This systematic review was heterogeneous, with 14 articles using qualitative methods in the study, 4 cross-sectional articles, and 2 cohort articles. Results from selected articles using single and combined survey interventions. The sample varies because the journals selected are heterogeneous. The types of interventions provided are described in the following table:

1. Weibo friends with benefits for people living with HIV/AIDS? The implications of Weibo use for enacted social support, perceived social support, and health outcomes

With the development of social media. Online support is becoming increasingly important for people with health problems such as HIV/AIDS to seek social support. Where the platform generates discussions on various health topics and becomes a place to share information and experiences ⁵⁻⁶

2. Envisioning Women-Centered HIV Care: Perspectives from Women Living with HIV in Canada
A women-centered approach to HIV care is critical to guiding policy and practice to improve care and outcomes for women living with HIV ⁷.
3. Management of nursing care in HIV/AIDS from a palliative and hospital perspective
In the context of palliative care, actions to improve comfort and quality of life are emphasized in the physical, socio-cultural, and psycho-spiritual fields in an altruistic way, to put oneself in the position of others to jointly try to understand the suffering of patients and families ⁸
4. De-simplifying single-tablet antiretroviral treatments for cost savings in France: From the patient perspectives to a 6-month follow-up on generics
Simplification of ART treatment to generic drugs allows cost savings in health care costs ⁹
5. Investigating the contributing factors to HIV/AIDS infection from the perspective of HIV-infected patients
An important cause of the spread of HIV is the government's weakness in public health education, family and individual factors ¹⁰
6. Factors hindering the integration of care for non-communicable diseases within HIV care services in Dar es Salaam, Tanzania: The perspectives of health workers and people living with HIV
NCD service bundling in HIV care and treatment clinics has adequate treatment adherence support to improve quality of life, close monitoring of biomarkers that can predict the development of NCDs and social support providers ¹¹
7. Prospective association of social circumstance, socioeconomic, lifestyle, and mental health factors with subsequent hospitalization over 6-7 year follow up in people living with HIV
Current markers of socioeconomic disadvantage (non-employment, insecure housing situations, and financial difficulties) are strong predictors of hospitalization among PLWHA ¹²
8. Health-related quality of life of people living with HIV infection in Spain: a gender perspective
The stereotypic differences associated with each sex may explain the perceived differences in quality of life. Men adopt healthy behaviors, helping to improve physical and mental health, emotional status, and a better quality of life through physical exercise. While women appear to be affected by their quality of life by personal experiences of rejection, receiving social stigma from various community sources affects their physical health and coping strategies used in response to HIV-related stress ¹³⁻¹⁴
9. Facilitators and Barriers Affecting Adherence Among People Living With HIV/AIDS: A Qualitative Perspective
Quality nursing outcomes depend on patient medication adherence. Lack of social support and poor doctor-patient communication often force patients to bear the burden of illness that affects their social and psychological well-being ¹⁵

10. How can we achieve person-centered care for people living with HIV/AIDS?
A qualitative interview study with healthcare professionals and patients in Ghana PCC as care delivered in the community, free from stigma and discrimination, that addresses financial, human, and time resource issues¹⁶
11. Life Experiences in Parenting: The Perspective of Women with HIV-AIDS
The form of hope for negative HIV status from loved ones, the lamentation of wives with PLWHA status in the hope that they will remain healthy and hope that their children will not experience the same disease¹⁷
12. HIV Stigma and Discrimination: Perspectives and Personal Experiences of Healthcare Providers in Yogyakarta and Belu, Indonesia
Stigma and discrimination are major challenges facing PLWHA globally because of their HIV status. Lack of knowledge about HIV, fear of being infected, personal values, religious thoughts, and socio-cultural values and norms are the driving forces for HIV-related stigma¹⁸
13. Advocacy for HIV/TB co-infection collaborative policy and service delivery in India: a civil society perspective
HIV prevention is a priority for TB control, and TB care and prevention is a priority for HIV/AIDS programs¹⁹
14. Care sharing for people with HIV/AIDS: a look targeted at young adults
The Health Care Network (HCN) is an organization that regulates health services, including technology, which are interconnected by support systems, seeking to ensure comprehensive care²⁰
15. Difficulties of living with HIV/Aids: Obstacles to quality of life
The impact of living with a chronic disease that still carries a lot of stigma and prejudice is the biggest obstacle between these individuals and their quality of life²¹
16. Social Capital Associated with Quality of Life among People Living with HIV/AIDS in Nanchang, China
Bonding social capital score has a positive effect on mental health summary and physical health summary²²
17. Stakeholders' perspectives on integrating the management of depression into routine HIV care in Uganda: qualitative findings from a feasibility study
Integrating depression management into routine HIV care is acceptable among key stakeholders²³
18. Perceptions of Intersectional Stigma among Diverse Women Living with HIV in the United States
Women living with HIV experience interrelated forms of stigma. The complex social environment in which HIV-positive women navigate shapes their realities of life, life opportunities, and well-being²⁴
19. "My determination is to live": Narratives of African-American Women Who Have Lived with HIV for 10 or More Years
Public health programs for long-term survivors, including ways to promote resilience, create opportunities for meaningful work, build skills and disclosure

20. Learning to address multiple syndemics for people living with HIV through client perspectives on CBT

Future psychotherapeutic interventions for HIV syndromes and treatments address complex issues comprehensively²⁶

No	Article Title	Intervention	Result
1	Weibo friends with benefits for people living with HIV/AIDS? The implications of Weibo use for enacted social support, perceived social support, and health outcomes	Filling out online questionnaires for HIV/AIDS positive Weibo users	Better socioeconomic conditions and poorer physical conditions were associated with higher PSS-ON rates. PLWHA experiences a higher level of social support via Weibo than offline.
2	Envisioning WomenCentered HIV Care: Perspectives from Women Living with HIV in Canada	Conduct focus group discussions with semistructured interviews to listen to the experiences of women living with HIV before shifting to women's perspectives on a women-centered	Women envision the 3 main characteristics of women-centered HIV care: coordinated and integrated services that address HIV and women's health care priorities that protect against exclusion from

		approach to HIV care	treatment because of stigma, care that recognizes and responds to structural barriers, and cares that encourages peer support and leadership. same age
3	Management of nursing care in HIV/AIDS from a palliative and hospital perspective	Conduct semi-structured interviews in two sample groups with 17 participants	Generating an understanding of the relationship between nurse interactions in the management of nursing care for people who have clinical complications caused by AIDS and its relationship to palliative care

4	De-simplifying singletablet antiretroviral treatments for cost savings in France: From the patient perspectives to a 6-month follow-up on generics	98 patients received a two-tablet treatment regimen containing the generic drug. 60 received Eviplera® and 38 received Triumeq®, six months later, a second survey was administered to patients	45 patients received treatment that did not refuse simplified, 37 refused and 16 did not decide and followed the decision offered by their doctor
5	Investigating the contributing factors to HIV/AIDS infection from the perspective of HIV-infected patients	150 patients with HIV were interviewed and filled out a medication adherence questionnaire and a mental health standard questionnaire	The results revealed that the main cause of HIV infection in men was an injection of drugs and in women was sexual intercourse with an infected individual.
6	Factors hindering the integration of care for non-communicable diseases within HIV care services in Dar es Salaam, Tanzania: The perspectives of health workers and people living with HIV	A guided checklist was observed for 5 days at the HIV care and treatment clinic on 130 PLWHA who provided PTM treatment services. Then conducted interviews with 5 CTC managers, 9 health care providers, and 27 PLWHA	There is a positive attitude between PLWHA and health workers towards the integration of noncommunicable disease services in HIV care and treatment clinics. This is due to the perceived benefits associated with the integration of PTM services in one clinic
7	Prospective association of social circumstance, socioeconomic, lifestyle, and mental health factors with subsequent hospitalization over 6-7	Conducted the ASTRA study by recruiting PLWHA from 8 HIV outpatient clinics with a total of 798 participants and asking them to fill	Socio-economic losses, social conditions, evidence of mental health disorders, and adverse lifestyle factors are hospitalization for PLWHA
	year follow up in people living with HIV	out a baseline questionnaire from February 2011 to June 2018	

8	Health-related quality of life of people living with HIV infection in Spain: a gender perspective	Evaluated gender differences in healthrelated quality of life in 744 PLHIV drawn from 32 Spanish hospitals offering HIV care, and participants were asked to complete the study assessment protocol with the supplied questionnaire.	Women who showed more emotional distress and negative moods related to their HIV condition used less positive coping strategies and experienced more rejection and stigma than men.
9	Facilitators and Barriers Affecting Adherence Among People Living With HIV/AIDS: A Qualitative Perspective	Conduct in-depth interviews with 18-yearold patients diagnosed with HIV/AIDS about medication adherence	Anxiety and stigma, the negative perceived image of HIV diagnosis, poor support from the community, and side effects of treatment are factors for treatment nonadherence.
10	How can we achieve person-centered care for people living with HIV/AIDS? A qualitative interview study with healthcare professionals and patients in Ghana	Conducted semi-structured interviews with PLWHA and health professionals at the community clinic aimed at participants' perspectives on their illness, access to care, involvement in decisionmaking, their symptoms, concerns, and preferences for care.	This study provides new insights into what patientcentered care (PCC) is all about. People with HIV understand PCC as care that involves them in their care decisions.
11	Life Experiences in Parenting: The Perspective of Women with HIV-AIDS	Conducted interviews with wives who were diagnosed with HIV/AIDS and registered by the HIV-AIDS program at the Serang City Health Center, to explore the phenomenon of mothers with HIV/AIDS in caring for their children.	The results showed that the mother wanted her child or husband to have a negative HIV status.

12	HIV Stigma and Discrimination: Perspectives and Personal Experiences of	Conducted interviews with 92 PLWHA and 20 health care providers regarding stigma and	Perspectives and experiences of health care providers regarding HIV stigma and discrimination
	Healthcare Providers in Yogyakarta and Belu, Indonesia	discrimination against PLWHA	still occur in families and communities. Reflected in negative labeling by others
13	Advocacy for HIV/TB co-infection collaborative policy and service delivery in India: a civil society perspective	Conducted interviews with 26 participants consisting of representatives of advocacy organizations, health activists and PLWHA	Civil society prioritizes HIV/TB coinfection is an important public health issue and advocates identifying common indicators of coinfection for better program management
14	Care sharing for people with HIV/AIDS: a look targeted at young adults	Conducting interviews with 19 participants with HIV/AIDS who were divided into 2 groups was shown to understand the sharing of care for PLWHA	City governments have transitioned into a model of health care for people with HIV/AIDS, whose care is shared within the health care network.
15	Difficulties of living with HIV/Aids: Obstacles to quality of life	Conducted interviews with 26 PLWHA about quality of life	There are difficulties faced by PLWHA. The main difficulties cited are prejudice experienced in family and social contexts, managing affective and sexual partnerships, and achieving quality of life
16	Social Capital Associated with Quality of Life among People Living with HIV/AIDS in Nanchang, China	225 PLWHA in Nanchang, China was asked to fill out a questionnaire to identify social capital factors that identify the quality of life in PLWHA	People living with HIV have low social capital and poor quality of life in Nanchang. Social bonding capital has a positive effect on the quality of life of PLWHA

17	Stakeholders' perspectives on integrating the management of depression into routine HIV care in Uganda: qualitative findings from a feasibility study	Conducted interviews with 11 PLWHA patients, doctors, supervisors, and mental health specialists and group-focused discussions by lay health workers. Interviews were conducted to evaluate HIV+D Interventions	Patients highlight the benefits of treating depression in the context of HIV care, including improved adherence to antiretroviral therapy, coping with sleeplessness, and suicidal ideation, and regaining a sense of self-efficacy.
18	Perceptions of Intersectional Stigma among Diverse Women Living with HIV in the United States	Conducted interviews with 76 women living with HIV from various socioeconomic backgrounds	Women living with HIV feel many forms of stigma apart from the stigma associated with HIV. The most common forms of stigma and discrimination are related to their gender,
			race, and income level.
19	"My determination is to live": Narratives of African-American Women Who Have Lived with HIV for 10 or More Years	Conducting interviews with women living with HIV for 10 years	The need for specific programs for long-term survivors, including resilience training, educational programs on dating and disclosure, and opportunities to engage in meaningful work or volunteer initiatives
20	Learning to address multiple syndemics for people living with HIV through client perspectives on CBT	Conduct interviews with multiple comorbid PLWHA who complete cognitive-behavioral therapy (CBT)	Mental health issues are interrelated and provide an informative insight into how participants perceive the interconnections between their mental health diagnoses, and how they behave in their health.

5. Discussion

The findings from this systematic review found that the perspective of PLWHA on stigma where stigma is still a major problem in improving their quality of life. Research by Jesus GJ, et al. (2017) the impact of living with a chronic disease that still carries a lot of stigma and prejudice is the biggest obstacle to PLWHA regarding their quality of life⁷. It is difficult for PLWHA to improve their quality of life in community stigma that still exists today so that they do not receive

social support in carrying out their lives. Through social support, especially family members, it is very important to help PLWHA accept their diagnosis and choose a healthy life and find ways to grow²⁵.

Many PLWHA expressed their perspective that it was difficult for them to be involved in a social environment because of the stigmatization of their HIV/AIDS status and made it difficult for them to seek social support from friends and family members. With the social conditions of PLWHA like that, they also lack emotional support so this can have an impact on their clinical condition. In this digital era, the development of information technology and social media makes PLWHA try to find alternatives in overcoming their social problems. PLWHA can seek support through online communities that they can access via the internet, where they can get information about their illness and find a community that suits them, so they can share experiences and their daily lives. In this way, PLWHA can get the social support that has an impact on their quality of life-related to their emotional needs. In the research of X. Han et al. (2018) with the development of social media in the PLWHA community, they can receive social support from online communities, where they can exchange information and share their experiences, as well as seek the needed emotional support²⁷.

In addition to the social support they seek through online communities, there is also a need for programs from the government that help PLWHA reduce the stigma they receive so that they can receive social support to improve their quality of life and their economic status. Several studies discuss the perspective of PLWHA about the need for programs to address social problems that occur. In the study of M. AbboahOffei et al. (2019) PLWHA understands patient-centered care (PCC) programs not only involve them in making decisions about their care but also as a way to deal with broader social problems such as living a normal life like other people, getting married, having children and being employed¹⁶.

The limitations and weaknesses of this research lie in the process and systematic review, where the researcher realizes that in a study there must be many obstacles and obstacles. One of the factors that became an obstacle in this study was the limitation of language and time, where the researcher used only a literature review that used English where had an effect on the results obtained.

6. Conclusion

In this digital era, the perspective of PLWHA on stigma is still the main problem they face. However, with the development of technology and social media, they can get support from the online community they follow, so they can share experiences in improving their quality of life.

7. References

1. Mak WWS, Mo PKH, Ma GYK, Lam MYY. Meta-analysis and systematic review of studies on the effectiveness of HIV stigma reduction programs. *Soc Sci Med*. 2017;188:30-40. doi:10.1016/j.socscimed.2017.06.045
2. Harding R, Simms V, Alexander C, et al. Can palliative care integrated within HIV outpatient settings improve pain and symptom control in a low-income

- country? A prospective, longitudinal, controlled intervention evaluation. *AIDS Care*. 2013;25(7):795-804. doi:10.1080/09540121.2012.736608
3. Harding R, Simms V, Penfold S, et al. Quality of life and wellbeing among HIV outpatients in East Africa: A multicentre observational study. *BMC Infect Dis*. 2014;14(1):1-10. doi:10.1186/s12879-014-0613-1
4. Macapagal KR, Ringer JM, Woller SE, Lysaker PH. Personal Narratives, Coping, and Quality of Life in Persons Living With HIV. *J Assoc Nurses AIDS Care*. 2012;23(4):361-365. doi:10.1016/j.jana.2011.08.011
5. Nambisan P. Information seeking and social support in online health communities: Impact on patients' perceived empathy. *J Am Med Informatics Assoc*. 2011;18(3):298-304. doi:10.1136/amiajnl-2010-000058
6. Robillard JM, Johnson TW, Hennessey C, Beattie BL, Illes J. Aging 2.0: Health Information about Dementia on Twitter. *PLoS One*. 2013;8(7):1-5. doi:10.1371/journal.pone.0069861
7. O'Brien N, Greene S, Carter A, et al. Envisioning Women-Centered HIV Care: Perspectives from Women Living with HIV in Canada. *Women's Heal Issues*. 2017;27(6):721-730. doi:10.1016/j.whi.2017.08.001
8. Cunha GH da, Galvão MTG, Pinheiro PN da C, Vieira NFC. Health literacy for people living with HIV/Aids: an integrative review. *Rev Bras Enferm*. 2017;70(1):180-188. doi:10.1590/0034-7167-2015-0052
9. Giraud JS, Doisne M, Wai ACH, et al. De-simplifying single-tablet antiretroviral treatments for cost savings in France: From the patient perspectives to a 6-month followup on generics. *PLoS One*. 2020;15(9 September):1-13. doi:10.1371/journal.pone.0239704
10. Mehraeen M, Heydari M, Lankarani KB, Joulaei H, Faghih M. Investigating the contributing factors to HIV/AIDS infection from the perspective of HIV-infected patients. *BMC Psychol*. 2021;9(1):1-9. doi:10.1186/s40359-021-00513-w
11. Haregu TN, Setswe G, Elliott J, Oldenburg B. Integration of HIV/AIDS and noncommunicable diseases in developing countries: rationale, policies and models. *Int J Healthc*. 2015;1(1). doi:10.5430/ijh.v1n1p21
12. Rein SM, Smith CJ, Chaloner C, et al. Prospective association of social circumstance, socioeconomic, lifestyle and mental health factors with subsequent hospitalisation over 6–7 year follow up in people living with HIV. *EClinicalMedicine*. 2021;31:100665. doi:10.1016/j.eclinm.2020.100665
13. Nosrat S, Whitworth JW, Ciccolo JT. Exercise and mental health of people living with HIV: A systematic review. *Chronic Illn*. 2017;13(4):299-319. doi:10.1177/1742395317694224
14. Norcini Pala A, Steca P. Illness perceptions and coping strategies among individuals diagnosed with HIV. *J Behav Med*. 2015;38(4):620-631. doi:10.1007/s10865-015-9639-0

15. Ahmed SI, Farooqui M, Syed Sulaiman SA, Hassali MA, Lee CKC. Facilitators and Barriers Affecting Adherence Among People Living With HIV/AIDS: A Qualitative Perspective. *J Patient Exp.* 2019;6(1):33-40. doi:10.1177/2374373518770805
16. Abboah-Offei M, Bristowe K, Koffman J, et al. How can we achieve personcentred care for people living with HIV/AIDS? A qualitative interview study with healthcare professionals and patients in Ghana. *AIDS Care - Psychol Socio-Medical Asp AIDS/HIV.* 2020;32(12):1479-1488. doi:10.1080/09540121.2019.1698708
17. Ernawati E, Rahmawati D, Wulandari I, Afriyani F. Life experiences in parenting: The perspective of women with hiv-aids. *Open Access Maced J Med Sci.* 2021;9(T4):35-43. doi:10.3889/oamjms.2021.5760
18. Fauk NK, Ward PR, Hawke K, Mwanri L. HIV Stigma and Discrimination: Perspectives and Personal Experiences of Healthcare Providers in Yogyakarta and Belu, Indonesia. *Front Med.* 2021;8(May):1-11. doi:10.3389/fmed.2021.625787
19. Reddy KS, Sahay S. Advocacy for HIV/TB co-infection collaborative policy and service delivery in India: A civil society perspective. *HIV AIDS Rev.* 2018;17(4):259-266. doi:10.5114/hivar.2018.80257
20. Coelho B, Meirelles BHS. Care sharing for people with HIV/AIDS: a look targeted at young adults. *Rev Bras Enferm.* 2019;72(5):1341-1348. doi:10.1590/0034-7167-2018-0248
21. Article O. Difficulties of living with HIV / Aids : Obstacles to quality of life. 2017;30(3):301-308.
22. Xie F, Zheng H, Huang L, Yuan Z, Lu Y. Social capital associated with quality of life among people living with HIV/AIDS in nanchang, China. *Int J Environ Res Public Health.* 2019;16(2):1-13. doi:10.3390/ijerph16020276
23. Rutakumwa R, Ssebunnya J, Mugisha J, et al. Stakeholders' perspectives on integrating the management of depression into routine HIV care in Uganda: qualitative findings from a feasibility study. *Int J Ment Health Syst.* 2021;15(1):1-15. doi:10.1186/s13033-021-00486-8
24. Watkins-Hayes C, Pittman-Gay LS Da, Beaman J. "Dying from" to "living with": Framing institutions and the coping processes of African American women living with HIV/AIDS. *Soc Sci Med.* 2012;74(12):2028-2036. doi:10.1016/j.socscimed.2012.02.001
25. Cherry S, De Marrais K, Keita C, Davis M, Lee J. "My determination is to live": Narratives of African-American women who have lived with HIV for 10 or more years. *Qual Rep.* 2019;23(10):2490-2510. doi:10.46743/2160-3715/2018.3389

26. Perry NS, Remmert JE, Psaros C, Pinkston M, Safren SA. Learning to address multiple syndemics for people living with HIV through client perspectives on CBT. *Psychother Res.* 2019;29(4):492-502. doi:10.1080/10503307.2017.1380863
27. Han X, Li B, Qu J, Zhu Q. Weibo friends with benefits for people live with HIV/AIDS? The implications of Weibo use for enacted social support, perceived social support and health outcomes. *Soc Sci Med.* 2018;211(June):157-163. doi:10.1016/j.socscimed.2018.06.016