Thyroid hormone axis and anthropometric recovery of children/adolescents with overweight/obesity: a scoping review

Carlos Ramos Urrea^{1*}, Amanda Paula Pedroso¹, Fernanda Thomazini¹, Andreia Cristina Feitosa do Carmo², Mônica Marques Telles¹, Ana Lydia Sawaya¹, Maria do Carmo Pinho Franco¹, Eliane Beraldi Ribeiro¹

¹ Universidade Federal de São Paulo, Escola Paulista de Medicina, Department of Physiology, Rua Botucatu, 862, Biomedical Sciences Building, 2° floor CEP 04023-060 São Paulo, SP, Brazil; carlosramos2690@gmail.com (C.R.U.); amandapedroso.bio@hotmail.com (A.P.P.); fernanda.thomazini@unifesp.br (F.T.); mmtelles@unifesp.br (M.M.T.); al.sawaya@yahoo.com.br (A.L.S.); mariadocarmo.franco@gmail.com (M.C.P.F.); eliane.beraldi@unifesp.br (E.B.R.)

² Universidade Federal de São Paulo, Escola Paulista de Medicina, Campus São Paulo Library; carmo@unifesp.br (A.C.F.C.)

* Correspondence: carlosramos2690@gmail.com

Abstract: Thyroid hormones play multiple physiological effects essential for the maintenance of basal metabolic rate (BMR), adaptive thermogenesis, fat metabolism, and appetite. The links between obesity and the hormones of the thyroid axis, i.e., triiodothyronine (T3), thyroxine (T4), thyrotropin-releasing hormone (TRH), and thyrotropin (TSH), are still controversial, especially when considering children and adolescents. This population has high rates of overweight and obesity and several treatment approaches, including nutritional, psychological, and physical exercise interventions have been used. Understanding the importance of the hormones of the thyroid axis in the recovery from overweight and obesity may help directing measures to the maintenance of a healthy body composition. The present scoping review was carried out to analyze studies evaluating these hormonal levels throughout interventions directed at treating overweight and obesity in children and adolescents. The main purpose was to ascertain whether the hormones levels vary during weight loss. We selected for analysis 16 studies published between 1999 and 2019. Most of the studies showed that the changes in body composition parameters in response to the multidisciplinary interventions correlated positively with free T3 (fT3)/ total T3 (TT3)/TSH. With respect to free T4 (fT4)/ total T4 (TT4), the most common finding was of unchanged levels and hence, no significant association with weight loss. Importantly, the response to the intervention has even been found to not be affected by fT4 supplementation. Further studies are necessary to elucidate the relevance of the variations in hormone levels to the establishment of overweight/obesity and to the recovery from these conditions in children/adolescents.

Keywords: Thyroid hormones; multidisciplinary intervention; obesity

1. Introduction

Globally, obesity is a well-recognized public health problem affecting both adults and children(1). In children/adolescents, the prevalence of overweight/obesity is high (2) and associates with increased risk to develop diabetes and other co-morbidities (3).

The pathophysiology of overweight/obesity includes genetic, environmental, behavioral, metabolic, psychological factors, and hormonal factors. Thyroid hormones play multiple physiological effects essential for the maintenance of basal metabolic rate (BMR), adaptive thermogenesis, fat metabolism, and appetite (4). The participation of the levels of the hormones of the thyroid axis, i.e., triiodothyronine (T3), thyroxine (T4), thyrotropin-releasing hormone (TRH), and thyrotropin (TSH), has been studied with no conclusive results, especially when considering children and

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

adolescents. They have indicated either that thyroid-hormones resistance is a causal factor of obesity or that elevated hormone levels may represent an adaptive response to obesity (5).

The treatment of children and adolescents with overweight or obesity is a very relevant issue, and several approaches, including nutritional, psychological, and physical exercise interventions have been used (6). Understanding the importance of the hormones of the thyroid axis in the recovery from overweight and obesity may help directing measures to the maintenance of a healthy body composition.

The present scoping review was carried out to analyze studies evaluating these hormonal levels throughout interventions directed at treating overweight and obesity in children and adolescents.

2. Materials and Methods

This scoping review was registered on the International Prospective Register of Systematic Reviews (PROSPERO, CRD42020203359) and performed with the recommendations of Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR).

2.1. Eligibility criteria

Original articles published in peer-reviewed journals, written in English, Portuguese, or Spanish, that had children and/or adolescents with overweight or obesity as participants and that performed some intervention for weight management, including, nutritional and/or psychological, and/or medical, and/or exercise.

The exclusion criteria were systematic reviews, studies in animals or adults, use of growth hormone or steroid hormones, diagnostic of thyroid kidney, heart, or neurological illness.

2.2. Literature search

The electronic search in the databases was carried out based on a search strategy aiming at locating both published and unpublished studies. Data collection and analysis were performed in September and October 2020. Electronic searches were conducted using the following databases: MEDLINE way Pubmed, Latin American and Caribbean Literature in Health Sciences (LILACS), Scopus (Elsevier) and Cochrane Library.

The following descriptors were extracted from the Health Science Descriptors database: obesity, overweight, obese, excess weight, weight gain, malnutrition, thyroid hormones, thyroid concentrations, thyroid stimulating hormone,

triiodothyronine, thyroxine, thyroid gland, child, children, adolescents. The development of the search strategy followed the recommendations of the checklist Peer Review of Electronic Search Strategies (PRESS) (7).

2.3. Study selection and appraisal

The selection and analysis of the studies were carried out by two independent authors using the Rayyan tool. The first selection was based on the title and summary of the studies. Duplicates and articles whose full texts were not available were excluded. Conflicts were resolved by consensus. After selection according to the inclusion criteria, two reviewers independently analyzed the texts in full to identify the relevant outcomes.

Data extraction and synthesis of results

To characterize the findings, the following variables were considered: age, type of intervention, effect of the intervention on body composition, and hormone levels, both at baseline and after the intervention. The articles were grouped by type of comparisons (in-group or between obese and eutrophic). Two studies involving thyroxine supplementation were constituted a third category.

3. Results

A total of 1031 articles were screened, leading to 20 eligible articles, of which 4 were excluded due to absence of full text. Sixteen articles were thus included in the qualitative synthesis. Figure 1 describes the selection process and Table 1 shows the results of each selected article.

Figure 1. Flow diagram of the studies selection.

Table 1. Summary of the characteristics and results of the 16 studies included in the analysis.

Reference	Sample (Age)	Intervention group (Gender)	Control group	Intervention type (Time)	Effect of Intervention*	Hormones		Authors' conclusion	Our observation
						Baseline	After intervention		
Studies conducted with in-group comparisons									
Kiortsis et al. 1999 (8)	64 (10 – 14y)	64 OB (BW>120%) (22 M and 42 F)		Caloric restriction, including a 24h recall method (6 weeks)	BMI decreased *		TT3: decreased TT4: unchanged TSH: unchanged	The decline in TT3 levels seems to play an important role in the decrease of RMR in children	TT3 levels decreased along with weight loss
Rijks et al. 2017 (9)	330 (2.6 – 18.9y)	66 OW (BMI Z-score >75<90), 148 OB (BMI Z-score=90<97), 115 MO (≥97) 142 M; 188 F 99 out of 330 children were evaluated 1-year post-intervention		Nutritional education and physical activity (8 weeks) Clinical assessment 1-year post-intervention	BMI Z-score decreased *	fT4: no differences among the subgroups OW, OB and MO TSH: no differences among the subgroups OW, OB and MO	fT4: decreased in children with decreased BMI Z-score (n = 62), but unchanged in children with increased BMI Z-score (n = 37) TSH: unchanged	In OW and OB, TSH is positively associated with CVD markers. Changes in TSH are also associated with changes in lipid concentrations in children showing weight loss	fT4 levels decreased along with weight loss
Aeberli et al. 2010 (10)	206 (10 – 18y)	206 OB (BMI-SDS >98 percentile) (119 M; 87 F) 197 OB completed the intervention		Hypocaloric diet, physical activity, and psychological treatment. (8 weeks)	BMI- SDS decreased *	fT3: normal range fT4: normal range TSH: normal range (but 52% in the high normal range and 1.9% had	fT3: decreased fT4: unchanged TSH: decreased (but 44% in the high normal range and 0.5% had HTTR)	TSH concentrations tend to be higher in OB	fT3 and TSH levels decreased along with weight loss

					HTTR)				
Shalitin et al. 2009 (11)	207 (5 – 18y)	207 OB (BMI-SDS >95 percentile) (97 M, 110 F) 142 subjects completed the intervention	Hypocaloric diet (12 weeks)	BMI-SDS decreased *	TT3: normal range	TT3: not reported	In OB, high TSH levels with normal fT4 levels appears to be frequent	No significant differences regarding fT4 and TSH between the children showing decreased BMI-SDS with those showing no BMI-SDS changes	
					fT4: normal range	fT4: positive Δ (0.18 ± 1.9) in children showing decreased BMI-SDS; negative Δ (-0.6 ± 1.97) in children showing no BMI changes			
						TSH: 161 subject showed normal range levels; 46 subjects showed levels above the normal range			TSH: not reported
									fT3: decreased
Bas et al. 2013 (12)	150 (3 – 17y)	150 OB (BMI Z-score >percentile 95) (67 M; 83 F) - Only 21 OB who had TSH levels above the normal range at the baseline were evaluated after the	Nutritional education and physical activity (6 months)	BMI Z-score decreased BMI Z-score unchanged	fT4: normal range	fT4: positive Δ (0.18 ± 1.9) in children showing decreased BMI-SDS; negative Δ (-0.6 ± 1.97) in children showing no BMI changes	In OB with high TSH levels, the fT3 and TSH levels decreased along with weight loss		
					fT4: normal range	fT4: positive Δ (0.18 ± 1.9) in children showing decreased BMI-SDS; negative Δ (-0.6 ± 1.97) in children showing no BMI changes			
						TSH: 161 subject showed normal range levels; 46 subjects showed levels above the normal range			TSH: not reported
									fT3: decreased

intervention					unchanged			
Wolters et al. 2013 (13)	477 (10.6 ± 2.7y)	477 OB (BMI-SDS >97 percentile) (219 M; 258 F)	Physical exercise, nutrition education, and psychological therapy (12 months)	BMI- SDS decreased *	ft3: normal range	ft3: unchanged	Moderately increased TSH and ft3 concentrations in OB, which normalized after substantial weight loss	No changes in ft3, ft4 and TSH levels along with weight loss
					ft4: normal range	ft4: unchanged		
					TSH: High range	TSH: unchanged		
Licenziati et al. 2019 (14)	96 (4.7 – 16.4y)	96 OB (BMI-SDS >75 percentile) (49 M; 47 F)	Dietary recommendations, physical activity, and behavioral strategies (0.8 ± 0.3 year)	BMI-SDS decreased *	ft4: normal range	ft4: unchanged	The alterations of thyroid function and structure in OB are reversible after weight loss (thyroid function significantly improved along with weight loss, as documented by a significant reduction of TSH without a change in ft4)	OB with alterations of the thyroid structure showed decreased TSH levels along with weight loss
					TSH: High range	TSH: decreased		
					- TSH levels exceeded the normal range in 83.3% OB	- TSH levels exceeded the normal range in 57.3% of OB		
Bouglé et al. 2014 (15)	528 (4.1 – 17.9y)	528 OW + OB (BMI Z-score >2 SDS) (238 M; 290 F) - 79 subjects completed the intervention	Nutritional education and physical activity (52 ± 15 weeks)	BMI Z-score decreased	ft3: normal range	ft3: unchanged	Increased TSH may be predictive of decreased insulin resistance; ft4 was associated with a low metabolic risk. Changes in thyroid function could protect against	Non-significant decrease of ft3, ft4, TSH and BMI Z-score
					ft4: normal range	ft4: unchanged		
					TSH: normal range levels, but 69 showed levels above the normal range	TSH: in the group with normal TSH at baseline, 26 and 42 subjects showed decrease and		

						increased TSH levels, respectively TSH decreased in the whole group of patients with initial values above the normal range	obesity-associated metabolic diseases	
						ft3: normal range ft4: normal range TSH: normal range (17.2% high range)	ft3: decreased ft4: unchanged TSH: decreased. 6.2% of the children showed a TSH above the normal range	A yet slightly reduced body fat can be followed by an improvement in biochemical parameters such as TSH and ft3
Radetti et al. 2012 (16)	72 (8 – 14y)	72 OW and OB (BMI-SDS >85 percentile) (41 M; 31 F)		Meetings with instructions on nutrition and physical activity at 3-month intervals (1.8 ± 1.0 year)	BMI-SDS decreased			ft3 and TSH levels decreased along with a non-significant weight loss
Studies conducted with comparisons between obese and eutrophic groups								
Marras et al. 2010 (17)	520 (3.7 – 17.9y)	468 OB (BMI-SDS >95 percentile) (213 M; 255 F) - 63 OB (43 M; 20 F) completed the intervention	52 EUT (24 M; 28 F)	Nutritional education and physical activity (6 months)	BMI-SDS decreased in the OB group	ft3: no difference (17.9% above the normal range) ft4: no differences (1.28% above normal range) TSH: no differences	ft3: no difference ft4: no difference TSH: no difference	In OB, an increased ft3 concentration is the most frequent thyroid function abnormality. Serum ft3 and TSH correlate with BMI-SDS. Moderate weight loss frequently restores these

						(3.2% above normal range)		abnormalities	
						Between-groups comparison (n = 50)			
						ft3: no differences			
						ft4: no differences between the boy groups, but OB girls had higher levels than EUT girls			
						TSH: no differences			
						In-group comparison (n = 36)			
						ft3: unchanged			
						ft4: no changes among the boys. OB girls showed decreased levels (became like the EUT girls)			
						TSH: unchanged			
Cayir et al. 2014 (18)	109 (5 – 18y)	85 (OW: 28; OB: 29; MO: 29) (No information on criteria) (43 M; 42 F)	24 age matched EUT (13 M; 11 F)	All patients received a diet list based on their ideal body weight. Recommendation of 30 min aerobic exercise per day (6 months)	BMI of OW, OB and MO subgroups decreased *	Normal thyroid function in OB, not influenced by a prolonged period of caloric restriction ft4 levels were higher in OB girls than in EUT girls, reaching normal levels along with weight loss induced by caloric restriction			
Reinehr, Souza, Andler	317 (9.5 – 12y)	246 OB (BMI Z-score >97 percentile) (109 M;	71 EUT	Physical exercise, nutrition education, and	BMI Z-score decreased (with	ft3: OB had higher levels than EUT	ft3: OB with substantial weight loss	ft3 and TSH were moderately increased in OB	Substantial weight loss decreased ft3 and TSH

2006 (19)	137 F)	behavior therapy (12 months)	substantial weight loss n=49)	(normal range)	showed decreased levels (no differences between OB with substantial weight loss and OB without substantial weight loss)	and weight loss led to a reduction. The elevation of these hormones seems to be rather a consequence of obesity than a cause of obesity
				fT4: no differences (normal range)	fT4: unchanged. (No differences between OB with substantial weight loss and OB without substantial weight loss.)	
				TSH: OB had higher levels than EUT (17% above normal range)	TSH: OB with substantial weight loss showed decreased levels (no differences between OB with substantial weight loss and OB	

without substantial weight loss.)									
Reinehr & Andler 2002 (20)	225 (4.5 – 16y)	118 OB (BMI Z-score >97 percentile) (63M; 55 F) - 68 OB completed the intervention	107 EUT (61 M; 46 F)	Physical exercise, nutrition education, and psychological therapy (12 months)	BMI Z-score decreased (n=55)		<u>TT3: decreased</u> <u>TT4: decreased</u> TSH: unchanged	TT3, TT4 and TSH levels were moderately increased in OB at baseline. A normal energy diet induces a long-term decrease in the peripheral thyroid hormones as opposed to TSH	TT3, TT4 and TSH levels moderately increased in OB. TT3 and TT4 levels decreased along with intervention-induced weight loss
Butte et al. 2015 (21)	16 (12 – 17y)	5 MO (BMI >50 or ≥40 with comorbidities) (3 M; 2 F)		Roux-en-Y gastric bypass surgery (RYGB). The subjects were reevaluated 12 months after surgery	BMI decreased after RYGB*		<u>TT3: decreased</u> <u>ft3: unchanged</u> <u>TT4: unchanged</u> <u>ft4: unchanged</u> TSH: decreased	Energy adaptations that occur in adolescents following RYGB possibly involves TT3 mediation	TT3 and TSH levels decreased along with BMI decrease induced by RYGB
Studies involving thyroxine supplementation									
Eliakim et al. 2006 (22)	196 (5 – 17y). - 41 OB showed HTTR and were	26 OB (BMI >85 percentile), with HTTR, with thyroxine supplementation	15 OB, with HTTR, without thyroxine supplementation	Dietary and exercise program for 12 months followed by thyroxine supplementation (1-2 µg/kg) for 6	BMI decreased in both supplemented and not supplemented groups	ft4: normal range	ft4: no differences between supplemented and not supplemented groups	HTTR is relatively common in OB. TSH levels returned to normal in the majority of	The TSH levels decreased significantly independent of ft4

Table 2. Correlations between anthropometric recovery and hormonal levels.

Reference	Correlation results	
	Baseline	After intervention
Marras et al. 2010 (17)	TSH and fT3 correlated positively with BMI-SDS	
Bouglé et al. 2014 (15)	TSH correlated positively with BMI Z-score	
Aeberli et al. 2010 (10)	Body weight correlated negatively with fT3 Percentage body fat correlated positively with fT4	Delta fT3 correlated positively with body weight, BMI-SDS, fat mass, and percentage body fat and negatively with lean body mass. Delta fT4 correlated negatively with percentage body fat
Licenziati et al. 2019 (14)	TSH correlated positively with BMI-SDS	TSH correlated positively with BMI-SDS
Bas et al. 2013 (12)		TSH and fT3 correlated positively with BMI Z-score
Reinehr & Andler 2002 (20)		TT3 and TT4 correlated positively with BMI Z-score
Butte et al. 2015 (21)		TT3 and TSH correlated positively with BMI

3.1. Description of the included studies

The literature search identified 1049 references (Figure 1) and 16 articles met all the inclusion criteria.

Table 1 summarizes the characteristics of the studies included in the analysis. There are studies performed with subjects from Italy (14,16,17), France (8,15), Israel (11,22), Turkey (12,18), Germany (13,19,20), the Netherlands (9), Swiss (10), United States (21), and India (23).

Twelve studies analyzed only obese children and adolescents (8,10–14,17,19–23) while the other 4 studies targeted both overweight and obese children and/or adolescents (9,15,16,18).

Table 2 describes the correlations between the thyroid hormonal axis and body composition parameters in the studies that performed this calculation. At baseline, body weight was negatively correlated with fT3 (10), the percentage of fat mass was positively correlated with fT4 (10), and BMI was positively correlated with TSH (14,15,17) and fT3 (17). After intervention, BMI correlated positively with TSH, TT3, and TT4 (14,20,21). When considering the baseline-after intervention changes, the authors reported positive correlations of BMI and TT3 (8), and fT3 (10,12), and TSH (12). fT3 also correlated positively with body weight, fat mass, and percentage body fat (10). Negative correlations were seen between lean body mass and fT3 and percentage body fat and fT4 (10).

3.2. Interventions

The duration of the interventions varied from 6 weeks to 18 months. One study utilized only a nutritional intervention [8], 6 studies performed nutritional intervention plus exercise intervention (11,12,15–18). One study allied nutritional and exercise interventions to administration of thyroxine (22).

In 5 studies, a psychological intervention was added to the nutritional plus exercise intervention (10,13,14,19,20). One study performed the 3 interventions plus levothyroxine supplementation (23).

One study performed psychological and exercise interventions (9) and 1 study used only a surgical intervention (21).

3.3. Nutritional Interventions

Fourteen studies used nutritional interventions. Eight studies used a nutritional education (12–16,19,20,23), 5 studies used a calorie restriction approach, with energy levels ranging from 900 to 2000 kcal/d and with varied macronutrient combinations and 1 study used the Mediterranean diet (17).

3.4. Exercise Interventions

Thirteen studies utilized physical exercise interventions. Nine studies conducted supervised training sessions with no specific routine or duration, with varying types and intensities (9,12–16,19,20,23), 1 involved two daily group endurance exercise sessions to improve aerobic performance, with a typical session lasting 60–90 min (10), and 3 performed aerobic exercise 3–5 times/week for at least 45–60 min (17), 30 minutes per day (18), or 90 minutes per day (11).

3.5. Psychological intervention

This type of intervention was used in 6 studies, including individual psychological care of the child/adolescent (10,14) or of the child/adolescent and their family (9,13,19,20). In the 2 studies detailing the psychological intervention, it consisted of techniques focusing on increasing self-esteem, responsibilities, and problem-solving strategies (9) and relaxation techniques and breathing therapy (10).

4. Discussion

All the 16 studies included in this analysis, published between 1999 and 2019, reported anthropometric recovery of the overweight/obese children/adolescents in response to the interventions, which, as depicted in table 1, varied largely with respect to the type and duration.

Concerning the baseline levels of the thyroid hormones (fT3, fT4, TT3 or TT4), 6 studies did not report these data (8,9,12,20,21,23). Among the 10 studies in which this information was available, the majority (9 studies) reported no significant alterations, either in relation to the normality ranges (10,11,13–16,19,22) or in comparison to eutrophic individuals (17,18), although one of these latter studies reported higher levels of fT4 in obese than in eutrophic girls (18) and one study reported small percentages of subjects with levels of fT3 (17.9%) or fT4 (1.28%) above normal range. Only one study reported higher mean fT3 values in the obese than in the eutrophic subjects, although still in the normal range (19). These results show that the most common status of thyroid hormones falls into normal levels. TSH levels at baseline were not reported in 5 studies (8,9,12,20,21) while 2 studies selected only individuals with hyperthyrotropinemia (22,23). Among the remaining 9 studies, the mean levels were normal in 6 studies (10,11,15–19), although some of these studies found mean values in the high normal range and highlighted the presence of elevated levels in variable percentages of their cohorts, namely 1.9% (10), 28.6% (11), 13.1% (15), 17.2% (16), and 3.2% (17). Elevated baseline levels of TSH were reported in 3 studies (14,19,17). These data demonstrate that the most common status of TSH among the studies analyzed fell into normal levels, although the finding of values in the high normal range was frequent.

Four studies reported correlations between body measures and hormone levels at baseline. One study reported a positive association of fT3 with BMI-SDS (17) while another study found a positive association of fT3 and body weight and a negative association of fT4 and percentage body fat (10). In 3 studies, TSH correlated positively with BMI-SDS (14,17) and BMI Z-score (15).

We searched other studies reporting levels of the hormones of the thyroid axis in children/adolescents with overweight/obesity. In one study, no differences were found in the levels of fT4 and TSH between children/adolescents with excess weight and the eutrophic ones (13). Many studies showed that these levels fell into the normal range, although a common finding was that they were higher than those of eutrophic children/adolescents, concerning TT3 (24), TT4 (25), and TSH (24–28), fT3 and fT4 (5).

Similar findings have been found in adults, with respect to fT3 (29) and TT4 (29,30), i.e., levels in the normal range but higher than the eutrophic levels. There are also reports that the hormone levels were in the normal range but lower in obese than in eutrophic adults, concerning TSH (29–31), fT3 and fT4 (31).

Examining studies performing correlation analysis of hormone levels and body parameters in overweight/obese children/adolescents, we observed one study reporting no significant associations of fT3, fT4, and TSH levels with body composition parameters (32). In contrast, we found reports of a positive correlation between fT3 and BMI (5) and of a negative correlation of fT4 and BMI (25). There are also studies showing a positive correlation of TSH and body measures (5,33). These latter results agree with the findings of the studies analyzed in this scoping review.

These findings are like some reports in adults, finding positive associations between BMI and TT3 (34) and fT3 (29,31,34–38), and a negative correlation with fT4 (38). With respect to TSH, the studies found achieved a negative (30) or positive (38–41) association of BMI and TSH.

Concerning the response to the interventions, 10 of the studies included in this scoping review reported a decrease in at least one thyroid hormone measured (TT3, TT4, fT3, fT4) in relation to the respective baseline values (8–12,16,17,19–21). Five studies did not find any type of significant change between the baseline and the post-intervention values (13–15,18,23). One study did not report these data (22).

In relation to TSH, 9 studies reported a decrease after the intervention (10,12,14–16,19,21–23) and 5 studies showed no changes (8,9,13,18,20). It is important to point out that 2 of these studies included only subjects with hyperthyrotropinemia. In 2 studies, this information was not reported (11,17).

The relation of hormonal levels on anthropometric recovery was evaluated in 5 studies by a correlation analysis. Positive associations were found between delta of fT3 and body weight, BMI-SDS, fat mass, and percentage of fat mass (10), fT3 and BMI Z-score (12) and TT3 and BMI (21). TT4 correlated positively with BMI Z-score (20) but delta fT4 correlated negatively with percentage of fat mass (10). TSH correlated positively with BMI-SDS (14), BMI Z-score (12) and BMI (21). Studies performed in adults submitted to multidisciplinary interventions to treat obesity corroborate the above results, as they have found positive associations of fT3/TT3 with BMI and body weight (34,42,43) and of TSH with body weight (43–46). However, we found one study in which body and fat mass losses were not accompanied by in TSH levels (47).

The main purpose of this scoping review was to ascertain whether the hormones of the thyroid axis vary during weight loss in overweight/obese children/adolescents. The examination of the 16 selected studies allowed us to conclude that most of the results pointed to the absence of elevated levels at baseline, in agreement with a previous review (5). Also, most of the studies showed that the changes in body composition parameters in response to the multidisciplinary interventions correlated positively with fT3/TT3/TSH. Further studies are necessary to elucidate the relevance of the variations in hormone levels to the establishment of overweight/obesity and to the recovery from these conditions in children/adolescents. With respect to fT4/TT4, the most common finding was of unchanged levels and hence, no significant association with weight loss. Importantly, the response to the intervention has even been found to not be affected by fT4 supplementation.

Author Contributions: C.R.R. and E.B.R.; methodology, C.R.U., E.B.R., A.C.F.C. and M.C.P.F.; review per peers, C.R.U., A.P.P., F.T., and E.B.R.; formal analysis, C.R.U., A.P.P. F.T., M.M.T., A.L.S., and E.B.R.; writing—original draft preparation, C.R.U., E.B.R. and M.C.P.F.; writing—review and editing, C.R.U., E.B.R. and M.M.T.. All authors have read and agreed to the published version of the manuscript.”

Funding: This research was funded by Coordenação de Aperfeiçoamento de Pessoal do Nível Superior (CAPES), this study was approved by the Research Ethics Committee of the Federal University of São Paulo, (CAAE = 17459618.0.0000.5505). CAPES has no role in the design, analysis or writing of this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. World Health Organization (WHO). Guideline: assessing and managing children at primary health-care facilities to prevent overweight and obesity in the context of the double burden of malnutrition. Updates for the Integrated Management of Childhood Illness (IMCI). [Internet]. 2017 [cited 2019 Jul 18]. Available from: <http://apps.who.int/bookorders>.
2. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight,

and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* [Internet]. 2017 [cited 2019 Mar 11];390:2627–42. Available from: <http://dx.doi.org/10.1016/>

3. Abdullah A, Aceh B, Courten M De, Stevenson CE, Walls H. The number of years lived with obesity and the risk of all-cause and cause-specific mortality. *Int J Epidemiol*. 2011;50(2):985–96. 202
4. Mullur R, Liu YY, Brent GA. Thyroid hormone regulation of metabolism. *Physiol Rev*. 2014;94(2):355–82. 204
5. Witkowska-Sędek E, Małgorzata Rumińska AK. Thyroid dysfunction in obese and overweight children. *Endokrynol Pol*. 2017;68(1):54–60. 205
6. Hoelscher DM, Kirk S, Ritchie L, Cunningham-Sabo L. Position of the Academy of Nutrition and Dietetics: Interventions for the Prevention and Treatment of Pediatric Overweight and Obesity. *J Acad Nutr Diet*. 2013;113:1375–94. 207
7. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. GUIDELINE STATEMENT PRESS Peer Review of Electronic Search Strategies : 2015 Guideline Statement. *J Clin Epidemiol*. 2016;75:40–6. 210
8. Kiortsis DN, Durack I, Turpin G. Effects of a low-calorie diet on resting metabolic rate and serum tri-iodothyronine levels in obese children. *Eur J Pediatr*. 1999;158(6):446–50. 212
9. Rijks JM, Plat J, Dorenbos E, Penders B, Gerver WJM, Vreugdenhil ACE. Association of TSH with cardiovascular disease risk in overweight and obese children during lifestyle intervention. *J Clin Endocrinol Metab*. 2017;102(6):2051–8. 214
10. Aeberli I, Jung A, Murer SB, Wildhaber J, Wildhaber-Brooks J, Knöpfli BH, et al. During Rapid Weight Loss in 217

Obese Children, Reductions in TSH Predict Improvements in Insulin Sensitivity Independent of Changes in Body Weight or Fat. *J Clin Endocrinol Metab.* 2010;95(12):5412–8.

11. Shalitin S, Yackobovitch-Gavan M, Phillip M. Prevalence of thyroid dysfunction in obese children and adolescents before and after weight reduction and its relation to other metabolic parameters. *Horm Res.* 2009;71(3):155–61. 220
12. Baş VN, Ayca Z, Ağladioğlu SY, Kendirci HNP. Prevalence of hyperthyrotropinemia in obese children before and after weight loss. *Eat Weight Disord.* 2013;18(1):87–90. 223
13. Wolters B, Lass N, Reinehr T. TSH and free triiodothyronine concentrations are associated with weight loss in a lifestyle intervention and weight regain afterwards in obese children. *Eur J Endocrinol.* 2013;168(3):323–9. 225
14. Licenziati MR, Valerio G, Vetrani I, Maria G De, Liotta F, Radetti G. Altered Thyroid Function and Structure in Children and Adolescents Who Are Overweight and Obese: Reversal After Weight Loss. *J Clin Endocrinol Metab.* 2019;104(7):2757–65. 227
15. Bouglé D, Morello R, Brouard J. Thyroid function and metabolic risk factors in obese youth. Changes during follow-up: a preventive mechanism? *Exp Clin Endocrinol diabetes.* 2014;122(9):548–52. 230
16. Radetti G, Longhi S, Baiocchi M, Cassar W, Buzi F. Changes in lifestyle improve body composition, thyroid function, and structure in obese children. *J Endocrinol Invest.* 2012;35(3):281–5. 232
17. Marras V, Casini MR, Pilia S, Carta D, Civolani P, Porcu M, et al. Thyroid function in obese children and adolescents. *Horm Res Paediatr* [Internet]. 2010;73(3):193–7. Available from: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-77951276542&doi=10.1159%2F000284361&partnerID=40> 234

&md5=12905e0603e2ec9b14938e68ac15537d

18. Cayir A, Doneray H, Kurt N, Orbak Z, Kaya A, Turan MI, et al. Thyroid functions and trace elements in 238
pediatric patients with exogenous obesity. *Biol Trace Elem Res*. 2014;157(2):95–100.
19. Reinehr T, Sousa G De, Andler W. Hyperthyrotropinemia in Obese Children Is Reversible after Weight Loss 240
and Is Not Related to Lipids. *J Clin Endocrinol Metab*. 2006;91(8):3088–91.
20. Reinehr T, Andler W. Thyroid hormones before and after weight loss in obesity. *Arch Dis Child*. 242
2002;87(4):320–3.
21. Butte NF, Brandt ML, Wong WW, Liu Y, Mehta NR, Wilson TA, et al. Energetic adaptations persist after 244
bariatric surgery in severely obese adolescents. *Obesity*. 2015;23(3):591–601.
22. Eliakim A, Barzilai M, Wolach B, Nemet D. Should we treat elevated thyroid stimulating hormone levels in 246
obese children and adolescents? *Int J Pediatr Obes*. 2006;1(4):217–21.
23. Kumar S, Dayal D, Attri SV, Gupta A, Bhalla AK. Levothyroxine supplementation for obesity-associated 248
thyroid dysfunction in children: A prospective, randomized, case control study. *Pediatr Endocrinol Diabetes
Metab*. 2019;25(3):107–13.
24. Soydan L, Ozturk E, Onal Z, C N. Associations of Thyroid Volume and Function with Childhood Obesity. *Acta 251
Endocrinol*. 2019;15(1):123–8.
25. Stichel H, L'allemand D, Grüters A. Thyroid function and obesity in children and adolescents. *Horm Res*. 253
2000;54(1):14–9.

26. Lobotková D, Staníková D, Staník J, Červeňová O, Bzdúch V, Tichá L. Lack of association between peripheral activity of thyroid hormones and elevated TSH levels in childhood obesity. *J Clin Res Pediatr Endocrinol.* 2014;6(2):100–4. 255
27. Ayala-Moreno M del R, Guerrero-Hernández J, Vergara-Castañeda A, Salazar-Aceves G, Cruz-Mercado DE. Thyroid function in pediatric population with different nutritional status. *Bol Med Hosp Infant Mex.* 2018;75(5):279–86. 258
28. Ghergherehchi R, Hazhir N. Thyroid hormonal status among children with obesity. *Ther Adv Endocrinol Metab.* 2015;6(2):51–5. 261
29. Wang Y, Dong X, Fu C, Su M, Jiang F, Xu D. Thyroid Stimulating Hormone (TSH) Is Associated With General and Abdominal Obesity: A Cohort Study in School-Aged Girls During Puberty in East China. *Front Endocrinol (Lausanne).* 2020;11(9):1–9. 263
30. Zhang L, Wang G. Association Between Body Mass Index and Thyroid Function in Euthyroid Chinese Adults. *Med Sci Monit.* 2021;27(27):1–8. 266
31. Mwafy S, Yassin M, Mousa R. Thyroid hormones, lipid profile and anthropometric changes after programmed weight loss in Palestinian obese adult females. *Diabetes Metab Syndr Clin Res Rev.* 2018;12(3):269–73. 268
32. Sari R, Balci MK, Altunbas H, Karayalcin U. The effect of body weight and weight loss on thyroid volume and function in obese women. *Clin Endocrinol (Oxf).* 2003;59(2):258–62. 270
33. Dursun F, Atasoy Öztürk T, Seymen Karabulut G, Kırmlızbekmez H. Obesity-related thyroiditis in childhood: Relationship with insulin resistance. *J Pediatr Endocrinol Metab.* 2019;32(5):471–8. 272

34. Joshi SR. Thyro-weight: Unlocking the link between thyroid disorders and weight. J Assoc Physicians India. 274
2018;66(March):70–3.
35. Liu G, Liang L, Bray GA, Qi L, Hu FB, Rood J, et al. Thyroid hormones and changes in body weight and 276
metabolic parameters in response to weight loss diets: The POUNDS LOST trial. Int J Obes. 2017;41(6):878–86.
36. Chen H, Zhang H, Tang W, Xi Q, Liu X, Duan Y, et al. Thyroid function and morphology in overweight and 278
obese children and adolescents in a Chinese population. J Pediatr Endocr Met 2013; 2013;26(5–6):489–96.
37. Roef GL, Rietzschel ER, Van Daele C, Taes YE, De Buyzere M, Gillebert TC, et al. Triiodothyronine and free 280
thyroxine levels are differentially associated with metabolic profile and adiposity-related cardiovascular risk
markers in euthyroid middle-aged subjects. Thyroid. 2014;24(2):1–31.
38. Natah TM, Wtw MA, Hadi MA, Farhood HF. Thyroid Metabolic Hormones and its Correlation with BMI and 283
Lipid Profile in Healthy People. Food Sci Qual Manag. 2013;18:18–25.
39. Fontenelle LC, Feitosa MM, Severo JS, Freitas TEC, Morais JBS, Torres-Leal FL, et al. Thyroid Function in 285
Human Obesity: Underlying Mechanisms. Horm Metab Res. 2016;48(12):787–94.
40. Bastemir M, Akin F, Alkis E, Kaptanoglu B. Obesity is associated with increased serum TSH level, independent 287
of thyroid function. Swiss Med Wkly [Internet]. 2007 [cited 2018 Sep 14];137(29–30):431–4. Available from:
<https://smw.ch/en/resource/jf/journal/file/view/article/smw/en/smw.2007.11774/smw.2007.11774.pdf/>
41. Hoermann R, Midgley JEM, Larisch R, Dietrich JW. Homeostatic Control of the Thyroid–Pituitary Axis: 290
Perspectives for Diagnosis and Treatment. Front Endocrinol (Lausanne) [Internet]. 2015 Nov 20 [cited 2018
Nov 26];6:177. Available from: <http://journal.frontiersin.org/Article/10.3389/fendo.2015.00177/abstract>

42. Fox CS, Pencina MJ, D'Agostino RB, Murabito JM, Seely EW, Pearce EN, et al. Relations of Thyroid Function to 293
Body Weight. Arch Intern Med [Internet]. 2008;168(6):587–92. Available from:
<https://jamanetwork-com.rsm.idm.oclc.org/journals/jamainternalmedicine/fullarticle/414105>
43. Agnihothri R V, Courville AB, Linderman JD, Smith S, Brychta R, Remaley A, et al. Moderate Weight Loss Is 296
Sufficient. Thyroid. 2014;24(1).
44. Marzullo P, Minocci A, Mele C, Fessehatsion R, Tagliaferri M, Pagano L, et al. The relationship between resting 298
energy expenditure and thyroid hormones in response to short-term weight loss in severe obesity. PLoS One
[Internet]. 2018 [cited 2018 Nov 26];13(10):1–12. Available from: <https://doi.org/10.1371/journal.pone.0205293>
45. Rosenbaum M, Hirsch J, Murphy E, Leibel RL. Effects of changes in body weight on carbohydrate metabolism , 301
catecholamine excretion , and thyroid function 1 – 4. Am J Clin Nutr. 2018;71(March):1421–32.
46. Muller M, Enderle J, Pourhassan M, Braun W, Eggeling B, Lagerpusch M. Metabolic adaptation to caloric 303
restriction and subsequent refeeding : the Minnesota Starvation Experiment revisited 1 , 2. Am J Clin Nutr.
2015;102:1–13.
47. Johnstone AM, Murison SD, Duncan JS, Rance KA, Speakman JR. Factors influencing variation in basal 306
metabolic rate include fat-free mass , fat mass , age , and circulating thyroxine but not sex ,. Am J Clin Nutr.
2005;82(March):941–8.

Figure 1. Flow diagram of the studies selection.