mRNA-COVID19 vaccination can be considered safe and tolerable for frail patients

Running title: **Safety and tolerability of SARS-COV-2-vaccine in frail patients**

An analysis from the Italian, multicentric, observational, prospective trial VAX4FRAIL study.

Maria Teresa Lupo Stanghellini^{1*}, Serena Di Cosimo^{2*}, Massimo Costantini³, Sara Monti⁴, Renato Mantegazza⁵, Alberto Mantovani⁶⁻⁷⁻⁸, Carlo Salvarani⁹⁻¹⁰, Pier Luigi Zinzani¹¹⁻¹², Matilde Inglese¹³⁻¹⁴, Fabio Ciceri¹⁻¹⁵, Giovanni Apolone¹⁶, Gennaro Ciliberto¹⁷, Fausto Baldanti¹⁸⁻¹⁹, Aldo Morrone²⁰, Valentina Sinno²¹, Franco Locatelli²²⁻²³, Stefania Notari²⁴, Elena Turola²⁵, Diana Giannarelli²⁶, Nicola Silvestris²⁷⁻²⁸, on behalf of the VAX4FRAIL Study Group

1 Hematology and Bone Marrow Transplantation Unit, IRCCS San Raffaele Scientific Institute, Milano, Italy;

2, Biomarkers Unit, Department of Applied Research and Technological Development, Fondazione IRCCS Istituto Nazionale dei Tumori, Milano, Italy;

3 Scientific Directorate, Azienda USL-IRCCS, Reggio Emilia, Italy;

4 Department of Rheumatology, Policlinico San Matteo IRCCS Fondazione, University of Pavia, Pavia, Italy;

5 Neuromuscular Diseases and Neuroimmunology Unit, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milano, Italy;

6 Humanitas Scientific Directorate, IRCCS Humanitas, Clinical and Research Center, Rozzano, Italy

7 Department of Biomedical Sciences, Humanitas University, Milano, Italy;

8 William Harvey Research Institute, Queen Mary University, London, United Kingdom;

9 Unità di Reumatologia, Azienda USL-IRCCS, Reggio Emilia, Italy;

10 Unità di Reumatologia, Università degli Studi di Modena e Reggio Emilia, Modena;

11 Istituto di Ematologia “Seràgnoli”, Azienda Ospedaliero-Universitaria di Bologna - IRCCS, Bologna, Italy;

12 Dipartimento di Medicina Specialistica, Diagnostica e Sperimentale Università di Bologna, Bologna, Italy;

13 Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health and Center of Excellence for Biomedical Research, University of Genoa, Genoa, Italy;

14 IRCCS Ospedale Policlinico San Martino, Genoa, Italy;

15 University Vita-Salute San Raffaele, Milan, Italy;

16 Scientific Directorate, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Italy;

17 Scientific Directorate IRCCS Regina Elena, National Cancer Institute, Istituti Fisioterapici Ospitalieri (IFO), Rome, Italy;

18 Molecular Virology Unit, Fondazione IRCCS Policlinico San Matteo;

19 Department of Clinical, Surgical, Diagnostics and Pediatric Sciences, University of Pavia, Pavia, Italy;

20 Scientific Directorate, San Gallicano Dermatological Institute IRCCS, Rome, Italy;

21 Department of Oncology and Hematology, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Italy;

22 Department of Pediatric Hematology and Oncology and Cell and Gene Therapy, IRCCS Ospedale Pediatrico Bambino Gesù;

23 Department of Gynecology-Obstetrics and Pediatrics, University “La Sapienza”, Roma, Italy;

24 Cellular Immunology Laboratory, National Institute for Infectious Diseases L Spallanzani – IRCCS, Rome, Italy;

25 Infrastruttura ricerca e statistica, Azienda USL-IRCCS, Reggio Emilia, Italy;

26 Biostatistical Unit, Istituto Nazionale Tumori Regina Elena IRCCS - IFO, Rome, Italy;

27 Medical Oncology Department, IRCCS Istituto Tumori “Giovanni Paolo II”, Bari, Italy;

28 Department of Biomedical Sciences and Human Oncology, University of Bari “Aldo Moro”, Bari, Italy.

* Equal contribution as first author

Corresponding author:

Nicola Silvestris

Medical Oncology Department,

IRCCS Istituto Tumori “Giovanni Paolo II”

DIMO – University of Bari

70124 - Bari, Italy

Mail: n.silvestris@oncologico.bari.it

Key Points

Question

Can m-RNA-COVID19 vaccination be considered safe for frail patients?

Findings

In this national, multicentric, observational, prospective trial (NCT04848493) that included 566 frail patients, the occurrence of both local and systemic adverse events was manageable and did not negatively impact on the general treatment program.

Meaning

mRNA-COVID19 vaccination is safe among frail immunocompromised patients.

Importance

Our study reports the safety analysis of the trial VAX4FRAIL confirming that mRNA-COVID-19 vaccination is safe in frail immunocompromised patients: expected side effects were manageable and had a minimum impact on patient care path.

Objective

To evaluate the safety of mRNA-COVID-19 vaccination in vulnerable patients.

Design

VAX4FRAIL is a national, multicentric, observational, prospective trial (start date March 3rd, 2021 – primary completion date September 2nd, 2021).

Setting

Multicenter prospective trial.

Participants

Frail patients were defined per protocol as patients under treatment with solid tumors (191), immune-rheumatological diseases (86), hematological malignancies (131), and neurological diseases (158), including multiple sclerosis and generalized myasthenia.

Exposure

Overall, 105 received the mRNA-1273 vaccine and 461 the BNT162b2 vaccine.

Main Outcome

The occurrence of adverse events after 1st and 2nd m-RNA-COVID-19 vaccination was analyzed. Adverse events were collected through a questionnaire comprising both open and closed questions.

Results

The most frequently reported moderate-severe adverse events were pain at the injection site (60.3% after the first dose, 55.4% after the second), fatigue (30.1% - 41.7%), bone pain (27.4% - 27.2%) and headache (11.8% - 18.9%).

Risk factors associated with the occurrence of severe symptoms after vaccine administration were identified through a multivariate logistic regression analysis: age was associated with severe fever presentation (younger patients vs. middle-aged vs. older ones), females presented a higher probability of severe pain at the injection site, fatigue, headache, and bone pain; the mRNA-1237 vaccine was associated with a higher probability of severe pain at the injection site and fever. Patients presenting a severe symptom after the first dose were at a relevant risk of recurrence of the same severe symptom after the second one.

Overall, 11 patients (1.9%) after the first dose and 7 (1.2%) after the second one was required to postpone or suspend their disease-specific treatment. Finally, 2 fatal events occurred among our 566

patients, and these two events were due to disease progression and considered unrelated to the vaccine.

Conclusion and Relevance

Our study reports that mRNA-COVID-19 vaccination is safe also in frail patients as expected side effects were manageable and had a minimum impact on patient care path.

Study Registration

A National, Multicentric, Observational, Prospective Study to Assess Immune Response to COVID-19 Vaccine in Frail Patients (VAX4FRAIL). NCT04848493

<https://clinicaltrials.gov/ct2/show/NCT04848493>

Background

Frail patients are considered at relevant risk of complications due to COVID-19 infection and, for this reason, are prioritized candidates for vaccination. As these patients were originally not included in the registration trials, fear related to vaccine side-effects and disease worsening was one of the reasons for vaccine hesitancy. Herein we report the safety profile of the prospective, multicenter, national VAX4FRAIL study (NCT04848493) to evaluate vaccines in a large trans-disease cohort of patients with solid or hematological malignancies, neurological and rheumatological diseases.

Methods

Between March 3rd and September 2nd, 2021, 566 patients were evaluable for safety endpoint: 105 received the mRNA-1273 vaccine and 461 the BNT162b2 vaccine.

Frail patients were defined per protocol as patients under treatment with hematological malignancies (131), solid tumors (191), immune-rheumatological diseases (86), and neurological diseases (158), including multiple sclerosis and generalized myasthenia.

The impact of the vaccination on the health status of patients was assessed through a questionnaire focused on the first week after each vaccine dose.

Results

The most frequently reported moderate-severe adverse events were pain at the injection site (60.3% after the first dose, 55.4% after the second), fatigue (30.1% - 41.7%), bone pain (27.4% - 27.2%) and headache (11.8% - 18.9%).

Risk factors associated with the occurrence of severe symptoms after vaccine administration were identified through a multivariate logistic regression analysis: age was associated with severe fever presentation (younger patients vs. middle-aged vs. older ones), females presented a higher probability of severe pain at the injection site, fatigue, headache, and bone pain; the mRNA-1237 vaccine was associated with a higher probability of severe pain at the injection site and fever. After the first dose, patients presenting a severe symptom were at a relevant risk of recurrence of the same severe symptom after the second one.

Overall, 11 patients (1.9%) after the first dose and 7 (1.2%) after the second one required to postpone or suspend the disease-specific treatment. Finally, 2 fatal events occurred among our 566 patients. These two events were considered unrelated to the vaccine.

Conclusions

Our study reports that mRNA-COVID-19 vaccination is safe also in frail patients as expected side effects were manageable and had a minimum impact on patient care path.

Keywords: Frail patients; SARS-CoV2 mRNA vaccine; safety

Introduction

The currently authorized mRNA-COVID-19 vaccines – m-RNA-1237 Moderna (1) and BNT-162b2 Pfizer BioNTech (2) - have been evaluated in clinical trials that excluded in accordance with the current regulations, immunocompromised subjects, and restricted participation to healthy volunteers. Fragile patients were not considered in such pivotal studies despite being the subjects at greatest risk of COVID-19 complications and with the potential greatest advantage.

Patients diagnosed with solid or hematological malignancy or under immunosuppressive treatment due to rheumatological or neurological diseases are considered at high risk of COVID-19 complications and categorized as fragile (3-7). The intended acceptance of the COVID-19 vaccine in fragile patients confirms the positive attitudes towards vaccination, but questions arise around the safety of these vaccines in the setting of immune alterations engendered by their diseases and/or therapies (8-14). One of the most typical reasons for vaccine hesitancy has been fear related to vaccine side-effects and underlying disease worsening. An additional effort through public education campaigns specific to fragile patients to clarify vaccine's safety, long-term effects, and potential health implications are still needed.

Over the past eight months, several groups have tried to answer questions about the efficacy and safety of COVID vaccination in different cohorts of fragile subjects. No safety concerns emerged, confirming the profile described in the series of healthy subjects (14-43) and pointing out an acceptable safety profile. VAX4FRAIL (44) study aimed at assessing immune responses to vaccination in a large trans-disease cohort of patients with hematological malignancies, solid tumors, neurological and rheumatological diseases. The study main objective is to assess prospectively the immunologic response to the COVID-19 vaccination in these specific subgroups, characterizing the kinetics of the immune response to the vaccination and its persistence over time. Longitudinal, prospective evaluation of the safety profile was part of this trans-disease study. Herein we report safety profile results as outlined in VAX4FRAIL trial.

Methods

Safety analysis was performed among patients enrolled in the VAX4FRail trial between t0 and t2 according to the protocol, being t0 “time point 0” at first dose of vaccine and t2 “time point 2” the blood sampling 2-4 weeks after the second dose of vaccine (44). This is a national, multicentric observational prospective study conducted in Italy with the primary aim of assessing the immune response of COVID-19 vaccination in frail, immunocompromised patients.

Patients were considered for the current safety analysis if they met the general inclusion criteria: being ≥ 18 years of age, having received COVID-19 vaccination with mRNA vaccines (BNT-162b2 Pfizer-BioNTech or m-RNA-1237 Moderna vaccine), and having completed the health status assessment questionnaire after one of the two vaccine doses. Frail patients under evaluation were diagnosed with hematological malignancies, solid tumors, immune-rheumatological disease, and neurological disease. Detailed inclusion and exclusion criteria were previously reported (44) and disease stratification according to diagnosis and treatment of the primary disease.

The impact of the vaccination on the health status of patients was assessed through a questionnaire focused on the first week after each vaccine dose. The questionnaire was administered to the patients after the second vaccine dose (3-4 weeks after the first dose) and at the subsequent timing according to VAX4FRail protocol (2-4 weeks after the second dose).

The questionnaire assessed how much the patient was troubled by eight symptoms (pain or swelling at the injection site, fatigue, headache, bone pain, fever, enlarged lymph nodes, skin rash, insomnia, diarrhea, and nausea or vomiting). (See supplementary; Appendix 1) The answers were graded according to a 5-level scale (not at all, slightly, moderately, severely, and overwhelmingly). The patient was also asked to report and grade other symptoms possibly occurring after each vaccine dose and if he/she had to postpone or suspend therapies due to symptoms related to vaccination.

Statistical methods

For this analysis, we grouped answers to the questionnaire in three levels of troubles: not at all, moderate (slightly or moderately), and severe (severely or overwhelmingly).

We estimated the probability of the occurrence of a severe symptom after the second dose of the vaccine according to the occurrence of a severe symptom after the first dose using Positive and Negative Predictive Values (PPV and NPV, respectively).

The probability of occurrence of a severe symptom after one of the two doses of the vaccine according to age, sex, main diagnosis, and type of vaccine was estimated in a multivariate logistic regression analysis adjusting for all variables.

Results

Patients

Between March 3rd and September 2nd, 2021, 566 patients were enrolled in the study VAX4FRAIL and were eligible for this analysis. Among 566 patients who received the first dose of the mRNA vaccine, 105 received the mRNA-1273 and 461 the BNT162b2. Among the 566 patients who received the first dose, they also received the second dose (**Figure 1**).

Overall, 488 patients after the 1st dose and 489 patients after the 2nd dose were evaluable for safety evaluation having complete clinical evaluation and questionnaires. Sixty-eight patients after the 1st dose and 57 after the 2nd dose did not fill the questionnaires. Consent withdrawn occurred in 8 cases after the 1st dose and 10 cases after the 2nd one.

Patient characteristics are reported in **Table 2**. Most patients (No.=299, 52.8%) were aged between 51 and 70 years, and 317 (56.0%) were females. The cohort of patients with solid tumors comprised 191 patients (33.7%), the one with neurological diseases included multiple sclerosis and generalized myasthenia 158 patients (27.9%); overall, 131 patients were diagnosed with hematological malignancies (23.1%) and 86 with immunopharmacological diseases (15.2%).

Vaccine-related adverse events

Overall, 438 (77.3%) patients reported any grade adverse events after the first dose (62, 11.0% reported as severe), and 373 (65.9%) after the second dose (87, 15.4% reported as severe). Detailed analysis of adverse events and severity is reported in **Table 3**.

After the first dose (both mRNA-1273 and BNT162b2) 53.9% of patients reported moderate pain at the injection site and 6.4% severe pain: this was the most frequently reported complaint after vaccine administration. Interestingly, after the second dose - t2 - 50.3% and 5.1% reported this adverse event as moderate and severe, respectively.

At t1 25.6% of patients reported fatigue as a moderate event, 4.5% as a severe one. We observed a slight increase at t2, with 35% of patients reporting fatigue as a moderate event and 6.7% as a severe one.

Bone pain and headache were reported as moderate by 13.7% and 9.8% of patients at t1, while 2.7% and 2.0% respectively reported the symptoms as severe. After the second dose bone pain was

reported as moderate by 23.3% and severe by 3.9% of patients, while headache was reported as moderate by 16% and severe by 2.9% of patients.

Only 5.9% of patients reported fever as relevant symptoms at t1 (moderate 4.7% - severe 1.2%), while at t2 the percentage of patients reporting this symptom slightly increased: 15.9% overall, with 10.0% moderate and 5.9% severe. Only a minority of patients – less than 2% - reported nausea, diarrhea, insomnia, skin rash or enlarged lymph nodes as severe manifestation both at t1 and t2.

Among unsolicited adverse events, after the first dose 5 patients reported chills (5/5 moderate), 3 itching (2 moderate, 1 severe), 2 gastro-intestinal pain (2/2 moderate), 5 myalgia (5/5 moderate), 6 dizziness (3 moderate, 3 severe), 1 drowsiness of moderate degree and 2 sweats (1 patient moderate and 1 severe). Similarly, after the second dose, 12 patients reported chills (11 moderate, 1 severe), 7 itching (5 moderate, 2 severe), 1 a moderate gastrointestinal pain, 8 myalgia (6 moderate, 2 severe), 9 dizziness (8 moderate, 1 severe), 1 drowsiness of moderate degree and 2 sweats (1 patient moderate and 1 severe). After the second dose, one patient reported mental confusion and one dysesthesia. Among the 11 patients with hematological malignancies that received an allogeneic stem cell transplantation, *graft versus host disease* (GvHD) occurrence or reactivation was not reported after the first and/or the second dose.

Of note, the absence of severe symptoms after the first dose strongly predicts the high probability of the absence of the same severe symptoms after the second dose (**Table 3**). Patients reporting severe fever after the first dose were more prone to develop the same severe symptom after the second one.

Multivariate model for occurrence of severe symptoms.

A multivariate logistic regression – adjusted by age, sex, diagnosis, and vaccine – was performed to identify risk factors associated with the occurrence of severe symptoms after vaccine administration (**Table 4**).

Age was associated with severe fever presentation: younger patients (age 18-50 years) had a higher probability of occurrence of severe fever than middle-aged patients (age 51-70 – odd ratio 0.3, 95% confidence interval [CI] 0.1-0.7) or older ones (≥ 71 – odd ratio 0.1, 95% CI 0.1-0.7) – p 0.003.

Female patients presented higher probability of severe pain at the injection site (odd ratio 2.9, 95% CI 1.4-6.2, p 0.005), severe fatigue (odd ratio 2.3, 95% CI 1.2-4.5, p 0.016), severe headache (odd ratio 7.6, 95% CI 1.7-33.6, p 0.007), and severe bone pain (odd ratio 2.7, 95% CI 1.1-6.5, p 0.028).

Diagnosis (hematological diseases, solid tumors, neurological diseases, rheumatological diseases) has no impact on the occurrence of severe fatigue, or headache, or bone pain or fever, but was associated with severe pain at the injection site: patients with a diagnosis of solid tumors reported severe pain less frequently than patients with hematological diseases (odds ratio 0.3, 95% CI 0.1-1.0), patients with a diagnosis of neurological or rheumatological diseases have a higher probability of occurrence of severe pain at the injection site in comparison with patients with the hematological disease (odd ratio for neurological diseases 4, 95% CI 1.4-11; the odds ratio for rheumatological diseases 2.8, 95% CI 1.1-7.5) – p <0.001.

Finally, the conditional contribution made by the vaccine (mRNA-1237 versus BNT162b2) clearly showed that mRNA-1237 was associated with a higher probability of severe pain at the injection site (odds ratio 0.3, 95% CI 0.1-0.6 – p 0.001) and a higher probability of severe fever (odds ratio 0.3, 95% CI 0.1-0.7 – p 0.004).

Impact on disease and therapy

Overall, 11 patients (1.9%) after the first dose and 7 (1.2%) after the second dose, required to postpone or suspend the disease-specific treatment due to the occurrence of symptoms associated with the vaccine. Among the 11 patients who required delays after the first dose, 7 were diagnosed with hematological disease and 4 with a solid tumor. Among the 7 patients who required delays after the second dose, 3 were diagnosed with the hematological disease and 4 with a solid tumor.

No patients from the neurological diseases' cohort or the rheumatological diseases' cohort required treatment delay.

Fatal events, long-term sequelae, or hospitalization related to the vaccine were not registered. Two patients died from disease progression; the events were clearly reported by the investigator as not related to the vaccine administration.

Discussion

Questionnaires based upon open and /or closed questions administered through the web-based app, phone interview, self-reporting forms, and face-to-face visits were all utilized to collect adverse events after COVID-19 vaccinations in the effort to clarify the safety of vaccines better. Absence of a univocal method to grade and collect this information make it almost impossible to directly compare the different experiences, but, of note, all the studies – both retrospective and prospective – are concordant in underlying the general perception of the absence of safety issue among distinct cohorts of frail patients (**Table 1**).

Providing information on the safety of COVID-19 vaccination in high-risk fragile populations is a duty of the international scientific community. To date, several groups are trying to answer two fundamental questions: i) if and how frail patients develop an effective response to the vaccine; ii) if the safety profile is confirmed valid in this category both with reference to toxicity and with reference to the maintenance of control of the underlying disease.

Safety profile is confirmed acceptable in all the experience reported so far in fragile patients.

More systemic and local side-effects were observed after the second dose of vaccine than after the first dose. The most common local side-effects were pain at the injection site, local rash, and local swelling, whereas the most common systemic side-effects were muscle pain, fatigue, headache, fever, chills, gastrointestinal complications, and flu-like symptoms (15-16-19-20-21-23-24-25-27-28-33-34-35). Overall, the incidence of severe symptoms was low - <2.5% (16, 36, 38) – with authors reporting no severe grade 3-4 adverse events (17-18-19-21-30-35-39-40-41-42-43). In a national prospective cohort study evaluating outcomes in patients with hematological malignancies in Lithuania (22) the authors reported that adverse events were more common after the second dose, with fatigue being the most prevalent symptom (13%). No grade 4 adverse events were reported.

Among patients with multiple sclerosis, Lotan and colleagues (23) outlined how 15% of the participants reported new or worsening neurological symptoms following the vaccination, the most frequent being sensory disturbances (58.3%). Most symptoms occurred within the first 24 h after vaccination and resolved within 3 days. A total of 28 participants (77.8%) did not require any medication to treat their symptoms. Of note, no increased risk of relapse activity was noted across patients with multiple sclerosis (23-24) or autoimmune inflammatory rheumatic diseases (26-28): Boekel and coworkers clearly showed how multivariable logistic regression analyses showed similar odds for any adverse event, systemic adverse events, or moderate or severe adverse events

between patients and controls, which was consistent when patients with rheumatoid arthritis or multiple sclerosis were compared with healthy controls (33).

In our experience, cohort composition was clearly defined at enrolment of patients, identifying a subset of high-risk subjects for severe COVID-19. The monitoring strategy was defined per protocol and homogeneous across the 4 categories of diseases (hematological disease, solid tumors, neurological conditions, and rheumatological diseases).

Overall, our study confirms the positive safety profile reported by other authors in both prospective and retrospective analysis (**Table 1**). Incidence of severe adverse events after vaccine administration was generally low and <3%. Most frequent complaints were pain at the injection site (severe 6.3%) and fatigue (severe 4.5%) after the first dose; pain at the injection site (severe 5.1%), fatigue (severe 6.8%), bone pain (severe 3.9%) and fever (severe 6%) after the second dose.

Patients experiencing a severe symptom after the first dose were more likely to report the same after the second one. Similarly, the absence of a severe symptom after the first dose was significantly associated with an absence of the same symptoms also after the second dose. This observation should be taken in consideration during the counseling of patients: patients should be reassured about the possibility of adopting a preventive strategy to reduce the burden of symptoms and on the not-unexpected onset of the symptom itself.

Moreover, the definition of predisposing factors to an increased possibility of presenting specific adverse events (e.g., younger patients are more likely to experience fever) can help both in the implementation of preventive measures (e.g., pre-emptive administration of painkiller drugs or antifever), in the optimization of counseling (e.g., if a patient experienced adverse events after the first shot he should be advised that the recurrence of the same after the second is very likely and that this is not unexpected, moreover strategies to counterbalance this inconvenience can be applied), and in the planning of therapies and/or vaccination itself.

Of note, less than 2% of patients required a delay or a suspension of the ongoing or planned treatment of the underlying diseases due to vaccination. This pointed out the positive safety profile of the vaccine strategy and – furthermore – this aspect should be discussed with patients to confirm the absence of impact on the whole therapy program. Whatever the underlying disease (a hematological cancer, a solid tumor, a rheumatological disease or a neurological disease) the mRNA-COVID-19 vaccine should be considered a crucial step to allow a safe program of treatment

more than a possible obstacle or danger to pursue the control of the disease, being the safety profile reassuring. Fatal events or hospitalization related to the vaccine were not registered in our cohort of high-risk frail patients.

Despite the absence of safety concerns emerging from our study, we believe it is mandatory to maintain constant and careful surveillance concerning post-vaccination adverse events. This can only contribute to the reliability that the vaccination strategy pursues and represents a cornerstone of general medical practice.

As outlined by several authors (3-9), one of the most common reasons for vaccine hesitancy was fear related to vaccine side-effects and disease worsening: we can confirm that vaccine-side effects were both manageable and in line with previously reported in the general population, without the occurrence of unexpected events and no concern related to worsening of the underlying disease or the need to delay treatment.

Conclusion

Frail patients' candidates to mRNA-COVID-19 vaccination should be reassured about the safety profile of vaccine strategy: adverse events were in line with the report from the healthy cohort of subjects and national observatories, no evidence of worsening of the underlying disease was reported and no concern on the adherence to the treatment program of the disease itself emerged from our prospective multicenter national study. Pursuing careful and timely monitoring of expected and unexpected adverse events represents a gold standard of modern medicine and can only support evidence-based medicine.

Article Information

Corresponding Author: Nicola Silvestris

Author Contributions: MC, VS, and ET had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: MTLP, SDC, MC, SM, NS, GA, AM.

Acquisition, analysis, or interpretation of data: MTLs, SDC, MC, SM, NS.

Drafting of the manuscript: MTLs, SDC, SM, MC, NS.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: MC, DG, ET.

Administrative, technical, or material support: VS, ET.

Conflict of Interest Disclosures: None reported.

Funding/Support: This study has been financed by Italian Ministry of Health within Ricerca Corrente 2021-Special Projects-Vax4Frail

Additional Contributions: We are indebted for their precious support to this study the Research Director Dr. Giuseppe Ippolito, and the Deputy General Director for Health Research and Innovation Dr. Gaetano Guglielmi, from the Italian Ministry of Health. We would also like to thank the patients who will be enrolled in the VAX4FRAIL study and their families, and all those who will be actively involved in their continuous care, study data collection and analysis, and ultimately in the scientific production that will result from this research.

THE VAX4FRAIL STUDY GROUP:

PRINCIPAL INVESTIGATORS (alphabetical order): Giovanni Apolone (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano); Alberto Mantovani (IRCCS Istituto Clinico Humanitas, Milano).

SCIENTIFIC COORDINATOR: Massimo Costantini (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano).

STEERING COMMITTEE (alphabetical order): Chiara Agrati (IRCCS Istituto per le Malattie Infettive Lazzaro Spallanzani, Roma); Giovanni Apolone (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano); Fabio Ciceri (IRCCS Ospedale San Raffaele, Milano); Gennaro Ciliberto (IRCCS Istituto Nazionale Tumori Regina Elena, Roma); Massimo Costantini (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano); Franco Locatelli (Università La Sapienza, Roma); Alberto Mantovani (IRCCS Istituto Clinico Humanitas, Milano); Fausto Baldanti (Fondazione IRCCS Policlinico San Matteo di Pavia); Aldo Morrone (Istituto Dermatologico San Gallicano IRCCS, Roma); Carlo Salvarani (Azienda USL-IRCCS Reggio Emilia); Nicola Silvestris (IRCCS Istituto Tumori “Giovanni Paolo II”, Bari); Fabrizio Tagliavini (Fondazione IRCCS Istituto Neurologico Carlo Besta, Milano); Antonio Uccelli (Ospedale Policlinico San Martino IRCCS, Genova); Pier Luigi Zinzani (IRCCS Azienda Ospedaliero-Universitaria di Bologna).

DISEASE GROUPS

1. HAEMATOLOGICAL MALIGNANCIES Referent: Paolo Corradini (Fondazione IRCCS Istituto Nazionale dei Tumori, Milano);
2. SOLID TUMORS Referent: Gennaro Ciliberto (IRCCS Istituto Nazionale Tumori Regina Elena, Roma);
3. IMMUNORHEUMATOLOGICAL DISEASES Referent: Carlo Salvarani (Azienda USL IRCCS Reggio Emilia);
4. NEUROLOGICAL DISEASES: Referent: Antonio Uccelli (Ospedale Policlinico San Martino IRCCS, Genova); Renato Mantegazza (Fondazione I.R.C.C.S Istituto Neurologico Carlo Besta (INCB), Milano).

IMMUNOLOGICAL GROUP

REFERENTS: Chiara Agrati (IRCCS Istituto per le Malattie Infettive Lazzaro Spallanzani, Roma); Maria Rescigno (IRCCS Istituto Clinico Humanitas, Milano); Daniela Fenoglio (Ospedale Policlinico San Martino IRCCS, Genova);

PARTICIPANTS: Roberta Mortarini (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano); Cristina Tresoldi (IRCCS Ospedale San Raffaele, Milano); Laura Conti (IRCCS Istituto Nazionale Tumori Regina Elena, Roma); Stefania Croci (Azienda USL IRCCS Reggio Emilia); Fausto Baldanti (Fondazione IRCCS Policlinico San Matteo di Pavia); Vito Garrisi (IRCCS Istituto Tumori “Giovanni Paolo II”, Bari); Fulvio Baggi (Fondazione IRCCS Istituto Neurologico Carlo Besta, Milano); Francesca Bonifazi (IRCCS Azienda Ospedaliero-Universitaria di Bologna); Fulvia Pimpinelli (Istituto Dermatologico San Gallicano IRCCS, Roma).

INMI CENTRALIZED LABORATORY (INMI Lazzaro Spallanzani – IRCCS, Roma) (alphabetical order)

Enrico Girardi (Scientific Director), Aurora Bettini; Veronica Bordoni; Concetta Castillett; Eleonora Cimini; Rita Casetti; Francesca Colavita; Flavia Cristofanelli; Massimo Francalancia; Simona Gili; Giulia Gramigna; Germana Grassi; Daniele Lapa; Sara Leone; Davide Mariotti; Giulia Matusali; Silvia Meschi; Stefania Notari; Enzo Puro; Marika Rubino; Alessandra Sacchi; Eleonora Tartaglia

CLINICAL TASK FORCE

Paolo Corradini, Silvia Damian, Filippo de Braud (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano); Maria Teresa Lupo Stanghellini, Lorenzo Dagna, Francesca Ogliari, Massimo Filippi (IRCCS Ospedale San Raffaele: Milano); Giulia Piaggio (IRCCS Istituto Nazionale Tumori Regina Elena, Roma); Elena Azzolini, Chiara Pozzi, Luca Germagnoli, Carlo Selmi, Maria De Santis, Carmelo Carlo-Stella, Alexia Bertuzzi, Francesca Motta, Angela Ceribelli (IRCCS Istituto Clinico Humanitas, Milano);

Fondazione IRCCS Policlinico San Matteo di Pavia: Sara Monti,
Istituto Dermatologico San Gallicano IRCCS, Roma;

Azienda USL-IRCCS Reggio Emilia: Maria Grazia Catanoso,
Rosa Divella, Antonio Tufaro, Vito Garrisi, Sabina Delcuratolo, Mariana Miano (IRCCS Istituto Tumori “Giovanni Paolo II”, Bari);

Fondazione IRCCS Istituto Neurologico Carlo Besta, Milano;
Ospedale Policlinico San Martino IRCCS, Genova;

Marco Fusconi, Vittorio Stefoni, Maria Abbondanza Pantaleo (IRCCS Azienda Ospedaliero-Universitaria di Bologna).

STATISTICAL COMMITTEE

Diana Giannarelli (IRCCS Istituto Nazionale Tumori Regina Elena, Roma).

e-CRF AND MONITORING REFERENT:

Valentina Sinno, Serena Di Cosimo (Fondazione IRCCS Istituto Nazionale dei Tumori di Milano).

PROJECT MANAGERS OF THE STUDY:

REFERENCES: Elena Turola, Azienda USL-IRCCS di Reggio Emilia.

PARTICIPANTS: Iolanda Pulice, Roberta Mennitto Fondazione IRCCS Istituto Nazionale dei Tumori, Milano); Stefania Trinca (IRCCS Ospedale San Raffaele, Milano); Giulia Piaggio (IRCCS Istituto Nazionale Tumori Regina Elena, Roma); Chiara Pozzi (IRCCS Istituto Clinico Humanitas, Milano);

Irene Cassaniti (Fondazione IRCCS Policlinico San Matteo, Pavia); Alessandro Barberini (Istituto Dermatologico San Gallicano IRCCS, Roma); Arianna Belvedere (Azienda USL-IRCCS Reggio Emilia);

Sabina Del Curatolo (IRCCS Istituto Tumori “Giovanni Paolo II”, Bari); Rinaldi Elena, Federica Bortone (Fondazione IRCCS Istituto Neurologico Carlo Besta, Milano); Maria Giovanna Dal Bello (Ospedale Policlinico San Martino IRCCS, Genova); Silvia Corazza (IRCCS Azienda Ospedaliero-Universitaria, Bologna).

Institutional Review Boards:

The study, according to the National COVID-19 procedures, was approved by the Italian Regulatory Agency (AIFA) and by the Ethics Committee of IRCCS L. Spallanzani (code 304,2021).

References

- 1 Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med*. 2021 Feb 4;384(5):403-416. doi: 10.1056/NEJMoa2035389. Epub 2020 Dec 30. PMID: 33378609; PMCID: PMC7787219.

- 2 Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020 Dec 31;383(27):2603-2615. doi: 10.1056/NEJMoa2034577. Epub 2020 Dec 10. PMID: 33301246; PMCID: PMC7745181.

- 3 Lee LYW, Cazier JB, Starkey T, et al. COVID-19 prevalence and mortality in patients with cancer and the effect of primary tumour subtype and patient demographics: a prospective cohort study. *Lancet Oncol*. 2020 Oct;21(10):1309-1316. doi: 10.1016/S1470-2045(20)30442-3. Epub 2020 Aug 24.

- 4 Lee LY, Cazier JB, Angelis V, et al. COVID-19 mortality in patients with cancer on chemotherapy or other anticancer treatments: a prospective cohort study. *Lancet*. 2020 Jun 20;395(10241):1919-1926. doi: 10.1016/S0140-6736(20)31173-9. Epub 2020 May 28.

- 5 Chaudhry F, Bulka H, Rathnam AS, et al. COVID-19 in multiple sclerosis patients and risk factors for severe infection. *J Neurol Sci*. 2020 Nov 15;418:117147. doi: 10.1016/j.jns.2020.117147. Epub 2020 Sep 19. PMID: 32980780; PMCID: PMC7834402.

- 6 Salter A, Fox RJ, Newsome SD, et al. Outcomes and Risk Factors Associated With SARS-CoV-2 Infection in a North American Registry of Patients With Multiple Sclerosis. *JAMA Neurol*. 2021 Jun 1;78(6):699-708. doi: 10.1001/jamaneurol.2021.0688. Erratum in: *JAMA Neurol*. 2021 Jun 1;78(6):765. PMID: 33739362; PMCID: PMC7980147.

- 7 Goldman JD, Robinson PC, Uldrick TS, Ljungman P. COVID-19 in immunocompromised populations: implications for prognosis and repurposing of immunotherapies. *J Immunother Cancer*. 2021 Jun;9(6):e002630. doi: 10.1136/jitc-2021-002630. PMID: 34117116; PMCID: PMC8206176.

- 8 Rodríguez-Blanco N, Montero-Navarro S, Botella-Rico JM, Felipe-Gómez AJ, Sánchez-Más J, Tuells J. Willingness to Be Vaccinated against COVID-19 in Spain before the Start of Vaccination:

A Cross-Sectional Study. *Int J Environ Res Public Health*. 2021 May 15;18(10):5272. doi: 10.3390/ijerph18105272. PMID: 34063476; PMCID: PMC8155897.

9 Chan WL, Ho YT, Wong CK, Choi HC, Lam KO, Yuen KK, et al. Acceptance of COVID-19 Vaccination in Cancer Patients in Hong Kong: Approaches to Improve the Vaccination Rate. *Vaccines (Basel)*. 2021 Jul 16;9(7):792. doi: 10.3390/vaccines9070792. PMID: 34358208; PMCID: PMC8310340

10 Brodziak A, Sigorski D, Osmola M, Wilk M, Gawlik-Urban A, Kiszka J, et al. Attitudes of Patients with Cancer towards Vaccinations-Results of Online Survey with Special Focus on the Vaccination against COVID-19. *Vaccines (Basel)*. 2021 Apr 21;9(5):411. doi: 10.3390/vaccines9050411. PMID: 33919048; PMCID: PMC8142983.

11 Hwang JK, Zhang T, Wang AZ, Li Z. COVID-19 vaccines for patients with cancer: benefits likely outweigh risks. *J Hematol Oncol*. 2021 Feb 27;14(1):38. doi: 10.1186/s13045-021-01046-w. PMID: 33640005; PMCID: PMC7910769.

12 Uhr L, Mateen FJ. COVID-19 vaccine hesitancy in multiple sclerosis: A cross-sectional survey. *Mult Scler*. 2021 Jul 27;13524585211030647. doi: 10.1177/13524585211030647. Epub ahead of print. PMID: 34313513.

13 Ehde DM, Roberts MK, Humbert AT, Herring TE, Alschuler KN. COVID-19 vaccine hesitancy in adults with multiple sclerosis in the United States: A follow up survey during the initial vaccine rollout in 2021. *Mult Scler Relat Disord*. 2021 Jul 22;54:103163. doi: 10.1016/j.msard.2021.103163. Epub ahead of print. PMID: 34325399; PMCID: PMC8295053.

14 Gaur P, Agrawat H, Shukla A. COVID-19 vaccine hesitancy in patients with systemic autoimmune rheumatic disease: an interview-based survey. *Rheumatol Int*. 2021 Sep;41(9):1601-1605. doi: 10.1007/s00296-021-04938-9. Epub 2021 Jul 2. PMID: 34213580; PMCID: PMC8249840.

15 Waissengrin B, Agbarya A, Safadi E, Padova H, Wolf I. Short-term safety of the BNT162b2 mRNA COVID-19 vaccine in patients with cancer treated with immune checkpoint inhibitors. *Lancet Oncol*. 2021 May;22(5):581-583. doi: 10.1016/S1470-2045(21)00155-8. Epub 2021 Apr 1

16 So ACP, McGrath H, Ting J, Srikantharajah K, Germanou S, Moss C, et al. COVID-19 Vaccine Safety in Cancer Patients: A Single Centre Experience. *Cancers (Basel)*. 2021 Jul 16;13(14):3573. doi: 10.3390/cancers13143573. PMID: 34298785; PMCID: PMC8304256

17 Ram R, Hagin D, Kikozashvili N, Freund T, Amit O, Bar-On Y, et al. Safety and Immunogenicity of the BNT162b2 mRNA COVID-19 Vaccine in Patients after Allogeneic HCT or CD19-based CART therapy-A Single-Center Prospective Cohort Study. *Transplant Cell Ther*. 2021 Jun 30:S2666-6367(21)01027-7. doi: 10.1016/j.jct.2021.06.024. Epub ahead of print. PMID: 34214738; PMCID: PMC8242200.

18 Monin L, Laing AG, Muñoz-Ruiz M, McKenzie DR, Del Molino Del Barrio I, Alaguthurai T, et al. Safety and immunogenicity of one versus two doses of the COVID-19 vaccine BNT162b2 for patients with cancer: interim analysis of a prospective observational study. *Lancet Oncol*. 2021 Jun;22(6):765-778. doi: 10.1016/S1470-2045(21)00213-8. Epub 2021 Apr 27. PMID: 33930323; PMCID: PMC8078907.

19 Mahil SK, Bechman K, Raharja A, Domingo-Vila C, Baudry D, Brown MA, et al. The effect of methotrexate and targeted immunosuppression on humoral and cellular immune responses to the COVID-19 vaccine BNT162b2: a cohort study. *Lancet Rheumatol*. 2021 Jul 8. doi: 10.1016/S2665-9913(21)00212-5. Epub ahead of print. PMID: 34258590; PMCID: PMC8266273.

20 Pimpinelli F, Marchesi F, Piaggio G, Giannarelli D, Papa E, Falcucci P, et al. Fifth-week immunogenicity and safety of anti-SARS-CoV-2 BNT162b2 vaccine in patients with multiple myeloma and myeloproliferative malignancies on active treatment: preliminary data from a single institution. *J Hematol Oncol*. 2021 May 17;14(1):81. doi: 10.1186/s13045-021-01090-6. PMID: 34001183; PMCID: PMC8128283.

21 Chevallier P, Coste-Burel M, Le Bourgeois A, Peterlin P, Garnier A, Béné MC, et al. Safety and immunogenicity of a first dose of SARS-CoV-2 mRNA vaccine in allogeneic hematopoietic stem-cells recipients. *EJHaem*. 2021 Jun 1;10.1002/jha2.242. doi: 10.1002/jha2.242. Epub ahead of print. PMID: 34226903; PMCID: PMC8242867.

22 Maneikis K, Šablauskas K, Ringelevičiūtė U, Vaitekėnaitė V, Čekauskienė R, Kryžauskaitė L, et al. Immunogenicity of the BNT162b2 COVID-19 mRNA vaccine and early clinical outcomes in patients with haematological malignancies in Lithuania: a national prospective cohort study. *Lancet Haematol*. 2021 Jul 2:S2352-3026(21)00169-1. doi: 10.1016/S2352-3026(21)00169-1. Epub ahead of print. PMID: 34224668; PMCID: PMC8253543

23 Lotan I, Wilf-Yarkoni A, Friedman Y, Stiebel-Kalish H, Steiner I, Hellmann MA. Safety of the BNT162b2 COVID-19 vaccine in multiple sclerosis (MS): Early experience from a tertiary MS center in Israel. *Eur J Neurol*. 2021 Jul 21. doi: 10.1111/ene.15028. Epub ahead of print. PMID: 34288285.

24 Achiron A, Dolev M, Menascu S, Zohar DN, Dreyer-Alster S, Miron S, et al. COVID-19 vaccination in patients with multiple sclerosis: What we have learnt by February 2021. *Mult Scler*. 2021 May;27(6):864-870. doi: 10.1177/13524585211003476. Epub 2021 Apr 15. PMID: 33856242; PMCID: PMC8114441.

25 Allen-Philbey K, Stennett A, Begum T, Johnson AC, Dobson R, Giovannoni G, et al. Experience with the COVID-19 AstraZeneca vaccination in people with multiple sclerosis. *Mult Scler Relat Disord*. 2021 Jul;52:103028. doi: 10.1016/j.msard.2021.103028. Epub 2021 May 18. PMID: 34049216; PMCID: PMC8129799.

26 Furer V, Eviatar T, Zisman D, Peleg H, Paran D, Levartovsky D, et al. Immunogenicity and safety of the BNT162b2 mRNA COVID-19 vaccine in adult patients with autoimmune inflammatory rheumatic diseases and in the general population: a multicentre study. *Ann Rheum Dis*. 2021 Jun 14;annrheumdis-2021-220647. doi: 10.1136/annrheumdis-2021-220647. Epub ahead of print. PMID: 34127481; PMCID: PMC8206170.

27 Rotondo C, Cantatore FP, Fornaro M, Colia R, Busto G, Rella V, et al. Preliminary Data on Post Market Safety Profiles of COVID 19 Vaccines in Rheumatic Diseases: Assessments on Various Vaccines in Use, Different Rheumatic Disease Subtypes, and Immunosuppressive Therapies: A Two-Centers Study. *Vaccines (Basel)*. 2021 Jul 2;9(7):730. doi: 10.3390/vaccines9070730. PMID: 34358147; PMCID: PMC8310114.

28 Braun-Moscovici Y, Kaplan M, Braun M, Markovits D, Giryès S, Toledano K, et al. Disease activity and humoral response in patients with inflammatory rheumatic diseases after two doses of the Pfizer mRNA vaccine against SARS-CoV-2. *Ann Rheum Dis*. 2021 Jun 18;annrheumdis-2021-220503. doi: 10.1136/annrheumdis-2021-220503. Epub ahead of print. PMID: 34144967; PMCID: PMC8214990.

29 Felten R, Kawka L, Dubois M, Ugarte-Gil MF, Fuentes-Silva Y, Piga M, Arnaud L. Tolerance of COVID-19 vaccination in patients with systemic lupus erythematosus: the international VACOLUP study. *Lancet Rheumatol*. 2021 Jul 21. doi: 10.1016/S2665-9913(21)00221-6. Epub ahead of print. PMID: 34312612; PMCID: PMC8294805.

30 Cherian S, Paul A, Ahmed S, Alias B, Manoj M, Santhosh AK, et al. Safety of the ChAdOx1 nCoV-19 and the BBV152 vaccines in 724 patients with rheumatic diseases: a post-vaccination cross-sectional survey. *Rheumatol Int*. 2021 Aug;41(8):1441-1445. doi: 10.1007/s00296-021-04917-0. Epub 2021 Jun 17. PMID: 34142203; PMCID: PMC8211311.

31 Connolly CM, Ruddy JA, Boyarsky BJ, Avery RK, Werbel WA, Segev DL, et al. Safety of the first dose of mRNA SARS-CoV-2 vaccines in patients with rheumatic and musculoskeletal diseases. *Ann Rheum Dis*. 2021 Mar 19;annrheumdis-2021-220231. doi: 10.1136/annrheumdis-2021-220231. Epub ahead of print. PMID: 33741555.

32 Boekel L, Kummer LY, van Dam KPJ, Hooijberg F, van Kempen Z, Vogelzang EH, et al. Adverse events after first COVID-19 vaccination in patients with autoimmune diseases. *Lancet Rheumatol*. 2021 Aug;3(8):e542-e545. doi: 10.1016/S2665-9913(21)00181-8. Epub 2021 Jun 18. PMID: 34179831; PMCID: PMC8213359.

33 Sattui SE, Liew JW, Kennedy K, Sirotich E, Putman M, Moni TT, et al. Early experience of COVID-19 vaccination in adults with systemic rheumatic diseases: results from the COVID-19 Global Rheumatology Alliance Vaccine Survey. *RMD Open*. 2021 Sep;7(3):e001814. doi: 10.1136/rmdopen-2021-001814. PMID: 34493645; PMCID: PMC8424419.

34 Ali A, Dwyer D, Wu Q, Wang Q, Dowling CA, Fox DA, et al. Characterization of humoral response to COVID mRNA vaccines in multiple sclerosis patients on disease modifying therapies.

Vaccine. 2021 Oct 1;39(41):6111-6116. doi: 10.1016/j.vaccine.2021.08.078. Epub 2021 Sep 2. PMID: 34483021; PMCID: PMC8411370

35 Cavanna L, Citterio C, Biasini C, Madaro S, Bacchetta N, Lis A, et al. COVID-19 vaccines in adult cancer patients with solid tumours undergoing active treatment: Seropositivity and safety. A prospective observational study in Italy. *Eur J Cancer*. 2021 Sep 2;157:441-449. doi: 10.1016/j.ejca.2021.08.035. Epub ahead of print. PMID: 34601285; PMCID: PMC8410513.

36 Peeters M, Verbruggen L, Teuwen L, Vanhoutte G, Vande Kerckhove S, Peeters B, et al. Reduced humoral immune response after BNT162b2 coronavirus disease 2019 messenger RNA vaccination in cancer patients under antineoplastic treatment. *ESMO Open*. 2021 Sep 8;6(5):100274. doi: 10.1016/j.esmoop.2021.100274. Epub ahead of print. PMID: 34597941; PMCID: PMC8423808.

37 Shroff RT, Chalasani P, Wei R, Pennington D, Quirk G, Schoenle MV, et al. Immune responses to two and three doses of the BNT162b2 mRNA vaccine in adults with solid tumors. *Nat Med*. 2021 Sep 30. doi: 10.1038/s41591-021-01542-z. Epub ahead of print. PMID: 34594036.

38 Di Noia V, Pimpinelli F, Renna D, Barberi V, Maccallini MT, Gariazzo L, et al. Immunogenicity and safety of COVID-19 vaccine BNT162b2 for patients with solid cancer: a large cohort prospective study from a single institution. *Clin Cancer Res*. 2021 Sep 28;clincanres.2439.2021. doi: 10.1158/1078-0432.CCR-21-2439. Epub ahead of print. PMID: 34583970.

39 Linardou H, Spanakis N, Koliou GA, Christopoulou A, Karageorgopoulou S, Alevra N, et al. Responses to SARS-CoV-2 Vaccination in Patients with Cancer (ReCOVer Study): A Prospective Cohort Study of the Hellenic Cooperative Oncology Group. *Cancers (Basel)*. 2021 Sep 15;13(18):4621. doi: 10.3390/cancers13184621. PMID: 34572848; PMCID: PMC8466969.

40 Lasagna A, Agustoni F, Percivalle E, Borgetto S, Paulet A, Comolli G, et al. A snapshot of the immunogenicity, efficacy and safety of a full course of BNT162b2 anti-SARS-CoV-2 vaccine in cancer patients treated with PD-1/PD-L1 inhibitors: a longitudinal cohort study. *ESMO Open*. 2021 Sep 1;6(5):100272. doi: 10.1016/j.esmoop.2021.100272. Epub ahead of print. PMID: 34543863; PMCID: PMC8407964.

41 Shmueli ES, Itay A, Margalit O, Berger R, Halperin S, Jurkowicz M, et al Efficacy and safety of BNT162b2 vaccination in patients with solid cancer receiving anticancer therapy - a single centre prospective study. *Eur J Cancer*. 2021 Sep 8;157:124-131. doi: 10.1016/j.ejca.2021.08.007. Epub ahead of print. PMID: 34508994; PMCID: PMC8424105.

42 Goshen-Lago T, Waldhorn I, Holland R, Szwarcwort-Cohen M, Reiner-Benaim A, Shachor-Meyouhas Y, et al. Serologic Status and Toxic Effects of the SARS-CoV-2 BNT162b2 Vaccine in Patients Undergoing Treatment for Cancer. *JAMA Oncol*. 2021 Jul 8:e212675. doi: 10.1001/jamaoncol.2021.2675. Epub ahead of print. PMID: 34236381; PMCID: PMC8267843.

43 Delvino P, Bozzalla Cassione E, Biglia A, Quadrelli VS, Bartoletti A, et al. Safety of BNT162b2 mRNA COVID-19 vaccine in a cohort of elderly, immunocompromised patients with systemic vasculitis. *Clin Exp Rheumatol*. 2021 Sep 16. Epub ahead of print. PMID: 34528506.

44 Agrati C, Di Cosimo S, Fenoglio D, Apolone G, Ciceri F, Ciliberto G, et al. COVID-19 Vaccination in Fragile Patients: Current Evidence and an Harmonized Transdisease Trial. *Front. Immunol.*, 10 August 2021 | <https://doi.org/10.3389/fimmu.2021.704110>.

Figure legends:

Figure 1 – Patients’ disposition

Cohort	N of pts	Control group	Vaccine	Methods for evaluation of AE	AE reported	Severe AE
Allo-HSCT ^{17,21}	192	26 healthy subjects	BNT162b2	Questionnaire within 7 days following the first vaccination ²¹ . Interview and clinical evaluation, grading CTC AE v5 ¹⁷	Only grade 1 or 2 AEs were observed. Cytopenia 12% after the 1st dose and 10% after the 2nd dose ¹⁷ . 3 cases of GvHD exacerbation after each dose ¹⁷ .	There were no grade 3 or 4 non-hematologic adverse events. A single case of impending graft rejection was summarized as possibly related.
Oncohematolog ^y ¹⁹⁻²²	977	67 healthy subjects + 36 elderly w/o cancers	BNT162b2	Questionnaires adapted from the original phase 3 BNT162b2 trial ²² Registration of local and systemic side effects ¹⁹	AEs were more common after the 2nd dose ²² . Mild pain at the injection site most common side effect in all the cohorts ¹⁹ .	No grade 4 adverse events were reported
Cancer ^{16, 18, 35-39, 41, 42}	2387	739 healthy subjects	BNT162b2 mRNA-1273 ChAdOx1	Solicited and unsolicited AE collection through face-to-face or telephone consultations ¹⁶ . Weekly telephone consultation ¹⁸ . Patient-reported outcomes between 1-2-4 weeks after each dose ³⁵ . Web-based electronic platform for home toxicity monitoring ³⁶ . AE graded according to patients interview within 2-4-week windows ³⁷ . Patient-reported scale (very mild / mild / moderate / severe) ³⁸ .	The most frequently reported local AE was mild-to-moderate pain at the injection site ³⁶ . AEs occurred between 39% and 56% patients after the 1 st dose and between 29% and 58% after the 2 nd ^{18,38,41} . 1st dose AEs: local side-effects 29.8% - 31.5%, fatigue 8.3% - 7%, headaches 3.9% - 8.17% and arthralgia/myalgia 3.7% - 2.71% ³⁵⁻³⁸ . 2nd dose AEs: fever 5.8% - 14.1%, local reactions 12.6% - 33.46%, fatigue 10.4% - 8.9%, arthralgia/myalgia 5.2% ³⁸ . Overall: 45% - 69% injection site pain, myalgia 9.5% - 15%, bone	No grade 3 / 4 side effects were recorded ^{18, 35, 37, 39, 41, 42} . Low rate of grade 3 / 4 AE: 2.1% vaccine related AE grade 3 ¹⁶ , 3.3% of patients after the 1st inoculum and 1.4% after the booster ³⁸ . One sudden death occurred after 5 days of the vaccine in a hospitalized patient who received active chemotherapy for advanced bladder cancer. The post-mortem examination revealed pulmonary embolism ³⁸ . Three cancer-related deaths were considered unrelated to the BNT162b2 vaccine ³⁶ . One patient died due to myocardial infarction - the authors cannot complete rule out that this event was vaccine

				Questionnaire ^{39, 41} . Review of medical records ⁴² .	pain 5%, headache 10% - 24.3%, fatigue 10% - 28.6%, chills 10%, appetite loss 5% ¹⁶⁻³⁷⁻³⁹⁻⁴² . AEs were more common after the 2nd vaccine dose than that with the 1st (33.3% vs 7%) ⁴¹ .	related ³⁶ . Severe reactogenicity occurred in 1% of the patients following the priming dose and in 3% of the patients following the booster dose ³⁶ .	
cancers inhibitors) ^{15, 40}	(checkpoint	222	2241 healthy subjects	BNT162 b2	Solicited and unsolicited AE collection Telephone questionnaires 17-21 days after the first vaccine dose and median 19 days after the second	1st dose AEs pain at the injection site 21%-28.57% ¹⁵⁻⁴⁰ . 2nd dose AEs fever 6.94% ⁴⁰ ; pain at the injection site 63%, local rash 2%, local swelling 9%; muscle pain 34%, fatigue 34%, headache 16%, fever 10%, chills 10%, gastrointestinal complications 10%, flu-like symptoms 2%. ¹⁵ Systemic events more frequent after the 2nd dose, while local effects were less common. ⁴⁰	No thrombosis, hypersensitivity adverse events or vaccine-related anaphylaxis were signaled. One patient has reported two immune-related side-effects (hepatitis G3 and colitis G3) 10 days after the first dose of vaccine.
		896	7 healthy subjects	BNT162 b2 ChAdOx 1 mRNA-1273	Questionnaires ^{23,25} Telemedicine and face-to-face evaluation (7-21 days after each dose) ²⁴	Between 56.9% and 94% reported AEs ^{23, 25} . AEs: sore arm 70%, flu-like symptoms 64%, fever 21%, fatigue 27%, headache 21% ^{24, 25, 34} .	No severe adverse effects occurred ²⁵ 15.1% of participants reported new or worsening neurological symptoms ²³ . Acute relapse 2.1% and 1.6% following the first and second doses, respectively ²⁴ . 2 people reported new or worsening MS symptoms ³⁴
Systemic autoimmune disease 14, 19, 26-33, 43		649 2	394 healthy subjects + 26 COVID recoverers ²⁸	BNT162 b2 ChAdOx 1 mRNA-1273 BBV152 CoronaVac	Registration of local and systemic side effects ¹⁹ Questionnaire distributed via e-mail ³² , phone ²⁶ , web-based ²⁹ 7th days after vaccination collection of data ³⁰ Questionnaire within the 1st week after	Absence of AEs 20% of patients ¹⁹ . Overall occurrence of AEs 35% - 60.5% ^{14, 30, 32, 33,43} AEs: local reactions 17% - 89%; systemic symptoms 69%, headache 12% - 20%; muscle sore 10% - 22.8% fatigue 7.4% - 33.4% ^{19,27, 28, 31, 33} . 1st dose AEs 45% - 53%, 2nd dose	No participants reported severe adverse events ^{19, 29-31, 43} . Severe adverse events 1% ³² . Major adverse events: death (2) several weeks after the 2nd dose, non-disseminated herpes zoster (6), uveitis (2), and pericarditis (1) ²⁶ . Postvaccination disease activity remained stable in the majority of patients ²⁶ . Relapses of the underlying disease 2.2% of

Ad26CO V2.S	vaccination ³¹ Questionnaires ^{27, 28, 33, 43}	AEs 26% - 53% ^{27, 29} . Patients who received both vaccine doses and reported side-effects after the 1st dose were more likely to report side-effects after the 2nd dose than those who did not (relative risk 2.30, 95% CI 1.88–2.82; p<0.0001) ²⁹ .	patients after the first dose of vaccine ²⁷ . 11% of patients reported that their psoriasis worsened after vaccination ¹⁹ . Immunosuppressive treatment was postponed because of COVID-19 vaccination in 6% patients ³² Four patients reported flare of arthritis that resolved within 5 days ³⁰ . One patient with Familial Mediterranean fever reported new-onset arthritis 2 weeks after the 1st dose. No flare-up of the underlying IRD occurred in any other patient ²⁸ . Flares of existing systemic rheumatic disease 13.4%, 4.6% requiring treatment ³³ .

Table 1 – Literature review

Table 2 – Patients’ characteristics

	N (%)
Age	
18-50 years	160 (28.3)
51-70 years	299 (52.8)
>= 71 years	107 (18.9)
Patient sex	
Males	249 (44.0)
Females	317 (56.0)
Main diagnosis	
Haematological malignancies	131 (23.1)
Solid tumours	191 (33.7)
Immunorheumatological diseases	86 (15.2)
Neurological diseases	158 (27.9)
Vaccine	
mRNA-1237 Moderna	105 (18.6)
BNT-162b2 Pfizer BioNTech	461 (81.4)
All patients	566 (100)

Table 3 – Occurrence of symptoms / adverse events over the week after vaccine administration

	t1- One week after the first dose			t2 - One week after the second dose			PPV	NPV
	N 488			N 489				
	Not at all (%)	Moderate (%)	Severe (%)	Not at all (%)	Moderate (%)	Severe (%)	(%)	(%)
<i>Pain</i>	194 (39.8)	263 (53.9)	31 (6.4)	218 (44.6)	246 (50.3)	25 (5.1)	17.2	96.2
<i>Fatigue</i>	341 (69.9)	125 (25.6)	22 (4.5)	285 (58.3)	171 (35.0)	33 (6.7)	28.6	93.9
<i>Headache</i>	430 (88.1)	48 (9.8)	10 (2.0)	397 (81.2)	78 (16.0)	14 (2.9)	25.0	97.7
<i>Pain bone</i>	408 (83.6)	67 (13.7)	13 (2.7)	356 (72.8)	114 (23.3)	19 (3.9)	25.0	96.8
<i>Fever</i>	459 (94.1)	23 (4.7)	6 (1.2)	411 (84.0)	49 (10.0)	29 (5.9)	83.5	94.8
<i>Enlarged lymph nodes</i>	487 (99.8)	-	1 (0.2)	478 (97.8)	10 (2.0)	1 (0.2)	NE	99.8
<i>Skin rash</i>	471 (96.5)	14 (2.9)	3 (0.6)	458 (93.7)	22 (4.5)	9 (1.8)	NE	98.4
<i>Insomnia</i>	464 (95.1)	22 (4.5)	2 (0.4)	443 (90.6)	41 (8.4)	5 (1.0)	NE	99.1
<i>Diarrheal</i>	461 (94.5)	25 (5.1)	2 (0.4)	454 (92.8)	32 (6.5)	3 (0.6)	NE	99.6
<i>Nausea or vomiting</i>	471 (96.5)	15 (3.1)	2 (0.4)	457 (93.5)	29 (5.9)	3 (0.6)	NE	99.6

PPV (positive predictive value) probability of occurrence of a severe symptom after the second dose according to the occurrence of the same symptom after the first dose.

NPV (negative predictive value) probability of absence of a severe symptom after the second dose according to the not occurrence of the same after the first dose.

NE not evaluable

“Moderate” and “Severe” include patients who have been troubled slightly or moderately (moderate) and severely or overwhelmingly (severe), respectively.

Table 4 - Probability of occurrence of a severe symptom after one of the two doses of vaccine estimated through a multivariate logistic regression adjusted by age, sex, diagnosis, and vaccine

	Severe pain			Severe fatigue		Severe headache		Severe pain bone		Severe fever	
	N.	N. (%)	Odd Ratio (95% CI)	N. (%)	Odd Ratio (95% CI)	N. (%)	Odd Ratio (95% CI)	N. (%)	Odd Ratio (95% CI)	N. (%)	Odd Ratio (95% CI)
Age											
18-50	160	19 (11.9)	Ref.	17 (10.7)	Ref.	12 (7.5)	Ref	8 (5.0)	Ref	17 (10.7)	Ref
51-70	299	26 (8.7)	1.1 (0.6-2.2)	25 (8.4)	1.0 (0.5-2.0)	7 (2.3)	0.5 (0.2-1.3)	14 (4.7)	1.1 (0.4-2.8)	11 (3.7)	0.3 (0.1-0.7)
71 +	107	6 (5.6)	0.6 (0.2-1.6)	7 (6.5)	0.7 (0.3-1.9)	3 (2.8)	0.5 (0.1-2.0)	7 (6.5)	1.5 (0.5-4.4)	2 (1.9)	0.1 (0.1-0.7)
p-value		0.401		0.740		0.315		0.725		0.003	
Sex											
Males	249	10 (4.0)	Ref.	13 (5.2)	Ref.	2 (0.8)	Ref	7 (2.8)	Ref	9 (3.6)	Ref
Females	317	41 (12.9)	2.9 (1.4-6.2)	36 (11.4)	2.3 (1.2-4.5)	20 (6.3)	7.6 (1.7-33.6)	22 (6.9)	2.7 (1.1-6.5)	21 (6.6)	1.6 (0.7-3.8)
p-value		0.005		0.016		0.007		0.028		0.253	
Diagnosis											
Haematological	131	6 (4.6)	Ref.	9 (6.9)	Ref.	4 (3.1)	Ref	7 (5.3)	Ref	7 (5.3)	Ref
Solid tumours	191	4 (2.1)	0.3 (0.1-1.0)	10 (5.2)	0.7 (0.3-1.7)	2 (1.0)	0.2 (0.1-1.4)	8 (4.2)	0.7 (0.2-1.9)	10 (5.2)	0.8 (0.3-2.2)
Rheumatological	86	15 (17.4)	4.0 (1.4-11.0)	10 (11.6)	1.5 (0.6-4.0)	3 (3.5)	0.8 (0.2-3.7)	4 (4.7)	0.7 (0.2-2.6)	5 (5.8)	1.0 (0.3-3.5)
Neurological	158	26 (16.5)	2.8 (1.1-7.5)	20 (12.7)	1.7 (0.7-4.1)	13 (8.2)	1.7 (0.5-5.8)	10 (6.3)	1.1 (0.4-3.0)	8 (5.1)	0.5 (0.2-1.6)

<i>p-value</i>		)									
Vaccine		<0.001		0.108		0.090		0.761		0.646	
<i>m-RNA-1237</i>	10	19	Ref.	9	Ref.	6	Ref	6	Ref	12	Ref
	5	(18.1	)	(8.6)		(5.7)		(5.7)		(11.4)	
<i>BNT-162b2</i>	46	32	0.3	40	1.1	16	0.7	23	0.9	18	0.3
	1	(6.9)	(0.1-0.6)	(8.7)	(0.5-2.4)	(3.5)	(0.3-2.0)	(5.0)	(0.4-2.5)	(3.9)	(0.1-0.7)
<i>p-value</i>		0.001		0.826		0.510		0.923		0.004	
Ref. is the referent strata											