Use of magnetic resonance imaging in neuroprognostication after pediatric cardiac arrest: Survey of current practices

Juan A. Piantino, MD^{1*}; Christopher M. Ruzas, MD^{2*}; Craig A. Press, MD, PhD³; Subramanian Subramanian, MD⁴; Binod Balakrishnan MD⁵; Ashok Panigrahy, MD⁴; David Pettersson, MD⁶; John A. Maloney, MD⁷; Arastoo Vossough, MD⁸; Alexis Topjian, MD, MSCE⁹; Matthew P Kirschen, MD, PhD⁹; Lesley Doughty, MD¹⁰; Melissa G. Chung, MD¹¹, David Maloney, BS¹²; Tamara Haller, BS¹³; Anthony Fabio, PhD, MPH¹³; Ericka L. Fink, MD, MS¹² for the POCCA Investigators.

* Dual first authors

¹ Department of Pediatrics, Division of Child Neurology, Doernbecher Children's Hospital, Oregon Health and Science University, Portland, OR, USA

² Department of Pediatrics, Division of Pediatric Critical Care Medicine, University of Colorado School of Medicine, Children's Hospital Colorado, Aurora, CO, USA

³ Department of Pediatrics, Division of Neurology, University of Pennsylvania Perelman School of Medicine, Children's Hospital of Philadelphia, Philadelphia, PA USA

⁴ Department of Radiology, UPMC Children's Hospital of Pittsburgh, Pittsburgh, PA, USA

⁵ Department of Pediatrics, Division of Pediatric Critical Care Medicine, Children's Wisconsin, Medical College of Wisconsin, Milwaukee, WI, USA

⁶ Department of Diagnostic Radiology, Division of Neuroradiology, Doernbecher Children's Hospital, Oregon Health & Science University, Portland, OR, USA

⁷ Department of Radiology, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, CO, USA

⁸ Department of Radiology, Children's Hospital of Philadelphia, Philadelphia, PA, USA

⁹ Department of Anesthesiology and Critical Care Medicine, University of Pennsylvania Perelman School of Medicine, Children's Hospital of Philadelphia, Philadelphia, PA USA

¹⁰ Division of Critical Care Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

¹¹ Department of Pediatrics, Divisions of Critical Care Medicine and Pediatric Neurology, Nationwide Children's Hospital, The Ohio State University, Columbus, OH, USA

¹² Department of Critical Care Medicine, UPMC Children's Hospital of Pittsburgh, Pittsburgh, PA USA

¹³ Department of Epidemiology, University of Pittsburgh, Pittsburgh, PA, USA

Corresponding author:

Ericka L. Fink, MD, MS

Division of Pediatric Critical Care Medicine

UPMC Children's Hospital of Pittsburgh

4401 Penn Avenue, Faculty Pavilion, 2nd floor

Pittsburgh, PA 15224

Phone: 412-692-5164 Fax: 412-692-6076

Email: finkel@ccm.upmc.edu

Keywords: Child; Cardiac Arrest; Neuroimaging; MRI; Hypoxia-Ischemia, Brain; Surveys and Questionnaires

Funding:

Juan Piantino: National Heart, Lung, and Blood Institute (NHLBI) K23HL150217

Ericka Fink: National Institute of Neurological Disorders and Stroke (NINDS) R01NS096714

Declarations of Interest:

Dr. Fink receives funding from National Institute of Neurological Disorders and Stroke (NINDS) R01NS096714; Neurocritical Care Society Incline Grant

Dr. Piantino receives support from the National Institute of Health, National Heart, Lung, and Blood Institute; grant K23HL150217

Dr. Press has performed paid expert testimony and provides consulting for Marinus Pharmaceuticals
The remainder of the authors have no relevant interests to disclose.

Word Count: 3261

Abstract

Background: Use of MRI as a tool to aid in neuroprognostication after cardiac arrest (CA) has been described, yet details of specific indications, timing, and sequences are unknown. We aim to define the current practices in use of brain MRI in prognostication after pediatric cardiac arrest.

METHODS: A survey was distributed to pediatric institutions participating in three international studies. Survey questions related to center demographics, clinical practice patterns of MRI after CA, neuroimaging resources, and details regarding MRI decision support.

RESULTS: Response rate was 31% (44/143). Thirty-four percent (15/44) of centers have a clinical pathway informing the use of MRI after CA. Fifty percent (22/44) of respondents reported that an MRI is obtained in nearly all CA patients and 32% (14/44) obtain an MRI in those who did not return to baseline neurological status. Sixty-four percent of centers (28/44) obtain an MRI greater than 72 hours after return of spontaneous circulation. Poor neurologic exam was reported as the most common factor (91% [40/44]) determining the timing of the MRI. Conventional sequences (T1, T2, FLAIR, and diffusion weighted imaging/apparent diffusion coefficient) are widely used. Advanced imaging techniques, such as MR spectroscopy, diffusion tensor imaging and resting state functional MRI are less commonly used.

CONCLUSIONS: Conventional brain MRI is a common practice for prognostication after CA. Advanced imaging techniques, are used infrequently. The lack of standardized clinical pathways and variability in

reported practices support a need for higher-quality evidence regarding the indications, timing, and acquisition protocols of clinical MRI studies.

INTRODUCTION

Pediatric cardiac arrest (CA) is a relatively uncommon event yet continues to be a major public health concern. Epidemiological studies have described a prevalence of nontraumatic out-of-hospital CA (OHCA) of 8 per 100,000 person years and a prevalence of in-hospital CA (IHCA) of 0.77 per 1,000 hospital admissions. Mortality rates range from 40-60% for IHCA with greater than 90% for OHCA.¹⁻⁵ In those who survive to hospital discharge, rates of survival with favorable neurologic outcome are 4-16%^{6,7} for OHCA and 80-96%^{3, 5} for IHCA.

Multiple monitoring modalities including electroencephalography (EEG), laboratory tests, and neuroimaging, are used to aid management and prognostication after CA.⁸⁻¹¹ However, no single modality has sufficient accuracy to facilitate neurological prognostication after return of spontaneous circulation (ROSC).¹² In recent years, brain magnetic resonance imaging (MRI)-based biomarkers have been increasingly recognized as promising tools to evaluate children after CA. Recent studies have shown that conventional MRI sequences (T1, T2, and diffusion-weighted imaging [DWI]) can accurately identify hypoxic-ischemic injury and can be used to assist in prognostication.¹³⁻¹⁹ Additionally, more advanced MRI techniques, such as MR spectroscopy (MRS), are able to measure the concentration of metabolites that serve as markers of injury and may also have prognostic value.^{10,20,21} Quantitative whole brain white matter fractional anisotropy (FA) measurements utilizing diffusion tensor imaging (DTI) are associated with long term neurological outcome in patients with cardiac arrest.¹⁸ Resting state functional MRI (rs-fMRI) has also been shown to have prognostic information in cardiac arrest patients who are comatose and have indeterminate prognosis and patients with higher functional connectivity recovered consciousness.¹⁹

Despite these advances, there is a lack of evidence-based consensus regarding the indications, timing, and MRI sequences most useful in this population, and little is known about current practice. The objective of this study was to assess current practices of post-CA MRI for neuroprognostication at

international tertiary care pediatric centers. To achieve our objective, we surveyed pediatric centers to assess practices in four areas: neuro-imaging resources, rationale for MRI, timing of MRI acquisition following CA, and specific MRI sequences used.

METHODS

Study design

This was a survey of current practices regarding brain MRI use in children with post-CA. The study was approved by the University of Pittsburgh Institutional Review Board as “Not Human Research” (STUDY20010055). The survey was conducted between May and August 2020. Pediatric hospitals participating in the Personalizing Outcomes after Child Cardiac Arrest (POCCA) study, the Approaches and Decisions after Pediatric Traumatic Brain Injury (ADAPT) trial, and the Prevalence of Acute critical Neurological disease in children: a Global Epidemiological Assessment (PANGEA) study were invited to participate in the survey. All participating sites have expertise in critical care and imaging-related research.²²⁻²⁴

Survey development and distribution

The 16-item survey was developed in consultation with experts in the field of Pediatric Critical Care and Neuroradiology at the University of Pittsburgh and Oregon Health & Science University. After the initial survey was developed, the remainder of the study group provided feedback. The final survey consisted of a multiple-choice questionnaire divided into two sections. The first section contained general information about each institution, the characteristics of each intensive care unit (total number of pediatric intensive care unit [PICU] beds, annual number of pediatric CA), and neuro-imaging resources (number and type of MRI scanners, availability of experts in neuroradiology). The second section

focused on MRI practices and presence of an institutional clinical pathway to support clinical decisions surrounding indications of MRI after CA, timing, and MRI sequences used in these patients.

The survey was distributed by the POCCA data coordinating center to all the participating sites. Each site's medical director or principal investigator was contacted via email and asked to complete the survey, with the assistance of the institution's neuroradiologist if needed. Follow-up emails were sent every two weeks up to four times. A single survey was completed per institution. If an individual site participated in more than one of the studies used to generate the survey distribution list, the site principal investigator of the most contemporary study was selected to receive this survey. Site principal investigators were physicians in Pediatric Neurology or Pediatric Critical Care.

Statistical analysis

Descriptive statistics were summarized as frequencies for categorical data and median and inter-quartile-range (IQR) for continuous data. Data from survey results were dichotomized based on the presence or absence of a clinical pathway for use of brain MRI as a prognostication tool following pediatric CA. Submitted surveys that did not include a response to this question were excluded from data analysis. Statistical analyses were performed using SAS® software, Version 9.4 of the SAS System for Windows. Copyright © 2015, Cary, NC, USA.

RESULTS

Cohort characteristics

The survey was distributed to 143 individual institutions. Of those, 32% (46/143) completed the survey. Two surveys did not include a response to the survey question: "Does your center have a clinical pathway for the use of brain MRI as a prognostication tool following pediatric cardiac arrest?" and thus were excluded. The respondents included 27 institutions from the US (61.4%) and 17 international

institutions (38.6%). Fifty-nine percent (26/44) of respondents were from freestanding children's hospitals, 36% (16/44) were from hospitals annexed to an adult hospital, and 5% (2/44) responded "other" regarding hospital type. Participating sites had a median of 23 (IQR 12-36) pediatric ICU beds. A minority of sites (34% [15/44]) reported a clinical pathway to guide MRI use in CA. The number of CA cases at responding sites varied substantially (Table 1).

Table 1. Site characteristics.

n(%) or median(25 th , 75 th percentiles)	All (n=44)	Clinical Pathway for MRI	
		No (n=29)	Yes (n=15)
Center's country	n=44	n=29	n=15
United States	27 (61.3)	18 (62)	9 (60)
Canada	3 (6.8)	3 (10.3)	0 (0)
United Kingdom	2 (4.5)	1 (3.4)	1 (6.67)
Spain	3 (6.8)	2 (6.9)	1 (6.6)
Australia	3 (6.8)	2 (6.9)	1 (6.6)
Other	6 (13.6)	3 (10.3)	3 (20)
Type of hospital	n=44	n=29	n=15
Freestanding children's hospital	26 (59)	15 (51.7)	11 (73.3)
Children's hospital annexed to an adult hospital	16 (36.3)	13 (44.8)	3 (20)
Other	2 (4.55)	1 (3.45)	1 (6.67)
Number of PICU beds	n=43	n=28	n=15
	23 (12 - 36)	19 (13.5 - 32.5)	36 (10 - 48)
Estimated no. pediatric in-hospital cardiac arrests per year	n=43	n=29	n=14
< 10	10 (23.2)	8 (27.5)	2 (14.2)
10-19	11 (25.5)	9 (31.0)	2 (14.2)
20-29	12 (27.9)	7 (24.1)	5 (35.7)
30-39	6 (13.9)	2 (6.9)	4 (28.0)
>= 40	3 (6.9)	2 (6.9)	1 (7.1)
Unknown	1 (2.3)	1 (3.4)	0 (0.0)

n(%) or median(25 th , 75 th percentiles)	All (n=44)	Clinical Pathway for MRI	
		No (n=29)	Yes (n=15)
Estimated no. pediatric out-of-hospital cardiac arrests per year	n=43	n=28	n=15
< 10	11 (25.5)	8 (28.5)	3 (20.0)
10-19	13 (30.2)	9 (32.1)	4 (26.6)
20-29	13 (30.2)	7 (25.0)	6 (40.0)
30-39	1 (2.3)	1 (3.5)	0 (0.0)
>= 40	4 (9.3)	2 (7.1)	2 (13.3)
Unknown	1 (2.3)	1 (3.5)	0 (0.0)

Neuroimaging resources

The majority (75% [33/44]) of respondents reported that their institution employs a neuroradiologist to interpret brain MRIs (Table 2). Of centers with neuroradiology expertise, 55% (18/33) report these subspecialists are available to interpret MRIs 24/7. A higher percentage of institutions with a clinical pathway for post-CA MRI reported having 24/7 Neuro-radiology availability (57% vs. 34%). Eighty-one percent (35/43) of responding centers have a 3 Tesla MRI available, though 19% (8/43) are limited to 1.5 Tesla MRI.

Table 2. Neuroimaging resources.

n(%) or median(25 th , 75 th percentiles)	All (N=44)	Clinical Pathway for MRI	
		No (N=29)	Yes (N=15)
Neuroradiologist for MRI interpretation available	n=44	n=29	n=15
Yes	33 (75.0)	23 (79.3)	10 (66.7)
Neuroradiologist available 24/7	n=43	n=29	n=14
Yes	18 (41.9)	10 (34.5)	8 (57.1)
MRI scanner within the ICU	n=44	n=29	n=15
Yes	1 (2.3)	0 (0.0)	1 (6.7)
MRI Scanner brand	n=43	n=28	n=15

Siemens only	16 (37.2)	11 (39.3)	5 (33.3)
Philips only	6 (14.0)	3 (10.7)	3 (20.0)
GE only	5 (11.6)	4 (14.3)	1 (6.7)
More than one brand among Siemens, Philips, GE, and Other	16 (37.2)	10 (35.7)	6 (40.0)
Platform			
MRI field strength	n=43	n=28	n=15
1.5 Tesla only	8 (18.6)	7 (25.0)	1 (6.7)
3.0 Tesla only	3 (7.0)	2 (7.1)	1 (6.7)
More than one field strength among 1.5T, 3.0T, and Other	32 (74.4)	19 (67.9)	13 (86.7)

Practices regarding MRI use in cardiac arrest

Of the participating institutions, 45% (20/44) have a dedicated Pediatric Neurocritical Care team/service (Table 3). Institutions with a pediatric neurocritical care team/service were almost twice as likely to have a clinical pathway for MRI after CA (66.7% vs. 34.5%). Fifty percent of institutions (22/44) obtain an MRI in nearly all pediatric CA patients and 31.8% (14/44) of institutions reported performing MRI in patients who are not back to their baseline. Centers with a clinical pathway for MRI use after CA were less likely to obtain an MRI on a “case by case” basis (6.7% [1/15]) compared to centers without a clinical pathway (24.1% [7/29]).

The majority of institutions (63.6% [28/44]) perform an MRI after 72 hours of return of spontaneous circulation (ROSC) but before transfer to the inpatient ward, 27.3% of institutions (12/44) obtain an MRI within 72 hours of ROSC and (9% (4/44) report the timing of MRI for prognostication after CA as “other.” All centers with a clinical pathway reported obtaining an MRI either within the first 72 hours after ROSC (40% [6/15]) or greater than 72 hours after ROSC and before transfer to the inpatient ward (60% [9/15]). None of the centers indicating the timing of MRI as “other” reported having a clinical pathway. When asked to indicate what factors are used to determine timing of brain MRI for prognostication, 90% of centers (40/44) reported poor neurologic exam as the main factor determining the timing of the MRI.

Other clinical factors determining timing of imaging for prognostication included intubation status (70% [31/44]) and favorable neurologic exam (61% [27/44]). All centers with a clinical pathway for MRI after CA indicated at least one clinical factor is used to determine timing of imaging for prognostication. Only one center reported using no clinical factors to determine timing of MRI.

Table 3. Clinical practices regarding the use of MRI in pediatric CA.

n(%) or median(25 th , 75 th percentiles)	All (N=44)	Pathway	
		No (N=29)	Yes (N=15)
Pediatric neurocritical care team/service	n=44	n=29	n=15
Yes	20 (45.5)	10 (34.5)	10 (66.7)
Typical practice for MRI prognostication after CA	n=44	n=29	n=15
Brain MRI is performed for nearly all patients (with some exception such as DNR)	22 (50.0)	15 (51.7)	7 (46.7)
Brain MRI is performed for patients who are not back to their baseline neurological status	14 (31.8)	7 (24.1)	7 (46.7)
Brain MRI is performed for patients only on a "case by case" basis	8 (18.2)	7 (24.1)	1 (6.7)
Typical timing of MRI for prognostication after CA	n=44	n=29	n=15
Within the first 72 hours after ROSC (acute)	12 (27.3)	6 (20.7)	6 (40.0)
After 72 hours from ROSC, but prior to transfer to ward (subacute)	28 (63.6)	19 (65.5)	9 (60.0)
Prior to hospital discharge or transfer to inpatient rehabilitation	0 (0)	-	-
Other	4 (9.1)	4 (13.8)	0 (0.0)
Factors determining timing of MRI			
Intubation status: MRI timing is adjusted to accommodate/prioritize extubation	n=44	n=29	n=15
Yes	31 (70.5)	20 (69.0)	11 (73.3)
Poor neurologic exam: The team decides to obtain MRIs earlier	n=44	n=29	n=15
Yes	40 (90.9)	25 (86.2)	15 (100.0)
Favorable neurologic exam: The team waits longer to obtain MRI	n=44	n=29	n=15
Yes	27 (61.4)	16 (55.2)	11 (73.3)

Types of MRI sequences used in children with cardiac arrest

Sites were asked to indicate which sequences were included in the imaging protocol used to evaluate children with CA. The percentage of centers using each specific MRI sequence, stratified by centers with and without a pathway, is illustrated in Figure 1. Over 97% (43/44) of institutions use T1, T2, and fluid-attenuated inversion recovery (FLAIR). All centers reported obtaining diffusion-weighted imaging (DWI). Sequences sensitive to hemorrhage such as gradient-echo (GRE) and susceptibility weighted imaging (SWI) are used in 43% and 67% of centers, respectively. Less than half of the institutions (40% and 31% of institutions with and without a pathway, respectively) report use of spectroscopy (MRS). Other techniques, such as functional MRI, MR venogram, and post-contrast T1 are used by less than 30% of institutions, regardless of the presence or absence of a pathway.

Figure 1. Type of MRI sequences used in pediatric cardiac arrest

Figure 1 Definitions: DWI-Diffusion weighted imaging; ADC-Apparent diffusion coefficient; FLAIR-Fluid-attenuated inversion recovery; GRE-Gradient Echo; SWI-Susceptibility weighted imaging; CBFP-Cerebral blood flow/perfusion; MRS-Magnetic resonance spectroscopy; DTI-Diffusion tensor imaging; MRA-Magnetic resonance angiogram; MRV-Magnetic resonance venogram; fMRI-Resting state functional MRI.

DISCUSSION

Our study describes current practices in use of brain MRI as a neuroprognostic tool in children with CA from 44 national and international pediatric tertiary care centers. The results suggest that clinicians often perform MRI as part of their routine care in children with CA, with half of responding institutions indicating that they obtain MRI in nearly all patients post-CA, and an additional 30% indicating that they obtain an MRI in those who are not back to their neurological baseline. Nearly all institutions routinely incorporate conventional sequences (i.e., T1, T2, DWI and FLAIR). Reported use of sequences sensitive to hemorrhage are 43% (GRE) and 67% (SWI). These rates appear low, but it is likely that centers choose to utilize a single sequence to evaluate for hemorrhage, resulting in little overlap in these values, thus indicating that this practice is also widely used at surveyed centers. However, there is still significant heterogeneity and less use of advanced MRI sequences post CA. We found that only a minority of the protocols incorporate advanced imaging sequences such as MRS. There are several potential reasons or explanations that may contribute to these observations. First, although promising, data regarding the clinical value of sequences such as MRS is limited to single-center, retrospective studies. Second, the interpretation of these images requires particular expertise. In our study, 25% of centers do not have a member of the radiology team with specific training in neuroradiology.

In our study, only 34% of centers indicated that they have an established clinical pathway for the use of MRI in pediatric CA. Institutions with CA pathways informing MRI utilization reported less variation in patient selection or timing of MRIs. The lack of pathway-driven practices is likely due to a number of factors including: sparse and low quality evidence in pediatrics and practice guidelines

developed by data extrapolated from adult patients or by expert opinion.³¹ Additionally, an increasing number of institutions have incorporated pediatric neurocritical care teams to build institutional expertise and provide standardized, evidence-based care for children with acute brain injury.^{32, 33} While less than half of responding institutions have a Pediatric Neurocritical Care team available, the presence of a Pediatric Neurocritical Care team doubled the likelihood of pathway driven care.

Taken together, these observations further highlight the need for evidence to guide clinical practice in the area of pediatric CA. Recent guidelines published by the American Heart Association (AHA) cite insufficient evidence to support the use of MRI as a standard of care in pediatric CA and state that "MRI during the first seven days post-arrest may be valuable" in assisting with prognostication.¹² Beyond this, no recommendations are made regarding indications, optimal timing, or acquisition protocol. The guidelines cite retrospective study design, small sample size, selection bias, lack of blinding in imaging interpretation, variable outcome measures, and follow-up times as limitations to the current evidence.

Conventional MRI protocols designed for clinical use include sequences such as T1 and T2-weighted images, FLAIR and diffusion-weighted imaging.²⁵ Prior studies have investigated the utility of these sequences in the assessment of children with CA. In a single-center study of 28 children who received a brain MRI within the first 14 days post-ROSC, T2 signal intensity abnormalities in the basal ganglia and diffusion restriction in the cerebral lobes were associated with unfavorable outcomes.¹⁵ A separate study evaluated the correlation between imaging abnormalities and unfavorable outcomes in 23 children with CA who underwent MRI within 14 days after ROSC.¹⁰ The injured areas most commonly identified by T2-weighted imaging were the lentiform nucleus (68%), the caudate (55%), and the thalamus (50%). T2 abnormalities in the lentiform nuclei were associated with unfavorable outcomes at hospital discharge. Diffusion restriction was most commonly seen in the lentiform nucleus (41%), thalamus (32%), frontal (30%), and parietal (30%) lobes. More recently, a single center study of 77

pediatric patients after OHCA who had an MRI at a median of day 4 post-CA found that the extent of diffusion restriction and T2/Flair abnormalities were associated with unfavorable outcomes with good discrimination (AUROC of 0.96 (95% CI 0.91, 0.99) for diffusion and 0.92 (0.85, 0.97) for T2/FLAIR injury).¹⁷ In addition to T2-weighted and DWI, more advanced imaging modalities such as MRS may have prognostic value in this population. Increased lactate concentrations in the parietooccipital gray matter and decreased NAA in the parietal white matter using MRS have been observed in children with CA and unfavorable outcomes.¹⁰ Decreased NAA and increased lactate were also observed in a study of children with acute brain injury (including CA) and unfavorable outcomes.²⁰

Given the evolution of the appearance of brain lesions on MRI, both the composition of the MRI protocol and the timing of the initial MRI after presentation are critical to capture the extent of brain injury accurately. MRI sequences sensitive to ischemia and cytotoxic edema such as DWI/apparent diffusion coefficient (ADC) often identify lesions in children with CA. Unlike abnormalities on T1 and T2 MRI sequences, diffusion abnormalities can be seen within minutes in acute ischemia (i.e. arterial ischemic stroke). However, diffusion abnormalities can also be delayed in CA possibly due to the variability of hemodynamic compromise over the course of the resuscitation and post resuscitation period.²⁶⁻²⁹ Thus, to ensure appropriate evaluation of the presence or absence of hypoxic ischemic brain injury after CA, the timing of MRI is very important. The AHA guidelines recommend considering use of MRI "within the first seven days" after injury."¹² In our survey, most centers stated that they perform MRIs after 72 hours post-ROSC. Ninety percent stated that poor neurologic exam is a main determinant for obtaining an MRI early and 61% stated that a favorable neurologic exam would lead to clinical teams waiting longer to obtain an MRI.

Our survey has a number of limitations that should be considered when interpreting these results. First, our survey response rate was relatively low at 30%. Second, a limitation common to survey studies with a single respondent per center is that the responses are based on the individual's

understanding of current institutional practices and may represent her/his opinion and practice over fact. Third, the surveyed institutions represent tertiary care facilities with an interest in clinical research in pediatric cardiac arrest which may not be generalizable to all pediatric intensive care settings. Imaging resources and practices may differ from institutions not participating in large research consortia, limiting the external validity of our findings. Of note, the cohort was not limited to large centers and included institutions with a broad spectrum of size and number of CA cases per year. Lastly, the correlation between the indications for MRI, timing, and acquisition protocols and patient outcomes was beyond the scope of this survey. Prospective, adequately powered, multi-center studies with standardized, patient-oriented outcomes are needed to bridge this knowledge gap.

Our survey reveals prevalent use of MRI for neuroprognostication in pediatric CA. Responses suggest a wide variability in the timing, MRI sequences, and neuroradiology resources, as well as a dearth of centers with clinical pathways to guide MRI utilization after pediatric CA. Studies to ultimately determine the optimal use of MRI in pediatric CA will need to account for and implement strategies reduce this variation.

Acknowledgements. We acknowledge members of the **POCCA Steering Committee**: Patrick Kochanek, MD, MCCM; Robert Clark, MD; Hulya Bayir, MD; Ashok Panigrahy, MD; Rachel Berger, MD, MPH; Sue Beers, PhD; Tony Fabio, PhD, MPH; **POCCA Investigators**: Karen Walson, MD (Children's Healthcare of Atlanta); Alexis Topjian, MD, MSCE (Children's Hospital of Philadelphia); Christopher JL Newth, MD, FRCPC (Children's Hospital of Los Angeles); Elizabeth Hunt, MD, MPH, PhD, Jordan Duval-Arnould, DrPH, MPH (Johns Hopkins Children's Center); Binod Balakrishnan, MD, Michael T. Meyer, MD, MS, FCCM (Children's Hospital of Wisconsin); Melissa G. Chung, MD (Nationwide Children's Hospital); Anthony Willyerd, MD (Phoenix Children's Hospital); Lincoln Smith, MD, Jesse Wenger, MD (Seattle Children's); Stuart Friess, MD, Jose Pineda, MD, MS (St. Louis Children's); Ashley Siems, MD, Jason Patregnani, MD, John Diddle, MD (Children's National Hospital); Aline Maddux, MD, MSCS and Craig Press, MD, PhD (Children's Hospital of Colorado); Lesley Doughty, MD (Cincinnati Children's Hospital Medical Center); Juan Piantino, MD (Doernbecher Children's Hospital); **and the POCCA Coordinators**: David Maloney, BS, Pamela Rubin, RN (UPMC Children's Hospital of Pittsburgh); Beena Desai, BS, CCRC, Maureen G. Richardson, BSN, RN, CPN, Cynthia Bates, CCRP (Children's Healthcare of Atlanta); Darshana Parikh, Janice Prodel, Maddie Winters, Katherine Smith, MPH, BSN, RN, CPN (Children's Hospital of Philadelphia); Jeni Kwok, JD, Adriana Cabrales, BA (Children's Hospital of Los Angeles); Ronke Adewale, Pam Melvin (Johns Hopkins Children's Center); Sadaf Shad, Katherine Siegel, Katherine Murkowski, Mary Kasch (Children's Hospital of Wisconsin); Josey Hensley RN, BSN, Lisa Steele, RN, BSN (Nationwide Children's Hospital); Danielle Brown, Brian Burrows, Lauren Hlivka (Phoenix Children's Hospital); Deana Rich (Seattle Children's Hospital); Amila Tutundzic, Tina Day, Lori Barganier (St. Louis Children's Hospital); Ashley Wolfe, Mackenzie Little, Elyse Tomanio, Neha Patel, Diane Hession (Children's National Hospital); Yamila Sierra MPH, CCRP (Children's Hospital of Colorado); and Rhonda Jones, Laura Benken (Cincinnati Children's).

We also thank Jonathan Elmer, MD, MS; Subramanian Subramanian, MD; Srikala Narayanan, MD; Julia Wallace; Tami Robinson; Andrew Frank; Stefan Bluml, PhD; Jessica Wisnowski, PhD; Keri Feldman; Avinash Vemulapalli; Linda Ryan; Scott Szypulski, MBA; Christopher Keys, and all of the children and families who generously participated in POCCA. We are grateful to the all ICU staff, nurses, and physicians for their efforts in study recruitment and provision of excellent clinical care to families.

Bibliography

- 1 Atkins D. L., Everson-Stewart S, Sears GK, Daya M, Osmond MH, Warden CR, *et al.* Epidemiology and outcomes from out-of-hospital cardiac arrest in children: The resuscitation outcomes consortium epistry-cardiac arrest. *Circulation* **119**, 1484-1491, doi:10.1161/CIRCULATIONAHA.108.802678 (2009).
- 2 Topjian AA & Berg RA. Pediatric out-of-hospital cardiac arrest. *Circulation* **125**, 2374-2378, doi:10.1161/CIRCULATIONAHA.111.071472 (2012).
- 3 Berg RA, Sutton RM, Holubkov R, Nicholson CE, Dean JM, Harrison R, *et al.* Ratio of PICU versus ward cardiopulmonary resuscitation events is increasing. *Crit Care Med* **41**, 2292-2297, doi:10.1097/CCM.0b013e31828cf0c0 (2013).
- 4 Knudson JD, Neish SR, Cabrera AG, Lowry AW, Shamazad P, Morales DLS, *et al.* Prevalence and outcomes of pediatric in-hospital cardiopulmonary resuscitation in the United States: an analysis of the Kids' Inpatient Database*. *Crit Care Med* **40**, 2940-2944, doi:10.1097/CCM.0b013e31825feb3f (2012).
- 5 Girotra S, Spertus JA, Li Y, Berg RA, Nadkarni VM, Chan PS, *et al.* Survival trends in pediatric in-hospital cardiac arrests: an analysis from Get With the Guidelines-Resuscitation. *Circulation. Cardiovascular quality and outcomes* **6**, 42-49, doi:10.1161/circoutcomes.112.967968 (2013).
- 6 Meert KL, Telford R, Holubkov R, Slomine BS, Christensen JR, Dean JM, *et al.* Pediatric Out-of-Hospital Cardiac Arrest Characteristics and Their Association with survival and neurobehavioral outcome. *Pediatric Critical Care Medicine*. **17**:e543-e550, doi:10.1097/PCC.0000000000000969 (2016).

- 7 Donoghue AJ, Nadkarni V, Berg RA, Osmond MH, Wells G, Nesbitt L, et al. Out-of-Hospital Cardiac Arrest: An Epidemiologic Review and Assessment of Current Knowledge. *Annals of Emergency Medicine*. **46**:512-522, doi: 10.1016/j.annemergmed.2005.05.028 (2005).

- 8 Fink EL, Berger RP, Clark RSB, Watson RS, Angus DC, Panigrahy A, et al. Exploratory study of serum ubiquitin carboxyl-terminal esterase L1 and glial fibrillary acidic protein for outcome prognostication after pediatric cardiac arrest. *Resuscitation* **101**, 65-70, doi:10.1016/j.resuscitation.2016.01.024 (2016).

- 9 Fink EL, Berger RP, Clark RSB, Watson RS, Angus DC, Richichi R, et al. Serum biomarkers of brain injury to classify outcome after pediatric cardiac arrest*. *Crit Care Med* **42**, 664-674, doi:10.1097/01.ccm.0000435668.53188.80 (2014).

- 10 Fink EL, Wisnowski J, Clark R, Berger RP, Fabio A, Furtado A, et al. Brain MR imaging and spectroscopy for outcome prognostication after pediatric cardiac arrest. *Resuscitation* **157**, 185-194, doi:10.1016/j.resuscitation.2020.06.033 (2020).

- 11 Abend NS, Topjian AA, Kessler SK, Gutierrez-Colina AM, Berg RA, Nadkarni V, et al. Outcome prediction by motor and pupillary responses in children treated with therapeutic hypothermia after cardiac arrest. *Pediatric critical care medicine : a journal of the Society of Critical Care Medicine and the World Federation of Pediatric Intensive and Critical Care Societies* **13**, 32-38, doi:10.1097/PCC.0b013e3182196a7b (2012).

- 12 Topjian AA, de Caen A, Wainwright MS, Abella BS, Abend NS, Atkins DL, et al. Pediatric Post-Cardiac Arrest Care: A Scientific Statement From the American Heart Association. *Circulation* **140**, e194-e233, doi:10.1161/cir.0000000000000697 (2019).

- 13 Högler S, Sterz F, Sipos W, Schratte A, Weihs W, Holzer M, *et al.* Distribution of neuropathological lesions in pig brains after different durations of cardiac arrest. *Resuscitation* **81**, 1577-1583, doi:10.1016/j.resuscitation.2010.07.005 (2010).
- 14 Back, T., Hemmen, T. & Schöler, O. G. Lesion evolution in cerebral ischemia. *J Neurol* **251**, 388-397, doi:10.1007/s00415-004-0399-y (2004).
- 15 Fink EL, Panigrahy A, Clark RSB, Fitz CR, Landsittel D, Kochanek PM & Zuccoli. Regional brain injury on conventional and diffusion weighted MRI is associated with outcome after pediatric cardiac arrest. *Neurocrit Care* **19**, 31-40, doi:10.1007/s12028-012-9706-0 (2013).
- 16 Christophe C, Fonteyne C, Ziereisen F, Christiaens F, Deltenre P, De Maertelaer V & Dan B. Value of MR imaging of the brain in children with hypoxic coma. *AJNR Am J Neuroradiol* **23**, 716-723 (2002).
- 17 Kirschen MP, Licht DJ, Faerber J, Antara M, Graham K, Winters M, *et al.* Association of MRI Brain Injury With Outcome After Pediatric Out-of-Hospital Cardiac Arrest. *Neurology* **96**, e719-e731, doi:10.1212/WNL.00000000000011217 (2021).
- 18 Velly L, Perlberg V, Boulrier T, Adam N, Delphine S, Luyt CE, *et al.* Use of brain diffusion tensor imaging for the prediction of long-term neurological outcomes in patients after cardiac arrest: a multicentre, international, prospective, observational, cohort study. *Lancet Neurol* **17**, 317-326, doi:10.1016/S1474-4422(18)30027-9 (2018).
- 19 Pugin D, Hofmeister J, Gasche Y, Vulliemoz S, Lovblad KO, Ven De Ville D, & Haller S. Resting-State Brain Activity for Early Prediction Outcome in Postanoxic Patients in a Coma with Indeterminate Clinical Prognosis. *AJNR Am J Neuroradiol* **41**, 1022-1030, doi:10.3174/ajnr.A6572 (2020).
- 20 Ashwal S, Holshouser BA, Tomasi LG, Shu S, Perkin RM, Nystrom GA, & Hinshaw DB. 1H-magnetic resonance spectroscopy-determined cerebral lactate and poor neurological outcomes

- in children with central nervous system disease. *Annals of neurology* **41**, 470-481, doi:10.1002/ana.410410410 (1997).
- 21 Barkovich AJ, Baranski K, Vigneron D, Partridge JC, Hallam DK, Hajnal BL, & Ferriero DM. Proton MR spectroscopy for the evaluation of brain injury in asphyxiated, term neonates. *AJNR Am J Neuroradiol* **20**, 1399-1405 (1999).
- 22 Fink EL, Clark RSB, Panigrahy A, Berger R, Wisnowski J, Bluml S, *et al.* Personalising Outcomes after Child Cardiac Arrest (POCCA): design and recruitment challenges of a multicentre, observational study. *BMJ Open* **10**, e039323, doi:10.1136/bmjopen-2020-039323 (2020).
- 23 Larsen GY, Schober M, Fabio A, Wisniewski SR, Grant MJC, Shari N, *et al.* Structure, Process, and Culture Differences of Pediatric Trauma Centers Participating in an International Comparative Effectiveness Study of Children with Severe Traumatic Brain Injury. *Neurocrit Care* **24**, 353-360, doi:10.1007/s12028-015-0218-6 (2016).
- 24 Fink EL, Kochanek PM, Tasker RC, Beca J, Bell MJ, Clark RSB, *et al.* International Survey of Critically Ill Children With Acute Neurologic Insults: The Prevalence of Acute Critical Neurological Disease in Children: A Global Epidemiological Assessment Study. *Pediatr Crit Care Med* **18**, 330-342, doi:10.1097/pcc.0000000000001093 (2017).
- 25 Arbelaez A, Castillo M, & Mukherji SK. Diffusion-weighted MR imaging of global cerebral anoxia. *AJNR Am J Neuroradiol* **20**, 999-1007 (1999).
- 26 Beauchamp NJ, Jr & Bryan RN. Acute cerebral ischemic infarction: a pathophysiologic review and radiologic perspective. *AJR. American journal of roentgenology* **171**, 73-84, doi:10.2214/ajr.171.1.9648768 (1998).
- 27 Chien D, Kwong KK, Gress DR, Buonanno FS, Buxton RB, & Rosen BR. MR diffusion imaging of cerebral infarction in humans. *AJNR Am J Neuroradiol* **13**, 1097-1102; discussion 1103-1095 (1992).

- 28 Gray L & MacFall J. Overview of diffusion imaging. *Magnetic resonance imaging clinics of North America* **6**, 125-138 (1998).
- 29 Reith W, Hasegawa Y, Latour LL, Dardzinski BJ, Sotak CH, & Fisher M. Multislice diffusion mapping for 3-D evolution of cerebral ischemia in a rat stroke model. *Neurology*. **45(1)**:172–177 (1995).
- 30 Andrews PJ, Piper IR, Dearden NM & Miller JD. Secondary insults during intrahospital transport of head-injured patients. *Lancet* **335**, 327-330, doi:10.1016/0140-6736(90)90614-b (1990).
- 31 Cronin M & Wainwright MS. Underpowered and Too Heterogenous: A Humbling Assessment of the Literature Supporting Neuroprognostication After Pediatric Cardiac Arrest. *Pediatr Crit Care Med* **21**, 915-916, doi:10.1097/pcc.0000000000002546 (2020).
- 32 Wainwright MS, Grimason M, Goldstein J, Smith CM, Amlie-Lefond C, Revivo G, *et al.* Building a pediatric neurocritical care program: a multidisciplinary approach to clinical practice and education from the intensive care unit to the outpatient clinic. *Semin Pediatr Neurol* **21**, 248-254, doi:10.1016/j.spen.2014.10.006 (2014).
- 33 Wainwright MS, Hansen G, & Piantino J. Pediatric neurocritical care in the 21st century: from empiricism to evidence. *Curr Opin Crit Care* **22**, 106-112, doi:10.1097/MCC.0000000000000288 (2016).