Metagenomic analysis of antimicrobial resistance genes in wastewaters in Benin and Burkina Faso indicates a serious health risk from untreated hospital wastewaters in low-income countries

Melina A. Markkanen^{a,b} #, Kaisa Haukka^{a,g}, Katariina M. M. Pärnänen^a, Victorien Tamegnon Dougnon^c, Isidore Juste O. Bonkougou^d, Zakaria Garba^e, Halidou Tinto^e, Anniina Sarekoski^a, Antti Karkman^{a,b}, Anu Kantele^{b,f,g} *, Marko P. J. Virta^{a,b} #*

^a Department of Microbiology, University of Helsinki, Finland

^b Multidisciplinary Center of Excellence in Antimicrobial Resistance Research, Finland

^c Research Unit in Applied Microbiology and Pharmacology of Natural Substances, Polytechnic School of Abomey-Calavi, University of Abomey-Calavi, Benin

^d Department of Biochemistry and Microbiology, University Joseph KI-ZERBO, Ouagadougou, Burkina Faso

^e Clinical Research Unit Nanoro, Institute for Research in Health Sciences, National Center for Scientific and Technological Research, Burkina Faso

^f Meilahti Infectious Diseases and Vaccine Research Center, MeiVac, Department of Infectious Diseases, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

^g Human Microbiome Research Program, Faculty of Medicine, University of Helsinki, Helsinki, Finland

Corresponding authors:

Marko P. J. Virta, marko.virta@helsinki.fi

Melina A. Markkanen, melina.markkanen@helsinki.fi

* Marko P. J. Virta and Anu Kantele contributed equally to this work.

Running title

HWW resistomes in Benin, Burkina Faso, and Finland

Abstract

Antimicrobial resistance (AMR) is one of the greatest global threats to human health, but substantial gaps in AMR data exist in West African countries. We explored the presence of antimicrobial resistance genes (ARGs) in the hospital wastewaters (HWW) of nine hospitals in Benin and Burkina Faso and, for comparison, four hospitals in Finland using shotgun metagenomic sequencing. The highest relative sum abundance of ARGs in HWWs was observed in Benin and the lowest in Finland. HWW resistomes resembled each other more in Benin and Burkina Faso than in Finland. Different carbapenemases were detected in varying abundances, especially in HWWs from Burkina Faso and Finland. The most widespread carbapenemase gene in the Beninese hospitals, *bla*GES, was also found in water used for handwashing. *bla*NDM was present in the HWW of one Beninese hospital and was also detected in the stools of a hospitalized patient. Mobile colistin resistance genes (*mcr*) were detected in the HWWs of all the three studied countries, with *mcr-5* as the most common variant. These and other *mcr* genes were observed in high abundance in treated wastewater released into rivers in Burkina Faso. In Benin, only little is done

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

to treat wastewaters, including HWWs. The results provide evidence for public health decision-makers for the dire need to increase wastewater treatment capacity, with particular attention to HWWs.

Keywords

antimicrobial resistance (AMR)

West Africa

hospital wastewater (HWW)

carbapenemase

colistin resistance

metagenomes

1. Introduction

The AMR crisis is a global issue with multifaceted effects on human, animal, and environmental health, and it comes with substantial economic losses (1). Resistance to almost all antibiotics, including the last resort regimen reserved to treat the most severe infections, has tended to emerge only a couple of years after introducing a new antibiotic to the global market (2, 3). Due to limitations in diagnostic testing in low-resource settings in low-income countries (LICs), broad-spectrum antibiotics are often prescribed empirically without microbiological verification of the causative pathogen or its sensitivity to different antibiotics (4). In addition, unregulated access to antimicrobials allows their wide use in these countries (5–7).

Bacterial resistance to antimicrobial compounds may be intrinsic or acquired from other bacteria or the surrounding environment via horizontal gene transfer of mobile genetic elements (MGE) (8–10). Acquired, potentially mobile ARGs are considered to have greater clinical relevance and impact the current AMR problem induced by human activity (10, 11). Class 1 integrons are strongly linked to the dissemination of clinically relevant acquired ARGs (12, 13). Although integrons are not mobile themselves, integron-carried multidrug-resistance gene cassettes can be transmitted to new hosts, for example, via plasmids (14). The gene encoding the integron integrase (*intI*), among other associated genes, is involved in the recombination of incoming genetic material into the gene cassette, and *intII* is typically used as a marker for class 1 integrons (13, 15). Over time, quaternary ammonium compound (QAC) resistance genes, such as *qacEΔ*, have established their position in the structure of a typical class 1 integron gene cassette whose spread is induced by human activity (15, 16). It has been speculated that increasing the use of such disinfectant compounds in hospitals and community environments, especially during the COVID-19 pandemic, would further exacerbate the problem of multidrug resistance mediated by class 1 integrons (17).

The high-risk ESKAPEc bacteria (9) (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Enterobacter* spp., and *Escherichia coli*) (18) are top priority pathogens for AMR research and the development of new antibiotics (9, 19). They can feature an impressive resistance profile in terms of both intrinsic and acquired ARGs and often exhibit multidrug-resistant phenotypes. However, investigating only bacterial isolates may result in a limited understanding of ARGs

occurring in a particular hospital. In contrast, shotgun metagenomic sequencing enables the analysis of resistomes and mobilomes (all ARGs or MGEs, respectively) as well as the microbial communities present in the studied environment (20). Hence, as opposed to PCR-based methods, there is no requirement for a priori knowledge of the resistome (20).

As a consequence of the increased prevalence of extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE), the use of broad-spectrum antibiotics has increased in clinical practice (2, 21, 22). Tracking down the spread of ESBL genes is, in many situations, no longer feasible or relevant, and attention has been directed to plasmid-mediated genes that encode carbapenemases. The carbapenemases *bla*GES, *bla*IMP, *bla*KPC, *bla*NDM, *bla*OXA-48, *bla*OXA-58, and *bla*VIM carried by plasmids have emerged and spread around the world during the past three decades (23–26).

Metronidazole is an example of an empirically widely used first-line treatment in many African countries (27). This antimicrobial drug is effective against various protozoan parasites and anaerobic bacteria (28). Although acquired resistance towards metronidazole (mediated mainly by *nim* genes) is considered to be relatively rare, an increasing trend in its emergence has been reported (28). Colistin is a last-resort antibiotic used for treating infections caused by multidrug and extensively resistant bacteria (29). The rapid emergence of mobile colistin resistance mediated by *mcr* genes threatens the efficacy of colistin in clinical use (30, 31).

Although AMR is of global concern, the crisis affects most dramatically LICs, such as those in West Africa (3, 4, 32–34). Lack of research data is a major factor hindering the development of solutions to tackle the AMR problem in these countries (4, 32, 35, 36). Despite gaps in resistance surveillance data, it is well known that the level of AMR is highest in many LICs, such as in many African countries (21, 33, 37, 38). By contrast, in Northern European countries, such as Finland, AMR occurrence is among the lowest globally, both in the community (38, 39) and healthcare settings (40).

As hospital wastewater (HWW) from healthcare facilities is considered to be at the frontline of AMR emergence and spread (41), we set out to investigate the AMR situation and the characteristics of the resistomes in nine hospitals in two West African countries, Benin and Burkina Faso, where prior data was scarce (21). For comparison, we analyzed samples from four hospitals in Finland, where the level of AMR was expected to be low (40). In this study, we applied shotgun metagenomic sequencing as it allows the analysis of the total catalog of genes present in the studied environment.

2. Results

2.1. Features of hospital wastewaters in Benin, Burkina Faso, and Finland

We analyzed hospital wastewater-associated resistomes from Benin, Burkina Faso, and Finland. On average, 31 million sequence reads per sample were analyzed. HWWs from Benin showed the highest, and from Burkina Faso the second-highest abundance of all detected ARGs normalized to bacterial 16S rRNA genes, here referred to as

the relative sum abundance of ARGs. The lowest relative sum abundance was detected in HWWs from Finland (Figure 1A, Table S3A). No significant differences were observed in the relative ARG sums among the different hospitals in each country or between the hospital sections or wards within the sampled hospitals (Table S3A). Country-wise differences for the relative abundances of class 1 integron gene (*intI1*) (Figure 1C, Table S3C) followed the trend observed for relative ARG abundances (Figure 1A). The observed high Pearson correlation of 0.80 ($p = 3.2 \times 10^{-16}$) between ARGs and *intI1* genes across all 67 HWW samples indicated an association between the genes. For all MGEs, including also other than class 1 integron associated genes, a similar trend was not observed (Figure 1B, Table S3B).

The resistomes of HWWs from Benin and Burkina Faso resembled each other more than those of the Finnish HWWs (Figure 2A). The resistomes in Benin and Burkina Faso shared 384 ARGs, while Finland and Benin shared only 15 ARGs, and Finland and Burkina Faso shared 134 ARGs (Figure S4). The resistome compositions differed the most between the Beninese and Finnish HWW samples (Figure 2A). The country explained the differences in the resistomes by 29.5–33.7 % in the Benin-Finland comparison and 24.5–25.6 % in the Burkina Faso-Finland comparison (Table S1). In addition, while the ARGs found uniquely in HWWs from Burkina Faso made up approximately 14 % of all the detected ARGs in this study, the corresponding percentage for HWWs from Benin was only 2.5 % (Figure S4).

As the resistomes of West African countries and Finland were divergent, the drivers for the differences were subsequently investigated. The country explained less of the variance between taxonomical compositions than resistomes (Figure 2B, Table S1). A set of samples from a single hospital (Hospital I, Figure S3B) seemed to drive the distinct clustering seen in the microbial community in HWWs from Burkina Faso.

Figure 1. Relative sum abundance of ARGs, MGEs, and class 1 integrons. A generalized linear model (GLM) was fitted to 16S rRNA, and gene length normalized A) ARG B) MGE, and C) *intI1* count data to model the differences in the levels of these genes in the three countries. The vertical line shows the median in boxplots, while the lower and higher hinges denote the 25 % and 75 % quantiles, respectively. In addition, Wilcoxon signed-rank test significances are shown for pair-wise comparisons.

Figure 2. PCoA plots describe the ordinations based on A) resistomes, B) taxonomical compositions, and C) mobilomes in HWWs from Benin, Burkina Faso, and Finland. All 67 HWW samples were included in the plots. However, due to unequal group sizes, groups of random sample sets were applied for testing the significance of the clustering based on the country using adonis. The tests were performed in triplicate with different samples to account for variation caused by sample selection. The R^2 values are shown in Supporting Information, Table S1. Confidence ellipses represent 90 % confidence levels.

2.2. Clinically relevant taxa and associated ARGs

Taxonomical composition present in septic tanks and sumps in Benin and Burkina Faso differed from those found in hospital sewers in Finland. In Benin and Burkina Faso, the 15 most abundant genera present in HWW included *Desulfobulbus*, *Sulfurospirillum*, *Thauera*, and *Geobacter*, while in Finnish HWW, *Bifidobacterium* and *Faecalibacterium* were among the most abundant genera (Table S4). Genera known for opportunistically pathogenic species such as *Escherichia* and *Aeromonas* were detected among the 15 most abundant genera in the HWWs of all three countries (Table S4, Figure S5). *Klebsiella pneumoniae* was among the 15 most abundant species in Benin and Burkina Faso in terms of relative abundance (Table S4, Figure S5). In contrast, at the species level, *Escherichia coli*, *Enterococcus faecium*, and species of *Enterobacter cloacae* and *Pseudomonas aeruginosa* complexes were present but rare in all the countries. *Staphylococcus aureus* was detected exclusively in Burkina Faso samples (Figure S5). Other species in the genus *Staphylococcus* were more commonly detected in HWWs from Benin, and Burkina Faso compared to Finland (Tables S5A and S6A). However, knowing the limitations of species-level taxa identification using shotgun metagenomics (42), these results must be evaluated with some caution.

There was no significant difference (adjusted p -value < 0.05) in the abundance of the genus *Acinetobacter* between the HWWs from the three countries, but the abundances of different species of *Acinetobacter* varied significantly. *A. johnsonii* and carbapenemases in the *bla*OXA-211-like group (for example, *bla*OXA-211, -212, -280, and -281) that are intrinsically carried by this species were significantly more abundant in HWWs from Finland compared to either Benin or Burkina Faso (Figure 4, Figure S5 and Tables S5A, S6A, S7, and S8). Certain species of *Acinetobacter* that are not intrinsically carbapenem-resistant, such as *A. townerii*, were detected at significantly higher relative abundances in HWWs from Benin and Burkina Faso compared to Finland (Tables S5A, and S6A).

2.3. Carbapenem resistance

The presence of seven acquired carbapenemases (*bla*GES, *bla*IMP, *bla*KPC, *bla*NDM, *bla*OXA-48, *bla*OXA-58, and *bla*VIM) was analyzed in detail due to their ability to cause complex infections with minimal treatment options when in a relevant host. One of the Finnish (hospital K) samples possessed the highest relative abundance of these carbapenemases (Figure 3A). This hospital is the University Central Hospital of the capital of Finland. In HWWs from Benin, the carbapenemase genes in the *bla*GES family seemed to dominate over other carbapenemases in all four hospitals (Figure 3A). Instead, most of the other carbapenemases were present at similar prevalence levels in HWWs from Burkina Faso and Finland (Figure 3A), in line with the observation of the lower number of different ARGs in Benin versus Burkina Faso (Figure S4).

*bla*VIM, *bla*IMP, and *bla*NDM were detected in Burkinabe and Finnish HWW samples, but they were more abundant in HWWs from Burkina Faso (Figure 3A). Among the studied acquired oxacillinases, *bla*OXA-48 was detected solely in two samples from Burkina Faso (Figure 3A). Instead, *bla*OXA-58 was present in the majority of Burkinabe and Finnish samples (Figure 3A). Interestingly, most *bla*KPC genes were exclusively detected in HWWs from Finland (Figure 3A, Table S11).

This suggests a stronger class I integron association of ARGs in Beninese HWWs than in the other two countries. The detection of some of these ARGs in contigs carrying several other acquired ARGs (Figure S6) or class 1 integron-associated genes (Table S13) suggests that the genes are linked to class 1 integrons. For example, the plasmid-mediated ARGs *qnrVC4* and *qnrVC5* strongly correlated with *intI1* and *qacEΔ*. The genes were located on contigs with other ARGs such as aminoglycoside ARGs and either carbapenemase (Figure S6) or ESBL genes in the *blaGES* gene family (Figure S6B). While *qnrVC4* and *qnrVC5* were more abundant in HWWs from Benin and Burkina Faso than in those from Finland, quinolone ARGs unassociated with integrons, such as *qnrB* variants, were instead more common in HWWs from Finland (Figure 4, and Tables S7, and S8).

Oxacillinases constituted half of the top 20 most abundant ARGs in HWWs from Finland (Figure 4). Most of these oxacillinases (*blaOXA-211*, -212, -334, -373, -309, -280, and -281) belong to the group of *blaOXA-211*-like carbapenemases typically intrinsically carried by certain *Acinetobacter* species and showed no significant correlation with *intI1* or *qacEΔ* (Figure 4, Tables S12A and S12B). In contrast, certain genes of oxacillinases from other *blaOXA* gene families, which were significantly more abundant in HWWs from Benin or Burkina Faso, showed significant correlation with *intI1* and *qacEΔ* (Figure 4, Tables S12A and S12B) and were found in shared contigs with the *intI1* gene and other ARGs (Table S13, Figure S6C).

Figure 4. Differently abundant ARGs in HWWs from Benin and Burkina Faso and their putative associations with class 1 integrons. 20 significantly more abundant ARGs in HWWs of each country in pair-wise comparisons (Benin-Finland, Burkina Faso-Finland) are shown. Pearson correlations between each ARG and the genes *intI1* and *qacEA* associated with class 1 integrons (Supporting Information, Tables S12A, and S12B) are denoted with color-coding. The red color indicates a positive and blue negative correlation. Non-significant ($p > 0.05$) correlations are marked as blank (white) squares.

2.5. Mobile colistin resistance genes

Mobile colistin resistance genes (*mcr* genes) were observed in several HWW samples in Benin, Burkina Faso, and Finland (Table S16). Variants of *mcr-5* (*mcr-5.1* and *mcr-5.2*) were the most common as they were found in HWWs from all except one hospital (hospital F) in the three countries. We were able to confirm the location of *mcr-5* within a Tn3-like transposase element in the metagenome-assembled contigs (Figure S7, Table S14).

In addition to the HWW samples, high relative abundances of *mcr-5* genes were also detected in the immediate and more distant surroundings of the hospitals, such as in the street gutter water 100 m away from hospital B and in the soil between two HWW septic tanks in Benin (Table S15A). For hospital C, the relative abundance of *mcr-5* in a puddle surrounding the surgery room HWW septic tank was greater than the average relative sum abundance of *mcr-5* in the actual HWW septic tanks of that hospital (Figure S2D, Table S15A).

In Burkina Faso, very high relative sum abundances (2.8×10^{-2}) of *mcr-5* were observed in a municipality channel leading out from hospital I after passing through a biological treatment process (Table S15B). Furthermore, the presence of *mcr-5* was also detected in a nearby wetland where the processed wastewater is released (Table S15B). Similarly, in another Burkinabe city, *mcr-5* was found in a sample before wastewater treatment and the river receiving the water after treatment (Table S15B).

The next most prevalent *mcr* genes, namely *mcr-3*, *mcr-10*, and *mcr-7*, were detected in certain cases at very high relative abundances in the HWW samples (Table S16). The gene *mcr-3* was also present in the water used for handwashing at hospital C (Figure S2E, Table S15A). The gene *mcr-2* was not detected, and the lowest possible average relative sum abundance was detected for the gene *mcr-1* (Table S16).

2.6. Other than hospital wastewater samples

Other sample types within and outside the studied hospitals were investigated to obtain a more comprehensive understanding of ARG prevalence. Metronidazole resistance genes, such as *nimA* and *nimE*, were among the top 15 most abundant ARGs in the fecal samples of the four in-patients at a Beninese hospital B (Table S17). These ARGs were also significantly more abundant in HWWs from Benin, and Burkina Faso compared to Finland (Figure 4, Tables S7 and S8). *blaNDM* was detected in one of the four fecal samples (Figure 3B) and was also present in the HWWs of hospital B (Figure 3A). Other high abundance ARGs in the fecal samples included ESBL and broad-spectrum beta-lactamase genes of the *cfxA* family (Table S17).

As in the HWWs from Benin (Figure 3A), *bla*GES carbapenemases were also prevalent in other sample types from a hospital environment, such as in the water intended for handwashing (Figure S2E) for hospital C, the soil between HWW septic tanks in the yard of hospital B, and street gutter water 100 meters away from hospital B (Figure 3B). *bla*GES carbapenemase genes were also found in various waters used for drinking, including river waters in a remote countryside village (Figure 3B). However, the relative sum abundance of ARGs was low in the three rivers in central Benin (Figure 3B). We also detected *bla*NDM in well water used for drinking in Benin (Figure 3B). Albeit at low relative abundances, *bla*GES, *bla*VIM, and *bla*OXA-58-like carbapenemases were detected in natural waters receiving treated HWWs from Burkina Faso (Figure 3B). Significantly higher abundances of these ARGs were found in biologically treated wastewater before releasing into the river (Figure 3B).

3. Discussion

We characterized the bacterial community composition and resistome in 67 HWWs and 16 related samples from Benin and Burkina Faso and compared them to HWWs from Finland. This is the first study investigating hospital wastewaters in the two West African countries using a metagenomic approach. Due to the lack of systematic AMR surveillance in these countries, available data on bacterial resistance are patchy and heterogeneous (1, 4, 31, 34, 37). Thus, the magnitude of the resistance problem and the specific ARG reservoirs in many African countries are yet to be unraveled. The relative sum abundance of all ARGs detected in HWWs was the highest in Benin, yet the number of different ARGs detected there was the lowest. The differences in resistomes between the countries seemed to be only partially explained by variation in taxonomical composition, as the clustering of taxonomical composition was not as strongly correlated by country compared to clustering of resistomes. This suggests that despite differences in the HWW collection systems in the three countries (in Benin and Burkina Faso, HWW is relatively stagnant in septic tanks or sumps, while in Finland it flows in HWW drains), the microbial community compositions appear to be similar. However, of the most abundant taxa, we found many environmental bacteria in HWWs from Benin and Burkina Faso but not in Finland. This may be due to the closer contact of water with environmental surfaces in septic tanks and sumps compared to the isolated wastewater systems in Finland. Bacterial genera typically found in soil and aquatic environments (44) have also been previously described to dominate HWWs from Benin and Burkina Faso (45). In contrast, the 15 most abundant genera in Finnish HWWs included less environmental bacteria and more bacteria typical for human gut communities, such as *Bifidobacterium* (46) and *Faecalibacterium* (47). Another factor possibly explaining the differences in the dominant taxa present in the HWWs from Benin and Burkina Faso in comparison to Finland could arise from the fact that in Finland, the hospital toilet waters containing human fecal material are directed to HWWs, while for the majority of the West African HWWs studied here, this was not the case.

The ARGs that were more abundant in Benin's HWWs than in Finland appeared to be more commonly carried by class 1 integrons. They were strongly correlated with class 1 integron-associated genes (*intI1* and *qacEA* (15)) and, in certain cases, detected in shared contigs with these genes. Moreover, we could detect gene cassettes consisting of several different acquired ARGs linked to each other, likely referring to gene cassettes carried by

class 1 integrons (48). Furthermore, a higher abundance of ARGs intrinsically carried by specific species, such as *blaOXA-211* and *blaOXA-299* genes by *A. johnsonii* and *A. bouvetii* (49, 50), as well as *cphA* intrinsically carried by *Aeromonas* spp. (51) were distinctive features of the HWW resistomes from Finland. Intrinsic resistance is often chromosomally encoded and rarely mobile or related to horizontal gene transfer (52), whereas acquired resistance is typically carried by MGEs, such as class 1 integrons. As expected, no significant correlations between these intrinsic *blaOXA* genes and class 1 integron-associated genes were found, supporting the observation that ARGs putatively carried by class 1 integrons had a smaller impact on resistomes in HWWs from Finland compared to Benin. Based on these results, we speculate that the high sum abundance of ARGs seen in HWWs from Benin may, in part, be explained by the mobilome, more specifically, the prevalence of class 1 integrons. However, correlation does not necessarily indicate causation, and in this study, quantitative inferences cannot be made for metagenome-assembled contigs and the putative co-localization of ARGs and class 1 integron-associated genes. MGEs other than class 1 integron-associated genes did not exhibit a significant correlation with ARGs. This is in line with the current understanding of the role of class 1 integrons, in particular, among all MGEs in spreading acquired ARGs (13).

Interestingly, the ARGs observed in HWWs from Benin were the highest in number but probably lower in clinical importance compared to those in the HWWs from Burkina Faso and Finland. While the most significant carbapenemase genes *blaKPC*, *blaNDM*, *blaVIM*, *blaIMP*, *blaOXA-48*, and *blaOXA-58* were found in several HWW samples from Burkina Faso and Finland at varying abundances, the abundance of many of them was low, or they were not detected at all in HWWs from Benin. The *blaGES* carbapenemase genes were an exception. They were the most prevalent of the seven carbapenemases in Beninese HWWs and were also found in samples unrelated to hospitals.

We suspect that as *blaGES* carbapenemases are typically carried by class 1 integrons, as confirmed by previous studies (53), a co-selection phenomenon (3, 54) might play a role in increasing their spread also in here. In this scenario, residues of antibiotics other than carbapenems would cause the co-selection of *blaGES*. We also found a high relative abundance of plasmid-mediated quinolone ARGs in the HWWs from Benin. These *qnrVC* genes showed strong correlations to class 1 integron-associated genes and were in some cases found to be located in the same contigs with *blaGES-5* carbapenemase genes (Figure S6B). Hence, selection targeting quinolone ARGs could co-select for carbapenem resistance encoded by the gene *blaGES-5*. Altogether our observations of ARGs putatively carried by class 1 integrons are worrisome as new ARG combinations among integron gene cassettes enable their selection by a broader set of antimicrobials or biocides such as quaternary ammonium compounds (QACs) (15, 55, 56).

Despite the lowest relative sum abundance of ARGs detected in HWWs from Finland, some of the seven carbapenemase genes of particular interest seemed to have even higher abundances in Finland than in the surveyed West African countries. However, the occurrence of these ARGs was not homogenous among the Finnish samples. The highest relative abundance seen in especially one sample from the University Central Hospital of the capital of Finland (hospital K) was likely explained by the high patient turnover compared to other hospitals in Finland and Benin and Burkina Faso studied here. Furthermore, a high relative abundance *blaKPC* was observed in only

one of the Finnish samples. Furthermore, the overall occurrence of *blaKPC* across all HWWs from Benin, Burkina Faso, and Finland studied here aligned with many local and global reports and systematic reviews (57–60), according to which *blaKPC* carbapenemases are spreading more profusely in Europe and North America than in Africa. Similarly, the relative abundance of *blaOXA-58* was higher among Finnish than Beninese or Burkinabe HWW samples.

Disconcertingly, *blaNDM* was found in the fecal sample of an in-patient staying at a Beninese hospital. *blaNDM* was also detected in the wastewaters of the hospital, indicating the sensitivity of wastewater-based monitoring of bacterial resistance carried by hospital in-patients. *blaNDM* carried by *A. baumannii* has recently, in 2021, been reported in clinical settings in Benin (61).

Our observation of the high prevalence of metronidazole ARGs in Benin and Burkina Faso but not in Finland most likely reflects differences in the consumption of this drug. In LICs, diarrheal infections caused by pathogens such as *Clostridioides difficile* and other anaerobic bacteria, as well as protozoa, are common. Therefore, the consumption of metronidazole commonly used to treat these infections may be high in these countries as these drugs are being prescribed, for example, both with and without a positive diagnostic test for *C. difficile* (62). However, multidrug-resistant strains of *C. difficile* seem to already be prevalent in hospitals in certain parts of Africa as a consequence of the heavy use of antimicrobials among in-patients(62).

Compared to many HMICs such as those in Northern Europe, antimicrobial usage in agricultural (6, 7) and clinical (4) settings are less controlled – or even entirely unregulated – in many African countries. For example, while banned in many other countries, the use of colistin as a feed additive is allowed in many LICs (63), such as those in Africa (7). In contrast to previous reports on the prevalence of mobile colistin resistance globally (30) and in Africa (31, 64), *mcr-5* was the most commonly detected gene in the HWWs from our study. Instead, our observations are in line with the results obtained by using both metagenomics and qPCR (65). This discrepancy may be due to differences between the methodologies used to screen for colistin resistance. Colistin ARGs other than the best-known genes *mcr-1*, -2, and -3 are rarely targeted when screening for mobile colistin resistance with conventional PCR (65). Our study shows that to obtain a more realistic view of mobile colistin resistance genes in Africa, screening should also be conducted for different *mcr* genes, as was recently done by Ngbede and colleagues (66).

The *mcr-5* gene detected here was embedded in a Tn3-like transposase similar to previous reports for *Salmonella enterica* (67, 68) and *Escherichia coli* (69) plasmids and the chromosome of *Cupriavidus gilardii* (67). These Tn3-like elements are flanked by inverted repeats, which enable translocation and putatively a wide host range for the *mcr-5.1* harboring element (67–69). Based on our results, the occurrences of *mcr-5* and other mobile colistin resistance genes in Benin and Burkina Faso were not restricted to HWW septic tanks. For example, at a Beninese hospital, a water puddle in the hospital yard had a higher relative abundance of *mcr* genes compared to the actual HWW septic tanks for that hospital. Furthermore, these genes were also detected in water intended for handwashing of visitors to the hospital. Thus, the transmission of these clinically critical ARGs within hospitals could be enabled via hands contaminated by this water. Similarly, Jacobs and colleagues have drawn attention to

the potentially inferior microbiological water quality in similar hand washing water tanks, which are very common in West Africa (4).

In Burkina Faso, the relative abundance of *mcr-5* released into natural waters after wastewater treatment was high (2.36×10^{-4}). For comparison, a similar relative abundance of *mcr-5* genes (3.0×10^{-4}) was reported for untreated municipal wastewater in Germany (65). Although we cannot confirm the association between the *mcr-5* genes detected in the natural waters and the hospital-associated wastewater in the Burkinabe hospital, our study suggests the inability of the currently used wastewater treatment processes in Burkina Faso to remove colistin resistance genes. Wastewater treatment systems have also earlier been described to be inadequate in Benin and Burkina Faso (70). This increases the risk of the further spread of colistin and other ARGs as the inadequately treated water is released into natural waters and used for various purposes by local people. Furthermore, in many African countries, untreated wastewater is used for irrigation in urban agriculture, possibly enabling the dissemination of ARGs to fresh produce (71, 72). The situation appears to be especially critical in Cotonou, where the hospitals involved in the study did not use any kind of wastewater treatment. Groundwater in the city is extracted from the shallow aquifer, which is polluted due to unauthorized waste deposits, inadequate toilets, pit latrines, and septic tanks prone to leakage and hydraulic failure (73). Our study shows that untreated hospital wastewaters, which can leak into groundwater and the environment, carry clinically dangerous antimicrobial-resistant bacteria and resistance genes.

Taken together, based on our results from nine hospitals, we can state that clinically important ARGs are circulating in West African hospitals and can spread to their surroundings. There was no significant difference in the relative sum abundances of ARGs based on the hospital section or different hospitals within these countries. Various carbapenem ARGs were found in HWWs from Burkina Faso and Finland, while among these *bla*GES seem to dominate in the studied hospitals in Benin. Our results suggest that class 1 integron mediated co-selection could play an important role in shaping the resistome in the Beninese samples. In addition to HWWs, carbapenemase genes and mobile colistin resistance genes were detected in the vicinity as well as locations more distant from the hospitals. Finally, there were differences between the HWW resistomes in Benin and Burkina Faso concerning the relative sum abundance of ARGs and the prevalence of clinically relevant ARGs. Therefore, it is important to consider the differences between African countries in AMR surveillance.

4. Material and Methods

4.1. Sample description

Hospital wastewater (HWW) samples were collected in Benin from four (hospitals A-D) and in Burkina Faso from five different hospitals (hospitals F-J) in November and December of 2019. In Finland, HWW samples were collected from four different hospitals (hospitals K-N) in January 2020. In Burkina Faso, the samples were mainly from septic tanks or sewers in the hospital area. In Benin, none of the hospitals was connected to a sewer system, and the samples were from septic tanks or sumps (unstructured wastewater wells), which were never emptied to our knowledge. In most cases, the toilet water was not directed into these sumps. For comparison, some other

types of samples were included: samples from river waters used for drinking in a remote countryside village in the community of Savalou in central Benin (n=3); river or wetland water receiving treated wastewater in Burkina Faso (n=2, hospitals I and J); and fecal samples of in-patients in Benin (n=4, hospital B). In addition, the following samples were analyzed; well water (hospital A), treated tap water (community of Savalou in central Benin), water for handwashing (hospital C), water in a municipality channel receiving biologically treated wastewater (hospital I), water and sediment from a street gutter next to the hospital (hospital B), an empty HWW septic tank (hospital A), and the soil between HWW septic tanks (hospital B). The four fecal samples were obtained from patients staying at hospital B with diagnoses mainly associated with musculoskeletal injuries. No clinical data concerning bacterial infections were reported for these patients, and we have no information on their possible use of antibiotics. Patient age varied from 24 to 50 years old, and at the time of sampling, patients had stayed in the hospital for one to two months. All sample details are provided in the Supporting Information, Table S2A. Selected pictures and a map indicating sample collection regions are shown in Figures S1 and S2. Letters A–N were used to denote the hospitals/sampling locations.

4.2. DNA extraction and metagenomic sequencing

Water samples were collected into 1-liter bottles and transported to a laboratory on ice. They were kept at +4 °C until processed within 24 hours. A volume of 50–100 ml was filtered through a 0.2 µm polycarbonate filter (Whatman™, GE Healthcare Life Sciences) using a portable vacuum pump (Millivac-Mini Vacuum Pump XF54, Millipore, Merck). DNA was extracted from the filters using the Qiagen DNeasy PowerWater DNA Kit for the water samples and Qiagen DNeasy PowerSoil DNA Kit for the soil samples, following manufacturer instructions. The concentration and quality of the extracted DNA were determined with a NanoDrop spectrophotometer. Altogether 93 samples were subjected to shotgun metagenomic sequencing using Illumina Novaseq5000 with Nextera XT library preparation at the Institute of Biotechnology, University of Helsinki.

4.3. Bioinformatic analyses

All samples were analyzed using an in-house Snakemake workflow (74). Briefly, the quality control steps included in the workflow were performed using FastQC (75) and MultiQC (76), with adapter and low-quality read removal using Cutadapt (77). Nucleotide sequence reads were mapped using Bowtie2 (78) against the ResFinder database (version 3.2, downloaded on 2020-06-28) (79). The reads were sorted and filtered using SAMtools (80) such that the reads mapping as pair or alone were calculated as a single count. Mobile genetic elements were identified by mapping the reads against the MobileGeneticElementDatabase (<https://github.com/KatariinaParnanen/MobileGeneticElementDatabase>, downloaded on 2020-06-28) consisting of 2714 unique MGE sequences including transposases; integrons of classes 1, 2, and 3; the integron associated disinfectant resistance gene *qacEΔ*; and IS and ISCR type insertion sequences (46). Class 1 integrons were studied using mapping results obtained for the *intI* gene, which encodes an integron integrase (*IntI*). Taxonomic profiling was performed using both Metaphlan3 (81) and Metaxa2 (82). The counts for bacterial 16S rRNA from Metaxa2 were used to normalize ARG and MGE counts. The lengths of the functional genes and the 16S rRNA SSU gene were taken into account in the normalization.

The accuracy of the normalization method was validated by comparing two distinct methods using the 16S rRNA and *rpoB* housekeeping genes. The *rpoB* counts were obtained using the translated sequence reads described above as input for hmmsearch (83) against a *rpoB* specific HMM profile (pf04563) from the Pfam-A database (84). Hits for forward reads (R1) were counted. The correlation between the ARGs normalized to both 16S rRNA, and *rpoB* was significant ($R^2 = 0.61$, $p = 4.8 \times 10^{-8}$), and we proceeded to use 16S rRNA genes for normalization.

To study mobile colistin resistance genes (*mcr* genes) and their genetic environments for their theoretical mobility potential, a subset of the samples (samples BH02, BH48, BFH19, BFH26, BFH41, FH1, and FH9) was assembled using MEGAHIT (v1.2.8) (85). The minimum contig length was set to 1,000 bp. Contigs harboring mobile colistin resistance genes were identified using a BLAST (86) search against *mcr* genes in the ResFinder database (n=54). Similarly, contigs containing class 1 integrons were identified and screened for putative ARGs located in the same contig.

Visualizing the genetic environments of contigs harboring *mcr-5.1* colistin ARGs was performed using the anvi'o (v7) (87) platform tool 'anvi-interactive'. For this, single profiles were formed for the samples BH02, BH48, BFH26, BFH41, and FH9 with the option '-cluster-contigs'. Then, a BLAST nr search of the contigs against the ResFinder and MobileGeneticElementDatabase was performed. The resulting annotations were imported into the individual sample profiles containing the contigs database and open reading frames predicted by Prodigal (v2.6.3) (88) with the command 'anvi-import-functions'. The other predicted genes, sharing the same genetic environment with the *mcr-5.1* gene, were identified by inspecting the resulting annotations from the BLAST nr search on the web BLAST 'Graphics' view.

To simplify the presentation of the relative sum abundance results of the mobile colistin resistance genes, the *mcr* genes found in ResFinder database (79) were clustered based on 90 % similarity in their sequence identity using CD-HIT (v4.8.1) (89).

To visualize the co-localization of multiple ARGs with Bandage (90), the assembled contigs were converted into assembly graphs using the MEGAHIT command 'megahit_toolkit contig2fastg'. The Bandage program was run with a BLAST (86) query of a custom database consisting of ARGs present in the ResFinder database and the integron sequence-related genes *qacEA* and *intI1*.

4.4. Statistical analyses

All statistical analyses described below were performed for all 67 HWW samples described in the 'Sample details' section and Supporting Information Table S2B. The various wards, clinics, and other units typically had their own septic tanks or sumps in the Beninese and Burkinabe hospitals. The samples were designated accordingly as 'surgery', 'maternity', 'pediatry', or 'other'. In addition, samples from septic tanks or sumps receiving wastewater from different sources were designated as 'other'.

To model the relative sum abundances of ARGs and MGEs in HWWs from Benin, Burkina Faso, and Finland, a generalized linear model (GLM) with gamma distribution and logarithmic link was fitted to 16S rRNA normalized count data. The effects of the country, hospital and hospital section on the abundance of ARGs and MGEs were studied with the function ‘glm’ in the package MASS (v7.3-53.1) (91).

The ‘ordinate’ function in the Phyloseq package (v1.30.0) (92) was used for principal coordinate analysis (PCoA) of resistomes, mobilomes, and taxonomical compositions. Country-related differences in the prevalence of specific resistance genes were examined using DESeq2 (v1.26.0) (93).

The significance of the distinct clustering between the HWWs from different countries drawn using Phyloseq package (v1.30.0) (92) was tested using pair-wise adonis in the package vegan (v2.5-7) (94) with equal sample groups. For this, eight samples per country were selected randomly among the HWWs. Unique and shared ARGs were visualized with the function ‘draw.triple.venn’ in the package VennDiagram (v1.6.20) (95).

Pearson correlation coefficients and adjusted *p*-values for each ARG for *intI1* and *qacEA* genes were obtained with the function ‘corr.test’ in the package psych (v2.1.3) (96) and visualized using package ggcorrplot (v0.1.3) (97). The correlation tests for different normalization methods (*rpoB* and 16S rRNA) were performed similarly.

DESeq2 (v1.26.0) (93) was used to study the differentially abundant ARGs between the HWWs from the three different countries. ARGs filtered using a variance threshold of 50 were screened for significantly ($p < 0.05$) different relative abundances represented by log-2-fold changes across the HWWs of the three countries in a pair-wise manner (Benin-Finland, Burkina Faso-Finland). The DESeq2 algorithm is designed for unnormalized count data from gene expression studies. Thus, to take into account the amount of bacterial DNA in the samples (counts of detected 16S rRNA), the relative abundances were multiplied by 10^5 and a pseudo count of one was added to avoid zero values unsuitable for integer transformation. A similar analysis using DESeq2 was performed to study differentially abundant taxa found in the HWWs from Benin or Burkina Faso versus Finland. The relative abundances obtained from Metaphlan3 (81) were multiplied with the corresponding 16S rRNA counts for each sample from the Metaxa2 (82) output. Analyses for differentially abundant genera and species were conducted separately.

Statistical analyses were performed in R (v3.6.3), and the results were visualized using ggplot2 (v3.3.3) (98), pheatmap (v 1.0.12) (99), and patchwork (v1.1.0.9000) (100). Vector maps were drawn using the package rnatuarearth (v0.1.0) (101). Detailed descriptions for the bioinformatic and statistical analyses can be viewed on the project’s GitHub page.

Ethical permission

The ethical permissions for the project were received from “Comité National d’Ethique pour la Recherche en Santé” under the Health Ministry in Benin and “Comité d’Ethique pour la Recherche en Santé” under the Health Ministry in Burkina Faso.

Supporting Information

The supporting information referred to within the text as Figures S1-S7 and Tables S1-S18 can be found in the two separate files “Supporting_Information_1” (Figures S1-S7, Table S1) and “Supporting_Information_2” (Tables S2-S18).

Acknowledgments

We thank the staff in all the participating hospitals for their collaboration in sampling. We also thank the laboratory personnel of the URMPha at the University of Abomey-Calavi in Benin and the Clinical Research Unit of Nanoro (CRUN) in Burkina Faso for sample processing. Finally, we acknowledge CSC, IT Center for Science, Finland, for providing the computational resources for the study.

Funding Sources

This work was supported by the Academy of Finland (Grant Numbers 346125, 318643 and 316708) and University of Helsinki HiLife Grand Challenges funding.

Data availability

The data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under the accession number PRJEB47975 and will be public upon article publication in a peer-reviewed journal.

Code availability

All custom codes used for the analyses are available from <https://github.com/melinamarkkanen/AMRIWA> upon article publication in a peer-reviewed journal.

References

1. 2015. Global Action Plan On Antimicrobial Resistance WHO Library Cataloguing-in-Publication Data Global Action Plan on Antimicrobial Resistance <https://doi.org/10.13140/RG.2.2.24460.77448>.
2. Ventola CL. 2015. The Antibiotic Resistance Crisis Part 1: Causes and Threats. *Physichal Therapy* 40:277–83.
3. Laxminarayan R, Duse A, Wattal C, Zaidi AKM, Wertheim HFL, Sumpradit N, Vlieghe E, Hara GL, Gould IM, Goossens H, Greko C, So AD, Bigdeli M, Tomson G, Woodhouse W, Ombaka E, Peralta AQ, Qamar FN, Mir F, Kariuki S, Bhutta ZA, Coates A, Bergstrom R, Wright GD, Brown ED, Cars O. 2013. Antibiotic resistance-the need for global solutions. *Lancet Infect Dis* 13:1057–1098.
4. Jacobs J, Hardy L, Semret M, Lunguya O, Phe T, Affolabi D, Yansouni C, Vandenberg O. 2019. Diagnostic Bacteriology in District Hospitals in Sub-Saharan Africa: At the Forefront of the Containment of Antimicrobial Resistance. *Frontiers in Medicine* 6:205.
5. Belachew SA, Hall L, Selvey LA. 2021. Non-prescription dispensing of antibiotic agents among community drug retail outlets in Sub-Saharan African countries: a systematic review and meta-analysis. *Antimicrobial Resistance and Infection Control*. BioMed Central Ltd.
6. Butcher A, Cañada JA, Sariola S. 2021. How to make noncoherent problems more productive: Towards an AMR management plan for low resource livestock sectors. *Humanities and Social Sciences Communications* 8:287.
7. Van TTH, Yidana Z, Smooker PM, Coloe PJ. 2020. Antibiotic use in food animals worldwide, with a focus on Africa: Pluses and minuses. *Journal of Global Antimicrobial Resistance* 20:170–177.
8. Hu Y, Gao GF, Zhu B. 2017. The antibiotic resistome: gene flow in environments, animals and human beings. *Frontiers of Medicine* 11:161–168.
9. Tacconelli E, Carrara E, Savoldi A, Kattula D, Burkert F. 2017. Global Priority List Of Antibiotic-Resistant Bacteria To Guide Research, Discovery, And Development Of New Antibiotics.

10. Martínez JL, Coque TM, Baquero F. 2014. What is a resistance gene? Ranking risk in resistomes. *Nature Reviews Microbiology* 13:116–123.
11. Zhang A-N, Gaston JM, Dai CL, Zhao S, Poyet M, Groussin M, Yin X, Li L-G, van Loosdrecht MCM, Topp E, Gillings MR, Hanage WP, Tiedje JM, Moniz K, Alm EJ, Zhang T. An omics-based framework for assessing the health risk of antimicrobial resistance genes <https://doi.org/10.1038/s41467-021-25096-3>.
12. Karkman A, Berglund F, Flach CF, Kristiansson E, Larsson DGJ. 2020. Predicting clinical resistance prevalence using sewage metagenomic data. *Communications Biology* 3:711.
13. Gillings MR, Gaze WH, Pruden A, Smalla K, Tiedje JM, Zhu Y-G. 2015. Using the class 1 integron-integrase gene as a proxy for anthropogenic pollution. *The ISME Journal* 9:1269–1279.
14. Li Y, Yang X, Zhang J, Yang S, Zhang S, Chen M, Xue L, Ding Y, Zeng H, Gu Q, Zhang Y, Wei X, Wang J, Wu Q. 2021. Molecular characterisation of antimicrobial resistance determinants and class 1 integrons of *Salmonella enterica* subsp. *enterica* serotype Enteritidis strains from retail food in China. *Food Control* 128:108191.
15. Gillings MR. 2014. Integrons: Past, Present, and Future. *Microbiology and Molecular Biology Reviews* 78:257–77.
16. Gillings M, Boucher Y, Labbate M, Holmes A, Krishnan S, Holley M, Stokes HW. 2008. The Evolution of Class 1 Integrons and the Rise of Antibiotic Resistance. *JOURNAL OF BACTERIOLOGY* 190:5095–5100.
17. Chen B, Han J, Dai H, Jia P. 2021. Biocide-tolerance and antibiotic-resistance in community environments and risk of direct transfers to humans: Unintended consequences of community-wide surface disinfecting during COVID-19? *Environmental Pollution* 283:117074.
18. de Angelis G, Fiori B, Menchinelli G, Tiziana D'inzeo & Flora & Liotti M, Morandotti GA, Sanguinetti M, Posteraro B, Spanu T. 2007. Incidence and antimicrobial resistance trends in bloodstream infections caused by ESKAPE and *Escherichia coli* at a large teaching hospital in Rome, a 9-year analysis (2007-2015). *European Journal of Clinical Microbiology & Infectious Diseases* 37:1627–1636.
19. Alemayehu T, Hailemariam M. 2020. Prevalence of vancomycin-resistant enterococcus in Africa in one health approach: a systematic review and meta-analysis. *Scientific Reports* 10:20542.
20. Karkman A, Do TT, Walsh F, Virta MPJ. 2018. Antibiotic-Resistance Genes in Waste Water. *Trends in Microbiology* 26:220–228.
21. World Health Organization. 2021. Global Antimicrobial Resistance and Use Surveillance System (GLASS) Report 2021. Geneva.
22. World Health Organization. 2018. WHO report on surveillance of antibiotic consumption: 2016-2018 early implementation.
23. Suay-García, Pérez-Gracia. 2019. Present and Future of Carbapenem-resistant Enterobacteriaceae (CRE) Infections. *Antibiotics* 8:122.
24. Poirel L, Marqué S, Héritier C, Segonds C, Chabanon G, Nordmann P. 2005. OXA-58, a novel class D β -lactamase involved in resistance to carbapenems in *Acinetobacter baumannii*. *Antimicrobial Agents and Chemotherapy* 49:202–208.
25. Poirel L, Weldhagen GF, Naas T, de Champs C, Dove MG, Nordmann P. 2001. GES-2, a class A β -lactamase from *Pseudomonas aeruginosa* with increased hydrolysis of imipenem. *Antimicrobial Agents and Chemotherapy* 45:2598–2603.
26. Nordmann P, Poirel L. 2013. The difficult-to-control spread of carbapenemase producers among Enterobacteriaceae worldwide. *Clinical Microbiology and Infection* 20:821–830.
27. Löfmark S, Edlund C, Erik Nord C. 2010. Metronidazole Is Still the Drug of Choice for Treatment of Anaerobic Infections. *Clinical Infectious Diseases* 1:S16–S23.
28. Alauzet C, Lozniewski A, el Ene Marchandin H. 2018. Metronidazole resistance and nim genes in anaerobes: A review. *Anaerobe* 55:40–53.
29. Falagas ME, Kasiakou Sofia. K. 2005. Colistin: The Revival of Polymyxins for the Management of Multidrug-Resistant Gram-Negative Bacterial Infections. *Clinical Infectious Diseases* 40:1333–41.
30. Ling Z, Yin W, Shen Z, Wang Y, Shen J, Walsh TR. 2020. Epidemiology of mobile colistin resistance genes mcr-1 to mcr-9. *Journal of Antimicrobial Chemotherapy* 75:3087–3095.
31. Anyanwu MU, Odilichukwu C, Okpala R, Chah KF, Shoyinka VS. 2021. Prevalence and Traits of Mobile Colistin Resistance Gene Harboursing Isolates from Different Ecosystems in Africa. *BioMed Research International* 22:6630379.
32. Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R. 2018. Anthropological and socioeconomic factors contributing to global antimicrobial resistance: a univariate and multivariable analysis. *Lancet Planet Health* 2:e398-405.

33. van Boeckel TP, Pires J, Silvester R, Zhao C, Song J, Criscuolo NG, Gilbert M, Bonhoeffer S, Laxminarayan R. 2019. Global trends in antimicrobial resistance in animals in low-and middle-income countries. *Science* 365:eaaw1944.
34. Iskandar K, Molinier L, Hallit S, Sartelli M, Craig Hardcastle T, Haque M, Lugova H, Dhingra S, Sharma P, Islam S, Mohammed I, Naina Mohamed I, Abi Hanna P, el Hajj S, Adilla Hayat Jamaluddin N, Salameh P, Roques C. 2020. Surveillance of antimicrobial resistance in low- and middle-income countries: a scattered picture. *Antimicrob Resist Infect Control* 10:63.
35. Ambler RP. 1980. The structure of beta-lactamases. *Philosophical transactions of the Royal Society of London Series B, Biological sciences* 289:321–331.
36. Bernabé KJ, Langendorf C, Ford N, Ronat J-B, Murphy RA. 2017. Antimicrobial resistance in West Africa: a systematic review and meta-analysis. *International Journal of Antimicrobial Agents* 50:629–639.
37. Mahamat OO, Kempf M, Lounnas M, Tidjani A, Hide M, Benavides JA, Carrière C, Bañuls A-L, Jean-Pierre H, Ouedraogo A-S, Dumont Y, Godreuil S. 2021. Epidemiology and prevalence of extended-spectrum β -lactamase-and carbapenemase-producing Enterobacteriaceae in humans, animals and the environment in West and Central Africa. *International Journal of Antimicrobial Agents* 57:106203.
38. Hendriksen RS, Munk P, Njage P, van Bunnik B, McNally L, Lukjancenko O, Röder T, Nieuwenhuijse D, Pedersen SK, Kjeldgaard J, Kaas RS, Clausen PTLC, Vogt JK, Leekitcharoenphon P, van de Schans MGM, Zuidema T, de Roda Husman AM, Rasmussen S, Petersen B, Amid C, Cochrane G, Sicheritz-Ponten T, Schmitt H, Alvarez JRM, Aidara-Kane A, Pamp SJ, Lund O, Hald T, Woolhouse M, Koopmans MP, Vigre H, Petersen TN, Aarestrup FM. 2019. Global monitoring of antimicrobial resistance based on metagenomics analyses of urban sewage. *Nature Communications* 10:1124.
39. Pärnänen KMM, Narciso-da-Rocha C, Kneis D, Berendonk TU, Cacace D, Do TT, Elpers C, Fatta-Kassinos D, Henriques I, Jaeger T, Karkman A, Martinez JL, Michael SG, Michael-Kordatou I, O'Sullivan K, Rodriguez-Mozaz S, Schwartz T, Sheng H, Sørum H, Stedtfeld RD, Tiedje JM, Giustina SV della, Walsh F, Vaz-Moreira I, Virta M, Manaia CM. 2019. Antibiotic resistance in European wastewater treatment plants mirrors the pattern of clinical antibiotic resistance prevalence. *Science Advances* 5:eaau9124.
40. European Centre for Disease Prevention and Control. 2020. Antimicrobial resistance in the EU/EEA (EARS-Net) - Annual Epidemiological Report 2019.
41. Buelow E, Rico A, Gaschet M, Lourenço J, Kennedy SP, Wiest L, Ploy M-C, Dagot C. 2020. Hospital discharges in urban sanitation systems: Long-term monitoring of wastewater resistome and microbiota in relationship to their eco-exposome. *Water Research X* 7:100045.
42. Thomas AM, Segata N. Multiple levels of the unknown in microbiome research <https://doi.org/10.1186/s12915-019-0667-z>.
43. Naas T, Oueslati S, Bonnin RA, Laura Dabos M, Zavala A, Dortet L, Retailleau P, Iorga BI, emy Bonnin RA. 2017. Beta-lactamase database (BLDB)-structure and function. *Journal of Enzyme Inhibition and Medicinal Chemistry* 32:917–919.
44. Thompson L, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill robert J, Tripathi A, Gibbons S, Ackermann G, Navas-molina JA, Janssen S, Kopylova E, Vázquez-baeza Y, González A, Morton JT, Mirarab S, Zech Xu Z, Jiang L, Haroon mohamed F, Kanbar J, Zhu Q, Jin Song S, Kosciolk T, Bokulich nicholas A, Lefler J, Brislawn colin J, Humphrey G. 2017. A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* 551:457–463.
45. Oyem IM, Oyem HH, Atuanya. 2020. Molecular characterisation of bacteria strains of septic tank sewage samples from related sites in Delta and Edo States of Nigeria using 16S rRNA denaturing gradient gel electrophoresis (DGGE) 14:132–138.
46. Pärnänen K, Karkman A, Hultman J, Lyra C, Bengtsson-Palme J, Larsson DGJ, Rautava S, Isolauri E, Salminen S, Kumar H, Satokari R, Virta M. 2018. Maternal gut and breast milk microbiota affect infant gut antibiotic resistome and mobile genetic elements. *Nature Communications* 9:389.
47. Laursen MF, Laursen RP, Larnkjaer A, Mølgaard C, Michaelsen KF, Frøkiaer H, Bahl MI, Licht TR. 2017. Faecalibacterium Gut Colonization Is Accelerated by Presence of Older Siblings. *mSphere* 2:e00448-17.
48. Collis CM, Hall RM. 1995. Expression of antibiotic resistance genes in the integrated cassettes of integrons. *Antimicrobial Agents and Chemotherapy* 39:155–162.
49. Périchon B, Goussard S, Walewski V, Krizova L, Cerqueira G, Murphy C, Feldgarden M, Wortman J, Clermont D, Nemec A, Courvalin P. 2014. Identification of 50 class D β -lactamases and 65 Acinetobacter-derived cephalosporinases in Acinetobacter spp. *Antimicrobial Agents and Chemotherapy* 58:936–949.

50. Figueiredo S, Bonnin RA, Poirel L, Duranteau J, Nordmann P. 2012. Identification of the naturally occurring genes encoding carbapenem-hydrolysing oxacillinases from *Acinetobacter haemolyticus*, *Acinetobacter johnsonii*, and *Acinetobacter calcoaceticus*. *Clin Microbiol Infect* 18:907–913.
51. Massidda O, Rossolini GM, Satta G. 1991. The *Aeromonas hydrophila* cphA gene: Molecular heterogeneity among class B metallo- β -lactamases. *Journal of Bacteriology* 173:4611–4617.
52. Cox G, Wright GD. 2013. Intrinsic antibiotic resistance: Mechanisms, origins, challenges and solutions. *International Journal of Medical Microbiology* 303:287–292.
53. Araújo S, Sousa M, Tação M, Baraúna RA, Silva A, Ramos R, Alves A, Manaia CM, Henriques I. 2021. Carbapenem-resistant bacteria over a wastewater treatment process: Carbapenem-resistant Enterobacteriaceae in untreated wastewater and intrinsically-resistant bacteria in final effluent. *Science of the Total Environment* 782:146892.
54. Zieliński W, Korzeniewska E, Harnisz M, Drzymala J, Felis E, Bajkacz S. 2021. Wastewater treatment plants as a reservoir of integrase and antibiotic resistance genes – An epidemiological threat to workers and environment. *Environment International* 156:106641.
55. Pal C, Bengtsson-Palme J, Kristiansson E, Larsson DGJ. 2015. Co-occurrence of resistance genes to antibiotics, biocides and metals reveals novel insights into their co-selection potential. *BMC Genomics* 16.
56. Gillings MR, Xuejun D, Hardwick SA, Holley MP, Stokes HW. 2009. Gene cassettes encoding resistance to quaternary ammonium compounds: a role in the origin of clinical class 1 integrons? *The ISME Journal* 3:209–215.
57. Olalekan A, Onwugamba F, Iwalokun B, Mellmann A, Becker K, Schaumburg F. 2020. High proportion of carbapenemase-producing *Escherichia coli* and *Klebsiella pneumoniae* among extended-spectrum β -lactamase-producers in Nigerian hospitals. *Journal of Global Antimicrobial Resistance* 21:8–12.
58. Manenzhe RI, Zar HJ, Nicol MP, Kaba M. 2015. The spread of carbapenemase-producing bacteria in Africa: a systematic review. *Journal of Antimicrobial Chemotherapy* 70:23–40.
59. van Duin D, Doi Y. 2017. The global epidemiology of carbapenemase-producing Enterobacteriaceae. *Virulence* 8:460–469.
60. Mushi MF, Mshana SE, Imirzalioglu C, Bwanga F. 2014. Carbapenemase Genes among Multidrug Resistant Gram Negative Clinical Isolates from a Tertiary Hospital in Mwanza, Tanzania. *BioMed Research International* 2014:303104.
61. Yehouenou C, Bogaerts B, Vanneste K, Roosens NHC, de Keersmaecker SCJ, Marchal K, Affolabi D, Soleimani R, Rodriguez-Villalobos H, van Bambeke F, Dalleur O, Simon A. 2021. First detection of a plasmid-encoded New-Delhi metallo-beta-lactamase-1 (NDM-1) producing *Acinetobacter baumannii* using whole genome sequencing, isolated in a clinical setting in Benin. *Annals of Clinical Microbiology and Antimicrobials* 20:5.
62. Mutai WC, Mureithi MW, Anzala O, Revathi G, Kullin B, Burugu M, Kyany'a C, Odoyo E, Otieno P, Musila L. 2021. High Prevalence of Multidrug-Resistant *Clostridioides difficile* Following Extensive Use of Antimicrobials in Hospitalized Patients in Kenya. *Frontiers in Cellular and Infection Microbiology* 10:897.
63. Olumuyiwa Olaitan A, Dandachi I, Alexandra Baron S, Daoud Z, Morand S, Rolain J-M. 2021. Banning colistin in feed additives: a small step in the right direction. *The Lancet Infectious Diseases* 21:29–30.
64. Olowo-Okere A, Yacouba A. 2020. Review Molecular mechanisms of colistin resistance in Africa: A systematic review of literature. *Germs* 10:367–379.
65. Kneis D, Berendonk TU, Heß S. 2019. High prevalence of colistin resistance genes in German municipal wastewater. *Science of the Total Environment* 694:133454.
66. Ngbede EO, Poudel A, Kalalah A, Yang Y, Adekanmbi F, Adikwu AA, Adamu AM, Mamfe LM, Daniel ST, Useh NM, Kwaga JKP, Adah MI, Kelly P, Butaye P, Wang C. 2020. Identification of mobile colistin resistance genes (*mcr-1.1*, *mcr-5* and *mcr-8.1*) in Enterobacteriaceae and *Alcaligenes faecalis* of human and animal origin, Nigeria. *International Journal of Antimicrobial Agents* 56:106108.
67. Borowiak M, Fischer J, Hammerl JA, Hendriksen RS, Szabo I, Malorny B. 2017. Identification of a novel transposon-associated phosphoethanolamine transferase gene, *mcr-5*, conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica* serovar Paratyphi B. *Journal of Antimicrobial Chemotherapy* 72:3317–3324.
68. Borowiak M, Hammerl JA, Deneke C, Fischer J, Szabo I, Malorny B. 2019. Characterization of *mcr-5*-Harboring *Salmonella enterica* subsp. *enterica* Serovar Typhimurium Isolates from Animal and Food Origin in Germany. *Antimicrobial Agents and Chemotherapy* 63:e00063-19.
69. Guo S, Tay MYF, Thu AK, Seow KLG, Zhong Y, Ng LC, Schlundt J. 2019. Conjugative IncX1 plasmid harboring colistin resistance gene *mcr-5.1* in *Escherichia coli* isolated from chicken rice retailed in Singapore. *Antimicrobial Agents and Chemotherapy* 63:e01043-19.

- 781 70. Peace Hounkpe S, Codjo Adjovi E, Crapper M, Awuah E. 2014. Wastewater Management in Third
782 World Cities: Case Study of Cotonou. *Benin Journal of Environmental Protection* 5:387–399.
- 783 71. Bougnom BP, Zongo C, McNally A, Ricci V, Etoa FX, Thiele-Bruhn S, Piddock LJV. 2019.
784 Wastewater used for urban agriculture in West Africa as a reservoir for antibacterial resistance
785 dissemination. *Environmental Research* 168:14–24.
- 786 72. Bougnom BP, Thiele-Bruhn S, Ricci V, Zongo C, Piddock LJV. 2020. Raw wastewater irrigation for
787 urban agriculture in three African cities increases the abundance of transferable antibiotic resistance
788 genes in soil, including those encoding extended spectrum β -lactamases (ESBLs). *Science of the Total*
789 *Environment* 698:134201.
- 790 73. Totin HS, Ernest A, Odoulami L, Edorh PA, Boukari M, Boko M. 2013. Groundwater pollution and the
791 safe water supply challenge in Cotonou town, Benin (West Africa). *IAHS-AISH Proceedings and*
792 *Reports* 361:191–196.
- 793 74. Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V, Forster J, Lee S,
794 Twardziok SO, Kanitz A, Wilm A, Holtgrewe M, Rahmann S, Nahnsen S, Köster J. 2021. Sustainable
795 data analysis with Snakemake. *F1000Research* 10:33.
- 796 75. Andrews S. 2010. FastQC A quality control tool for high throughput sequence data
797 <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- 798 76. Ewels P, Ns Magnusson M, Lundin S, Aller MK. Data and text mining MultiQC: summarize analysis
799 results for multiple tools and samples in a single report. *Bioinformatics* 32:3047–3048.
- 800 77. Marcel M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
801 *EMBnet.journal* 17:10–12.
- 802 78. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 9:357–
803 359.
- 804 79. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A, Allesoe RL, Rebelo
805 AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B, Werner G, Bender JK, Stingl K, Nguyen
806 M, Coppens J, Xavier BB, Malhotra-Kumar S, Westh H, Pinholt M, Anjum MF, Duggett NA, Kempf I,
807 Nykäsenoja S, Olkkola S, Wiecek K, Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C,
808 Lund O, Aarestrup FM. 2020. ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of*
809 *Antimicrobial Chemotherapy* 75:3491–3500.
- 810 80. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009.
811 The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079.
- 812 81. Beghini F, Mciver LJ, Blanco-mi A, Dubois L, Asnicar F, Maharjan S, Mailyan A, Manghi P, Scholz
813 M, Thomas AM, Valles-colomer M, Weingart G, Zhang Y, Zolfo M, Huttenhower C, Franzosa EA,
814 Segata N. 2021. Integrating taxonomic, functional, and strain-level profiling of diverse microbial
815 communities with bioBakery 3. *Elife* 4:1–42.
- 816 82. Bengtsson-Palme J, Hartmann M, Eriksson KM, Pal C, Thorell K, Larsson DGJ, Nilsson RH. 2015.
817 metaxa2: Improved identification and taxonomic classification of small and large subunit rRNA in
818 metagenomic data. *Molecular Ecology Resources* 15:1403–1414.
- 819 83. Eddy SR. 2011. Accelerated Profile HMM Searches. *Citation: Eddy SR* 7:1002195.
- 820 84. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ, Salazar
821 GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, Finn RD. 2018. The
822 Pfam protein families database in 2019. *Nucleic Acids Research* 47:427–432.
- 823 85. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node solution for
824 large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31:1674–1676.
- 825 86. Altschup SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic Local Alignment Search Tool. *J*
826 *Mol Biol* 215:403–410.
- 827 87. Murat Eren A, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef M,
828 Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD, Kivenson V, Graham
829 ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin A, Scott JJ, Vázquez-Campos X,
830 McKay LJ, McDaniel EA, R Stevens SL, Anderson RE, Fuessel J, Fernandez-Guerra A, Maignien L,
831 Delmont TO, Willis AD. 2021. Community-led, integrated, reproducible multi-omics with anvi'o.
832 *Nature Microbiology* 6:3–6.
- 833 88. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene
834 recognition and translation initiation site identification. *BMC bioinformatics* 11.
- 835 89. Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. Sequence analysis CD-HIT: accelerated for clustering the next-
836 generation sequencing data 28:3150–3152.
- 837 90. Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualization of de novo genome
838 assemblies. *Bioinformatics* 31:3350–3352.
- 839 91. Venables WN, Ripley B. 2002. *Modern Applied Statistics with S*.

840 92. McMurdie PJ, Holmes S. 2013. Phyloseq: An R Package for Reproducible Interactive Analysis and
841 Graphics of Microbiome Census Data. PLoS ONE 8.
842 93. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq
843 data with DESeq2. Genome Biology 15:550.
844 94. Oksanen J, Blanchet G, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB,
845 Simpson GL, Solymos P, Stevens MHM, Szoecs E, Wagner H. 2020. vegan: Community Ecology
846 Package.
847 95. Chen H. 2018. VennDiagram: Generate High-Resolution Venn and Euler Plots. R package version
848 1.6.20.
849 96. Revelle W. 2021. psych: Procedures for Psychological, Psychometric, and Personality Research.
850 97. Kassambara A. 2019. ggcorrplot: Visualization of a Correlation Matrix using "ggplot2."
851 98. Wickham H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.
852 99. Kolde R. 2019. pheatmap: Pretty Heatmaps. R package version 1.0.12.
853 100. Pedersen TL. 2020. patchwork: The Composer of Plots. R package version 1.1.1.
854 101. South A. rnaturalearth: World Map Data from Natural Earth. R package version 0.1.0.
855