Key factors for effective implementation of healthcare workers support interventions in health organisations after patient safety incidents: a protocol for a scoping review

Sofia Guerra Paiva^{a,b}, Maria João Lobão^{a,b,c}, João Diogo Simões^{a,d}, Helena Donato^e, José Joaquin Mira^{f,g}, Irene Carrillo^g, Paulo Sousa^{a,b}

Corresponding author:

Sofia Guerra Paiva^{a,b},

<https://orcid.org/0000-0002-4311-3799>

Avenida Padre Cruz 1600-560 Lisboa

sg.paiva@ensp.unl.pt

- a. Public Health Research Centre, National School of Public Health, NOVA University of Lisbon, Lisboa Portugal
- b. Comprehensive Health Research Center, NOVA Medical School, Lisboa, Portugal
- c. Internal Medicine Department, Hospital de Cascais, Cascais, Portugal
- d. Public Health Unit of ACES USP Almada-Seixal, Almada, Portugal
- e. Documentation and Scientific Information Service, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal
- f. Alicante-Sant Joan Healthcare District, 03013 Alacant, Spain
- g. Department of Health Psychology, Universidad Miguel Hernández de Elche

Word count: 3641 words (excluding title page, abstract, references, figures and tables)

ARTICLE SUMMARY:

Abstract:

Introduction: Health organisations should support healthcare workers who are physically and psychologically affected by patient safety incidents (second victims). There is a growing body of evidence which focusses on second victim support interventions. However, there is still limited research on the elements necessary to effectively implement and ensure the sustainability of these types of interventions. In this study, we propose to map and frame the key factors which underlie an effective implementation of healthcare worker support interventions in healthcare organisations when healthcare workers are physically and/or emotionally affected by patient safety incidents.

Methods and analysis: This scoping review will be guided by the established methodological Arksey and O'Malley framework, Levac and Joanna Briggs Institute (JBI) recommendations. We will follow the JBI three-step process: (1) a preliminary search conducted on two databases; (2) the definition of clear inclusion criteria and the creation of a list of search terms to be used in the subsequent running of the search on a larger number of databases; and (3) additional searches (cross-checking/cross-referencing of reference lists of eligible studies, hand-searching in target journals relevant to the topic, conference proceedings, institutional/organisational websites and networks repositories). We will undertake a comprehensive search strategy in relevant bibliographic databases (Pubmed/Medline, Embase, CINAHL, Web of Science, Scopus, PsycInfo, Epistemonikos, Scielo, Cochrane Library and Open Grey). We will use the Mixed Methods Appraisal Tool (MMAT) 2018 version for quality assessment of the eligible studies. Our scoping review will be guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis extension for Scoping Reviews (PRISMA-ScR).

Ethics and dissemination: This study will not require ethical approval. Results of the scoping review will be published in a peer-review journal and findings will be presented in scientific conferences as well as in international forums and other relevant dissemination channels.

This protocol is registered on the Open Science Framework (www.osf.io):
[10.17605/OSF.IO/WZSE9](https://doi.org/10.17605/OSF.IO/WZSE9)

Keywords: support programs; health personnel; second victim; clinical error; patient safety

Strengths and limitations of this study:

- To our knowledge, this is the first scoping review to map and frame the different organisational, operational and contextual factors which underlie the implementation of health worker support programmes after the occurrence of a patient safety incident.
- This scoping review uses a rigorous and transparent method for mapping the available evidence.
- Given the broad focus of this scoping review, we don't expect that eligible studies will show a direct relation between the key factors and the effectiveness of the implementation of the support interventions.
- For better interpretation of the results, we will provide a quality assessment of the included studies, although quality assessment is not mandatory to include in scoping reviews.
- We will involve key stakeholders as an additional source of information to complement the literature search.

INTRODUCTION

Patient safety incidents affect patients' lives at different levels and globally impact health organisations and their health care workers [1,2]. A patient safety incident is an unintended or unexpected healthcare event that either causes harm to one or more patients (adverse event) or that doesn't reach a patient but poses a risk of harm (near miss) [1,3].

Patients suffering direct harm caused by a healthcare incident are the 'first victims' of an adverse event. Approximately 1 in 10 patients admitted to a hospital will suffer an adverse event, which represents 4% to 17% of hospital admissions [4,5], as will 4 in 10 patients in primary and outpatient healthcare [6]. According to a meta-analysis published in 2019, half of patient harm is preventable, which means that 6% to 12% of the prolonged, permanent disability or death identified in this meta-analysis could be prevented [7].

In 2000, Albert Wu [8] drew attention to the 'second victim' phenomenon in an editorial entitled 'Medical error: the second victim. The doctor who makes the mistake needs help too' which focused on the need to support health professionals who suffered from involvement in an adverse event. There is a growing body of evidence which focusses on understanding physical and psychological consequences experienced by healthcare professionals who are involved in a patient safety incident which causes serious or minor harm or even did not cause direct harm to the patient (near miss) [9,10]. Anxiety about future adverse events, depression, shame [9], frustration, repetitive/intrusive memories, extreme fatigue, sleep disturbances [11] and burnout [12] are some of the most frequent consequences reported in the literature. These emotional and physical consequences can impact professional performance and self-confidence [9], which can increase the likelihood of being involved in future adverse events [9,13].

Several studies have mentioned that health professionals feel unprotected and that they lack support after involvement in a healthcare incident [9,14]. Healthcare worker support practices are still underdeveloped and underused in healthcare institutions [9,14]. Although growing attention is being paid to the importance of cultivating supportive and non-judgmental environments, the blame culture continues to be one of the most frequent problems in health organisations [15]. This culture may negatively impact healthcare workers that are experiencing the second victim phenomenon [16].

Health organisations should be responsible for providing tools and training to support healthcare workers after an incident occurs during healthcare, thereby contributing to a learning environment and safer healthcare [17]. Most of the support strategies that have been internationally adopted are programmes and interventions [17]. The common goals of established support programmes are to reduce the psychological distress of second victims due to incidents and to foster coping strategies [2]. We can find different formats for support interventions/programmes, such as online programmes, peer support programmes, specific support tools and resources for healthcare providers [18,19,20].

Improvements in patient safety culture [22], resilience and adaptative capacity [23] are some outcomes related to these types of interventions in healthcare organisations.

According to the Donabedian model for measuring quality in medical care [24], outcomes depend on factors related to structure and processes.

Some recent studies have shown the influence of organisational structures, processes and outcomes on the effectiveness of research implementation turned into practice [25,26].

The described healthcare organisational characteristics were clearly aggregated by Yano [25] into organisational structures, organisational processes and organisational outcomes (summarised in Table 1).

Table 1- Organisational structures, processes and outcomes adapted from Yano [21]

Organisational Structures	<p>Some examples:</p> <ul style="list-style-type: none"> • Size of the organisational unit(s) – number of facilities, beds, providers • Services – general and specialty services • Staffing characteristics and integrated networks • Leadership structure/authority • Resource allocation • Organisational culture • Work environment/organisational climate
Organisational Processes	<ul style="list-style-type: none"> • Management processes – practice arrangements, managerial coordination of services and follow up • Communication processes, procedures, quality of interactions • Relationships – nature of roles and responsibilities, interpersonal styles
Organisational Outcomes	<ul style="list-style-type: none"> • Process quality measures • Intermediate outcome measures • Global health status measures • Utilisation measures • Workflow or efficiency measures • Costs

Organisational structures are focused on ‘static’ resources, which means that the resources are related to infrastructures, tools, equipment, healthcare services, units and the staffing level of the function for managing and delivering services [27].

Organisational processes, on the other hand, tend to be more mutable as they refer to organisational actions, procedures and service coordination. Buse, Mays and Walt referred to processes as the way in which policies are initiated, developed or formulated, negotiated, communicated, implemented and evaluated [28].

Organisational outcomes are described as performance measures and include practice-level measures of effectiveness. As mentioned by Yano [25], ‘organizational outcomes are distinct only insofar as they represent what the entire practice or institution would experience as a whole once implementation is complete and are thus inter-related to other evaluation activities’.

According to San Martín-Rodríguez, Beaulieu, D'Amour and Ferrada-Videla [26], organisational factors combine institutional characteristics as well as communication and coordination mechanisms. Factors in epidemiology are considered to be ‘events, characteristics, or other definable entities that have the potential to bring about a change in a health condition or other defined outcome’ [29].

In accordance with the health policy implementation process, an effective implementation includes a complex set of interrelationships: content, processes, context and actors [28].

Beyond all the aforementioned domains associated with the implementation of interventions, healthcare workers that participate in the interventions are considered to be important stakeholders for evidence-based implementation. Participant contributions can lead to more effective implementation and to the facilitation of practical applications [30] which, in turn, may contribute to the sustainability of the interventions

Study rationale:

There is a growing body of published evidence that is already focused on second victim support programmes and other types of support interventions in health organisations for healthcare workers who are physically and/or emotionally affected by patient safety incidents. A preliminary

search related to this topic revealed two recently published scoping reviews [17,31], a systematic review [2] and a meta-analysis [32]. The main focus of the published studies has been to describe health professional support interventions and resources along with their benefits for second victims and challenges in their implementation processes. The studies also identify coping strategies used by second victims as well as peer support experiences.

Studies have shown the existence of barriers to the implementation of and concerns about the use of these types of support interventions or tools by healthcare teams (concerns about confidentiality of programmes, time investment, reluctance to show vulnerability and ask for help, limited awareness of the second victim phenomenon, lack of financial resources and a culture of blame) [2].

Although support interventions have demonstrated utility for health organisations, health worker wellbeing and patient safety, there is still limited research on the elements necessary for effective implementation of these types of interventions in order to overcome difficulties in the implementation process and ensure the sustainability of these initiatives in health organisations.

Our study is based on the knowledge from the implementation science and health policy implementation, and on the empirical recommendations and stated preferences from the participants involved in the evidence-based programs/interventions. We believe that this study will contribute to the future implementation of these types of interventions in health organisations.

Objectives:

In this scoping review, we propose to map and frame the key factors which underlie an effective implementation of healthcare workers support interventions in health organisations when healthcare workers are physically and/or emotionally affected by patient safety incidents.

Key factors will be organized in 5 domains: operational attributes, organisational and contextual factors, relevant actors and healthcare workers recommendations.

Our study was based on the knowledge from the implementation science and health policy implementation, and on the empirical recommendations and stated preferences from the participants involved in the evidence-based programs/interventions. We believe that this study will contribute to the future implementation of these types of interventions in health organisations.

METHODS AND ANALYSIS

Our scoping review protocol is registered on the Open Science Framework (www.osf.io): DOI 10.17605/OSF.IO/2W46H

A scoping review is a valid and comprehensive approach with a rigorous and transparent method for mapping the evidence available in a specific area in order to clarify key characteristics or factors related to a concept [33,34].

The methodology of this scoping review will be guided by the established Arksey and O'Malley methodological framework [35] as enhanced by Levac et al. [36] and Joanna Briggs Institute (JBI) recommendations [34]. In this study, we will follow the six-stages of: (1) identifying the research question(s); (2) identifying relevant studies; (3) selecting studies; (4) charting the data; (5) collating, summarising and reporting the results; and (6) completing a consultation exercise.

This study will be guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis extension for Scoping Reviews (PRISMA-ScR) to ensure transparency of the obtained results [37].

To develop this scoping review, a multidisciplinary team composed of healthcare workers, researchers, academics and a qualified librarian will be used.

We will now describe the different stages of this scoping review according to the Arksey and O'Malley methodological framework [35] for scoping reviews.

Stage 1: Research questions

Main research question:

What are the key factors for an effective implementation of healthcare worker support interventions in health organisations after the occurrence of patient safety incidents?

Secondary research questions:

What are the organisational factors that contribute to an effective implementation of a second victim support programme/other types of support intervention in health organisations when healthcare workers are physically and/or emotionally affected by patient safety incidents?

What are the operational attributes of a second victim support programme/other types of support intervention in health organisations when healthcare workers are physically and/or emotionally affected by patient safety incidents?

Who are the most relevant actors in second victim support programmes/other types of support intervention in health organisations when healthcare workers are physically and/or emotionally affected by patient safety incidents?

What are the contextual factors for second victim support programmes/other types of support intervention in health organisations when healthcare workers are physically and/or emotionally affected by patient safety incidents?

Stage 2: Identification of the relevant literature

Identification process:

In this study, we will follow the JBI three-step process: (1) a preliminary search conducted on at least two databases; (2) the definition of clear inclusion criteria and the creation of a list of search terms to be used in the subsequent running of the search on a larger number of databases; (3) possible additional searches (cross-checking/cross-referencing of reference lists of potentially eligible studies, hand-searching in target journals relevant to the topic).

Preliminary literature search:

The preliminary search strategy was based on a literature review previously developed to clarify our inclusion and exclusion criteria. Our search strategy structure was guided by the Population, Concept and Context (PCC) elements of the inclusion criteria recommended by JBI for scoping reviews [34]. We want to achieve as much sensitivity as possible in the scope of our research.

The preliminary literature search was tested in two databases that are widely used in the Health Sciences: Pubmed/ MEDLINE and Web of Science Core Collection.

For the search strategy, we used Mesh and natural language terms.

In table 2, we present the search strategy used in Pubmed/ MEDLINE and Web of Science Core Collection.

Table 2 –Preliminary literature search applied to Pubmed/ MEDLINE and Web of Science Core Collection databases

Search strategies applied during the preliminary search			Results
<i>Context</i>	#1	"Health Services" OR "Health Facilities" OR "Healthcare" OR "Primary health care" OR "General Practice" OR "Family practice" OR "Ambulatory Care" OR "Nursing Care" OR "Family unit" OR "Hospitals"	
<i>Content</i>	#2	"Program evaluation" OR "Support program" OR "Peer support" OR "Support strategies" OR "Organizational factors" OR "Organizational culture"	
<i>Population</i>	#3	("Health personnel" OR "Physicians" OR "Nurses" OR "Doctor" OR "Practitioner" OR "Medical students" OR "Medical residents" OR "Healthcare providers" OR "Healthcare worker" OR "Healthcare staff") AND ("Error" OR "Near miss" OR "Adverse Event" OR "Clinical Error" OR "Medical error" OR "Second victim" OR "Wounded caregiver" OR "Wounded healer" OR "Secondary trauma")	
Complete search on Web of Science	#1 AND #2 AND #3	"Health Services" OR "Health Facilities" OR "Healthcare" OR "Primary health care" OR "General Practice" OR "Family practice" OR "Ambulatory Care" OR "Nursing Care" OR "Family unit" OR "Hospitals" (Topic) and "Program evaluation" OR "Support program" OR "Peer support" OR "Support strategies" OR "Organizational factors" OR "Organizational culture" (Topic) and "Health personnel" OR "Physicians" OR "Nurses" OR "Doctor" OR "Practitioner" OR "Medical students" OR "Medical residents" OR "Healthcare providers" OR "Healthcare worker" OR "Healthcare staff" (Topic) and Error OR Near miss OR Adverse Event OR Clinical Error OR Medical error OR Second victim OR Wounded caregiver OR Wounded healer OR Secondary trauma (Topic) and Web of Science Core Collection (Database)	781 results
Complete search on Pubmed (Medline)		(((((("Health Services"[MeSH Terms]) OR ("Health Services"[Title/Abstract])) OR (("Health Facilities"[MeSH Terms])) OR ("Health Facilities"[Title/Abstract])) OR ("healthcare"[Title/Abstract])) OR ("Primary health care"[MeSH Terms])) OR ("Primary health care"[Title/Abstract])) OR ("General Practice"[MeSH Terms])) OR ("General Practice"[Title/Abstract])) OR ("Family practice"[MeSH Terms])) OR ("Family practice"[Title/Abstract])) OR ("Ambulatory Care"[MeSH Terms])) OR ("Ambulatory Care"[Title/Abstract])) OR ("Nursing Care"[Title/Abstract])) OR ("Family unit"[Title/Abstract])) OR ("Hospitals"[Title/Abstract])) AND (("Program evaluation"[Title/Abstract]) OR ("Support program"[Title/Abstract]) OR ("Peer support"[Title/Abstract]) OR ("Support strategies"[Title/Abstract]) OR ("Organizational factors"[Title/Abstract]) OR ("Organizational culture"[MeSH Terms]) OR ("Organizational culture"[Title/Abstract])))) AND (("Health personnel"[MeSH Terms]) OR ("Health personnel"[Title/Abstract]) OR ("Physicians"[Title/Abstract]) OR ("Nurses"[Title/Abstract]) OR ("Doctor"[Title/Abstract]) OR ("Practitioner"[Title/Abstract]) OR ("Medical students"[Title/Abstract]) OR ("Medical residents"[Title/Abstract]) OR ("Healthcare providers"[Title/Abstract]) OR ("Healthcare worker"[Title/Abstract]) OR ("Healthcare staff"[Title/Abstract])))) AND ((Error[Title/Abstract]) OR ((Near miss[Title/Abstract]) OR ("Adverse Event"[Title/Abstract]) OR ("Clinical Error"[Title/Abstract]) OR ("Medical error"[MeSH Terms]) OR ("Medical error"[Title/Abstract]) OR ("Second victim"[Title/Abstract]) OR ("Wounded caregiver"[Title/Abstract]) OR ("Wounded healer"[Title/Abstract]) OR ("Secondary trauma"[Title/Abstract]))))	688 results
<i>Language</i>		no language filter/restraint will be applied	
<i>Period</i>		no period filter/restraint will be applied	
<i>Exclusion</i>		Article types not included: editorial , letter to the editor, cases series,	

<i>criteria</i>	case reports, narrative review, commentary
-----------------	--

Structured search strategy:

With the support of a qualified research librarian, we will undertake a comprehensive search strategy in relevant bibliographic databases, particularly in peer-reviewed journals and grey literature databases: Pubmed/Medline, Embase, CINAHL, Web of Science, Scopus, PsycInfo, Epistemonikos, Scielo, Cochrane Library and Open Grey.

We will also hand search in the reference list of the included articles, conference proceedings, institutional/organisational websites and networks repositories: The European Researchers' Network Working on Second Victims (ERNST) website, Segundas y Terceras Víctimas Proyecto de Investigación, SARS-CoV-2 (COVID-19) Second Victims, Center for Patient Safety, Second Victim Support (UK) website, KU Leuven Research – Second Victim in Health Care, ForYOU team website, AHRQ website and World Health Organization (WHO) website.

We will conduct the searches from March to May 2022.

Stage 3: Study selection

After applying the search strategy, results will be collated and exported to EndNote web.

Duplicates will automatically be removed for the further screening stages of study titles and abstracts of the eligible studies.

To reduce potential selection bias, studies will be independently screened by two authors. A third author can be involved in the case of disagreements.

To assure the maximum sensitivity of the screening process, the review team will do a pilot test with 30 randomly selected studies. We will screen this random sample using the agreed inclusion and exclusion criteria, discuss discrepancies and make any necessary modifications.

For full-text screening, we will note specific reasons for exclusion. Full texts will be included if at least two reviewers consider them eligible.

Screening results will be summarised and also graphically presented in a PRISMA flow chart (the number of excluded and included articles at each stage of the screening process will be reported).

Inclusion criteria:

In our scoping review, we will consider the following criteria related to the typology of the studies:

- Study design/characteristics: original articles, systematic reviews and meta-analyses, grey literature (thesis, other documents);
- Period: no time filter/restraint will be applied; and
- Languages: no language filter/restraint will be applied.

We next describe the inclusion criteria according to Population, Concept and Context (PCC) elements.

Population

In this study, our target population is support interventions for healthcare workers who are involved in patient safety incidents.

The eligible interventions are focused on healthcare workers, defined as ‘people engaged in the promotion, protection, care or improvement of the health of population’, including health professionals and residents and other allied health professionals such as technicians and supply workers [38].

We will also include studies that use the term ‘second victims’, who are defined as ‘healthcare providers who are involved in an unanticipated adverse patient event, medical error and/or a patient related injury and become victimized in the sense that the provider is traumatized by the event’ [14]. However, there has been a discussion over the last few years about the adequacy of this term [39]. Therefore, in this scoping review, we will consider different terms for defining the second victim that have already been used in some of the literature, such as ‘secondary trauma’ [40], ‘wounded caregiver’ [41] and ‘wounded healer’ [42].

Concept

The concept of interest in this review is second victim support interventions/other types of support interventions in health organisations in which healthcare workers are physically and/or emotionally affected by patient safety incidents.

In this study, we intend to explore different types of healthcare worker support interventions: programmes, courses, pilot experiences and other tools and resources.

The development and/or the implementation and/or the evaluation of these types of support initiatives should be described in order to be eligible for this study.

We will focus on five key domains to study the implementation of these types of interventions (Figure 1).

We will in particular consider the following aspects based on a health policy implementation and implementation sciences background:

- **Organisational factors**

a) organisational structures such as infrastructures, resources, tools, equipment, units and staffing levels of function for managing and delivering services, leadership structure/authority and organisational culture;

b) organisational processes such as organisational actions, procedures, recruitment criteria, training, programme implementation, communication processes and quality of interactions and coordination during programme implementation and dissemination as well as the sustainability of the practice;

c) organisational outcomes such as implementation measures, process quality measures and utilisation measures; and effectiveness measures that assess the attainment of an end state, achievement of an objective, or creation of an effect in the healthcare organisation;

- **Operational attributes of the interventions** (format/type of programme, accessibility, usability and confidentiality of the programme/intervention);
- **Relevant actors** (the individuals and organisations that actively participate in the implementation);
- **Contextual factors** (type of healthcare setting and cultural context); and
- **Preferences and recommendations for the support programmes** – health care workers/second victim preference features and recommendations to improve the interventions.

Context

We will include all types of healthcare settings from high-, middle- and low-income countries in order to create a worldwide map of healthcare worker support interventions after the occurrence of a patient safety incident. According to the CDC Field Epidemiology Manual of the Center for Disease Control and Prevention, 'healthcare settings represent a broad of services and places where healthcare occurs, including acute care hospitals, urgent care centers, rehabilitation centers, nursing homes and other long-term care facilities, specialized outpatient services (e.g., hemodialysis, dentistry,...), and outpatient surgery centers' [43].

Exclusion criteria:

In this study, we will exclude private offices and particular services due to their potential structural differences, specific contexts, and characteristics. Also, we will exclude the following types of publication: editorials, letters to the editor, cases series, case reports, narrative review, commentaries. A table of excluded evidence will be presented, with detailed reasons for their exclusion in the final scoping review

Stage 4: Charting the data

This phase intends to give a logical and descriptive summary of eligible study characteristics and results. Arksey & O'Malley [35] referred to this phase as a 'basic numerical analysis of the extent, nature and distribution of the studies included'. This will enable the alignment of study data with the objectives and questions of the scoping review [34].

A data extraction template will be created to show the characteristics of the eligible studies (table 3):

Table 3- Data extraction template for charting the data

Authors	Study authors
Year	Year of publication
Country	Country where the programme was developed
Population and setting	Study population and setting where study was developed
Aim/propose	Aim/propose of the study
Methodology	Study design and types of methods used (survey, interview, observation, experiments...)
Outcome measures	Main results assessed in the study
Key findings	Results that relate to the scoping review questions

The content of the template was developed by means of a preliminary exercise by the research team.

As recommended in the literature, a trial of the data extraction form was conducted by the reviewers with respect to at least three studies to ensure that all relevant information will be extracted from the eligible studies [34].

Quality assessment

There is an ongoing debate about the inclusion of an assessment phase related to the quality of eligible studies in scoping reviews [44]. Although this phase is not mandatory in this type of study [33,35], the absence of quality assessment is usually considered a methodological limitation. Some evidence shows that quality assessment can provide a better interpretation of

results and an understanding of how the research was conducted, can clarify practice implications and can help to identify potential gaps in the evidence and the need for future research [33].

We will use the Mixed Methods Appraisal Tool (MMAT) 2018 version to conduct a quality assessment of the eligible studies. This is a validated tool to appraise the methodological quality of five categories of studies: qualitative research, randomised controlled trials, non-randomised controlled trials, quantitative descriptive studies and mixed methods studies [45]. Two independent reviewers will analyse the methodological quality of the included studies. A third reviewer will be involved in cases of disagreement in the quality assessment.

Information from each checklist item of the MMAT will be reported in a table format, described as 'yes', 'no' and 'can't tell'.

Stage 5: Collating, summarising and reporting the results

After charting the data, each of the elements will be divided into different conceptual categories. Data will be collected in an Excel table using a descriptive content analysis. In addition, the results will be described in a narrative summary.

With respect to each of the included studies, we will extract the data as detailed in table 4.

Table 4- Data extraction template according to scoping review research questions

Domains	Detailed description
Author, Year	Authors of the study, year of the study's publication
Type of intervention	Type of intervention described in the study (peer support programme, online programme, workshops, other)
Year of implementation	Year that support intervention was implemented
Duration of the intervention	Duration of the support intervention (in months)
Target population of the intervention	Health workers or medical residents/students (nurses, physicians, other allied health professionals such as technicians and supply workers,...)
Organisational factors	<p>Some examples to consider in data extraction:</p> <ul style="list-style-type: none"> a) organisational structures <ul style="list-style-type: none"> • Infrastructures (size of organisational or units, facilities, number of beds, ...) • Resources, tools, equipment • Staffing characteristics and integrated networks • Leadership structure • Organisational culture, work environment/organisational climate b) organisational processes <ul style="list-style-type: none"> • organisational procedures • communication processes, interactions, roles and responsibilities • management processes (practice arrangements, service coordination during programme implementation and follow-up) • dissemination and sustainability of the practice c) organisational outcomes <ul style="list-style-type: none"> • process quality measures • utilisation measures • effectiveness measures • global health status measures • efficiency measures • costs

Operational attributes of the programmes/interventions	Some examples to consider in data extraction: <ul style="list-style-type: none"> • intervention type • accessibility • usability • confidentiality issues • other relevant attributes
Relevant actors	Individuals and organisations that make up and implement the programmes
Contextual factors	Type of healthcare setting, healthcare services and units, cultural context
Healthcare workers/second victim recommendations	Participant preference features and recommendations for the support intervention

The content of the template was developed through a preliminary exercise by all the research team members. Data extraction can be updated during the data extraction process.

After collecting information in an Excel table (analytical framework), the information will be thematically organised according to the support intervention type provided to healthcare workers after the occurrence of an incident. In this phase, we will present information in a diagram using support intervention type as a primary unit of analysis.

Stage 6: Consultation exercise and stakeholder involvement

A consultation exercise has been suggested in the literature [35,36] as an opportunity to involve key stakeholders as an additional source of information to complement the literature search. This will bring a high level of meaning content and will enlarge the scope of the review. In this phase, we will involve some of the members of the European Researchers Network Working on Second Victims (ERNST) - COST Action 19113.

Ethics and dissemination:

This scoping review does not require ethical approval. Results and conclusions of the study will be published in a peer-reviewed journal and presented in scientific conferences as well as in international forums and through other relevant dissemination channels.

Author contributions:

SGP was involved in the writing of the article. SGP, MJL, PS and JJM contributed to the conceptualisation of the study. SGP, MJL and JDS analysed the data extraction templates, participated in the pilot testing stage and defined the inclusion criteria. HD contributed to the definition of a robust search strategy and participated in the preliminary search in relevant bibliographic databases, particularly in peer-reviewed journals and grey literature databases. PS, JJM and IC reviewed the article and contributed to the literature review. All authors read and approved the manuscript.

Funding statement:

This work was supported by FCT- Fundação para a Ciência e Tecnologia, I. P. under the PhD grant UI/BD/150875/2021. It is also partly supported by Comprehensive Health Research Centre (CHRC) and National School of Public Health, NOVA University of Lisbon (ENSP-NOVA), under the funding support for publications.

Competing interest statement:

This protocol is included in the PhD project of the corresponding author (SGP) previously approved by the National School of Public Health (ENSP- NOVA). Some of the research team members are part of the European Researchers Network Working on Second Victims (ERNST) - COST Action 19113. All the authors have completed the disclosure of interest form.

References

1. World Health Organization. Patient safety incident reporting and learning systems: technical report and guidance. Geneva: *World Health Organization*; 2020:1-51. <https://apps.who.int/iris/handle/10665/334323>.
2. Busch IM, Moretti F, Campagna I, et al. Promoting the Psychological Well-Being of Healthcare Providers Facing the Burden of Adverse Events: A Systematic Review of Second Victim Support Resources. *Int J Environ Res Public Health* 2021;18(10):5080. doi:10.3390/ijerph18105080.
3. NHS England. Report a patient safety incident. Accessed Jan 2021 <https://www.england.nhs.uk/patient-safety/report-patient-safety-incident/>.
4. Rafter N, Hickey A, Condell S, et al. Adverse events in healthcare: learning from mistakes. *QJM* 2014;108(4):273–7. <https://doi.org/10.1093/qjmed/hcu145>
5. Brennan TA, Hebert L, Localio AR, et al. Incidence of Adverse Events and Negligence in Hospitalized Patients. *N Engl J Med* 1991;324:370-376. doi: 10.1056/NEJM199102073240604.
6. Aaraaen A, Slawomirski L, Klazinga N. The economics of patient safety in primary and ambulatory care: Flying blind. OECD Health Working Papers, No. 106. Paris: *OECD Publishing*; 2018:106. Accessed Nov 2021 https://www.oecd-ilibrary.org/social-issues-migration-health/the-economics-of-patient-safety-in-primary-and-ambulatory-care_baf425ad-en.
7. Panagioti M, Khan K, Keers RN, et al. Prevalence, severity, and nature of preventable patient harm across medical care settings: systematic review and meta-analysis. *BMJ* 2019;366:l4185. doi: <https://doi.org/10.1136/bmj.l4185>.
8. Wu AW. Medical error: the second victim. The doctor who makes the mistake needs help too. *BMJ* 2000;320(7237):726-727. doi:10.1136/bmj.320.7237.726.

9. Ullström S, Andreen Sachs M, Hansson J, et al. Suffering in silence: a qualitative study of second victims of adverse events. *BMJ Qual Saf* 2014;23(4):325–31. doi: 10.1136/bmjqs-2013-002035.
10. Waterman AD, Garbutt J, Hazel E, et al. The Emotional Impact of Medical Errors on Practicing Physicians in the United States and Canada. *Jt Comm J Qual Patient Saf* 2007;33(8):467–76. doi: 10.1016/s1553-7250(07)33050-x.
11. Rinaldi C, Leigheb F, Vanhaecht K, et al. Becoming a “second victim” in health care: Pathway of recovery after adverse event. *Rev Calif Asist* 2016;31:11–9. doi: 10.1016/j.cali.2016.05.001.
12. Vogus TJ, Ramanujam R, Novikov Z, et al. Adverse Events and Burnout: The Moderating Effects of Workgroup Identification and Safety Climate. *Med Care* 2020;58(7):594–600. doi: 10.1097/MLR.0000000000001341.
13. Ozeke O, Ozeke V, Coskun O, et al. Second victims in health care: current perspectives. *Adv Med Educ Pract* 2019;10:593–603. doi: 10.2147/AMEP.S185912.
14. Scott SD, Hirschinger LE, Cox KR, et al. The natural history of recovery for the healthcare provider “second victim” after adverse patient events. *Qual Saf Health Care* 2009;18(5):325–30. doi: 10.1136/qshc.2009.032870.
15. Reis CT, Paiva SG, Sousa P. The patient safety culture: a systematic review by characteristics of Hospital Survey on Patient Safety Culture dimensions. *Int J Qual Health Care* 2018;30(9):660–677. doi: 10.1093/intqhc/mzy080.
16. Quillivan RR, Burlison JD, Browne EK, et al. Patient Safety Culture and the Second Victim Phenomenon: Connecting Culture to Staff Distress in Nurses. *Jt Comm J Qual Patient Saf* 2016;42(8):377–86. doi: 10.1016/s1553-7250(16)42053-2.
17. Quadrado ERS, Tronchin DMR, Maia FOM. Strategies to support health professionals in the condition of second victim: scoping review. *Rev Esc Enferm USP* 2021;55:e03669. <https://doi.org/10.1590/S1980-220X2019011803669>.
18. Mira J, Carrillo I, Guilbert M, et al. The Second Victim Phenomenon After a Clinical Error: The Design and Evaluation of a Website to Reduce Caregivers’ Emotional Responses After a Clinical Error. *J Med Internet Res* 2017;19(6):e203. doi: 10.2196/jmir.7840.
19. Edrees H, Connors C, Paine L, et al. Implementing the RISE second victim support programme at the Johns Hopkins Hospital: a case study. *BMJ Open* 2016;6(9):e011708. doi: 10.1136/bmjopen-2016-011708.
20. University of Missouri Health Care. forYOU Team 2021. Assessed Dec 2021 <https://www.muhealth.org/about-us/quality-care-patient-safety/office-of-clinical-effectiveness/foryou>.
21. El Hechi MW, Bohnen JD, Westfal M, et al. Design and Impact of a Novel Surgery-Specific Second Victim Peer Support Program. *J Am Coll Surg* 2020;230(6):926–33. doi: 10.1016/j.jamcollsurg.2019.10.015.
22. Wijaya MI, Mohamad AR, Hafizurrachman M. Second victim support program and patient safety culture: A quasi experimental study in Bali International Medical Centre (BIMC) Hospital. *Bali Med J* 2018;7(1):220–226. doi: 10.15562/bmj.v7i1.952.
23. Jones C, Robertson N, Chamdal V, et al. Second Victim Support Unit Scoping Project Final Report. AHSN Network Patient Safety. England: *NHS Improvement*; 2019:1–19.
24. Donabedian A. Evaluating the Quality of Medical Care. *Milbank Q* 2005;83(4):691–729. doi: 10.1111/j.1468-0009.2005.00397.x.
25. Yano EM. The role of organizational research in implementing evidence-based practice: QUERI Series. *Implement Sci* 2008;3(1):29. doi: 10.1186/1748-5908-3-29.
26. San Martín-Rodríguez L, Beaulieu MD, D’Amour D, et al. The determinants of successful collaboration: A review of theoretical and empirical studies. *J Interprof Care* 2005;19(sup1):132–47. doi: 10.1080/13561820500082677.
27. Donabedian A. Basic approaches to assessment: structure, process and

- outcome. In: Explorations in quality assessment and monitoring. Volume 1, The definition of quality and approaches to its assessment. Ann Arbor, Michigan: *Health Administration Press* 1981:77–128.
28. Buse K, Mays N, Walt G. The health policy framework: Context, process and actors. In: Black N, Raine R, editors. *Making Health Policy*. 1st edition. Maidenhead, Berkshire: McGraw-Hill Education: *University Press* 2005.
29. Harvard Catalyst Profiles. Epidemiologic Factors. Harvard University 2022. Accessed Dec 2021 <https://connects.catalyst.harvard.edu/Profiles/display/Concept/Epidemiologic%20Factors>.
30. Cook S, Mayers S, Goggins K, et al. Assessing research participant preferences for receiving study results. *J Clin Transl Sci* 2020;4(3):243–9. doi: 10.1017/cts.2019.427.
31. Wade L, Fitzpatrick E, Williams N, et al. Organizational Interventions to Support Second Victims in Acute Care Settings: A Scoping Study. *J Patient Saf* 2022;18(1):e61–e72. doi: 10.1097/PTS.0000000000000704.
32. Busch IM, Moretti F, Purgato M, et al. Dealing With Adverse Events: A Meta-analysis on Second Victims' Coping Strategies. *J Patient Saf* 2020;16(2):e51–e60. doi: 10.1097/PTS.0000000000000661.
33. Munn Z, Peters MDJ, Stern C, et al. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018;18(1):143. <https://doi.org/10.1186/s12874-018-0611-x>.
34. The Joanna Briggs Institute. The Joanna Briggs Institute Reviewers' Manual 2015 Methodology for JBI Scoping Reviews. South Australia: *The Joanna Briggs Institute*; 2015:1–24.
35. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8(1):19–32. <https://doi.org/10.1080/1364557032000119616>.
36. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5(1):69. <https://doi.org/10.1186/1748-5908-5-69>.
37. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169(7):467–73. doi: 10.7326/M18-0850.
38. Bragazzi NL, Dini G, Parodi V, et al. Protocol of a scoping review assessing injury rates and their determinants among healthcare workers in western countries. *BMJ Open* 2019;9(1):e023372. doi: 10.1136/bmjopen-2018-023372.
39. Clarkson MD, Haskell H, Hemmelgarn C, et al. Abandon the term “second victim”. *BMJ* 2019;364:l1233. doi: <https://doi.org/10.1136/bmj.l1233>.
40. Von Rueden KT, Hinderer KA, McQuillan KA, et al. Secondary Traumatic Stress in Trauma Nurses: Prevalence and Exposure, Coping, and Personal/Environmental Characteristics. *J Trauma Nurs* 2010;17(4):191–200. doi: 10.1097/JTN.0b013e3181ff2607.
41. Lents MB. The Wounded Caregiver: Understanding and Supporting the Second Victim. AORN Global Surgical Conference & Expo 2020. Accessed Nov 2021 <https://www.aorn.org/surgicalexpo/articles-and-resources/wounded-caregiver>.
42. Conti-O'Hare M. The Nurse as Wounded Healer: From Trauma to Transcendence. Massachusetts: *Jones & Bartlett Publishers*; 2002:1–196.
43. Christensen BE, Fagan RP. Healthcare Settings. In: The CDC Field Epidemiology Manual [Internet]. Atlanta, Georgia, United States: *Center for Disease Control and Prevention* 2018. Accessed Nov 2021 <https://www.cdc.gov/eis/field-epi-manual/chapters/Healthcare-Settings.html>.
44. Pham MT, Rajić A, Greig JD, et al. A scoping review of scoping reviews: advancing the approach and enhancing the consistency. *Res Synth Methods* 2014;5(4):371–85. doi: 10.1002/jrsm.1123.
45. Hong Q, Pluye P, Fàbregues S, et al. Mixed Methods Appraisal Tool (MMAT), version 2018: user guide. Quebec: *Mc Gill Department of Family Medicine*; 2018:1–10.

Legend of figure 1:

Figure 1: Five key domains for effective implementation of healthcare workers support interventions in health organisations