Prevalence mapping of Schistosomiasis among Pre-school aged children in Rwanda

Bayingana Jean Bosco^{1,2}, Nyandwi Elias³, Ntakarutimana Amans², Kagabo Joseph², Shema Eliah², Kanimba Philbert², Mbonigaba Jean Bosco⁴, Ruberanziza Eugene⁴ & Rujeni Nadine^{2#}

¹ University of Rwanda, Single Project Implementation Unit

² University of Rwanda, College of Medicine and Health Sciences, School of Health Sciences

³ University of Rwanda, College of Science and Technology, Centre for Geographic Information Systems and Remote Sensing (CGIS)

⁴ Ministry of Health, Rwanda Biomedical Center, NTDs & Other Parasitic Diseases Unit

#Corresponding author email address: nrujeni@gmail.com

Abstract

Schistosomiasis is endemic in Rwanda and control programs have been implemented with a special focus on school-aged children (SAC) ignoring pre-school aged children (pre-SAC) for which the actual prevalence of the disease is not well established. This study consisted of a cross-sectional quantitative mapping of the distribution of schistosomiasis and identification of associated risk factors among pre-SAC throughout the country. The study covered all the 17 Districts of Rwanda endemic for schistosomiasis with a total sample of 4675 children enrolled from 80 purposively selected villages. The Parasitological assessment of children's urine and stool samples was conducted using CCA and Kato Katz methods respectively for infection detection. A standard questionnaire was used to collect data on the risk factors and geospatial assessment was performed using tablets and GPS to record geographic coordinates for plotting locations on maps using ArcGIS software.

The overall prevalence of *S. mansoni* infection across the surveyed areas was 24% and 0.8 by CCA and Kato-Katz, respectively. Infection was significantly associated with bathing children in open water bodies. Furthermore, pre-SAC looked after by siblings (sisters) were twice as much likely to be infected compared to those looked after by mothers. Schistosomiasis control interventions are needed for pre-SAC to limit their exposure to open water bodies with expectations of adapted chemotherapy to be availed. Community based deworming campaigns may be the best way to ensure good treatment coverage of pre-SAC in Rwanda.

Key words: Schistosomiasis, pre-school aged children, risk factors, *Schistosoma mansoni*, Rwanda

39 Author summary

40 Schistosomiasis is one of the Neglected Tropical Diseases (NTD) of public health concern in
 41 Rwanda like in many tropical countries. The recently published NTD roadmap by the World
 42 Health Organization indicates that schistosomiasis is targeted for elimination as a public
 43 health problem worldwide by 2030. For this target to be achieved, all at risk populations
 44 should be reached by control programs' interventions such as preventive chemotherapy,
 45 health education as well as water, sanitation and hygiene (WASH) practices. However, pre-
 46 school children (pre-SAC) are among populations at risk for whom infection burden is not
 47 fully documented. Implementation of the recent WHO guideline on schistosomiasis control
 48 and elimination that recommend inclusion of pre-SAC in control programs will be informed
 49 by detailed assessment of the infection burden and distribution in endemic countries. This
 50 study is showing for the first time a high burden of schistosomiasis among pre-SAC in high-
 51 risk areas of Rwanda. With the imminent availability of the paediatric formulation of
 52 Praziquantel (the drug of choice against the disease), the findings should guide the country in
 53 implementation strategies that include these children in mass deworming. The findings also
 54 highlight a number of risk factors including the passive exposure of these young children by
 55 their siblings while bathing them in open waters and the lack of knowledge of
 56 parents/guardians on the disease. Additional integrated interventions such as health
 57 education, improvement of water supply and sanitation as well as snail surveillance will lead
 58 to more sustainable solutions in the march towards schistosomiasis elimination.

Introduction

Schistosomiasis is a chronic parasitic infection that is caused by trematodes of the genus *Schistosoma*, transmitted through fresh water snails. It can cause vascular damage in individuals and is associated with growth retardation and impaired metabolism and cognition (1,2). Schistosomiasis remains a public health concern and highly prevalent in low and middle-income countries, including Rwanda, mostly affecting poor communities with no access to potable water (3). Globally, more than 250 million people are affected by schistosomiasis and Sub Saharan Africa accounts for 90% of all these infection burdens (4,5). Control interventions mainly rely on regular targeted treatments (using Praziquantel) of populations in endemic areas coupled with health education. Although the disease is targeted (as per the WHO roadmap) for elimination as a public health problem (defined as prevalence of <1% of heavy intensity schistosomiasis infections) by 2030 (6), control interventions do not currently reach all populations in need in most endemic areas.

For many years, pre-school aged children (pre-SAC) have been left out in most schistosomiasis treatment programs. This was partly due to the assumption that they are not at great risk of contracting schistosomiasis. However, considerable advances have been made in this regard and a number of studies have shown that most of the pre-SAC in endemic areas are significantly exposed to schistosomiasis in many countries (7–14). This has prompted the WHO to formulate a new guideline including recommendations to include pre-SAC aged 2 years+ in preventive chemotherapy while those below 2 years should be treated on individual basis (15).

With the realization that pre-SAC are in need of treatment, came a number of challenges including the absence of an appropriate child friendly formulation of Praziquantel and the lack of information on the real burden of schistosomiasis in this young age group in some countries. Indeed, inclusion of pre-SAC in schistosomiasis control programs should be justified by quantification of the burden of infection in that age group in each country for optimal intervention. Furthermore, an assessment of their accessibility and local acceptability should guide implementation strategies.

Rwanda is among the countries where the burden of schistosomiasis among pre-SAC is not well documented. Our own study on Nkombo Island in the western part of the country reported an overall prevalence of 9.5% (9), but a countrywide assessment of the disease in this age group is necessary to better inform the control program. Nevertheless, the Rwandan Ministry of health has recently introduced a community-based deworming strategy whereby community health workers (CHWs) and community leaders coordinate MDAs with minimal supervision from the central level. Such a deworming strategy should be suitable for treating schistosomiasis among pre-SAC.

In light of the above, this study was conducted to estimate the prevalence, identify risk factors and produce an illustrative, countrywide map of schistosomiasis distribution in all moderate to high - risk areas of the country.

Methods

Study design, population, and sample size

This was a cross-sectional quantitative study that involved parasitological assessment of schistosomiasis among children aged 7 to 59 months. Structured interviews were also held with participants' parents/caregivers to assess risk factors of infection in this young population. The study covered all the 17 Districts of Rwanda at moderate to high risk of schistosomiasis (16,17) and targeted 40 Sectors reported with a prevalence of 10% and above among school children (based on incidence and mapping data). From each targeted sector, 2 villages were selected based on their proximity to existing water body and/or wetlands. The villages layer was overlaid with layers of lakes, multipurpose water dam, fish pond and/or important wetlands (wetland of ≥ 0.7 ha hosting a socio-economic activity, mostly irrigated agriculture). From eligible villages (risky villages), final selection was purposively decided considering the spatial distribution pattern analysed visually by researchers and complemented with field verification, making 80 study villages with proximity to open water sources and wetlands. A total of 4675 children were enrolled from the selected villages.

Parasitological assessment

Urine samples were collected and tested for schistosomiasis circulating antigen using the Point-Of-Care Circulating Cathodic Antigen (POC-CCA or CCA) following manufacturer's instructions. CCA test results were recorded as negative (- or trace) or positive (+; ++ or +++ according to the intensity of the test line in comparison to a test control). Stool samples were also collected and tested using the Kato Katz method for the detection of patient infection following published protocols (18). A single specimen was collected from which 2 slides were prepared and read by 2 laboratory technicians independently. Ten per cent (10%) of all

slides were re-tested by the National Reference Laboratory senior technicians for quality control.

Risk factors and spatial assessment

A standard questionnaire was designed in Kobo toolbox and used to collect data from parents/guardians by a well-trained data collector using a tablet. Geographic coordinates were recorded using tablets and ArcGIS software version 10.4 was used to produce maps.

Statistical analysis

Data collected were analysed by Stata 13 (STATA Corp, Lakeway, College Station, Texas, USA). Descriptive analysis was conducted. Categorical variables were presented as frequencies and their respective percentages. Numerical variables were presented as mean and standard deviations. For comparison purposes, cross-tabulation analyses were conducted and contingency tables were produced, while chi-square tests were used to compare proportions. To assess the association between variables, logistic regression was performed. For binary logistic regression, each risk factor was assessed with the outcome (infection status) separately while for multivariable logistic regression, a backward stepwise logistic regression was run and all risk factors with a p-value less or equal to 0.05 were considered. Odds ratios and the corresponding 95% confidence interval (CI) were reported. A two-tailed significance level of 0.05 was considered.

Ethical consideration

The current study was approved by the Institutional Review Board (IRB) of the University of Rwanda College of Medicine and Health Sciences (Approval Notice No 148/CMHS/IRB/2019) and endorsed by the Ministry of Health. Local authorities, community

health workers and parents/guardians were informed about the aims, voluntary nature of their participation, potential risks and benefits of the study. A signed consent was obtained from parents/guardians before enrolling their child(ren) in the study.

Results

The results of this research are presented by study sites and do not represent the administrative District. Study sites are those sites in proximity with open water bodies and vary in number per District from one to six. These sites are the base for defining the targeted areas of control interventions.

Demographic characteristics of the study participants

The total sample was 4675 pre-SAC with a mean age of 37.5 months (± 13.0), and a sex ratio M:F of 1.01: 1 (Table 1). Majority of the study participants (corresponding to 96.2%) were born in their respective villages (permanent residents).

172 **Table 1.** Characteristics of pre-school aged children enrolled in the study

	n	%
Age in months (Mean, SD)	(37.5, 13.0)	
Age categories (in months)		
12 months and less	73	1.6
13-24 months	842	18
25-36 months	1230	26.3
37-48 months	1359	29.1
49 months and above	1171	25
Total	4675	100
Sex		
Female	2317	49.6
Male	2358	50.4
Total	4675	100
Child born in village		
No	177	3.8
Yes	4498	96.2
Total	4675	100

173

174

Water contact activities (exposure) for children in study areas

More than 90% parents reported farming as their occupation and 83.3% (n= 3895) of the families use open water sources (lake, dam or swamp) for daily domestic activities. This result in more than 60% of enrolled children being regularly in contact with open water sources (Table 2).

Table 2. Water contact activities for parents and their children in the study areas

	n	%
Parent's occupation		
Casual worker	143	3.18
Farmer	4,060	90.26
Government employee	52	1.16
Self employed	157	3.49
Unemployed	86	1.91
Total	4,498	100
Fetching water in the open water source		
No	780	16.7
Yes	3895	83.3
Total	4675	100
Child taken to open water source		
No	1444	30.9
Yes	3222	69.1
Total	4666	100
Frequency child taken to lake		
Once a day	2055	63.4
Once a week	824	25.4
Once a month	185	5.7
Once a year	53	1.6
Once in his life	123	3.8
Total	3240	100
Child bathed in the lake		
No	1669	36.2
Yes	2937	63.8
Total	4606	100
Frequency child bathed IN the lake		
Once a day	1943	65.2

Once a week	761	25.5
Once a month	155	5.2
Once a year	38	1.3
Once in his life	83	2.8
Total	2980	100
Child ever bathed using FETCHED lake water		
No	947	20.3
Yes	3728	79.7
Total	4675	100

Prevalence and infection intensity of schistosomiasis among pre-school aged children

Prevalence of schistosomiasis among pre-school aged children was assessed using both KK and POC-CCA. The POC-CCA results yielded a prevalence of 24% among children tested (trace results considered negative), while the KK only yielded a 0.8% prevalence. The POC-CCA estimate was considered as the overall prevalence given the high sensitivity of the assay (16). For infection intensity, KK results were considered and the raw faecal worm egg count (FWEC) was multiplied by 24 to estimate the eggs per gram (epg) (18,19). Based on WHO guidelines (20), 16.2% (n= 6) of the 37 children who tested positive on Kato Katz had moderate infection while the rest had light infection (Table 3).

205 **Table 3. Prevalence and infection intensity of the study population**

	n	%
POC-CCA test		
Negative	3554	76
Positive	1121	24
Total	4675	100
KK test results		
Negative	4638	99.2
Positive	37	0.8
Total	4675	100
Intensity of infection		
Light	31	83.8
Moderate	6	16.2
Heavy	0	0
Total	37	100

206

207

Prevalence of schistosomiasis by age, gender, and location among pre-school children

The results on the prevalence of schistosomiasis infection by gender revealed no statistically significant difference in infection between males and females. However, a statistically significant difference in infection was observed across age groups (Table 4).

Table 4. Infection prevalence per age category and gender

Age		CCA – traces considered negative		KK	
		Negative	Positive	Negative	Positive
≤12 months	n	52	21	73	0
	%	1.5	1.9	1.6	0.0
13-24 months	n	617	225	841	1
	%	17.4	20.1	18.1	2.7
25-36 months	n	915	315	1222	8
	%	25.7	28.1	26.3	21.6
37-48 months	n	1068	291	1346	13
	%	30.1	26.0	29.0	35.1
≥ 49 months	n	902	269	1156	15
	%	25.4	24.0	24.9	40.5
		χ^2 (P-value) = 11.7493 (0.019)		χ^2 (P-value) = 9.7921 (0.044)	
Gender					
Female	n	1782	535	2298	19
	%	50.1	47.7	49.5	51.4
Male	n	1772	586	2340	18
	%	49.9	52.3	50.5	48.6
		χ^2 (P-value) = 1.9890 (0.158)		χ^2 (P-value) = 0.0478 (0.827)	

There were significant differences in infection prevalence across study sites, Gisagara District having the highest prevalence with 60.4%, followed by Nyagatare and Gasabo Districts respectively with 46.8% and 40%, while Kamonyi District had the lowest prevalence (7.1%). Infection distribution across study sites is illustrated on Fig 1.

Factors associated with Schistosomiasis infection

The logistic regression performed highlighted the age as a contributing factor to infection status as expected. In addition, two behavioural factors were significantly associated with the occurrence of infection: children taken care of by a sister and those who are bathed in the lake/pond are approximately 2 times more at risk of being infected (OR = 1.9, $p < 0.0001$ and OR=1.8, $p < 0.0001$ respectively) compared to their counterparts.

Table 6. Association between behavioural factors and schistosomiasis

CCA_Results	Odds Ratio	Std. Err.	P	[95% Conf. Interval]
Age group				
≤ 12 months	REF			
13-24 months	0.800925	0.209265	0.3960	0.479945 1.336572
25-36 months	0.7182087	0.185582	0.2000	0.432816 1.191784
37-48 months	0.5782747	0.149883	0.0350	0.347946 0.961074
≥ 49 months	0.6029743	0.157244	0.0520	0.361678 1.005253
Child carer				
Mum	REF			
Dad	0.8052959	0.133487	0.1910	0.581915 1.114427
Sister	1.905744	0.323173	0.0000	1.366845 2.657111
Brother	0.3749097	0.200035	0.0660	0.131754 1.066813
Other	0.5014567	0.05911	0.0000	0.398012 0.631787
Child bathed in Lake				
No	REF			
Yes	1.825002	0.147079	0.0000	1.558347 2.137286

Knowledge on Schistosomiasis and perception on community based deworming

Approximately 35% (n=1629) of parents/ caregivers had never heard of schistosomiasis (or Bilharzia) before. However, out of the 155 community health workers (CHWs) interviewed, 98.7% reported knowing the disease by name and 96.1% knew that it is contaminated through contact with water bodies and that pre-SAC are also at risk of contamination (Table 7).

Table 7. CHWs' general knowledge on schistosomiasis

	n	%
Ever heard of Bilharzia		
No	2	1.3
Yes	153	98.7
Total	155	100
How is schistosomiasis/bilharzia acquired?		
Don't know	6	3.9
Contact with water bodies	148	96.1
Total	154	100
Can children under 5 years be infected?		
No	1	0.6
Yes	151	97.4
Don't Know	3	1.9
Total	155	100

Regarding the implementation of a community-based mass drug administration (MDA), 62.1% (n=95) of CHWs reported that this is implemented at their respective villages, while 37.9% (n=58) reported that it is not. However, of those (n=58, i.e. 37.9%) who reported that MDA is not yet implemented at their villages, 87.9% (n=51) strongly support the idea of introducing it in their villages. Majority of the CHWs (99.3%, n=136) reported that they are willing to be involved in the implementation of community-based MDA campaigns in their village and 77.9% (n=113) suggested that the village leader would be the best person to coordinate this activity. CHWs also acknowledged some advantages of a community-based MDA such as full treatment coverage, time saving and an opportunity for health education

260 (Table 8). However, some challenges were also highlighted such as the lack of distribution
261 spaces, poor management of drugs and poor knowledge of CHWs at the village level.

Discussion

The present study aimed to estimate the burden of schistosomiasis among pre-SAC in Rwanda and risk factors for the infection as well as to map the distribution of infection throughout the country for that category of population. This is the first comprehensive mapping exercise conducted in this young age group that has long been neglected by the Rwanda schistosomiasis control program. The study covered all the 17 Districts that had been highlighted by the previous school-based mapping as endemic areas (16,17). The 80 study villages as minimum units of observation (sampling units) within these endemic Districts were purposively selected for their proximity to open water bodies to allow for the focal nature of the disease distribution. From the targeted districts, the results are presented at the level of study sites which have been extrapolated to administrative sectors based on the layout of open water bodies, environmental sanitary conditions and exposed population. Overall, the prevalence of schistosomiasis was 24% using the CCA diagnostic method and 0.8% using the KK method. This discrepancy was expected as the KK has shown a low sensitivity (21), especially in pre-SAC who are most likely carrying pre-patent and/or light infections (22). Nevertheless, the CCA prevalence detected was comparable to findings among SAC in the same areas (16), highlighting an urgent need to include these children in treatment campaigns. As expected, the distribution of infection was skewed, few Sectors harbouring most of the infection burden as highlighted by the significant differences between sites (i.e 60.5%, maximum and 7.1%, minimum). In agreement with the previous mapping among SAC (16,17), sectors of Nyagatare and Gisagara Districts showed the highest infection prevalence (in 4 sectors the prevalence exceeded 50%) but also the Districts of Ngoma, Bugesera, Gatsibo, Kayonza, Nyamasheke, Gicumbi and Karongi were heavily burdened. With regard to infection intensity, and based on WHO criteria (20), most of the pre-SAC showed light to moderate infection intensities.

288

289 In this study, 83.3% of parents/guardians fetched their water for domestic use from open
290 water sources and 63.4% confirmed taking their children to these water sites every day. Most
291 importantly, 63.8% of the parents/guardians interviewed bath their children in open water
292 bodies and our logistic regression analysis indicate that their children are twice as much
293 likely to be infected compared to those who are not bathed in these waters. Previous studies
294 elsewhere have also reported this passive nature of exposure to schistosomiasis among pre-
295 SAC (11,23,24). Interestingly, our logistic regression highlights that children who are looked
296 after by a sibling (a sister in this case) –most likely while their parents conduct
297 domestic/professional chores- are more likely to be infected compared to those who are
298 constantly with their mothers. Although we were not able to conduct a thorough
299 observational investigation, these findings suggest that older sisters play and/or bath their
300 younger siblings at water contact points that are far enough from where mothers wash
301 dishes/clothes, and therefore far from detergents that may destroy cercaria (25). The latter
302 may protect –to some extent- those children who are constantly with their mothers.

303

304 At the time of the study, 65% of the parents/caregivers had some vague information on
305 schistosomiasis “*yes I heard about schistosomiasis*” but no extensive assessment of the level
306 of knowledge was done. Nevertheless, more than 90% of CHWs reported that
307 “*schistosomiasis is transmitted through open waters*”. The study was however limited in
308 terms of depth of health education assessment as the main focus was on screening infection in
309 the child and interviewing the caregivers to assess exposure risk factors. Nevertheless, given
310 the findings on infection prevalence, schistosomiasis control interventions are specifically
311 needed that focus on health education for household members’ behaviour change and
312 practices to (a) limit the children’s exposure to open water bodies, and (b) improve children’s

environmental sanitary conditions (26). Additional interventions such as provision of potable water supplies to these vulnerable communities as well as public latrines may provide longer term solutions. Indeed, more than 90% of the interviewed parents/guardians are farmers and most of our study sites were in close proximity to wetlands that are exploited for agriculture. Farmers generally spend several hours in the fields and, in the absence of latrines, are likely to defecate in these fields and contaminate the water bodies that are later used for bathing children.

The Rwandan Ministry of health has recently started implementing a community-based deworming that take place during the mother-child health (MCH) week. Our study involved interviews of CHWs to find out how they perceive this deworming approach and findings indicate that they strongly support it and highlight several advantages. This should be the best way to reach pre-SAC once the pediatric formulation of Praziquantel is availed.

Taken together, the present study findings stress that pre-SAC in Rwanda are part of the population at risk of schistosomiasis and control interventions of the disease in Rwanda must include them. In particular, we provide a detailed map of pre-SAC infection distribution that should guide intervention strategies countrywide. Further assessment of the parents/CHW knowledge, attitudes and practices towards schistosomiasis and WASH recommended practices is warranted.

Acknowledgements

This research was commissioned by the National Institute of Health Research (NIHR) Global Health Research programme (16/136/33) using UK aid from the UK Government. The authors would like to extend thanks to the study participants, the University of Rwanda, Rwanda Biomedical Center, Ministry of Health, district hospitals and health centers for all facilitations to the end of this research activity.

Conflict of interest

None

References

1. Gasparotto J, Senger MR, Telles de Sá Moreira E, Brum PO, Carazza Kessler FG, Peixoto DO, et al. Neurological impairment caused by *Schistosoma mansoni* systemic infection exhibits early features of idiopathic neurodegenerative disease. *The Journal of biological chemistry*. 2021;100979.
2. Osakunor DNM, Mduluzi T, Osei-Hyiaman D, Burgess K, Woolhouse MEJ, Mutapi F. *Schistosoma haematobium* infection is associated with alterations in energy and purine-related metabolism in preschool-aged children. *PLoS neglected tropical diseases*. 2020;14(12):e0008866.
3. Rujeni N, Morona D, Ruberanziza E, Mazigo HD. Schistosomiasis and soil-transmitted helminthiasis in Rwanda: An update on their epidemiology and control. *Infectious Diseases of Poverty*. 2017;6(1).
4. Engels D, Chitsulo L, Montresor A, Savioli L. The global epidemiological situation of schistosomiasis and new approaches to control and research. *Acta Tropica*. 2002;82(2):139–46.
5. Sokolow SH, Wood CL, Jones IJ, Swartz SJ, Lopez M, Hsieh MH, et al. Global Assessment of Schistosomiasis Control Over the Past Century Shows Targeting the Snail Intermediate Host Works Best. *PLoS Neglected Tropical Diseases*. 2016;10(7):1–19.
6. WHO. Ending-the-neglect-to-attain-the-SDGs--NTD-Roadmap. 2021;
7. Mutapi F, Rujeni N, Bourke C, Mitchell K, Appleby L, Nausch N, et al. *Schistosoma haematobium* treatment in 1-5 year old children: Safety and efficacy of the antihelminthic drug praziquantel. *PLoS Neglected Tropical Diseases*. 2011;5(5).
8. Imai N, Rujeni N, Nausch N, Bourke CD, Appleby LJ, Cowan G, et al. Exposure, infection, systemic cytokine levels and antibody responses in young children

- 385 concurrently exposed to schistosomiasis and malaria. *Parasitology*. 2011;138(12).
- 386 9. Rujeni N, Mazimpaka A, Tumusiime M, Nyandwi E, Rutayisire G, Kayiranga P, et al.
- 387 Pre-school aged children are exposed to *Schistosoma* through Lake Kivu in Rwanda.
- 388 AAS Open Research. 2019;
- 389 10. Stothard JR, Gabrielli AF. Schistosomiasis in African infants and preschool children:
- 390 to treat or not to treat? Vol. 23, *Trends in Parasitology*. 2007. p. 83–6.
- 391 11. Poole H, Terlouw DJ, Naunje A, Mzembe K, Stanton M, Betson M, et al.
- 392 Schistosomiasis in pre-school-age children and their mothers in Chikhwawa district,
- 393 Malawi with notes on characterization of schistosomes and snails. *Parasites and*
- 394 *Vectors*. 2014;
- 395 12. Sousa-figueiredo JC, Pleasant J, Day M, Betson M, Rollinson D, Montresor A, et al.
- 396 Treatment of intestinal schistosomiasis in Ugandan preschool children : best diagnosis
- 397 , treatment efficacy and side-effects , and an extended praziquantel dosing pole.
- 398 2010;2:103–13.
- 399 13. Ekpo UF, Oluwole AS, Abe EM, Etta HE, Olamiju F, Mafiana CF. Schistosomiasis in
- 400 infants and pre-school-aged children in sub-Saharan Africa: Implication for control.
- 401 *Parasitology*. 2012.
- 402 14. Ekpo UF, Laja-deile A, Oluwole AS, Sam-wobo SO, Mafiana CF. Urinary
- 403 schistosomiasis among preschool children in a rural community near Abeokuta,
- 404 Nigeria _ *Parasites & Vectors* _ Full Text. 2010. p. 1–5.
- 405 15. World Health Organization (WHO). WHO Guideline on control and elimination of
- 406 human schistosomiasis. Evidence-based recommendations. 2022.
- 407 16. Ruberanziza E, Wittmann U, Mbituyumuremyi A, Mutabazi A, Campbell CH, Colley
- 408 DG, et al. Nationwide remapping of *Schistosoma mansoni* infection in Rwanda using
- 409 circulating cathodic antigen rapid test: Taking steps toward elimination. *American*

- 410 Journal of Tropical Medicine and Hygiene. 2020;103(1):315–24.
- 411 17. Clark NJ, Umulisa I, Ruberanziza E, Owada K, Colley DG, Ortu G, et al. Mapping
412 Schistosoma mansoni endemicity in Rwanda: A critical assessment of geographical
413 disparities arising from circulating cathodic antigen versus Kato-Katz diagnostics.
414 PLoS Neglected Tropical Diseases. 2019;
- 415 18. Barbosa CS, Gomes ECS, Marcelino JMR, Cavalcante KRLJ, Nascimento WRC.
416 Quality control of the slides by Kato-Katz method for the parasitological diagnosis of
417 schistosomiasis infection by Schistosoma mansoni. Jornal Brasileiro de Patologia e
418 Medicina Laboratorial. 2017;53(2):110–4.
- 419 19. Leuenberger A, Nassoro T, Said K, Fenner L, Sikalengo G, Letang E, et al. Assessing
420 stool quantities generated by three specific Kato-Katz thick smear templates employed
421 in different settings. Infectious Diseases of Poverty. 2016;5(1):1–8.
- 422 20. Sturrock RF. Guidelines for the evaluation of soil-transmitted helminthiasis and
423 schistosomiasis at community level: A guide for managers of control programmes. A.
424 Montresor, DWT Crompton, DAP Bundy, A Hall & L. Savioli. Geneva: World Health
425 Organization, 1998. Transactions of the Royal Society of Tropical Medicine and
426 Hygiene. 1998;92(4):470–1.
- 427 21. Oliveira WJ, Magalh C, Mariana A, Elias S, Castro VN De, Favero V, et al. Evaluation
428 of diagnostic methods for the detection of intestinal schistosomiasis in endemic areas
429 with low parasite loads : Saline gradient , Helmintex , Kato-Katz and rapid urine test.
430 2018;1–22.
- 431 22. Stothard JR, Sousa-figuereido JC, Betson M, Adriko M, Arinaitwe M, Rowell C, et al.
432 Schistosoma mansoni Infections in Young Children : When Are Schistosome Antigens
433 in Urine , Eggs in Stool and Antibodies to Eggs First Detectable ? 2011;5(1).
- 434 23. Osakunor DNM, Woolhouse MEJ, Mutapi F. Paediatric schistosomiasis : What we

- 435 know and what we need to know. 2018;1–16.
- 436 24. Stothard JR; Gabrielli AF. Schistosomiasis in African infants and preschool children:
437 to treat or not to treat? Trends Parasitology. 2007;(23):83–6.
- 438 25. Okwuosa, V N; Osuala FO. Toxicity of washing soaps to Schistosoma mansoni
439 cercariae and effects of sublethal concentrations on infectivity in mice. Applied
440 Parasitology. 1993;34(1):69–75.
- 441 26. World Health Organization (WHO). Children’s Environmental Health Indicators
442 (CEHI): presenting regional successes learning for the future: summary.
443
444

446 Supporting information caption

447 **Fig 1. Map illustrating the distribution of schistosomiasis infection among pre-SAC in**

448 **Rwanda.** Infection prevalence was based on CCA findings (trace as negative); Geographical

449 coordinates were collected for each site (village) and were extrapolated to the lowest

450 administrative boundary (sector).

451