‘Anatomy of SARS-CoV-2 outbreak of ‘vaccinated’: An observational case-control study of Covid-19 breakthrough infections in medical college students at Rural Medical College, India

ABSTRACT:

Introduction

Breakthrough infections in reference to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) have been reported worldwide in both partially or completely vaccinated individuals irrespective of the type of vaccine. India, also emerged as the vaccine powerhouse producer, initiating the world’s one of the largest vaccination drives since January 16, 2021, with two vaccines named, BBV-152(COVAXIN™) and AZD1222 (COVISHIELD™). Breakthrough cases are reported from all over the globe for all kinds of vaccines. This is the investigation report into the outbreak of breakthrough covid infections at one of the medical colleges in Rural India unraveled in early October 2021. This report underscores the role of covid appropriate behavior along with vaccination and the role of IgG in evaluating immunity generated through vaccination.

Objective: The study objective was to 1. Clinically characterize the recent breakthrough infection of Covid-19 (SARS- CoV-2) infection among (students) vaccinated in BKL Walawalkar Rural Medical College. 2. To evaluate ‘Covid appropriate behavior’ (CAB) in cases and controls. 3. To evaluate Covid anti spike IgG in the cases in comparison with the control

Methods

A total of 74 students studying at BKL Walawalkar Rural Medical College and vaccinated for Covid were included in the study. RT-PCR diagnosis was done from 5 to 10 October 2021. The breakthrough infection in the cases was characterized using self-assessment questionnaires in comparison to the controls. The cases were assessed clinically and also using biochemical parameters. Both cases and controls were also assessed for their adherence to covid appropriate behavior using a separate semi-quantitative questionnaire and scoring system.

Results

In our study, out of the total subjects, 50% of Covaxin recipients had experienced vaccine breakthrough infection and 20% of Covishield recipients experienced breakthrough infection. Also, 6 out of the 35 cases were asymptomatic, and the rest were either having mild symptoms. None of them required any hospitalization or O2 therapy. The CAB score was lower in the cases when compared to controls. All the vaccine recipients are seroconvert. Anti-spike IgG antibodies titers are dynamic over time and across the clinically distinct groups. Irrespective of varying IgG titer, the vaccine protects against severity and possibly mortality.

Conclusion

The need for better awareness of covid appropriate methods as an alternative to and additive to vaccination is necessary to control the transmission of Covid infection and decrease the disease severity. Vaccines are effective against preventing severe disease and possibly mortality. The use of serum anti-spike covid IgG is restricted to know the status of SARS-CoV-2 seroconversion.

Keywords

Breakthrough infection; SARS-CoV-2; Covid-19 vaccination, covid appropriate behavior, Anti-spike-IgG seroconversion

Author details:

1. Miss Monika Chavan.
Department of Biochemistry, BKL Walawalkar Rural medical College, Maharashtra, India 415606
Email Id: monikachavan85@gmail.com
2. Dr. Sowmya Gayatri,
Department of Biochemistry, ESIC super-speciality Hospital, Sanath Nagar, Hyderabad, India
Email Id: gayatri20072008@gmail.com
3. Dr. Suvarna Patil
Department of Medicine, BKL Walawalkar Rural medical College, Maharashtra, India 415606
Email id : dr.suvarnanpatil@gmail.com
4. Dr. Janhavi Deshpande
Department of Medicine, BKL Walawalkar Rural medical College, Maharashtra, India 415606
Email id: jspjvd@gmail.com
5. Dr. Arvind Yadav.
Department of Biochemistry, BKL Walawalkar Rural medical College, Maharashtra, India 415606
Email id: asyadav13@gmail.com
6. Dr. Prasanna Nakate
VRDL Laboratory, Department of Microbiology, BKL Walawalkar Rural medical College, Maharashtra, India 415606
Email Id: prasannanakate@gmail.com
7. Dr Anup N Nillawar. (Corresponding Author)
Department of Biochemistry, BKL Walawalkar Rural medical College, Maharashtra, India 415606
Email Id: nilawaranup@gmail.com

Authors' Contribution:

MC and SG contributed equally and oversee the investigations and data collection and contributed in manuscript writing, JD and SP provided crucial clinical data about hospitalized patients, AY reviewed manuscript and contributed in data collection, PN gave RT-PCR inputs, AN conceptualized the paper, prepared study design and prepared CRF and did manuscript writing and oversee laboratory investigations.

INTRODUCTION

The vaccination campaign for COVID19 in India was started on January 16, 2021, using two vaccines; AZD1222- ChAdOx1-S (Covishield), manufactured in India by Serum Institute of India through a license from AstraZeneca-Oxford.¹ and BBV152 (Covaxin), an indigenous vaccine developed by Bharat Biotech in collaboration with the Indian Council of Medical Research (ICMR).² From May 1st onwards vaccination is open to all individuals (>18 yr of age) in India. The efficacy after administration of two doses of the vaccines irrespective of the interval between the doses has been reported as 63.1% and 78% for AZD1222 (ChAdOx1-S) and BBV152 respectively.¹³ Vaccines are widely regarded as the most effective means of halting and terminating the Covid-19 outbreak. However, due to a variety of factors vaccine efficacy in the real world may differ from that reported in clinical trials. Because no vaccine provides 100 % protection against the disease, and newer virus types periodically evolve mechanisms to evade the vaccine-induced antibody response, a small percentage of people will contract Covid-19 despite being fully vaccinated. Also, other preventive measures such as Covid appropriate behavior which determines the degree of exposure to the virus, play a major role in preventing breakthrough infections.⁴⁵ Further, strict adherence to such behavior would also allay the severity of infection. Breakthrough infections about Covid-19, refer to the incidence of SARS-CoV-2 infections in individuals who have already been partially or completely vaccinated with any authorized Covid-19 vaccine.⁶ According to the ICMR between 0.02 and 0.04% of infections have occurred after partial or complete vaccination with either BBV152 or AZD1222 (ChAdOx1-S).⁴ The current study attempts to characterize covid breakthrough infections to concerning clinical symptomatology, vaccination status, and inclination and awareness towards Covid appropriate behavior. This is pursued by using a self-assessment semi-quantitative questionnaire in simple understandable language. This study would provide crucial evidence in understanding not only the vaccine effectiveness but also the need for strict adherence to appropriate behavior in preventing transmission and decrease in the severity of the disease.

OBJECTIVES

1. To clinically characterize recent Covid 19 breakthrough infection of vaccinated subjects in BKL Walawalkar Rural Medical College during 5 to 10 October 2021.
2. To evaluate 'Covid appropriate behavior' (CAB) in cases and controls.
3. To evaluate Covid anti spike IgG in the cases in comparison with the control

MATERIAL AND METHODS

Study background:

BKL Walawalkar rural medical college and hostels are the closed knit campus situated in the rural Konkan region where college and hostel campus is situated away from village dwellings and with very little possible contact with the outsiders and even with the hospital premises. In such a campus, in the last week of September 2021, we found one index case having mild upper respiratory tract infection (URTI) and came covid positive on RT-PCR. This triggered a thought to check for all the students if they are having any spread as they live in closely placed hostel rooms and closed sitting arrangements in offline teaching classes. Boys and girls have separate hostel buildings and are situated far apart. All the students were vaccinated against SARS-CoV-2 almost at the same time period, around April 2021.

The rationale for selecting the study group:

The study group is in many ways a closed group. Being in the same age group and staying in the hostel campus before and after vaccination and during this investigation. Hence, there is less chance of bias on the exposure to external sources of infection. Hence, there is a considerable chance that the transmission was from the index case and later on the spread to the rest of the class and was also restricted only to the close contacts of the infected. In a way, this can be described as a point source outbreak. Also, all of them were vaccinated at around a similar time. (March-April 2021) This helps in understanding the clinical effectiveness of the vaccine. To assess the kinetics of antibody response, we evaluated titers of Covid anti-spike IgG.

LABORATORY TESTING: All nasopharyngeal swab samples were tested in an automated PCR machine, Roche LC-96. This targets two genes that are ORF-1 and E gene. Anti-spike

IgG antibodies against SARS-CoV-2 were quantitatively estimated in an automated chemiluminescence electro assay (CLIA) - based platform, ARCHITECT i1000SR (Abbott diagnostics) using a commercial quantitative kit SARS-CoV-2 IgG □ quant. The cut-off value is 50 AU/ml.

STUDY SUBJECTS: A total of 74 students studying at BKL Walawalkar rural medical college, Dervan, Maharashtra, India, who were all vaccinated for Covid 19 infection in March and April 2021 (first dose of vaccine) were included in the study. All the students who tested positive for Covid 19 by RT-PCR with/without clinical symptomatology and at least one dose of Covid vaccination were selected as cases. Similarly, all the RT-PCR negative tested students with no clinical symptomatology and with a history of at least one dose of vaccination were selected as controls. For finding out the best matching controls, we briefed the cases about the study design and asked them to choose the best matching control based on the contact time between them and amongst their peers. The study included 35 cases and 39 controls. Case record forms (CRF) were prepared to gather the following information: 1. Demographics, History of previous Covid 19 infection, and vaccination details (As per the government of India policy, two doses of either BBV152 (Covaxin) or AZD1222 (Covishield) vaccine at least 4 weeks apart were available for administration since January 2021 with no mixing of doses allowed) 2. Covid clinical symptomatology, assessed by a semi-quantitative ‘self-assessment clinical questionnaire’ and a self-assessment questionnaire on ‘Covid appropriate behavior (CAB)’ before and after this outbreak (CRF as Annexure). Both the questionnaires were prepared in an easily understandable manner and consisted of questions, each with a score from 0 to 10. The self-assessment questionnaire to assess ‘Covid appropriate behavior (CAB)’ included five topics (a. wearing of mask b. Use of hand sanitizer c. frequent hand washing d following social distancing e. following travel restrictions) with each topic having a maximum score of 10. The total maximum score possible in the clinical questionnaire is 230 and for CAB is 50. (Annexure). The study protocol was approved by the Institutional Ethics Committee. Written consent was obtained from the subjects before the initiation of the study.

STATISTICAL ANALYSIS

Continuous data has been presented as median (25th-75th centiles) while categorical data has been presented as percentages. Pearson's correlation was used to analyze the association between IgG titers and clinical scores. The independent-sample t- test was used to compare between two groups. Statistical analysis was performed using Statistical Package for the Social Sciences version 21.0 (SPSS, Inc.; Chicago, IL, USA) for Windows. p- Value <0.05 was considered statistically significant.

RESULTS

A total of 74 students studying at BKL Walawalkar rural medical college, were included in the study. Table 1 shows the basic characteristics of the study subjects. Among the breakthrough cases, 32 had received 2 vaccination doses (fully vaccinated according to vaccination schedule) while, 3 had received one dose of vaccine. Among the 39 matched controls, 33 had received 2 and 6 had received one vaccination dose respectively Out of the whole cohort, 6 had received AZD1222/Covishield while 68 had received BBV152/Covaxin. (Table no. 2).

1. Demography: The median age was 21.0 years. The median BMI of the study subjects was 21.02 kg/m². The median Concentration of IgG was 726.0 (AU/ml). The mean antibody IgG titer between cases and control were 2996.3 and 656.2 respectively. (Table No:) There is a significant difference in IgG titers between cases and controls (p=0.048).
2. 50% of Covaxin recipients had experienced vaccine breakthrough infection and 20% of Covishield recipients experienced breakthrough infection. The average duration between the second dose and confirmed SARS-CoV-2 infection by RT-PCR was 150 days; in Covaxin recipients and 88 days; in Covishield recipients.
3. Clinical: 6 out of 35 cases were asymptomatic, 29 positive cases with minimal symptomatology, 47% had anosmia and 41 % had a loss of taste. The most common symptom was runny nose and fever found in 50 % of patients. None had lower respiratory signs; none of them required hospitalizations or Oxygen support. All the symptoms were restricted to the upper respiratory tract. Out of 35, 10 were investigated for routine blood inflammatory markers and none had CRP>20 mg/L.

Only 2 cases had CRP at 10 mg/L. None had increased D-Dimer above normal levels.

The clinical score (calculated out of 23 symptoms, each symptom gauged on the scale of 0 to 10, on self-assessment questionnaire) is directly correlated with IgG titers ($r=0.293$, $p=0.011$). Subtly, this tells IgG titer increases with aggravation in clinical symptomatology.

4. Covid Appropriate Behavior (CAB) Score: (Score was calculated on 5 criterion, each carrying 10 points, maximum score possible 50). And to our surprise, the mean CAB scores for cases and control were 36.2 and 42.5 respectively. ($p=0.000$).

Both cases and controls improved on their CAB score post-outbreak.

5. Serum IgG titer: IgG antibodies against SARS-CoV-2 were quantitatively estimated in an automated chemiluminescence electro assay (CLIA) - based platform, ARCHITECT i1000SR (Abbott diagnostics). The present cohort (case; group2 and control; group3) was investigated for IgG after an average of 150 days of vaccination. Group 1 was vaccinated individuals with IgG measured at an average of 60 days of vaccination. Additionally, we compared IgG titers with a group (Group 4) having moderate to severe SARS-CoV-2, hospitalized, and needing oxygen support. (Table 3)

- a) All the vaccine recipients showed seropositivity, there was no significant difference in seropositivity between Covaxin and Covishield. The mean antibody IgG titer value for Covaxin and Covishield were 1719.0 AU/ml and 2261.8 AU/ml respectively. ($p=0.791$) (Table:2)
- b) In the control group of the present study (Group2, Table 5), mean IgG levels are = 656 AU/ml. This analysis is done after the average lapse of 150 days. IgG levels are significantly lower when done after a wider time-lapse and it indicates waning antibody titers. Even though the titers are done in 2 different groups, they are enough indicating about waning antibodies.
- c) Table 4 indicates, additional infection with SARS-CoV2 bolsters IgG titers in vaccinated individuals.

d) As the secondary objective was to know the SARS- CoV-2 anti-spike IgG titer and its association with immunity to SARS-CoV-2 re-infection. As shown in table 3, IgG titer varies across the groups, likewise; in ‘only vaccinated group’ (Group1), Group with mild breakthrough SARS-CoV-2 infection (Group 3), And patients with moderate to severe breakthrough infection (Group 4), who needed hospitalization and various levels of Oxygen support. Though the number in group 4 is less and no controlled matching was possible with the rest of the groups, the fact which remains assuring is that none of the patients with severe breakthrough infections also died. This trend needs to be ascertained and reinforced with more data. This table showed, though the IgG levels are varying across the groups and they are showing a waning trend, vaccination is helping to prevent severe disease and mortality. (Group 3 and 4 in Table 4)

Table1: Basic Characteristics of Cohort:

Age (years)*	21.0 (20.0-21.0)
BMI (kg/m ²)	21.02 (19.3-23.7)
Sex	
Male	38 (51.4%)
Female	36 (48.6%)
IgG (AU/ml)*	611.5 (349.0-1476.7)

Vaccine type received	
Covaxin	68 (91.9%)
Covishield	06 (8.1%)

Table 2: Demographic and Clinical Characteristics of breakthrough cases receiving Covaxin and Covishield

Variable	Covaxin recipients (n=68)		Significance	Covishield recipients (n= 6)		Significance
	Case (n=34)	Control (n=34)		Case (n=1)	Control (n=5)	
Age(median,IQR)	20	21	0.000	19	21	NA
Sex:						
Male	15	19		1	3	
Female	19	15		0	2	
BMI	22.0	20.1	0.059	23.4	22.2	NA
Symptomatic:						
(Yes)	28	NA				
If symptomatic, clinical Score on the scale of 230(mean)	33.9	NA		15	0	
Asymptomatic	6	NA		0	0	
Mean duration between second dose and Infection	150	88				
Antibody positive against S protein						
IgG titer(mean)	2993.5	444.5	p=0.036	3093	2095.6	NA
Covid appropriate behavior(On scale of 50)	36	43	p=0.000	44	39.6	NA

(Case Arm: At least 1 dose of Covid vaccine+ RT-PCR positive breakthrough covid infection)

(Control Arm: Age and ecology matching group with At least 1 dose of Covid vaccine+ RT-PCR Negative)

Table3: Comparison of IgG titers with additional data sets:

	Post-vaccination + 60 days healthy volunteers Grp1	Post vaccination + 150 days. ('control' group of the present study)Grp2	Post-vaccination +150 days+ mild breakthrough infection ('case' group of the present study) Grp3	Post-vaccination (7 to 30 days) + Moderate to severe covid Hospitalized patients Grp4
Number of cases (n)	n=31	n=39	n=35	n=10
Age:	42.8	21	19.9	47
IgG titer	1430.8	656.2	2996.3	7000
Resultant	No Covid infection reported till this 12 Dec 2021	No covid infection /RT –PCR negative till 12 Dec 2021	Mild breakthrough infection limited to URT with RT-PCR positive report.	No mortality

(Group 4, which is the cohort of Post-vaccination hospitalized patients who were suffering with moderate to severe covid disease, has IgG titers in the range of 2900 to 40000 AU/ml, with the mean of 7000 AU/ml. This is the group of 10 cases of severe disease, none died and they contracted infection within a short span of vaccination)

Table 4: Showing Bolstering of IgG titer after vaccine+ Infection:

	(Grp2) (With Only vaccination)	(Grp3) (With vaccination+ RT-PCR positive for SARS Cov-2)	p-Value
Number of cases (n)	39	35	
Age	21	19.9	
IgG titer (AU/ml)	656.2	2996.3	0.05
The duration between the second dose of vaccine and IgG Titer (Mean)	108	144	NA
Resultant	No Covid/ or covid Suspect	RT-PCR positive Covid	

Table 5: Showing waning of IgG titer:

	(Grp1) IgG Titer did after 60 Days	(Grp2) IgG Titer did after 150 Days	Significance
Number of cases (n)	31	39	
Age	42.8	21.0	p=0.000
Time since vaccination	60	150	
IgG titer Mean (AU/ml)	1430.8	656.2	p=0.034
Resultant	No Covid reported	No Covid reported	

Discussion:

Breakthrough infections were operationally defined as the occurrence of Covid-19 infection ≥ 14 days after administration of two doses of Covid-19 vaccine irrespective of the type. In the present study, 35 experienced a breakthrough infection after being administered both scheduled doses of either Covid-19 vaccine. Furthermore, nearly, 3 individuals reported the incidence of Covid-19 infection after receiving one dose of a Covid-19 vaccine but to before either complete vaccination or before 14 days post administration of the second vaccine dose. However, in our study, we also observed that despite the infection, the disease in the infected was often mild or sometimes with nil symptomatology. Though very few similar studies on breakthrough infections are reported from India.⁷⁸ Also, we observed that following a covid appropriate and covid preventive rules such as the use of face masks, maintaining social distancing, proper and regular hand washes, limiting social contact and gatherings, etc., had a huge impact on containing the transmission of infection among the exposed. It is beyond doubt that vaccines may have a greater impact on reducing viral circulation than naturally acquired immunity. The study by Tartof et al. observed that vaccines decrease disease severity among individuals who do acquire infection i.e., breakthrough infections.⁹ The mechanisms by which COVID-19 vaccination elicits disease attenuation are largely unknown but are likely due to the recall of immunologic memory responses that reduce viral replication and accelerate the elimination of virally infected cells. Studies of viral dynamics further suggest that while viral loads in breakthrough infections may be as high in vaccinated individuals as they are in unvaccinated individuals, viral loads in those who are vaccinated decline more rapidly, and the virus that they shed is less likely to be culture-positive than virus shed by unvaccinated individuals.¹⁰¹¹ This suggests that people who are fully vaccinated are less likely to become infected and if infected, will be contagious for shorter periods than unvaccinated.¹² Vaccines not only decrease transmission rates but also decrease disease severity among individuals who do acquire infection. This is very well supported by the present study by two means, one the

breakthrough infection found in vaccinated individuals after an average lapse of 160 days was found to be very mild. So also, in the small group of SARS Cov-2 hospitalized patients, those who contracted an infection after an average of 30 days of vaccination, had no mortality despite their needed hospitalizations and oxygen therapy. (Table No: 4) To concerning delta variant, when vaccinated subjects with breakthrough infections are compared with unvaccinated, are less likely to develop mild or severe symptoms, more likely to recover from their illness quickly, and much less likely to require hospitalization. Nonetheless, the fact that vaccinated individuals can still become ill enough with COVID-19 to be hospitalized is understandably concerning. Even though vaccines are widely regarded as the most effective means of halting and terminating the Covid-19 outbreak, vaccine efficacy differs in the real world due to a variety of factors than reported.¹³ Hence, other preventive measures such as Covid appropriate behavior which determines the degree of exposure to the virus also play a major role in preventing breakthrough infections. And our study strongly underscored the role of covid appropriate behavior and its additive effect on vaccine protection.⁴ A valuable new study by Tenforde and colleagues affirms the vaccine efficacy against hospitalization also depends on patients' immune status, age, vaccine preparation, time since vaccination, and Covid appropriate behavior among other parameters studied. The study further suggests that vaccines along with appropriate preventive measures are associated with a lower likelihood of infection and also transmission.¹⁴ Vaccination is the better effect when combined with better CAB, as all those who contracted covid breakthrough infections fared on the lower scale on CAB score in our study. This may have underlined the message that avoiding those contacts that are on the lower scale on CAB score (whose behavior is not in compliance with guidelines to prevent respiratory droplet infections) and avoiding close and crowded rooms is better for infection prevention. This warrants better awareness programs for propagating the importance of covid appropriate behavior. Of given the recent variants (omicron) and imminent variants of whatsoever nature, our study underlines the importance of vaccination and CAB.

Antibody response to SARS-Co-V 2: Evidence through multiple studies shows that protective antibody titers even though correlate well at the population level, are still

elusive at the individual level¹⁵. A positive antibody test is currently recommended only to detect an individual with prior infection but cannot be used to assess the level of protection provided by the immune system in post-vaccinated individuals. However, a study by Bates et al. suggests that even though the breakthrough infections in patients were mild, there is a substantial increase of humoral immunity similar to the findings in our study.¹⁶ Additionally, in the wake of newer variants of SARS-CoV-2 emerging at a rather fast rate, the breakthrough sera have also demonstrated variant cross-neutralization in the vaccinated individuals. This creates a scope for a broadened protective immune response if the existing 2 dose series are further supplemented by variant boosters with antigenic inserts similar to the emerging variants and further prolong the protection as seen in a study conducted by Falsey et al.¹⁷ Naturally, antibody response to infection takes several days to a week and level decreases over time. Quantitative SARS- Co-V-2 anti-spike IgG response is varied by prior infection status, age, sex, type of vaccine & number of doses received. Post-vaccine antibody response is increased over 2-4 weeks after 1st dose of vaccination. Vaccine dose and following breakthrough infection cause a surge in IgG response as reported by this study.¹⁸

The decline in IgG titers is expected; but how these titers relate to protection from subsequent re-infections (Symptomatic or severe covid disease and death) remain elusive. An individual with a low or undetectable level of SARS-COV -2 antibodies have an increased risk of shedding infectious viruses.¹⁹ Effectiveness of vaccination and antibody-based -correlates of this protection are a potential strength of serological testing to identify those who are immune. Though this is a humongous and complex but a very novel aim, for which larger and complex studies are required. Studies with 'Long longitudinal cohorts with baseline immunology' are needed. This small study throws some insights for future studies.

This present outbreak could be because a new unknown Covid variant that demands genomic sequencing and continuous surveillance. Unusual outbreaks like this and changing ecology of pandemic calls for deeper genomic and mechanistic studies.

The B.1.617.2 (Delta) variant of (SARS-CoV-2) was first identified in the state of

Maharashtra, adjoining to the place of the present study in late 2020, and spread throughout India, outcompeting pre-existing lineages including B.1.617.1 (Kappa) and B.1.1.7 (Alpha).²⁰¹³ This variant had higher replication and spike mediated entry and better host immune evasion. The latest variant omicron is allegedly milder because of its better availability to immune cells. The ongoing battle between the immune system and the natural selection of potential viral mutants is highlighted by the present study.

This study also demands real-time and representative studies from different geographies to understand the changing scene of emerging variants. Until now, Very few studies have been reported from India⁷⁸

Conclusion: Covid-19 vaccination may not be the silver bullet to defeat the novel SARS-CoV-2, but as of now, it is the most important weapon in the arsenal to prevent, manage and overcome the pandemic. The vaccine induces humoral immune response which is reflected in IgG response against spike protein and the effectiveness of the vaccine is shown in the prevention of infection, the mildness of infection, and likely protection against mortality. And if vaccination is supported by Covid-appropriate behavior, it increases the chances of scoring over the novel coronavirus. Since India is also facing the emergence of new variants and a surge in covid cases, it makes the case stronger for following the practice of hand hygiene, wearing of masks, and maintaining social distancing in public places. The two-pronged approach of vaccination and Covid-appropriate behavior would go a long way in fighting not only this wave but also future waves and variants of the pandemic.

Bibliography:

1. World health organization.COVID-19 vaccines.
<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/covid-19-vaccines>. Accessed January 11, 2022.
2. Ella R, Reddy S, Jogdand H, et al. Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBV152: interim results from a double-blind, randomised, multicentre, phase 2 trial, and 3-month follow-up of a double-blind, randomised phase 1 trial. *Lancet Infect Dis*. 2021;21(7):950-961. doi:10.1016/S1473-3099(21)00070-0
3. Covaxin efficacy 78% in Phase 3 analysis | Latest News India - Hindustan Times.
<https://www.hindustantimes.com/india-news/covaxin-efficacy-78-in-phase-3-analysis-101622227104005.html>. Accessed January 11, 2022.
4. Singh AK, Phatak SR, Singh R, et al. Antibody response after first and second-dose of

- ChAdOx1-nCOV (Covishield TM®) and BBV-152 (Covaxin TM®) among health care workers in India: The final results of cross-sectional coronavirus vaccine-induced antibody titre (COVAT) study. *Vaccine*. 2021;39(44):6492-6509. doi:10.1016/J.VACCINE.2021.09.055
5. COVID-19 Vaccine Breakthrough Case Investigation and Reporting | CDC. <https://www.cdc.gov/coronavirus/2019-ncov/php/hd-breakthrough.html>. Accessed January 11, 2022.
 6. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020;383(27):2603-2615. doi:10.1056/NEJMOA2034577
 7. Dash GC, Subhadra S, Turuk J, et al. Breakthrough SARS-CoV-2 infections among BBV-152 (COVAXIN®) and AZD1222 (COVISHIELD TM) recipients: Report from the eastern state of India. *J Med Virol*. 2021;1(5). doi:10.1002/JMV.27382
 8. Singh UB, Rophina M, Chaudhry R, et al. Variants of concern responsible for SARS-CoV-2 vaccine breakthrough infections from India. *J Med Virol*. November 2021. doi:10.1002/JMV.27461
 9. Tartof SY, Slezak JM, Fischer H, et al. Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated health system in the USA: a retrospective cohort study. *Lancet (London, England)*. 2021;398(10309):1407-1416. doi:10.1016/S0140-6736(21)02183-8
 10. Ke R, Martinez PP, Smith RL, et al. Longitudinal analysis of SARS-CoV-2 vaccine breakthrough infections reveal limited infectious virus shedding and restricted tissue distribution. *medRxiv Prepr Serv Heal Sci*. September 2021. doi:10.1101/2021.08.30.21262701
 11. Chia PY, Ong SWX, Chiew CJ, et al. Virological and serological kinetics of SARS-CoV-2 Delta variant vaccine breakthrough infections: a multicentre cohort study. *Clin Microbiol Infect*. November 2021. doi:10.1016/J.CMI.2021.11.010
 12. Shah ASV, Gribben C, Bishop J, et al. Effect of Vaccination on Transmission of SARS-CoV-2. *N Engl J Med*. 2021;385(18):1718-1720. doi:10.1056/NEJMC2106757
 13. Harder T, Külper-Schiek W, Reda S, et al. Effectiveness of COVID-19 vaccines against SARS-CoV-2 infection with the Delta (B.1.617.2) variant: second interim results of a living systematic review and meta-analysis, 1 January to 25 August 2021. *Eurosurveillance*. 2021;26(41). doi:10.2807/1560-7917.ES.2021.26.41.2100920
 14. Tenforde MW, Self WH, Adams K, et al. Association Between mRNA Vaccination and COVID-19 Hospitalization and Disease Severity. *JAMA*. 2021;326(20):2043-2054. doi:10.1001/JAMA.2021.19499
 15. National Center for Immunization and Respiratory Diseases (NCIRD) D of VD. Science Brief: SARS-CoV-2 Infection-induced and Vaccine-induced Immunity. *CDC COVID-19 Sci Briefs*. October 2021. <https://www.ncbi.nlm.nih.gov/books/NBK575088/>. Accessed

January 22, 2022.

16. Bates TA, McBride SK, Winders B, et al. Antibody Response and Variant Cross-Neutralization After SARS-CoV-2 Breakthrough Infection. *JAMA*. 2022;327(2). doi:10.1001/JAMA.2021.22898
17. Falsey AR, Frenck RW, Walsh EE, et al. SARS-CoV-2 Neutralization with BNT162b2 Vaccine Dose 3. *N Engl J Med*. 2021;385(17):1627-1629. doi:10.1056/NEJMC2113468
18. Wei J, Stoesser N, Matthews PC, et al. Antibody responses to SARS-CoV-2 vaccines in 45,965 adults from the general population of the United Kingdom. *Nat Microbiol*. 2021;6(9):1140-1149. doi:10.1038/S41564-021-00947-3
19. Glans H, Gredmark-Russ S, Olausson M, et al. Shedding of infectious SARS-CoV-2 by hospitalized COVID-19 patients in relation to serum antibody responses. *BMC Infect Dis*. 2021;21(1). doi:10.1186/S12879-021-06202-8
20. Mlcochova P, Kemp S, Dhar MS, et al. SARS-CoV-2 B.1.617.2 Delta variant replication and immune evasion. *Nature*. 2021;599(7883). doi:10.1038/S41586-021-03944-Y