1 **Liver Echinococcosis Lesion Classification Tool by** 2 **Deep Learning: development, deployment, and** 3 **validations**

4 Liang Huang¹, Xiaohong Xi², Zhu Chen¹, Yilan Zeng¹

5 ¹Chengdu Public Health Clinical Center, Chengdu City, Sichuan Province, China,

6 61000

7 ²Department of Sonography, Chenghua Women and Children's Hospital, Chengdu

8 City, Sichuan Province, China, 61000

9

10 Correspondence:

11 Yilan Zeng, Chengdu Public Health Clinical Center, 377 Jingming Road, Chengdu

12 City, Sichuan Province, China (hyuangx@qq.com).

13

14

15

16

17

18

19

20

21

22

23 Abstract

24 The endemic of Echinococcosis imposed heavy disease burden in some areas. The
 25 sonography for Echinococcosis lesions was essential to disease diagnosis and
 26 managements. Especially the biological typing of lesions was key to disease
 27 treatments. We used deep-learning tools to help sonographer to classify the lesion
 28 types. The model achieved 85%(302/376) accuracy, in contrast to senior sonographer
 29 achieved 72%(61/85) accuracy. The accuracy of AI model was higher than senior
 30 sonographer (p-value=0.01), could be a feasible method to help sonographer in
 31 remote area.

32

33

34

35

36

37

38

39

40 Introduction

41 Echinococcosis is a result of human infection by larval stages of taeniid cestodes of
 42 the genus *Echinococcus*. *Echinococcus granulosus* causes cystic echinococcosis(CE),
 43 *Echinococcus multilocularis* causes alveolar echinococcosis (AE), two separately
 44 diseases with different biological processes^{1,2}. The co-infection of the two pathogens
 45 was rare³. At least 58% of the population are at risk of acquiring echinococcosis,
 46 distributed in Europe, North and Central America, Africa, Australia and Western
 47 China⁴. It is a potential fatal zoonosis⁵, a major public health concern in
 48 Qinghai-Tibet Plateau due to the highly pandemic of both CE and AE. The
 49 echinococcosis DALYs in the Tibetan communities were 126,159 life years annually
 50 estimated by Wang⁶. The Tibet population in China holds 1.66% prevalence of both
 51 CE and AE, estimated 50 million people are at risk of infecting the disease in Western
 52 China⁷. China has been investing large amount of resources to control echinococcosis,
 53 launched national control program⁸. The strategies of the national control program
 54 majorly included definitive hosts deworming, slaughtering control, and intermediate
 55 hosts vacation². Ultrasound screening was included as an essential measure to find the
 56 patients and follow-up.

57 Echinococcosis is chronic parasites disease, liver echinococcosis were most common
 58 in clinical⁹. The treatments for it included surgically remove the lesion,
 59 PAIR(puncture, aspiration, injection, and respiration), albendazole or they

60 combined^{10,2}. The lesion classification for CE is the one of the key aspects in patients'
 61 management, since the classification was closely correlated to the underlying
 62 biological status of the larva.

63 For CE, the ultrasonic WHO classification is commonly used (CL, CE1, CE2, CE3,
 64 CE4, and CE5). The CE1-2 types of CE indicating active infection should be treated
 65 by Albendazole or/and surgical resection. The CE4-5 are considered degenerated
 66 lesion due to failure of natural infection or effects of treatments, should be closely
 67 monitored^{11,12}. The CE3 is considered the transitional phase of lesion. The CE1-5 are
 68 the continuous progress of “active-transitional-inactive”¹³ indicating importance for
 69 biological process and disease assessments.

70 The AE is a silent progressing infiltrative proliferation of the parasite, mimicking a
 71 malignancy¹⁴. For AE cases, the lesion classification are more complex and uncertain,
 72 research revealed that the clinical stage of AE was not correlated to ultrasound
 73 appearance¹⁵. WHO had proposed a PNM classification system for AE¹⁶. The PNM
 74 classification system (P=parasitic mass in the liver, N=involvement of neighboring
 75 organs, and M=metastasis) was designed to systematic evaluation for surgical purpose
 76 majorly. Besides PNM classification system, the Ulm university proposed a
 77 classification system for AE¹⁷, and five types of lesion proposed.

78 The PNM classification systems required extra examinations other than ultrasound,
 79 such as CT or MRI scan. However, the PNM system was not feasible for many

occasions. The accessibility of medical resources in Qinghai-Tibet Plateau is incomparable with urban area¹⁸ due to the high altitude and atrocious weather, the local paramedics are lack of professional training, especially for sonography.

This study purposed to develop deep learning-based application to automatically type lesions for both AE and CE, giving diagnostic assistant for local paramedics. In this study, we used 2820 ultrasound images training our model, and used external data to validate the model. It is the largest echinococcosis ultrasound training set so far and its application for was published on a web for open access.

Previous studies had developed deep learning system for hepatology by using medical images, pathology, clinical and laboratory data, natural languages, etc. They are majorly designed to address liver problems such as liver fibrosis(cirrhosis), liver cancer, portal hypotension, etc¹⁹. Relatively, less study were aimed to the liver echinococcosis, which is on the list of NZDs²⁰ by WHO. Before our study, Xin had built a segmentation and classification networks for echinococcosis, shedding light on potential development in this field. However, it used 160 CT scan images for training, and the classification of echinococcosis were not covered all types of both AE and CE²¹. Comparing with CT images, ultrasound was more widely used in filed echinococcosis diagnosis. In this study, we used 2820 ultrasound images for training covering all echinococcosis types.

Method and Material

100 **Patient Data**

101 Sonographies were retrospectively obtained from National Control Program for
 102 Echinococcosis Screening in Qinghai-Tibet Plateau²². The medical decisions were not
 103 based on results of this study. All the patients were previously diagnosed
 104 echinococcosis by systematic evaluations including medical images and serology tests,
 105 fulfilled the criteria of echinococcosis diagnosis criteria.
 106 In this study, we collected 3423 images of echinococcosis patients who were enrolled
 107 and routinely managed in national echinococcosis control program.
 108 Ethical approval for this retrospective study was obtained from the Ethics Committee
 109 of the Chengdu Public Health Clinical Center. The patient consents for inclusion were
 110 waived as the retrospective nature of this study and anonymous data use.

111

112 **Classification criteria**

113 In this study, we used CL-CE5 classification system for CE, which was recommended
 114 by WHO recommendation¹³. We used infiltrating, calcification, and necrotic lesion
 115 for AE classification, which was recommended by Chinses Guideline for AE²³.
 116 For CE lesions, the CL lesion, which was appears “cystic lesion”, which appears no
 117 different from simple cyst in liver, could not be confirmed by ultrasound image solely.
 118 The CE1 refers to a simple round or oval unilocular cyst with anechoic content and a
 119 visible double cystic wall. The CE2 cyst is filled with daughter vesicles. The CE3

120 cysts includes two stages, CE3a is characterized by the “water-lily” sign, CE3b
121 represented by floating membranes. The CE4 typically reveals coarse variable
122 hyperechogenic or hypoechogenic echotexture without daughter vesicles. The “ball of
123 wool” sign, corresponding to the detached endocyst as a hypoechoic folded structure
124 embedded in a hyperechoic matrix, is the key ultrasound sign. The CE5 cysts are
125 partially (with an egg-shell calcified wall) or completely calcified with shadowing¹³.

126 For AE lesions, in this study, we followed the Chinese Guideline for AE, which was
127 also applied in National Controlling Program in endemic areas. It proposed three
128 ultrasound types of lesions. The infiltrative lesion (AE1) appearing hailstorm pattern,
129 heterogeneously echogenic areas, in many cases scatter calcifications can be seen.
130 The necrosis lesion (AE2), appearing a pseudocystic pattern with an irregular
131 hyperechoic rim. The calcification lesion, appearing solid hyperechoic lesion (AE3).

132 **Data labelling**

133 Expert who had more than 15 years of echinococcosis ultrasound diagnosis and
134 ultrasound based echinococcosis patients managements had labeled the included
135 ultrasound images by using LabelMe²⁴, an open-source image label tool. The lesions
136 in the images were labeled by different color, and normal liver background was
137 labeled as black. The labeled images were automatically cropped into fixed size of
138 input images (512*512 pixels) for model learning and inference.

139 **Model constructions**

140 The U-net architecture achieves high performance on different biomedical images
 141 segmentation applications by using relatively less quantity of training images^{25,26}. In
 142 this study, we used two separated U-net models. Firstly, we constructed “Classifier”
 143 U-net model which was used generalized multiclass dice loss (GMD). The GMD
 144 penalizes the wrong classifications more aggressively and more effective for training
 145 if the unbalanced category of training samples used. However, it produces more noise
 146 (false positive predictions) produced. Secondly, the identical “Shaper” U-net using
 147 conventional binary dice loss function focusing on lesion shape learning black-white
 148 (lesion-background, values 0 and 1) outputs, omitting the lesion class information. It
 149 produced more accurate shape inference for lesions. The inference result produced by
 150 multiplication of two tensors of the models’ outputs (Figure 1).

151 The whole program was based on Python programming languages with PyTorch 1.3
 152 deep learning framework. The core construction of the models consist 10 of
 153 convectional layers including downsampling, upsampling, and attention gates were
 154 used in this model²⁷. The models were trained with mini-batch SGD optimizer with
 155 momentum on two Nvidia 1080TI GPUs. The learning rate was chosen to minimize
 156 the error in the tuning dataset. Early stop training strategies was used for preventing
 157 over fitting.

158 **Web-based user interface**

159 Trained weights files loaded in a web-based user interface under Flask framework for
160 internal logical process and HTML for appearances of the application of both
161 computer and mobile phone. The users will upload images transferred from
162 ultrasound equipment by computer or mobile phone. The web-based UI will
163 automatically calculate the hot-spot of the ultrasound images and crop them into
164 proper size and optimized location(Figure 2).

165 **Model validations**

166 We compared classification accuracies for human and model. The senior sonographer
167 has 15 years of working experiences. The performance of model was designed to
168 achieved equivalent accuracy with senior sonographer.

169 **Statistical analysis**

170 The statistical analysis was performed with the STATA/SE 14.1 software (StataCorp ,
171 4905 Lakeway Drive, College Station, Texas , USA) and R 4.0.2²⁸. The ggplot2²⁹ for
172 R was used for diagram plotting. Confusion matrix³⁰ was used for validation of the
173 accuracy of the model. The confusion matrix was plotted by ggplot2 packages. The α
174 was set to 0.05. $P < \alpha$ was statistically significant.

175 **Hardware and software**

176 The sonographies were obtained from various ultrasound equipment, including
177 Mindray™ M5, Seote MyLab™ Alpha. The server used for training model installed

178 two 1080TI GPUs with Ubuntu 18.04 LTS platform. A Nvidia AGX Xavier edge
179 computing device providing web-based interface and model inference. These devices
180 install Ubuntu 18.04 LTS with PyTorch 1.3 and Flask library installed.

181 **Result**

182 **Dataset**

183 According to our dataset, we had 2250 training images, 190 images validation images
184 and 376 testing images. In the training dataset, for CL and CE1-5 type, 140, 172, 198,
185 280, and 472 images were trained, respectively. For AE lesion, 420, 290, and 278
186 images were trained, respectively.

187 **Training the model and publishing**

188 Model was trained for about 10 hours to achieved convergence after 90 epochs
189 (multiple times of training for optimizing). The generalized dice score of Classifier
190 for validation set was 0.77, the Dice score of Shaper for validation set was, 0.86. The
191 web-based interface uses 3.1s for average for single image analysis in current
192 hardware configuration (each service has six times of inferences).

193 **Comparison with human sonographer**

194 Total of 376 samples were inputted into AI model, archived 85%(302/376) accuracy.
195 In contrast, total of 85 samples were classified by senior sonographer achieved

196 72%(61/85) accuracy. The accuracy of AI model was higher than senior sonographer
197 (p-value=0.01 by chi-square test)(Table 1).

198 However, the difference for each types of lesions were not statically significant
199 (except CE5, the human sonographer achieved extremely low accuracy). The result
200 were represented as confusion matrix(Figure 3 and Figure 4).

201 **Discussion**

202 **Model outperforms trained sonographers**

203 According to our testing, our deep learning model outperformed fast-trained
204 sonographer in echinococcosis classification task. The U-net based model has been
205 proven to be efficient for smaller size of training set. It could be highly useful for
206 local paramedics while managing echinococcosis patients.

207 **Background of the disease**

208 *Echinococcus granulosus* life cycle involves dogs and other canids as definitive hosts
209 for the intestinal tapeworm as intermediate hosts for metacestode (larval) stage. The
210 metacestode (echinococcal cyst). In humans, the slowly growing hydatid cysts can
211 attain a volume of several liters and contain many thousands of protoscolices³¹.
212 The life cycle of *E. multilocularis* involves small rodent intermediate hosts, such as
213 arvicolid, wild or domestic canid definitive hosts, such as red or arctic foxes, jackals,
214 wolves, or dogs. Humans are aberrant intermediate hosts acquiring the infection

215 through ingestion of eggs shed in the feces of definitive hosts. AE is of increasing
216 concern globally due to the geographical spread of the parasite, its increasing
217 prevalence in animals from endemic areas, the absence of a vaccine, and the lack of
218 active control measures to prevent the infection³².

219 **Web-based application improved accessibility of AI** 220 **tools**

221 Deep learning applications are provided as source code deposited on Github³³ or
222 similar platform for sharing. It is not feasible for many medical related applications,
223 since the privacy concern of the data. The run-time environments of the source code
224 were varied in wide range due to different platforms, different frameworks, and
225 versions used. Here we used open-access web-based tool available for PC and mobile
226 will significantly increase the accessibility for many remote areas.

227 **Clinical implications**

228 We demonstrated the AI model achieved better performance than sonographer, it
229 could help doctors in remoted area for echinococcosis lesion typing. The lesion typing
230 for echinococcosis was important for disease managements.

231 According to the consensus on cystic echinococcosis diagnosis and treatment, five
232 types of lesions were recommended for classification. A natural grouping of the cysts
233 into three relevant groups: active (CE1 and 2), transitional (CE3) and inactive (CE4

and 5). The lesion classification is critical to patients managements, according to the consensus, CE4 and 5 indicated an inactive infection, watch and wait was recommended, if patients with CE1-3, the consensus suggested an active larva infection, radical medical surgery or/and ABZ should be suggested^{11,2}.

Comparing CE, AE is more problematic, surgical resection, chemotherapy, early diagnosis, and multidisciplinary discussion contributed to the successful treatment for AE cases¹¹. AE lesions behave “cancer-like”³⁴, radical surgery is the first choice in all cases suitable for totally resection¹¹. Benzimidazoles should be used for all cases¹¹, and liver transplantation as an alternative to palliative surgery, however, has not been shown to be superior to long-term conservative therapy³⁵.

244

245

1. Moro, P. & Schantz, P. M. Echinococcosis: a review. *Int. J. Infect. Dis.* **13**, 125–133 (2009).
2. Wen, H. *et al.* Echinococcosis: Advances in the 21st Century. *Clin. Microbiol. Rev.* **32**, e00075-18, /cmr/32/2/CMR.00075-18.atom (2019).
3. Wang, X. Q., Han, X. M., Tian, Q. S., Zhao, S. Y. & A, J. D. [Hepatic cystic and alveolar echinococcosis co-infections: a report of 3 cases]. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi Chin. J. Schistosomiasis Control* **32**, 213–216 (2019).
4. Deplazes, P. *et al.* Global Distribution of Alveolar and Cystic Echinococcosis. in *Advances in Parasitology* vol. 95 315–493 (Elsevier, 2017).
5. Guo, B. *et al.* High endemicity of alveolar echinococcosis in Yili Prefecture, Xinjiang Autonomous Region, the People’s Republic of China: Infection status in different ethnic communities and in small mammals. *PLoS Negl. Trop. Dis.* **15**, e0008891 (2021).
6. Wang, Q. *et al.* Disease burden of echinococcosis in Tibetan communities—A significant public health issue in an underdeveloped region of western China. *Acta Trop.* **203**, 105283 (2020).

- 262 7. Shan, X. *et al.* Health-related quality of life (HRQoL) associated with
263 echinococcosis patients in Tibetan communities in Shiqu County, China: a
264 case-control study. *Qual. Life Res.* **29**, 1559–1565 (2020).
- 265 8. Nationa echinococcosis Control programe(in Chinese).
266 http://www.gov.cn/zhengce/zhengceku/2020-09/03/content_5539743.htm.
- 267 9. Eckert, J. & Deplazes, P. Biological, Epidemiological, and Clinical Aspects of
268 Echinococcosis, a Zoonosis of Increasing Concern. *Clin. Microbiol. Rev.* **17**, 107–135
269 (2004).
- 270 10. Gupta, N. *et al.* Hepatic Hydatid: PAIR, Drain or Resect? *J. Gastrointest. Surg.*
271 **15**, 1829–1836 (2011).
- 272 11. Brunetti, E., Kern, P. & Vuitton, D. A. Expert consensus for the diagnosis and
273 treatment of cystic and alveolar echinococcosis in humans. *Acta Trop.* **114**, 1–16
274 (2010).
- 275 12. Larrieu, E. *et al.* Epidemiology, diagnosis, treatment and follow-up of cystic
276 echinococcosis in asymptomatic carriers. *Trans. R. Soc. Trop. Med. Hyg.* **113**, 74–80
277 (2019).
- 278 13. Brunetti, E. *et al.* Ultrasound and Cystic Echinococcosis. *Ultrasound Int. Open*
279 **04**, E70–E78 (2018).
- 280 14. Casulli, A., Barth, T. F. E. & Tamarozzi, F. Echinococcus multilocularis. *Trends*
281 *Parasitol.* **35**, 738–739 (2019).
- 282 15. Sulima, M. *et al.* Ultrasound images in hepatic alveolar echinococcosis and
283 clinical stage of the disease. *Adv. Med. Sci.* **64**, 324–330 (2019).
- 284 16. Kern, P. *et al.* WHO classification of alveolar echinococcosis: Principles and
285 application. *Parasitol. Int.* **55**, S283–S287 (2006).
- 286 17. Kratzer, W. Proposal of an ultrasonographic classification for hepatic alveolar
287 echinococcosis: Echinococcosis multilocularis Ulm classification-ultrasound. *World J.*
288 *Gastroenterol.* **21**, 12392 (2015).
- 289 18. Yang, S.-J. *et al.* A remote management system for control and surveillance of
290 echinococcosis: design and implementation based on internet of things. *Infect. Dis.*
291 *Poverty* **10**, 50 (2021).
- 292 19. Ahn, J. C., Connell, A., Simonetto, D. A., Hughes, C. & Shah, V. H. The
293 application of artificial intelligence for the diagnosis and treatment of liver diseases.
294 *Hepatol. Baltim. Md* (2020) doi:10.1002/hep.31603.
- 295 20. Borhani, M. *et al.* Cystic echinococcosis in the Eastern Mediterranean region:
296 Neglected and prevailing! *PLoS Negl. Trop. Dis.* **14**, e0008114 (2020).
- 297 21. Xin, S. *et al.* Automatic lesion segmentation and classification of hepatic
298 echinococcosis using a multiscale-feature convolutional neural network. *Med. Biol.*
299 *Eng. Comput.* **58**, 659–668 (2020).
- 300 22. Yu, Q., Xiao, N., Han, S., Tian, T. & Zhou, X.-N. Progress on the national
301 echinococcosis control programme in China: analysis of humans and dogs population
302 intervention during 2004–2014. *Infect. Dis. Poverty* **9**, 137 (2020).

303 23. Sichuan Institute of Echinococcosis Clinical Research. Guide line for complex
304 alveolar echinococcosis. *Chin. J. Bases Clin. Gen. Surg.* **27**, 18 (23).
305 24. Kentaro, W. *labelme: Image Polygonal Annotation with Python*. (2016).
306 25. U-Net: Convolutional Networks for Biomedical Image Segmentation.
307 <https://lmb.informatik.uni-freiburg.de/people/ronneber/u-net/>.
308 26. Ronneberger, O., Fischer, P. & Brox, T. U-Net: Convolutional Networks for
309 Biomedical Image Segmentation. in *Medical Image Computing and*
310 *Computer-Assisted Intervention – MICCAI 2015* (eds. Navab, N., Hornegger, J.,
311 Wells, W. M. & Frangi, A. F.) vol. 9351 234–241 (Springer International Publishing,
312 2015).
313 27. Oktay, O. *et al.* Attention U-Net: Learning Where to Look for the Pancreas.
314 *ArXiv180403999 Cs* (2018).
315 28. *R Core Team* (2020). *R: A language and environment for statistical computing*. *R*
316 *Foundation for Statistical Computing, Vienna, Austria*. URL
317 <https://www.R-project.org/>.
318 29. Hadley, W. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New
319 York, 2016).
320 30. Foody, G. M. Status of land cover classification accuracy assessment. *Remote*
321 *Sens. Environ.* **80**, 185–201 (2002).
322 31. Moro, P. & Schantz, P. M. Echinococcosis: a review. *Int. J. Infect. Dis.* **13**,
323 125–133 (2009).
324 32. Casulli, A., Barth, T. F. E. & Tamarozzi, F. Echinococcus multilocularis. *Trends*
325 *Parasitol.* **35**, 738–739 (2019).
326 33. GitHub: Where the world builds software · GitHub. <https://github.com>.
327 34. Reinehr, M. *et al.* Pathology of Echinococcosis: A Morphologic and
328 Immunohistochemical Study on 138 Specimens With Focus on the Differential
329 Diagnosis Between Cystic and Alveolar Echinococcosis. *Am. J. Surg. Pathol.* **44**,
330 43–54 (2020).
331 35. Salm, L. A., Lachenmayer, A., Perrodin, S. F., Candinas, D. & Beldi, G. Surgical
332 treatment strategies for hepatic alveolar echinococcosis. *Food Waterborne Parasitol.*
333 **15**, e00050 (2019).

334

335

336

337 Table 1. Comparison of human and AI

338

	CL	CE1	CE2	CE3	CE4	CE5	AE1	AE2	AE3	total	acc.*	p-values#
CL-Human	2	2	0	0	0	0	0	0	0	4	0.50	0.27
CL-AI	38	4	4	0	4	0	0	0	0	50	0.76	
CE1-Human	1	11	0	0	0	0	0	0	0	12	0.92	0.68
CE1-AI	4	40	4	0	2	0	0	0	0	50	0.80	
CE2-Human	0	0	2	1	0	0	0	0	0	3	0.67	1.00
CE2-AI	0	0	38	9	3	0	0	0	0	50	0.76	
CE3-Human	0	0	2	8	1	0	0	0	0	11	0.73	1.00
CE3-AI	0	0	6	38	6	0	0	0	0	50	0.76	
CE4-Human	0	0	0	1	14	7	0	0	0	22	0.64	0.15
CE4-AI	0	0	8	2	40	0	0	0	0	50	0.80	
CE5-Human	1	0	0	0	2	0	1	0	0	4	0.00	0.00
CE5-AI	0	1	0	0	1	40	8	0	0	50	0.80	
AE1-Human	0	0	0	0	2	0	19	0	0	21	0.90	0.21
AE1-AI	0	0	0	0	0	0	49	0	1	50	0.98	
AE2-Human	0	0	0	0	0	0	1	3	0	4	0.75	1.00
AE2-AI	0	2	0	0	0	0	2	10	0	14	0.71	
AE3-Human	0	0	0	0	0	1	1	0	2	4	0.50	0.55
AE3-AI	0	0	0	0	0	2	1	0	9	12	0.75	

339 * acc. accuracy=right classification/total samples

340 # by using Fisher' exact tests.

341

342

Figure 1. The architecture of Two U-nets model.

Figure 2. Web Application structure of model

362
363
364
365
366
367
368
369

Figure 3.

370
371 The confusion matrix of AI prediction.

372
373
374
375
376
377
378
379
380
381
382
383
384
385
386

387
388
389
390
391
392
393
394
395

Figure 4.

396
397
398

The confusion matrix of human sonographer.