Breakthrough SARS-CoV-2 infections in MS patients on disease modifying therapies

Irene Schiavetti¹, Cinzia Cordioli², Maria Laura Stromillo³, Maria Teresa Ferrò⁴, Alice Laroni^{5,6}, Eleonora Cocco⁷, Gaia Cola⁸, Livia Pasquali⁹, Maria Teresa Rilla¹⁰, Elisabetta Signoriello¹¹, Rosa Iodice¹², Alessia Di Sapio¹³, Roberta Lanzillo¹⁴, Francesca Caleri¹⁵, Pietro Annovazzi¹⁶, Antonella Conte^{17,18}, Giuseppe Liberatore¹⁹, Francesca Ruscica²⁰, Renato Docimo²¹, Simona Bonavita²², Monica Ulivelli²³, Paola Cavalla²⁴, Francesco Patti²⁵, Diana Ferraro²⁶, Marinella Clerico²⁷, Paolo Immovilli²⁸, Massimiliano Di Filippo²⁹, Marco Salvetti^{30,18}, Maria Pia Sormani^{1,5} and the “Breakthrough infections in MS” study group

1. Department of Health Sciences, Section of Biostatistics, University of Genova, Italy
2. Centro Sclerosi Multipla ASST Spedali Civili di Brescia
3. Clinica Neurologica e Malattie Neurometaboliche, Università degli Studi di Siena
4. Neuroimmunology, Center for Multiple Sclerosis, Cardiocerebrovascular Department, Neurological Unit, ASST Crema
5. IRCCS Ospedale Policlinico San Martino, Genova, Italy
6. Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Italy
7. Centro Sclerosi Multipla Ospedale Binaghi Cagliari
8. "Department of Systems Medicine, Multiple Sclerosis Clinical and Research Unit, Tor Vergata, University, Rome, Italy"
9. Department of Clinical and Experimental Medicine, Neurology Unit, University of Pisa, Italy
10. Department of Neurology, Imperia Hospital, Imperia, Italy
11. II Division of Neurology, University of Campania Luigi Vanvitelli
12. Department of Neurosciences, Reproductive Sciences and Odontostomatology, University of Naples "Federico II", Naples, Italy
13. Department of Neurology, Regina Montis Regalis Hospital, Mondovì, Italy
14. Federico II University of Naples
15. Department of Neurology, MS Center, F. Tappeiner Hospital Meran (BZ), Italy
16. UOC Centro Sclerosi Multipla, ASST Valle-Olona, PO di Gallarate (VA)
17. Department of Human Neuroscience, Sapienza, University of Rome, Italy.
18. IRCCS Neuromed, Pozzilli (IS), Italy
19. Neuromuscular and Neuroimmunology Unit, IRCCS Humanitas Research Hospital, Rozzano, Italy
20. U.O.C. Neurologia e Centro SM - Fondazione Istituto G. Giglio, Cefalù (Italy)
21. Multiple Sclerosis Center, Aversa Hospital "San Giuseppe Moscati", ASL Caserta, Aversa (Ce) Italy
22. Dipartimento di Scienze Mediche e Chirurgiche avanzate, Università della Campania Luigi Vanvitelli, Naples-Italy
23. Department of Medical Sciences, Surgery and Neurosciences, University of Siena
24. Multiple Sclerosis Center and 1st Division of Neurology, Department of Neuroscience, City of Health and Science University Hospital of Turin, Italy

25. Department of Medical and Surgical Sciences and Advanced Technologies, GF Ingrassia, University of Catania, Catania, Italy.
26. Department of Biomedical, Metabolic and Neurosciences, University of Modena and Reggio Emilia, Modena, Italy.
27. Clinical and Biological Sciences Department, University of Torino
28. Guglielmo da Saliceto Hospital, Piacenza
29. Department of Medicine and Surgery, Section of Neurology, University of Perugia, Perugia, Italy
30. Department of Neuroscience, Mental Health and Sensory Organs, Sapienza University of Rome, Rome, Italy

Corresponding author: **Maria Pia Sormani**, Department of Health Sciences, Via Pastore 1, 16132, University of Genova, Italy

Email: mariapia.sormani@unige.it

Tel: +39-3669937472

Keywords: Multiple Sclerosis, Covid-19 vaccination, Breakthrough infections

Number of characters in the title: 80

Number of words: 1394

Number of figures: 1

Number of tables: 1

Number of references: 11

60 Abstract

61 **Background.** Patients with Multiple Sclerosis (pwMS) treated with anti-CD20 or fingolimod showed a
62 reduced humoral response to SARS-CoV-2 vaccines. In this study we aimed to monitor the risk of
63 breakthrough SARS-CoV-2 infection in pwMS on different Disease Modifying Therapy (DMT).

64 **Methods.** Data on number of vaccinated patients and of patients with a breakthrough infection were
65 retrospectively collected in 27 Italian MS centers. We estimated the rate of breakthrough infections and of
66 infection requiring hospitalization per DMT.

67 **Findings.** 19641 vaccinated pwMS were included in the database. After a median follow-up of 8 months, we
68 observed 137 breakthrough infections. As compared to the other DMTs, the rate of breakthrough infections
69 was significantly higher on ocrelizumab (0.57% vs 2.00%, RR=3.55,95%CI=2.74-4.58, p<0.001) and
70 fingolimod (0.58% vs 1.62%, RR=2.65,95%CI=1.75-4.00, p<0.001), while there were no significant
71 differences in any other DMT group. In the ocrelizumab group the hospitalization rate was 16.7% vs 19.4%
72 in the pre-vaccination era (RR=0.86,p=0.74) and it was 3.9% in all the other DMT groups vs 11.9% in the
73 pre-vaccination period (RR=0.33,p=0.02).

74 **Interpretation.** The risk of breakthrough SARS-CoV-2 infections is higher in patients treated with
75 ocrelizumab and fingolimod, and the rate of severe infections was significantly reduced in all the DMTs
76 excluding ocrelizumab.

77

Introduction

Several recent studies evaluated the effect of vaccination against SARS-CoV-2 in patients with multiple sclerosis (pwMS) treated with disease-modifying therapies (DMTs). There is wide consensus that the use of anti-CD20 monoclonal antibodies and fingolimod are associated with an impaired virus-specific humoral immune response as compared to all the other DMTs¹⁻⁴. On the other hand, there is also growing evidence that vaccinated pwMS treated with anti-CD20 generated robust virus specific CD4 and CD8 T cell responses⁴⁻⁵, while these are slightly reduced in fingolimod treated patients⁵. A preliminary follow-up study of 344 fully vaccinated pwMS on DMT reported 13 breakthrough infections, 10 of which were in patients on anti-CD20 therapy and the remaining 3 on fingolimod⁶, suggesting a relevant role of antibodies in preventing the infection. The French registry recently reported a case series of 18 pwMS who had Covid-19 after two doses of BNT162b2-vaccination, 13 of which treated with anti-CD20 and four with fingolimod⁷. Finally, the clinical follow up of the CovXiMS study¹ evaluating humoral response in 1705 pwMS who received two doses of mRNA vaccines⁸, reported 23 breakthrough infections over a 6 month follow up. The risk of infection was associated with lower SARS-CoV-2 antibody levels measured after 4 weeks from the second vaccine dose⁸.

Against this background and taking advantage of the large network of MS centers within the Italian Alliance against Covid-19 promoted by the Italian MS Society, we collected data from 27 Italian MS centers on the number of vaccinated patients and the number of patients who had a breakthrough infection in each DMT group, in the period preceding the spread of the Omicron variant, that started its massive diffusion in Italy after the December 2021 holiday season. Aim of this study is to estimate the rate of breakthrough infections per DMT class on a large sample of vaccinated pwMS and to compare the rates of severe infections to the rate observed in Italy in the pre-vaccination era⁹.

Patients and Methods

Study design and participants

This was a retrospective data collection conducted in 27 Italian MS centers on pwMS undergoing the SARS-CoV-2 vaccination. Each MS center was requested to report the number of pwMS vaccinated by two mRNA vaccine doses (BNT162b2 (Pfizer Inc, and BioNTech), or mRNA-1273 (Moderna Tx, Inc)) in each DMT group from March 2021 to December 25, 2021. Data cutoff was set before the spread of the Omicron variant in Italy, since on December 23, 2021 the percentage of Omicron infections was estimated to be 28% (<https://www.iss.it/primo-piano>, accessed on December 25, 2021). Breakthrough infections occurred within 8 months, defined as a PCR-confirmed test after 14 days from the second or the third vaccine dose, were extracted from the platform dedicated to Covid-19 data collection in pwMS (MuSC-19 database¹⁰) for the participating centers. The post-vaccination SARS-CoV-2 infection was recorded in a dedicated Case Report Form (CRF).

The study is done in compliance with the principles of the Declaration of Helsinki. The study was approved by the regional ethics committee of Liguria (University of Genoa; n 130/2020–DB id 10433) and at a national level by the Italian Medicines Agency. Written informed consent was obtained from all participants before starting any study procedures.

Primary Outcome: breakthrough infection

The primary objective of this analysis was to compare the incidence of breakthrough SARS-CoV-2 infections among the vaccinated pwMS in each DMT group. These conditions entail a PCR-confirmed swab, and a time lag of at least 14 days from a full vaccination cycle (after the second or third vaccination dose, or after the first dose following a Covid-19 infection).

Statistical analysis

The percentage of patients with a breakthrough infection in the different DMT groups was calculated. 95% Confidence Intervals (CI) were estimated using the normal approximation to the binomial calculation¹¹. Difference of rate of infections between DMT groups were estimated by Risk Ratios (RR) and evaluated by Chi-square tests. Difference of rate of infections in the first 4 months vs the second 4 months of follow up were estimated by ORs and evaluated by the McNamar test for paired data.

Funding

FISM [2021/Special-Multi/001].

Results

Data were collected between March 1, 2021 and December 24, 2021. 19641 pwMS who had a full vaccination cycle with an mRNA vaccine (2 or 3 vaccine dose, or 1 vaccine dose after Covid-19 infection) were included in the database. The number of vaccinated pwMS in each DMT group is reported in Table 1. The mean follow-up time was 249 days (range 99-354). Among them, 137 breakthrough infections were observed (26 (19.0%) after the third dose, 1 after Covid-19 infection and one dose) over a mean interval after the last vaccine dose of 142 days (range 14-262) (Table 1). Over the whole follow-up of about 8 months, we compared the proportion of patients with breakthrough infections in each DMT group to the pooled proportion of the patients on all the other DMTs (Figure 1, panel A). The rate of breakthrough infections was significantly higher in patients treated with ocrelizumab (2.00%, 95%CI=1.36-2.66) than in patients treated with all the other DMTs (0.57%, 95%CI=0.46%-0.68%) with a RR=3.55, 95%CI=2.74-4.58, $p<0.001$; the same was observed in patients treated with fingolimod who had a higher rate of breakthrough infections (1.62%, 95%CI= 1.02%-2.21%) than the patients treated with all the other DMTs (0.61%, 95%CI=0.50-0.72) with a RR=2.65, 95%CI= 1.75-4.00, $p<0.001$. Among the patients who had the SARS-CoV-2 infection, 10 (7.3%) had a severe disease course and were hospitalized. Six patients treated with ocrelizumab were hospitalized and in this group the rate of hospitalization was 16.7%, slightly lower but not significantly different than the pre-vaccination rate observed in Italy (19.4%) in the same DMT group⁷ (relative reduction=14%, RR=0.86, 95%CI=0.38-1.91, $p=0.74$). In the fingolimod group we observed just 1 hospitalized patient (3.6%). The rate of hospitalization was 3.9% in all the other DMT groups as compared to 11.9% in the pre-vaccination period⁷ (relative reduction=67%, RR=0.33, 95%CI=0.13-0.88, $p=0.02$). One patient in ocrelizumab was admitted to the Intensive Care Unit (ICU) and recovered.

Figure1, panel B, reports the rate of breakthrough infections in two time periods of equal duration: the first 4 months following the last vaccination dose vs the period 4-8 months after the last vaccination dose in patients treated with ocrelizumab, fingolimod and all the other DMTs. The rate in patients treated with

ocrelizumab and fingolimod was not significantly affected by the time since vaccination (ocrelizumab: 0-4 months after vaccination: 0.84%, 4-8 months after vaccination: 1.18%, OR=1.40, p=0.31; fingolimod: 0-4 months after vaccination: 0.86%, 4-8 months after vaccination: 0.76%, OR=0.88, p=0.75). In all the other DMT groups the rate is much lower (0-4 months after vaccination: 0.14%) and it was significantly increased after 4 months from the last vaccine dose (4-8 months after vaccination: 0.32%, OR=2.32, 95%CI=1.38-4.01, p<0.001).

Discussion

This study on a large sample of pwMS who received a full vaccination cycle confirms that the risk of contracting SARS-CoV-2 infection after Covid-19 m-RNA vaccines is higher in pwMS on anti-CD20 monoclonal antibodies or fingolimod. We observed just one admission to ICU and no deaths. Despite the small sample of 137 infections, two results emerge. First, in our cohort, among the infected patients after vaccination treated with ocrelizumab the hospitalization rate is very similar to the hospitalization rate of patients on the same treatment in the pre-vaccination era⁸, while it is reduced by 67% in pwMS in other DMTs. However, we must consider that this result can be confounded by an increased propensity of clinician to admit to hospital pwMS on ocrelizumab who develop Covid-19, because of previous studies showing that these patients are at a higher risk for a severe course⁸. Second, as expected⁹, the vaccine-induced protection from the disease is waning with time since vaccination, and this is more evident in patients treated with DMTs other than ocrelizumab and fingolimod, who already had low antibody levels soon after the vaccination. In fact, while the infection rate is similar in the first and in the second four months after vaccination in patients on ocrelizumab and fingolimod, and consistently higher than in patients on other DMTs, the initial protective effect is vanishing with time for patients in the other DMTs group, who had a good level of antibody response four weeks after vaccination¹. This study complements the information of previous studies reporting the antibody levels after anti-SARS-Cov-2 vaccination in pwMS on different DMTs¹⁻⁷, suggesting that antibodies play a dominant role in preventing Covid-19 infections and their severe consequences.

References

1. Sormani MP, Inglese M, Schiavetti I et al. Effect of SARS-CoV-2 mRNA vaccination in MS patients treated with disease modifying therapies. *EBioMedicine*. 2021 Oct;72:103581. doi: 10.1016/j.ebiom.2021.103581.
2. Achiron A, Mandel M, Dreyer-Alster S, et al. Humoral immune response to COVID-19 mRNA vaccine in patients with multiple sclerosis treated with high-efficacy disease-modifying therapies. *Ther Adv Neurol Disord*. 2021. doi:10.1177/17562864211012835
3. Tallantyre EC, Vickaryous N, Anderson V et al. COVID-19 Vaccine Response in People with Multiple Sclerosis. *Ann Neurol* 2021; Oct 22. doi: 10.1002/ana.26251. Online ahead of print
4. Tortorella C, Aiello A, Gasperini C et al. Humoral- and T-Cell-Specific Immune Responses to SARS-CoV-2 mRNA Vaccination in Patients With MS Using Different Disease-Modifying Therapies. *Neurology* in press
5. Apostolidis SA, Kakara M, Painter MM, et al. Cellular and humoral immune responses following SARS-CoV-2 mRNA vaccination in patients with multiple sclerosis on anti-CD20 therapy. *Nat Med*. 2021 Nov;27(11):1990-2001. doi: 10.1038/s41591-021-01507-2.
6. Rose DR, Mahadeem AZ, Carlson AK et al. Clinical features and outcomes of COVID-19 despite SARS-CoV-2 vaccination in people with multiple sclerosis. *Mult Scler J Exp Transl Clin*. 2021 Nov 26;7(4):20552173211057110. doi: 10.1177/20552173211057110. eCollection 2021 Oct.
7. Januel E , De Seze J, Vermersch P et al. Post-vaccine COVID-19 in patients with multiple sclerosis or neuromyelitis optica. *Mult Scler J*. 2022; doi: 10.1177/ 13524585211049737.
8. Sormani MP, Schiavetti I, Carmisciano L et al. COVID-19 Severity in Multiple Sclerosis: Putting Data Into Context. *Neurol Neuroimmunol Neuroinflamm*. 2021 Nov 9;9(1):e1105. doi: 10.1212/NXI.0000000000001105. Print 2022 Jan.
9. Sormani MP, Schiavetti I, Inglese M et al. Breakthrough SARS-CoV-2 infections after COVID-19 mRNA vaccination in MS patients on disease modifying therapies. . *medRxiv* 2021.12.23.21268177; doi: <https://doi.org/10.1101/2021.12.23.21268177>.

10. Sormani MP, De Rossi N, Schiavetti I, et al. Disease-Modifying Therapies and Coronavirus Disease 2019 Severity in Multiple Sclerosis. *Ann Neurol*. 2021;89(4):780-789. Doi:10.1002/ana.26028
11. Brown, Lawrence D.; Cai, T. Tony; DasGupta, Anirban (2001). "Interval Estimation for a Binomial Proportion". *Statistical Science*. 16 (2): 101–133.

Acknowledgements

Supported by FISM [2021/Special-Multi/001]

Disclosures

Annovazzi received honoraria for lecturing and participation in advisory boards, and/or travel expenses for attending congresses and meetings from Almirall, Biogen, BMS-Celgene, Merck, Novartis, Roche, Sanofi-Genzyme, Teva Italia, and Viatrix.

Bonavita received speaker and/or advisors board fee from Biogen, Novartis, Roche, viatrix, Merck serono.

Caleri received honoraria for advisory board and/or for public speaking, and/or travel grant, from Biogen, Merck, Teva, Novartis, Sanofi-Genzyme, Roche.

Cavalla has received advisory board membership, speaker honoraria or travel grants to attend national and international conferences from Biogen, Merck-Serono, Teva, Roche, Novartis, Cellgene-BMS and Sanofi-Genzyme.

Clerico received grants and consulting fees from Merck, Biogen, Novartis, Sanofi-Genzyme, Roche, Almirall.

Cola, Pasquali, Cocco, Ferrò, Liberatore, Rilla, Stromillo have nothing to disclose.

Conte reports speaking honoraria from Merck, Sanofi, Novartis, Biogen, Roche, Bristol Myers Squibb, Almirall and has received research support from Roche, Biogen and Merck.

Cordioli received personal compensation for advisory board and speaking for Merck Serono, Novartis, Almirall, Biogen and Roche.

Di Filippo participated on advisory boards for and received research support, speaker/writing honoraria and funding for traveling from Bayer, Biogen Idec, Genzyme, Merck, Mylan, Novartis, Roche, Teva and Viatrix

Di Sapio received personal compensation for speaking and consulting by Biogen, Novartis and Genzyme and has been reimbursed by Merck, Biogen, Genzyme and Roche for attending several conferences.

Docimo received grants from Roche, Novartis, Biogen, Merck, Viatrix, Genzyme.

Ferraro received travel grants and/or speaker/advisory board honoraria from Biogen, Roche, Novartis, TEVA, Sanofi Genzyme, Merck Serono.

Immovilli reports personal fees from Roche, Biogen, Merck.

Iodice received speaker honoraria and/or consultancy from Biogen, Teva, Genzyme, Merck, Almirall, Roche.

243 Lanzillo received personal compensations for speaking or consultancy from Biogen, Teva, Genzyme, Merck,
244 Mylan, Novartis and Roche.

245 Laroni received personal compensations from Merck, Biogen, Novartis, Roche, Almirall.

246 Patti received personal compensation for serving on advisory board Almirall, Bayer, Biogen, Bristol Meyers
247 Squibb, Merck, Novartis, Roche and Sanofi; he further received unrestricted research grants by Biogen,
248 Merck, Roche, FISM, University of Catania and Reload (onlus patients association).

249 Ruscica received speaker and/or advisors board fee from Merck, Novartis, Biogen, Genzyme.

250 Salvetti reports grants and personal fees from Biogen, Merck, Novartis, Roche, Sanofi, Teva, grants from
251 Italian Multiple Sclerosis Foundation, grants from Sapienza University of Rome.

252 Schiavetti received consulting fees from Hippocrates Research, NovaNeuro, Sakura Italia, ADL
253 Farmaceutici, Associazione Commissione Difesa Vista Onlus.

254 Signoriello received speaker honoraria and/or consultancy from Biogen, Teva, Genzyme, Merck, Novartis,
255 Almirall, Roche.

256 Sormani received consulting fees from Merck, Biogen, Novartis, Sanofi, Roche, Geneuro, GSK, Medday,
257 Immunic.

258 Ulivelli received consulting fees from Biogen, Novartis, Serono.

Table 1: Characteristics of patients with breakthrough infections (N = 137)

Female sex		85 (62.0)
Age, years		42.3 ± 10.70
BMI (kg/m²)		24.6 ± 4.95
MS phenotype	Primary progressive	9 (6.6)
	Relapsing remitting	116 (84.7)
	Secondary progressive	7 (5.1)
	Missing data	5 (3.6)
Disease duration, months		99.5 (44.0 - 182.0)
Last EDSS before Covid-19 infection		2.0 (1.0 - 3.5)
Relapse in the six months before Covid-19 infection		7 (5.1)
Number of breakthrough infections in each DMT/number of vaccinated patients, (%)	alemtuzumab	0/371 (0.0)
	azathioprine	0/298 (0.0)
	cladribine	4/570 (0.70)
	dimethyl fumarate	22/2668 (0.82)
	fingolimod	28/1733 (1.61)
	glatiramer acetate	4/1514 (0.26)
	interferon	7/2452 (0.29)
	natalizumab	15/1843 (0.81)
	ocrelizumab	36/1794 (2.00)
	rituximab	3/364 (0.82)
	teriflunomide	10/1379 (0.73)
	other	0/389 (0.0)
	untreated	8/4266 (0.19)
Boost COVID-19 vaccination		26 (19.0)
Heterologous vaccine		4 (2.9)
Covid severity	Asymptomatic, viral RNA detected	16 (11.8)
	Symptomatic, independent	95 (69.3)
	Symptomatic, assistance needed	16 (11.8)
	Hospitalized, no oxygen therapy	4 (2.9)
	Oxygen by mask or nasal prongs	5 (3.7)
	Intubation and mechanical ventilation, piO ₂ /FiO ₂ ≥150 or SpO ₂ /FiO ₂ ≥200	1 (0.7)

Results are expressed as count (%), mean ± Standard Deviation, or median [Inter Quartile Range], as appropriate.

MS=Multiple Sclerosis, BMI=Body Mass Index, EDSS=Expanded Disability Status Scale, DMT=Disease Modifying Therapy

261 Breakthrough infections in MS study group

262

Name Surname	Affiliation
Agostino Giulio Lanuto	Federico II university of Naples
Alessandro Giannotta	Department of Medical Sciences, Surgery and Neurosciences, University of Siena
Alessia Di Sapia	Department of Neurology, Regina Montis Regalis Hospital, Mondovì, Italy
Alice Laroni	1 - Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Italy AND IRCCS Ospedale Policlinico San Martino, Genova, Italy 2 - IRCCS Ospedale Policlinico San Martino, Genova, Italy
Anastasia Alteno	Multiple Sclerosis Center, Department of Neuroscience, City of Health and Science University Hospital of Turin, Italy
Antonella Conte	1- Department of Human Neuroscience, Sapienza, University of Rome, Italy. 2- IRCCS Neuromed, Pozzilli (IS), Italy
Carlo Serrati	Ospedale Civile Imperia
Caterina Lapucci	IRCCS Ospedale Policlinico San Martino, Genova, Italy
Chiara Rosa Mancinelli	Centro Sclerosi Multipla ASST Spedali Civili di Brescia
Cinzia Cordioli	Centro Sclerosi Multipla ASST Spedali Civili di Brescia
Clara Grazia Chisari	Department of Medical and Surgical Sciences and Advanced Technologies, GF Ingrassia, University of Catania, Catania, Italy.
Daiana Bezzini	Department of Life Sciences, University of Siena
Damiano Baroncini	UOC Centro Sclerosi Multipla, ASST Valle-Olona, PO di Gallarate (VA)
Diana Ferraro	Department of Biomedical, Metabolic and Neurosciences, University of Modena and Reggio Emilia, Modena, Italy
Donata Guidetti	Guglielmo da Saliceto Hospital, Piacenza
Doriana Landi	Department of Systems Medicine, Multiple Sclerosis Clinical and Research Unit, Tor Vergata University, Rome, Italy
Eduardo Nobile-Orazio	1. Neuromuscular and Neuroimmunology Unit, IRCCS Humanitas Research Hospital, Rozzano, Italy 2. Department of Medical Biotechnology and Translational Medicine, Milan University, Milan, Italy
Elena Di Sabatino	Department of Medicine and Surgery, Section of Neurology, University of Perugia, Perugia, Italy
Eleonora Cocco	Centro Sclerosi Multipla Ospedale Binaghi Cagliari
Elisabetta Signoriello	II Division of Neurology, University of Campania Luigi Vanvitelli
Elvira Sbragia	Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Italy AND IRCCS Ospedale Policlinico San

	Martino, Genova, Italy
Emanuele Cassano	Department of Neurosciences, Reproductive Sciences and Odontostomatology, University of Naples "Federico II", Naples, Italy.
Enri Nako	Department of Neurology, Regina Montis Regalis Hospital, Mondovì, Italy
Fabio Della Cava	Ospedale Civile Imperia
Flora Govone	Department of Neurology, Regina Montis Regalis Hospital, Mondovì, Italy
Francesca Bianchi	Department of Clinical and Experimental Medicine, Neurology Unit, University of Pisa, Italy
Francesca Caleri	Department of Neurology, MS Center, F. Tappeiner Hospital Meran (BZ), Italy
Francesca Ruscica	U.O.C. Neurologia e Centro SM - Fondazione Istituto G. Giglio, Cefalù (Italy)
Francesca Vitetta	Neurology Unit, Azienda Ospedaliero-Universitaria of Modena, Modena, Italy
Francesco Patti	Department of Medical and Surgical Sciences and Advanced Technologies, GF Ingrassia, University of Catania, Catania, Italy.
Gabriele Siciliano	Department of Clinical and Experimental Medicine, Neurology Unit, University of Pisa, Italy
Gaia Cola	Department of Systems Medicine, Multiple Sclerosis Clinical and Research Unit, Tor Vergata University, Rome, Italy
Giacomo Lus	II Division of Neurology, University of Campania Luigi Vanvitelli
Gianmarco Abbadessa	Dipartimento di Scienze Mediche e Chirurgiche avanzate, Università della Campania Luigi Vanvitelli, Naples-Italy
Gianmarco Bellucci	Department of Neuroscience, Mental Health and Sensory Organs Sapienza University, S. Andrea Hospital, Rome
Girolama Alessandra Marfia	Department of Systems Medicine, Multiple Sclerosis Clinical and Research Unit, Tor Vergata University, Rome, Italy
Giuditta Ilaria Scarano	Department of Psychology, F. Tappeiner Hospital Meran (BZ), Italy
Giuseppe Liberatore	Neuromuscular and Neuroimmunology Unit, IRCCS Humanitas Research Hospital, Rozzano, Italy
Giuseppina Miele	Dipartimento di Scienze Mediche e Chirurgiche avanzate, Università della Campania Luigi Vanvitelli, Naples-Italy
Graziella Callari	U.O.C. Neurologia e Centro SM - Fondazione Istituto G. Giglio, Cefalù (Italy)
Guido Urbano	Dipartimento di Neuroscienze "Rita Levi Montalcini, Università di Torino
Irene Schiavetti	Department of Health Sciences, Section of Biostatistics, University of Genova, Italy
Jessica Frau	Centro Sclerosi Multipla Ospedale Binaghi Cagliari
Leonardo Malimpensa	Department of Human Neuroscience, Sapienza, University of Rome, Italy
Lia Allegorico	Nola Hospital "Santa Maria della Pietà", ASL Napoli 3 Sud, Nola (Na) Italy
Livia Pasquali	Department of Clinical and Experimental Medicine, Neurology Unit, University of Pisa, Italy

Lorena Lorefice	Centro Sclerosi Multipla Ospedale Binaghi Cagliari
Lorenzo Gaetani	Department of Medicine and Surgery, Section of Neurology, University of Perugia, Perugia, Italy
Lucia Ruggiero	Department of Neurosciences, Reproductive Sciences and Odontostomatology, University of Naples "Federico II", Naples, Italy
Marco Salvetti	Department of Neuroscience, Mental Health and Sensory Organs. Sapienza University, S. Andrea Hospital, Rome
Marco Vercellino	Multiple Sclerosis Center and 1st Division of Neurology, Department of Neuroscience, City of Health and Science University Hospital of Turin, Italy
Maria Chiara Buscarinu	Department of Neuroscience, Mental Health and Sensory Organs Sapienza University, S. Andrea Hospital, Rome
Maria Francesca Creta	1. Neuromuscular and Neuroimmunology Unit, IRCCS Humanitas Research Hospital, Rozzano, Italy 2. Department of Medical Biotechnology and Translational Medicine, Milan University, Milan, Italy
Maria Laura Stromillo	Clinica Neurologica e Malattie Neurometaboliche, Università degli Studi di Siena
Maria Pia Sormani	1- Department of Health Sciences, Section of Biostatistics, University of Genova, Italy 2- IRCCS Ospedale Policlinico San Martino, Genova, Italy
Maria Rosaria Pennacchio	Multiple Sclerosis Center, Aversa Hospital "San Giuseppe Moscati", ASL Caserta, Aversa (Ce) Italy
Maria Teresa Ferrò	Neuroimmunology, Center for Multiple Sclerosis, Cardiocerebrovascular Department, Neurological Unit, ASST Crema
Maria Teresa Rilla	Ospedale Civile Imperia
Marinella Clerico	Dipartimento di Scienze Cliniche e Biologiche, Università di Torino
Massimiliano Di Filippo	Department of Medicine and Surgery, Section of Neurology, University of Perugia, Perugia, Italy
Matteo Scialabba	U.O.C. Neurologia e Centro SM - Fondazione Istituto G. Giglio, Cefalù (Italy)
Mauro Zaffaroni	UOC Centro Sclerosi Multipla, ASST Valle-Olona, PO di Gallarate (VA)
Monica Ulivelli	Department of Medical Sciences, Surgery and Neurosciences, University of Siena
Nicola De Stefano	Clinica Neurologica e Malattie Neurometaboliche, Università degli Studi di Siena
Paola Cavalla	Multiple Sclerosis Center and 1st Division of Neurology, Department of Neuroscience, City of Health and Science University Hospital of Turin, Italy
Paola De Mitri	Guglielmo da Saliceto Hospital, Piacenza
Paolo Immovilli	Guglielmo da Saliceto Hospital, Piacenza
Patrizia Sola	Neurology Unit, Azienda Ospedaliero-Universitaria of Modena, Modena, Italy
Pietro Annovazzi	UOC Centro Sclerosi Multipla, ASST Valle-Olona, PO di Gallarate (VA)
Raffaele Nardone	1. Paracelsus Medical University, Department of Neurology, Salzburg-AU 2. Department of Neurology, F. Tappeiner Hospital Meran (BZ), Italy

Renato Docimo	Multiple Sclerosis Center, Aversa Hospital "San Giuseppe Moscati", ASL Caserta, Aversa (Ce) Italy
Roberta Lanzillo	Federico II university of Naples
Rosa Iodice	Department of Neurosciences, Reproductive Sciences and Odontostomatology, University of Naples "Federico II", Naples, Italy
Rosanna Missione	II Division of Neurology, University of Campania Luigi Vanvitelli
Sarah Rasia	Centro Sclerosi Multipla ASST Spedali Civili di Brescia
Sebastiano Arena	Department of Medical and Surgical Sciences and Advanced Technologies, GF Ingrassia, University of Catania, Catania, Italy.
Simona Bonavita	Dipartimento di Scienze Mediche e Chirurgiche avanzate, Università della Campania Luigi Vanvitelli, Naples-Italy
Simona Rolla	Dipartimento di Scienze Cliniche e Biologiche, Università di Torino
Vincenzo Brescia Morra	Federico II University of Naples
Viola Baione	Department of Human Neuroscience, Sapienza, University of Rome, Italy.

Figure legends

Figure 1. Cumulative incidence of breakthrough infections in patients in each DMT group (A) and breakthrough infection rates according to time since vaccination in ocrelizumab, fingolimod and other DMTs (B).

**Ocrelizumab and fingolimod had a percentage of breakthrough infections that is significantly higher ($p < 0.001$) than the percentage in all the other DMT groups.*