Association study of HLA with the kinetics of SARS-CoV-2 spike specific IgG antibody responses to BNT162b2 mRNA vaccine

Seik-Soon Khor¹, Yosuke Omae¹, Junko S. Takeuchi², Ami Fukunaga³, Shohei Yamamoto³, Akihito Tanaka⁴, Kouki Matsuda⁵, Moto Kimura², Kenji Maeda⁵, Gohzoh Ueda⁶, Tetsuya Mizoue³, Mugen Ujiie⁷, Hiroaki Mitsuya⁵, Norio Ohmagari⁷, Wataru Sugiura⁸, Katsushi Tokunaga¹

Affiliations:

1. Genome Medical Science Project, National Center for Global Health and Medicine, Tokyo, Japan
2. Department of Academic-Industrial Partnerships Promotion, Center for Clinical Sciences, National Center for Global Health and Medicine, Tokyo, Japan
3. Department of Epidemiology and Prevention, Center for Clinical Sciences, National Center for Global Health and Medicine, Tokyo, Japan
4. Department of Laboratory Testing, Center Hospital of the National Center for the Global Health and Medicine, Tokyo, Japan
5. Department of Refractory Viral Infection, Research Institute, National Center for Global Health and Medicine, Tokyo, Japan
6. Division of Core Diagnostics, Abbott Japan LLC, Tokyo, Japan
7. Disease Control and Prevention Center, National Center for Global Health and Medicine, Tokyo, Japan
8. Center for Clinical Sciences, National Center for Global Health and Medicine, Tokyo, Japan

Abstract

BNT162b2, an mRNA-based SARS-CoV-2 vaccine (Pfizer-BioNTech), is one of the most effective COVID-19 vaccines and has been approved by more than 130 countries worldwide. However, several studies have reported that the COVID-19 vaccine shows high interpersonal variability in terms of humoral and cellular responses, such as those with respect to SARS-CoV-2 spike protein immunoglobulin (Ig)G, IgA, IgM, neutralizing antibodies, and CD4⁺ & CD8⁺ T cells. The objective of this study is to investigate the kinetic changes in anti-SARS-CoV-2 spike IgG (IgG-S) profiles and adverse reactions and their associations with HLA profiles among 100 hospital workers from the Center Hospital of the National Center for Global Health and Medicine (NCGM), Tokyo, Japan. DQA1*03:03:01 (P = 0.017; Odd ratio (OR) 2.80, 95%Confidence interval (CI) 1.05–7.25) was significantly associated with higher IgG-S production after two doses of BNT162b2 while DQB1*06:01:01:01 (P = 0.028, OR 0.27, 95%CI 0.05–0.94) was significantly associated with IgG-S declines after two doses of BNT162b2. No HLA alleles were significantly associated with either local symptoms or fever. However, C*12:02:02 (P = 0.058; OR 0.42, 95%CI 0.15–1.16), B*52:01:01 (P = 0.031; OR 0.38, 95%CI 0.14–1.03), DQA1*03:02:01 (P = 0.028; OR 0.39, 95%CI 0.15–1.00) and DPB1*02:01:02 (P = 0.024; OR 0.45, 95%CI 0.21–0.97) appeared significantly associated with protection against systemic symptoms after two doses of BNT162b2 vaccination. Further studies with larger sample sizes are clearly warranted to determine HLA allele associations with the production and long-term sustainability of IgG-S after COVID-19 vaccination.

Keywords: COVID-19, SARS-CoV-2, Pfizer-BioNTech, IgG, HLA, Japanese

1 Introduction

2 The concept of messenger RNA (mRNA) vaccination stems from 1987, when Robert
3 Malone confirmed that human cells can absorb cationic liposomes containing mRNA and can
4 create proteins from those mRNA sequences (1). However, over the years, academic
5 laboratories and companies working on mRNA had come to the consensus that mRNA is too
6 prone to degradation to be used effectively as a drug or a vaccine(2). From the end of the 20th
7 century, research into mRNA vaccines mainly focused on influenza disease (3) and cancer
8 (4), with testing in animal models yielding satisfactory results. Such results inspired CureVac,
9 BioNTech, and Moderna to focus on transforming mRNA into a drug platform (5-7). In 2005,
10 the Karikó and Weissman team performed a landmark experiment in which an mRNA
11 vaccine was used to successfully suppress RNA recognition by Toll-like receptors (TLRs)
12 using modified pseudouridine, a uridine analog (8). All this previous work contributed to the
13 rapid development of coronavirus disease 2019 (COVID-19) mRNA vaccines within days of
14 the genome for the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) being
15 released online(9).

16 BNT162b2, an mRNA-based SARS-CoV-2 vaccine (Pfizer-BioNTech), is one of the
17 most effective COVID-19 vaccines (10) and has been approved by more than 130 countries
18 worldwide. However, several studies have reported that the COVID-19 vaccine shows high
19 interpersonal variability in terms of humoral and cellular responses, such as those with
20 respect to anti-SARS-CoV-2 spike immunoglobulin (Ig)G (11-17), IgA (15, 16), IgM (16),
21 neutralizing antibodies (18-20), and CD4⁺ & CD8⁺ T cells (18, 19, 21). The long-term
22 sustainability of SARS-CoV-2 neutralizing antibodies is crucial for the protection against
23 infection and hospitalization.

24 COVID-19 mRNA vaccines work by introducing the SARS-CoV-2 spike protein (S-
25 protein) mRNA sequence and subsequently the translated protein to the human immune
26 system. This first prompts an innate (non-specific) immune response by triggering immune
27 cells such as natural killer cells, macrophages, neutrophils, dendritic cells, mast cells,
28 basophils, and eosinophils. The second line of immune response is the adaptive immune
29 response, in which T and B lymphocytes are activated to generate long-term specific immune
30 responses to the SARS-CoV-2 virus.

31 Human Leukocyte Antigen (HLA) allelic variation may be associated with vaccine
32 efficacy according to trials of hepatitis B vaccine (22, 23), influenza vaccine (24) and HIV-1
33 vaccine (25). However, data on possible associations with the kinetics of post-COVID-19
34 vaccination antibodies remain scarce. Here, we designed a longitudinal study aimed at
35 investigating the association of host HLA polymorphisms with kinetic changes in IgG and
36 adverse reactions during and after BNT162b2 vaccination among hospital workers at a
37 national medical institution in Japan.

Methods

Study design and participants

The National Center for Global Health and Medicine (NCGM) is a Japanese government-designated medical center for the treatment of COVID-19 patients. We recruited and monitored a total of 100 hospital workers ≥ 20 years of age from the period between March and June 2021. Details of the study design are explained in previous studies(26, 27). All participants were vaccinated with two doses of the BNT162b2 mRNA-based SARS-CoV-2 vaccine (Pfizer-BioNTech) according to the standard protocol (two doses of 30 μ g, administered 3 weeks apart) (18, 28). All participants received the first dose of the BNT162b2 vaccine in March 2021 and the second dose was administered 21 days after the first dose. Blood was drawn from volunteers on day 1 (immediately after the first dose). Samples were then collected on day 15, day 29 (7 days after the second dose) and day 61. None of the participants had any history of COVID-19 or administration of immunosuppressive medications. The study protocol was approved by the Ethics Committee of the NCGM, Japan (approval number: NCGM-A-004175-00). Written informed consent was obtained from all participants prior to enrollment.

Self-reported questionnaire

Participants were requested to complete a questionnaire regarding the adverse reactions experienced after each dose of vaccine. Based on the results, we classified side effects into local symptoms (pain at the injection site, swelling and redness) and systemic symptoms (fever, dizziness, fatigue, headache, chills, vomiting, diarrhea, muscle pain and joint pain) with reference to the U.S. Food and Drug Administration guidance. Symptoms were further graded into 4 categories: grade 0, no adverse effect; grade 1, adverse effect without interfering with daily activities; grade 2, adverse effect with some degree of interference with daily activities; and grade 3, adverse effect with considerable interference with daily activities. Redness or swelling at the side of infection was graded based on the size of the redness or swelling as: grade 0, 0–2.0 cm; grade 1, 2.1–5.0 cm; grade 2, 5.1–10.0 cm; and grade 3, >10.0 cm.

Serological assays

Antibodies against the SARS-CoV-2 receptor-binding domain (RBD) of the S1 subunit of the spike protein (IgG-S) (AdviseDx SARS-CoV-2 IgG II; Abbott, Chicago, IL) and SARS-CoV-2 nucleocapsid (IgG-N) (Abbott ARCHITECT® SARS-CoV-2 anti-N IgG; Abbott) were measured at each collection time point. For IgG-S, results > 50 AU/mL (as the cut-off set by the manufacturer) were considered indicative of seropositivity. For surrogate measurement of neutralizing antibody, a threshold of 4160 AU/mL was applied, as this threshold corresponds to a 95% probability of obtaining a positive result from plaque

reduction neutralization test (PRNT) for SARS-CoV-2 at 1:250 dilution (14). For IgG-N, results above the index value of 1.40 (as the cut-off set by the manufacturer) were considered indicative of seropositivity.

Next-generation sequencing (NGS)-based HLA genotyping

NGS HLA genotyping was performed using AllType™ NGS Assays (One Lambda, West Hills, CA) on an Ion GeneStudio S5 sequencing system (Thermo Fisher Scientific, Waltham, MA). HLA-A, -C, -B, DRB1, DQA1, DQB1, DPA1, DPB1 targeted gene amplification, HLA library preparation, HLA template preparation, and HLA library loading onto an ion 530v1 chip in an Ion Chef library preparation robot (Thermo Fisher Scientific) and final sequencing in an Ion GeneStudio S5 sequencer (Thermo Fisher Scientific) were performed following the instructions from the vendor.

HLA genotype assignment

Demultiplexing of barcodes and base-calling was carried out using Torrent Suite version 5.8.0 (Thermo Fisher Scientific). Raw fastq reads were extracted using the FileExporter function in Torrent Suite version 5.8.0. HLA genotype assignments were undertaken using HLATypeStream Visual (TSV v2.0; One Lambda, West Hills, CA) and NGSengine® (v2.22.0.22581; GenDX, Utrecht, Netherlands). The default analysis parameters and healthy metrics threshold were applied for TSV v2.0, while we applied the “ignore regions” function in NGSengine® to eliminate known sequencing error sites in the ion S5 system. Four-field resolution HLA alleles were obtained for HLA-A, -C, -B, DRB1, DQA1, DQB1, DPA1. Novel HLA alleles were subjected to Pacbio Sequel sequencing in collaboration with the H.U. Group Research Institute (Tokyo, Japan). Pacbio subreads and consensus reads were obtained from SMRT Link software and HLA calling was performed using NGSengine® software (v2.22.0.22581; GenDX).

Statistical analysis

Correlation coefficient tests and Kruskal-Wallis tests were applied to describe the correlation between IgG-S titers with age, sex, BMI of participants, and adverse reactions after two doses of vaccination. IgG-S levels in participants plateaued on day 29 (7 days after the second dose), and were further classified into low responders (top 25th percentile of the IgG-S distribution) and high responders (bottom <25th percentile of the IgG-S distribution). Strong decliners (top 25th percentile of Δ) and weak decliners (bottom 75th percentile of Δ) were defined as Δ of IgG-S level between day 29 (7 days after the second dose) and day 61 (39 days after second vaccination).

Case-control HLA allele association tests, HLA haplotype estimations and case-control HLA haplotype association tests were prepared and analyzed using the Bridging ImmunoGenomics Data Analysis Workflow Gaps (BIGDAWG) R package (29).

Results

Basic characteristics of participants

A total of 100 hospital workers at the NCGM were recruited, comprising 32 men and 68 women. Participants were further classified into 5 age strata to observe the changes of IgG-S in relation to age: 20–29 years old; 30–39 years old; 40–49 years old; 50–59 years old; and ≥ 60 years old (Table 1). The BMI of participants ranged from 17 kg/m² to 38 kg/m², with the highest BMI observed in the group ≥ 60 -years old (mean $\pm$ standard deviation, 26 $\pm$ 6.30 kg/m²).

Kinetics of antibodies activities

IgG-S

All participants tested negative for IgG-S, IgG-N and IgM at day 1 (before BNT162b2 vaccination). IgG-S levels increased exponentially after the first dose of BNT162b2 vaccination, plateauing on day 15. After the second dose of BNT162b2 vaccination, the IgG-S level peaked on day 29 (7 days after the second vaccination). Younger age (especially under 59 years old) (Kruskal-Wallis test, $P = 0.024$) (Supplementary Figure 1a) and female sex (Kruskal-Wallis test, $P = 0.034$) (Supplementary Figure 1b) were significantly associated with higher IgG-S production after two doses of BNT162b2 vaccinations (Supplementary Figure 1a–c). For the surrogate measurement of neutralizing antibody activities, we examined for IgG-S level > 4160 AU/mL, which corresponded to a 95% probability of obtaining a PRNT ID₅₀ (estimated number of virus particles required to produce infection in 50% of normal adult humans) at 1:250 dilution(14). Fifteen days after the first dose of BNT162b2 vaccination, none of the participants had reached a sufficient level of surrogate neutralizing activities, although 96% of participants obtained sufficient level of surrogate neutralizing activities against SARS-CoV-2 by day 29 (7 days after the second dose). However, the proportion showing sufficient level of surrogate neutralizing activities decreased to 81% on day 61.

At day 61 (39 days after second vaccination), we observed a significant correlation between age ($r = 0.325$, $P = 0.0012$) (Supplementary Figure 2a) and decreased IgG-S levels, but no associations with sex ($P = 0.0621$) (Supplementary Figure 2b) or BMI ($r = 0.173$, $P = 0.1065$) (Supplementary Figure 2c) were apparent for participants. However, although it is not statistically significant, strong decliners ($\Delta \geq 20,000$ AU/mL) were observed to be mainly

concentrated among the female group (n=15, 23%), compared with the male group (n=3, 9.4%).

IgG-N

None of the participants were seropositive for IgG-N across the five survey time points, indicating no prior SARS-CoV-2 infection. No infection was observed during or after the BNT162b2 vaccination.

Associations of HLA alleles with adverse reactions and IgG-S levels

We classified the self-reported body temperature into 4 categories: T1, < 37.5°C; T2, 37.5–37.9°C; T3, 38.0–38.4°C; and T4, 38.5–38.9°C. Systemic symptoms (grade 0–1 versus grade 2–3) showed a significant association (Kruskal-Wallis test, $P = 0.026$) with higher IgG-S level on day 29 (7 days after the second dose) (Supplementary Figure 3a). More specifically, fever (> 37.5°C) showed a strong association (Kruskal-Wallis test, $P = 0.009$) with higher IgG-S level on day 29 (Supplementary Figure 3b). However, local symptoms (Kruskal-Wallis test, $P = 0.465$) were not associated with IgG-S level on day 29 (Supplementary Figure 3c).

DQA1*03:03:01 ($P = 0.017$; OR 2.80, 95%CI 1.05–7.25) was significantly associated with higher IgG-S production after two doses of BNT162b2 (Table 2). Haplotype analysis of HLA-DQA1 and surrounding HLA genes did not reveal any significant associations with HLA haplotypes (DQA1*03:03:01-DQB1*04:01:01, $P = 0.072$, OR 2.45, 95%CI 0.92–6.52 and DRB1*04:05:01-DQA1*03:03:01-DQB1*04:01:01, $P = 0.072$, OR 2.45, 95%CI 0.92–6.52) (Table 3).

IgG-S levels declined after the second dose of BNT162b2 (day 61, 39 days after the second vaccination), showing a significant correlation with age ($r=0.325$, $P=0.012$) (Supplementary Figure 2a). DQB1*06:01:01:01 ($P = 0.028$, OR 0.27, 95%CI 0.05–0.94) was significantly associated with IgG-S declines after two doses of BNT162b2 (Table 4). Haplotype analysis of DQB1*06:01:01:01 revealed significant HLA haplotypes with DQA1, DQA1*01:03:01-DQB1*06:01:01 ($P = 0.028$; OR 0.27, 95%CI 0.05–0.94). However, the HLA haplotype did not extend to HLA-DRB1 with DRB1*15:02:01-DQB1*06:01:01 ($P = 0.392$; OR 0.57, 95%CI 0.16–2.05) or DRB1*15:02-DQA1*01:03:01-DQB1*06:01:01 ($P = 0.392$; OR 0.57, 95%CI 0.16–2.05) (Table 5). Due to the limited sample size, none of the abovementioned HLA alleles remained significant associated with IgG-S levels after multiple correction, but a potential correlation between individual HLA alleles and responsiveness to BNT162b2 vaccination was still suggested.

No HLA alleles were significantly associated with either local symptoms (Supplementary table 1) or fever (Supplementary table 2). However, C*12:02:02 ($P = 0.058$; OR 0.42, 95%CI 0.15–1.16), B*52:01:01 ($P = 0.031$; OR 0.38, 95%CI 0.14–1.03),

DQA1*03:02:01 ($P = 0.028$; OR 0.39, 95%CI 0.15–1.00) and DPB1*02:01:02 ($P = 0.024$; OR 0.45, 95%CI 0.21–0.97) appeared significantly associated with protection against systemic symptoms after two doses of BNT162b2 vaccination (Supplementary table 3).

Discussion

We have described in this study the kinetic changes in IgG-S profiles and adverse reactions and their associations with HLA profiles among hospital workers from a national medical institution in Japan. BNT162b2 was designed to present the entire spike glycoprotein of SARS-CoV-2 as a target to elicit neutralizing antibodies and so block viral entry through the angiotensin-converting enzyme 2 (ACE2) cellular receptor, particularly the viral sequences recognizing the receptor-binding domain (28). A structural study of SARS-CoV-2 proteins identified the S-protein as a major inducer of protective immunity (30). Assessment of changes in the kinetics of antibodies, adverse reactions after BNT162b2 vaccination, and possible associations between interpersonal differences and variations in HLA genes are thus important.

All participants in the present study displayed seroconversion on day 29 (7 days after the second dose) and displayed an increase of IgG-S against SARS-CoV-2. The youngest age group and female participants produced higher levels of IgG-S after two doses of BNT162b2 vaccinations, consistent with findings from previous studies (13, 31, 32). However, the IgG-S antibody response appears both time- and age-dependent (Figure 2) (11, 13, 16, 33). In this study, the female group produced significantly more IgG-S at the early stage of our longitudinal study, but we also observed a strong IgG-S decline ($\Delta 20,000$ AU/mL) in the female group. DQA1*03:03:01 ($P = 0.017$; OR 2.80, 95%CI 1.05–7.25) was significantly associated with strong responders for IgG-S. Interestingly, DQA1*03:03:01 is the second most common DQA1 allele in the Japanese population, with a frequency of 16.50% (34).

The long-term sustainability of IgG-S after COVID-19 vaccination is central to the protection afforded against SARS-CoV-2. This study did not identify any HLA alleles associated with strong declines in IgG-S levels. Instead, we identified an association between DQB1*06:01:01:01 ($P = 0.028$; OR 0.27, 95%CI 0.05–0.94) and protection against IgG-S declines.

The severity of systemic adverse symptoms after two doses of BNT162b2 vaccination was associated with the level of IgG-S production, consistent with previous research (35). We observed that carriers of C*12:02:02, B*52:01:01, DQA1*03:02:01 and DPB1*02:01:02 showed significant protection against systemic symptoms.

Data regarding long-term antibody kinetics in vaccinated subjects remain scarce. In a population of 33 healthy adults having received the Moderna mRNA-1273 vaccine and followed-up for 209 days, the estimated half-life of the antibody response was 52 days (95%CI 46–58 days) using an exponential decay model [13]. In a cohort of 188 unvaccinated

COVID-19 patients (mostly not hospitalized: 174/188) who were followed-up for as long as 8 months, the antibody half-life was 83 days (95%CI 62–126 days) [14]. In this study, we estimated that IgG-S levels would drop below 4,160 AU/mL after 101 days, suggesting a need for booster shots to sustain adequate levels of neutralizing antibody.

Research into the associations between HLA alleles and IgG-S antibody levels after vaccination remain limited, but a small Italian cohort (n=56) found no associations between HLA alleles and IgG-S levels after BNT162b2 vaccination. In the present study, we identified several HLA alleles showing potential associations with the kinetics of IgG-S antibody levels. Further studies with larger sample sizes are clearly warranted to determine HLA allele associations with the production and long-term sustainability of IgG-S antibody levels after COVID-19 vaccination.

Acknowledgments

We are grateful to the members of the working group for this study (Yoshimi Shigemori, Ayumi Nakayama, Yusuke Oshiro, Natsumi Inamura, Haruka Osawa, Maki Konishi, Azusa Kamikawa, and Yumiko Kito) for their support. We thank Dr. Ikue Ito and Dr. Tetsuya Sato from H.U. Group Research Institute for their technical support for Pacbio Sequel sequencing.

Role of the Funder/Sponsor

The funders did not play any role in the design and conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Conflict of Interests

All authors (except Gohzoh Ueda) declare no conflict of interest. Gohzoh Ueda is one of employees of Abbott Japan, which provided the antibody assay reagents and funding for the present study. The Role of the funder/sponsor is described above.

Figures legends

Supplementary Figure 1 SARS-CoV-2 spike-specific IgG production on day 29, stratified by age and sex

- a) SARS-CoV-2 spike-specific IgG production on day 29, stratified by 5 age groups: 20–29 years old (y.o), 30–39 y.o., 40–49 y.o. 50–59 y.o, and over 60 y.o
- b) SARS-CoV-2 spike-specific IgG production on day 29, stratified by sex
- c) SARS-CoV-2 spike-specific IgG production on day 29, stratified by 10 groups including the categories listed in (a) and (b)
- Violin plots showing median and 95% confidence interval (CI)

Supplementary Figure 2 SARS-CoV-2 spike-specific IgG decline on day 61, stratified by age, sex, and BMI of participants

- a) Scatter plot showing the correlation between age and the change in IgG-S between day 29 and day 61. Correlation coefficients and P-values are calculated. Color intensity of the heatmap indicates the accumulation of data points in a specific area.
- b) Violin plot showing the correlation between sex and the change in IgG-S between day 29 and day 61. Violin plots showing median and 95% confidence interval (CI)
- c) Scatter plot showing the correlation between the BMI of participants and the change in IgG-S between day 29 and day 61. Correlation coefficients and p-values are calculated. Color intensity of the heatmap indicates the accumulation of data points in a specific area

Supplementary Figure 3 Association between SARS-CoV-2 spike-specific IgG production on day 29 and adverse reactions after two doses of BNT162b2 vaccination

- a) Violin plot showing the distribution of IgG-S production on day 29 stratified by grades of systemic symptoms experienced by participants after two doses of BNT162b2 vaccination
- b) Violin plot showing the distribution of IgG-S production on day 29 stratified by fever experienced by participants after two doses of BNT162b2 vaccination. T1: < 37.5°C; T2: 37.5–37.9°C; T3: 38.0–38.4°C; and T4: 38.5–38.9°C.
- c) Violin plot showing the distribution of IgG-S production on day 29 stratified by the grade of local symptoms experienced by participants after two doses of BNT162b2 vaccination
- Violin plots showing median and 95% confidence interval (CI)

References

1. Malone RW, Felgner PL, Verma IM. Cationic liposome-mediated RNA transfection. *Proc Natl Acad Sci U S A*. 1989;86(16):6077-81.
2. Pardi N, Hogan MJ, Porter FW, Weissman D. mRNA vaccines - a new era in vaccinology. *Nat Rev Drug Discov*. 2018;17(4):261-79.
3. Conry RM, LoBuglio AF, Wright M, Sumerel L, Pike MJ, Johanning F, et al. Characterization of a messenger RNA polynucleotide vaccine vector. *Cancer Res*. 1995;55(7):1397-400.
4. Boczkowski D, Nair SK, Snyder D, Gilboa E. Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. *J Exp Med*. 1996;184(2):465-72.
5. Probst J, Weide B, Scheel B, Pichler BJ, Hoerr I, Rammensee HG, et al. Spontaneous cellular uptake of exogenous messenger RNA in vivo is nucleic acid-specific, saturable and ion dependent. *Gene Ther*. 2007;14(15):1175-80.
6. Kariko K, Kuo A, Barnathan ES, Langer DJ. Phosphate-enhanced transfection of cationic lipid-complexed mRNA and plasmid DNA. *Biochim Biophys Acta*. 1998;1369(2):320-34.
7. Kariko K, Kuo A, Barnathan E. Overexpression of urokinase receptor in mammalian cells following administration of the in vitro transcribed encoding mRNA. *Gene Ther*. 1999;6(6):1092-100.
8. Kariko K, Buckstein M, Ni H, Weissman D. Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA. *Immunity*. 2005;23(2):165-75.
9. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus associated with human respiratory disease in China. *Nature*. 2020;579(7798):265-9.
10. Dagan N, Barda N, Kepten E, Miron O, Perchik S, Katz MA, et al. BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. *N Engl J Med*. 2021;384(15):1412-23.
11. Favresse J, Bayart JL, Mullier F, Elsen M, Eucher C, Van Eeckhoudt S, et al. Antibody titres decline 3-month post-vaccination with BNT162b2. *Emerg Microbes Infect*. 2021;10(1):1495-8.
12. Zaffina S, Alteri C, Ruggiero A, Cotugno N, Vinci MR, Camisa V, et al. Induction of immune response after SARS-CoV-2 mRNA BNT162b2 vaccination in healthcare workers. *J Virus Erad*. 2021;7(2):100046.
13. Kageyama T, Ikeda K, Tanaka S, Taniguchi T, Igari H, Onouchi Y, et al. Antibody responses to BNT162b2 mRNA COVID-19 vaccine in 2,015 healthcare workers in a single tertiary referral hospital in Japan. *medRxiv*. 2021:2021.06.01.21258188.
14. Ebinger JE, Fert-Bober J, Printsev I, Wu M, Sun N, Prostko JC, et al. Antibody responses to the BNT162b2 mRNA vaccine in individuals previously infected with SARS-CoV-2. *Nat Med*. 2021;27(6):981-4.
15. Wisniewski AV, Campillo Luna J, Redlich CA. Human IgG and IgA responses to COVID-19 mRNA vaccines. *PLoS One*. 2021;16(6):e0249499.
16. Pratesi F, Caruso T, Testa D, Tarpanelli T, Gentili A, Gioe D, et al. BNT162b2 mRNA SARS-CoV-2 Vaccine Elicits High Avidity and Neutralizing Antibodies in Healthcare Workers.

Vaccines (Basel). 2021;9(6).

17. Ragone C, Meola S, Fiorillo PC, Penta R, Auriemma L, Tornesello ML, et al. HLA Does Not Impact on Short-Medium-Term Antibody Response to Preventive Anti-SARS-Cov-2 Vaccine. *Front Immunol*. 2021;12:734689.
18. Sahin U, Muik A, Vogler I, Derhovanessian E, Kranz LM, Vormehr M, et al. BNT162b2 vaccine induces neutralizing antibodies and poly-specific T cells in humans. *Nature*. 2021.
19. Kalimuddin S, Tham CYL, Qui M, de Alwis R, Sim JXY, Lim JME, et al. Early T cell and binding antibody responses are associated with COVID-19 RNA vaccine efficacy onset. *Med (N Y)*. 2021;2(6):682-8 e4.
20. Maeda K, Amano M, Uemura Y, Tsuchiya K, Matsushima T, Noda K, et al. Correlates of Neutralizing/SARS-CoV-2-S1-binding Antibody Response with Adverse Effects and Immune Kinetics in BNT162b2-Vaccinated Individuals. *medRxiv*. 2021:2021.07.27.21261237.
21. Martínez-Gallo M, Esperalba-Esquerria J, Pujol-Borrell R, Sandá V, Arrese-Muñoz I, Fernández Naval C, et al. T-cell responses as a correlate of COVID-19 vaccination. A pilot study in Health Care Workers. *medRxiv*. 2021:2021.03.31.21254472.
22. Nishida N, Sugiyama M, Sawai H, Nishina S, Sakai A, Ohashi J, et al. Key HLA-DRB1-DQB1 haplotypes and role of the BTNL2 gene for response to a hepatitis B vaccine. *Hepatology*. 2018;68(3):848-58.
23. Li ZK, Nie JJ, Li J, Zhuang H. The effect of HLA on immunological response to hepatitis B vaccine in healthy people: a meta-analysis. *Vaccine*. 2013;31(40):4355-61.
24. Moss AJ, Gaughran FP, Karasu A, Gilbert AS, Mann AJ, Gelder CM, et al. Correlation between human leukocyte antigen class II alleles and HAI titers detected post-influenza vaccination. *PLoS One*. 2013;8(8):e71376.
25. Gartland AJ, Li S, McNeven J, Tomaras GD, Gottardo R, Janes H, et al. Analysis of HLA A*02 association with vaccine efficacy in the RV144 HIV-1 vaccine trial. *J Virol*. 2014;88(15):8242-55.
26. Takeuchi JS, Fukunaga A, Yamamoto S, Tanaka A, Matsuda K, Kimura M, et al. SARS-CoV-2 specific T cell and humoral immune responses upon vaccination with BNT162b2. *medRxiv*. 2021:2021.11.06.21265632.
27. Yamamoto S, Fukunaga A, Tanaka A, Takeuchi JS, Inoue Y, Kimura M, et al. Association between reactogenicity and SARS-CoV-2 antibodies after the second dose of the BNT162b2 COVID-19 vaccine. *medRxiv*. 2021:2021.07.19.21260744.
28. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020;383(27):2603-15.
29. Pappas DJ, Marin W, Hollenbach JA, Mack SJ. Bridging ImmunoGenomic Data Analysis Workflow Gaps (BIGDAWG): An integrated case-control analysis pipeline. *Hum Immunol*. 2016;77(3):283-7.
30. Buchholz UJ, Bukreyev A, Yang L, Lamirande EW, Murphy BR, Subbarao K, et al. Contributions of the structural proteins of severe acute respiratory syndrome coronavirus to protective

immunity. *Proc Natl Acad Sci U S A*. 2004;101(26):9804-9.

31. Muller L, Andree M, Moskorz W, Drexler I, Walotka L, Grothmann R, et al. Age-dependent immune response to the Biontech/Pfizer BNT162b2 COVID-19 vaccination. *Clin Infect Dis*. 2021.
32. Terpos E, Trougakos IP, Apostolou F, Charitaki I, Skirou AD, Mavrianou N, et al. Age-dependent and gender-dependent antibody responses against SARS-CoV-2 in health workers and octogenarians after vaccination with the BNT162b2 mRNA vaccine. *Am J Hematol*. 2021;96(7):E257-E9.
33. Salvagno GL, Henry BM, di Piazza G, Pighi L, De Nitto S, Bragantini D, et al. Anti-SARS-CoV-2 Receptor-Binding Domain Total Antibodies Response in Seropositive and Seronegative Healthcare Workers Undergoing COVID-19 mRNA BNT162b2 Vaccination. *Diagnostics (Basel)*. 2021;11(5).
34. Ikeda N, Kojima H, Nishikawa M, Hayashi K, Futagami T, Tsujino T, et al. Determination of HLA-A, -C, -B, -DRB1 allele and haplotype frequency in Japanese population based on family study. *Tissue Antigens*. 2015;85(4):252-9.
35. Otani J, Ohta R, Sano C. Association between Immunoglobulin G Levels and Adverse Effects Following Vaccination with the BNT162b2 Vaccine among Japanese Healthcare Workers. *Vaccines (Basel)*. 2021;9(10).

Table 1. Basic characteristics of study participants					
			BNT162b2		
Characteristics	20-29 y.o	30-39 y.o	40-49 y.o	50-59 y.o	Over 60 y.o
Sample number	17	25	27	23	8
Male sex — n (%)	2 (11.8)	12 (48.0)	7 (25.9)	7 (30.4)	4 (50.0)
Age at vaccination — years					
Mean	26 ± 2.00	35 ± 3.00	44 ± 3.00	54 ± 3.00	64 ± 4.17
Median	26 (22-29)	35 (30-39)	45 (40-49)	53 (50-59)	63 (60-73)
BMI at vaccination					
Mean	21 ± 2.00	23 ± 4.00	22 ± 3.00	21 ± 2.00	26 ± 6.30
Median	21 (19-26)	22 (17-29)	22 (17-26)	22 (17-27)	24 (19-38)
Missing data	2	3	2	1	1
Baseline SARS-CoV-2 status — n (%)†					
Positive	0	0	0	0	0
Negative	17	25	27	23	8
Missing	0	0	0	0	0

† Positive severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) status is defined as positive IgG against N-protein at the time of 1st vaccination

Table 2 | Associational analysis of HLA-A, -C, -B, -DRB1, -DQB1, -DOA1 and -DPB1 alleles in 73 IgG-S low responders and 24 IgG-S high responders. Significant HLA alleles are highlighted. Abbreviations: OR, odds ratio; 95%CI, 95% confidence interval; binned, rare HLA alleles with expected count < 5 are combined into a common class.

Locus	Two-field Allele	IgG-S low responders (2n=146) %	IgG-S high responders (2n=48) %	OR (95%CI)	P-value	Three-field allele	IgG-S low responders (2n=146) %	IgG-S high responders (2n=48) %	OR (95%CI)	P-value	Four-field allele	IgG-S low responders (2n=146) %	IgG-S high responders (2n=48) %	OR (95%CI)	P-value
A	02:06	12.3	14.6	1.21 (0.40-3.32)	0.686	02:06:01	12.3	14.6	1.21 (0.40-3.32)	0.686					
A	11:01	11.0	12.5	1.16 (0.35-3.38)	0.770	11:01:01	11.0	12.5	1.16 (0.35-3.38)	0.770	11:01:01:01	11.0	12.5	1.16 (0.35-3.38)	0.770
A	24:02	34.2	33.3	0.96 (0.45-2.01)	0.908	24:02:01	34.2	33.3	0.96 (0.45-2.01)	0.908	24:02:01:01	32.2	31.3	0.96 (0.44-2.02)	0.903
A	binned	42.5	39.6	0.89 (0.43-1.81)	0.725	binned	42.5	39.6	0.89 (0.43-1.81)	0.725	binned	56.8	56.3	0.98 (0.48-2.00)	0.942
C	01:02	15.1	22.9	1.68 (0.67-4.00)	0.209	01:02:01	15.1	22.9	1.68 (0.67-4.00)	0.209					
C	03:03	10.3	14.6	1.49 (0.48-4.21)	0.414	03:03:01	10.3	14.6	1.49 (0.48-4.21)	0.414	03:03:01:01	10.3	14.6	1.49 (0.48-4.21)	0.414
C	07:02	10.3	14.6	1.49 (0.48-4.21)	0.414	07:02:01	10.3	14.6	1.49 (0.48-4.21)	0.414					
C	12:02	15.1	8.3	0.51 (0.12-1.63)	0.235	12:02:02	14.4	8.3	0.54 (0.13-1.73)	0.278	12:02:02:01	14.4	8.3	0.54 (0.13-1.73)	0.278
C	binned	49.3	39.6	0.67 (0.33-1.37)	0.241	binned	50.0	39.6	0.66 (0.32-1.33)	0.210	binned	75.3	77.1	1.10 (0.49-2.65)	0.807
B	52:01	15.1	8.3	0.51 (0.12-1.63)	0.235	52:01:01	15.1	8.3	0.51 (0.12-1.63)	0.235	52:01:01:02	15.1	8.3	0.51 (0.12-1.63)	0.235
B	binned	84.9	91.7	1.95 (0.61-8.20)	0.235	binned	84.9	91.7	1.95 (0.61-8.20)	0.235	binned	84.9	91.7	1.95 (0.61-8.20)	0.235
DRB1	09:01	15.8	6.3	0.36 (0.07-1.27)	0.094	09:01:02	15.8	6.3	0.36 (0.07-1.27)	0.094	09:01:02:01	15.8	6.3	0.36 (0.07-1.27)	0.094
DRB1	15:02	11.6	8.3	0.69 (0.16-2.28)	0.522	15:02:01	11.6	8.3	0.69 (0.16-2.28)	0.522					
DRB1	binned	72.6	85.4	2.21 (0.88-6.30)	0.072	binned	72.6	85.4	2.21 (0.88-6.30)	0.072	binned	84.2	93.8	2.80 (0.79-15.23)	0.094
DQA1	01:02	13.0	16.7	1.34 (0.47-3.50)	0.526	01:02:01	12.3	16.7	1.42 (0.50-3.75)	0.444					
DQA1	01:03	17.8	20.8	1.21 (0.48-2.90)	0.640	01:03:01	17.8	20.8	1.21 (0.48-2.90)	0.640	01:03:01:01	11.0	10.4	0.94 (0.26-2.91)	0.916
DQA1	03:01	12.3	10.4	0.83 (0.23-2.50)	0.722	03:01:01	12.3	10.4	0.83 (0.23-2.50)	0.722	03:01:01:01	12.3	10.4	0.83 (0.23-2.50)	0.722
DQA1	03:02	17.8	6.3	0.31 (0.06-1.09)	0.051	03:02:01	17.8	6.3	0.31 (0.06-1.09)	0.051	03:02:01:01	17.1	6.3	0.32 (0.06-1.14)	0.063
DQA1	03:03	9.6	22.9	2.80 (1.05-7.25)	0.017	03:03:01	9.6	22.9	2.80 (1.05-7.25)	0.017					
DQA1	binned	29.5	22.9	0.71 (0.30-1.59)	0.381	binned	30.1	22.9	0.69 (0.29-1.54)	0.336	binned	59.6	72.9	1.83 (0.85-4.08)	0.097
DQB1	03:01	12.3	8.3	0.65 (0.15-2.12)	0.449	03:01:01	12.3	8.3	0.65 (0.15-2.12)	0.449					
DQB1	03:02	12.3	10.4	0.83 (0.23-2.50)	0.722	03:02:01	12.3	10.4	0.83 (0.23-2.50)	0.722	03:02:01:01	12.3	10.4	0.83 (0.23-2.50)	0.722
DQB1	03:03	16.4	6.3	0.34 (0.06-1.20)	0.077	03:03:02	16.4	6.3	0.34 (0.06-1.20)	0.077	03:03:02:02	16.4	6.3	0.34 (0.06-1.20)	0.077
DQB1	06:01	17.1	20.8	1.27 (0.50-3.05)	0.562	06:01:01	17.1	20.8	1.27 (0.50-3.05)	0.562	06:01:01:01	17.1	20.8	1.27 (0.50-3.05)	0.562
DQB1	binned	41.8	54.2	1.65 (0.81-3.35)	0.134	binned	41.8	54.2	1.65 (0.81-3.35)	0.134	binned	54.1	62.5	1.41 (0.69-2.94)	0.309
DPA1	01:03	43.2	43.8	1.02 (0.50-2.08)	0.942	01:03:01	43.2	43.8	1.02 (0.50-2.08)	0.942	01:03:01:01	28.1	29.2	1.05 (0.47-2.27)	0.885
											01:03:01:05	11.6	8.3	0.69 (0.16-2.28)	0.522
DPA1	02:01	16.4	14.6	0.87 (0.29-2.28)	0.761	02:01:01	16.4	14.6	0.87 (0.29-2.28)	0.761	02:01:01:02	13.0	10.4	0.78 (0.21-2.33)	0.635
DPA1	02:02	40.4	41.7	1.05 (0.51-2.14)	0.878	02:02:02	40.4	41.7	1.05 (0.51-2.14)	0.878	02:02:02:01	40.4	41.7	1.05 (0.51-2.14)	0.878
											binned	6.8	10.4	1.58 (0.40-5.41)	0.422
DPB1	02:01	25.3	22.9	0.88 (0.37-1.98)	0.735	02:01:02	25.3	22.9	0.88 (0.37-1.98)	0.735	02:01:02:01	19.9	18.8	0.93 (0.36-2.25)	0.866
DPB1	04:02	11.6	8.3	0.69 (0.16-2.28)	0.522	04:02:01	11.6	8.3	0.69 (0.16-2.28)	0.522					
DPB1	05:01	34.2	39.6	1.26 (0.60-2.58)	0.503	05:01:01	34.2	39.6	1.26 (0.60-2.58)	0.503	05:01:01:01	27.4	35.4	1.45 (0.68-3.05)	0.290
DPB1	binned	28.8	29.2	1.02 (0.46-2.19)	0.958	binned	28.8	29.2	1.02 (0.46-2.19)	0.958	binned	52.7	45.8	0.76 (0.37-1.54)	0.406

Table 3 | HLA haplotype analysis for HLA-DRB1, -DQA1, and -DQB1 in IgG-S low responders versus high responders. Abbreviations: OR, odds ratio; 95%CI, 95% confidence Interval; binned, rare HLA alleles with expected count < 5 are combined into a common class.

Locus	Allele	IgG-S low responders (2n=146) %	IgG-S high responders (2n=48) %	OR (95%CI)	P-value
DRB1~DQA1~DQB1	09:01:02~03:02:01~03:03:02	14.4	6.3	0.40 (0.07-1.43)	0.138
DRB1~DQA1~DQB1	15:02:01~01:03:01~06:01:01	11.0	8.3	0.74 (0.23-2.33)	0.605
DRB1~DQA1~DQB1	15:01:01~01:02:01~06:02:01	8.2	10.4	1.30 (0.43-3.89)	0.641
DRB1~DQA1~DQB1	04:05:01~03:03:01~04:01:01	7.5	16.7	2.45 (0.92-6.52)	0.072
DRB1~DQA1~DQB1	08:03:02~01:03:01~06:01:01	6.2	10.4	1.77 (0.56-5.57)	0.329
DRB1~DQA1~DQB1	binned	52.7	47.9	0.82 (0.43-1.58)	0.562
DQA1~DQB1	01:03:01~06:01:01	17.1	20.8	1.27 (0.50-3.05)	0.562
DQA1~DQB1	03:01:01~03:02:01	12.3	10.4	0.83 (0.23-2.50)	0.722
DQA1~DQB1	03:02:01~03:03:02	16.4	6.3	0.34 (0.06-1.20)	0.077
DQA1~DQB1	01:02:01~06:02:01	8.2	10.4	1.30 (0.43-3.89)	0.641
DQA1~DQB1	03:03:01~04:01:01	7.5	16.7	2.45 (0.92-6.52)	0.072
DQA1~DQB1	binned	38.4	35.4	0.88 (0.45-1.74)	0.715

Table 4 | Associational analysis of HLA-A, -C, -B, -DRB1, -DQB1, -DOA1 and -DPB1 alleles in 75 IgG-S weak decliner and 22 IgG-S strong decliner. Abbreviations: OR, odds ratio; 95%CI, 95% confidence interval; binned, rare HLA alleles with expected count < 5 are combined into a common class.

Locus	Two-field allele	IgG-S Weak decliner (2n=150) %	IgG-S Strong decliner (2n=44) %	OR (95%CI)	P-value	Three-field allele	IgG-S Weak decliner (2n=150) %	IgG-S Strong decliner (2n=44) %	OR (95%CI)	P-value	Four-field allele	IgG-S weak decliner (2n=150) %	IgG-S strong decliner (2n=44) %	OR (95%CI)	P-value
A	02:06	12.7	13.6	1.09 (0.33-3.10)	0.866	02:06:01	12.7	13.6	1.09 (0.33-3.10)	0.866					
A	24:02	37.3	22.7	0.49 (0.20-1.12)	0.072	24:02:01	37.3	22.7	0.49 (0.20-1.12)	0.072	24:02:01:01	35.3	20.5	0.47 (0.19-1.10)	0.063
A	binned	50.0	63.6	1.75 (0.83-3.76)	0.111	binned	50.0	63.6	1.75 (0.83-3.76)	0.111	binned	64.7	79.5	2.12 (0.91-5.40)	0.063
C	01:02	18.7	11.4	0.56 (0.16-1.61)	0.257	01:02:01	18.7	11.4	0.56 (0.16-1.61)	0.257					
C	12:02	13.3	13.6	1.03 (0.31-2.90)	0.959	12:02:02	12.7	13.6	1.09 (0.33-3.10)	0.866	12:02:02:01	12.7	13.6	1.09 (0.33-3.10)	0.866
C	binned	68.0	75.0	1.41 (0.63-3.36)	0.375	binned	68.7	75.0	1.37 (0.61-3.27)	0.420	binned	87.3	86.4	0.92 (0.32-3.01)	0.866
B	52:01	13.3	13.6	1.03 (0.31-2.90)	0.959	52:01:01	13.3	13.6	1.03 (0.31-2.90)	0.959	52:01:01:02	13.3	13.6	1.03 (0.31-2.90)	0.959
B	binned	86.7	86.4	0.97 (0.34-3.18)	0.959	binned	86.7	86.4	0.97 (0.34-3.18)	0.959	binned	86.7	86.4	0.97 (0.34-3.18)	0.959
DRB1	09:01	13.3	13.6	1.03 (0.31-2.90)	0.959	09:01:02	13.3	13.6	1.03 (0.31-2.90)	0.959	09:01:02:01	13.3	13.6	1.03 (0.31-2.90)	0.959
DRB1	binned	86.7	86.4	0.97 (0.34-3.18)	0.959	binned	86.7	86.4	0.97 (0.34-3.18)	0.959	binned	86.7	86.4	0.97 (0.34-3.18)	0.959
DQA1	01:02	13.3	15.9	1.23 (0.41-3.33)	0.664	01:02:01	12.7	15.9	1.30 (0.43-3.55)	0.579					
DQA1	01:03	21.3	9.1	0.37 (0.09-1.14)	0.066	01:03:01	21.3	9.1	0.37 (0.09-1.14)	0.066					
DQA1	03:01	11.3	13.6	1.24 (0.37-3.58)	0.678	03:01:01	11.3	13.6	1.24 (0.37-3.58)	0.678	03:01:01:01	11.3	13.6	1.24 (0.37-3.58)	0.678
DQA1	03:02	14.7	15.9	1.10 (0.37-2.94)	0.839	03:02:01	14.7	15.9	1.10 (0.37-2.94)	0.839	03:02:01:01	14.7	13.6	0.92 (0.28-2.56)	0.864
DQA1	03:03	14.7	6.8	0.43 (0.08-1.53)	0.172	03:03:01	14.7	6.8	0.43 (0.08-1.53)	0.172					
DQA1	binned	24.7	38.6	1.92 (0.88-4.13)	0.069	binned	25.3	38.6	1.86 (0.85-3.97)	0.085	binned	74.0	72.7	0.94 (0.42-2.20)	0.866
DQB1	03:02	11.3	13.6	1.24 (0.37-3.58)	0.678	03:02:01	11.3	13.6	1.24 (0.37-3.58)	0.678	03:02:01:01	11.3	13.6	1.24 (0.37-3.58)	0.678
DQB1	03:03	13.3	15.9	1.23 (0.41-3.33)	0.664	03:03:02	13.3	15.9	1.23 (0.41-3.33)	0.664	03:03:02:02	13.3	15.9	1.23 (0.41-3.33)	0.664
DQB1	06:01	21.3	6.8	0.27 (0.05-0.94)	0.028	06:01:01	21.3	6.8	0.27 (0.05-0.94)	0.028	06:01:01:01	21.3	6.8	0.27 (0.05-0.94)	0.028
DQB1	binned	54.0	63.6	1.49 (0.71-3.20)	0.257	binned	54.0	63.6	1.49 (0.71-3.20)	0.257	binned	54.0	63.6	1.49 (0.71-3.20)	0.257
DPA1	01:03	41.3	50.0	1.42 (0.68-2.95)	0.308	01:03:01	41.3	50.0	1.42 (0.68-2.95)	0.308	01:03:01:01	27.3	31.8	1.24 (0.55-2.70)	0.562
DPA1	02:01	18.0	9.1	0.46 (0.11-1.43)	0.156	02:01:01	18.0	9.1	0.46 (0.11-1.43)	0.156	02:01:01:02	14.0	6.8	0.45 (0.08-1.63)	0.203
DPA1	02:02	40.7	40.9	1.01 (0.48-2.11)	0.977	02:02:02	40.7	40.9	1.01 (0.48-2.11)	0.977	02:02:02:01	40.7	40.9	1.01 (0.48-2.11)	0.977
DPA1											binned	18.0	20.5	1.17 (0.44-2.87)	0.713
DPB1	02:01	23.3	29.5	1.38 (0.59-3.07)	0.401	02:01:02	23.3	29.5	1.38 (0.59-3.07)	0.401	02:01:02:01	18.7	22.7	1.28 (0.50-3.05)	0.551
DPB1	05:01	36.0	34.1	0.92 (0.42-1.96)	0.816	05:01:01	36.0	34.1	0.92 (0.42-1.96)	0.816	05:01:01:01	29.3	29.5	1.01 (0.44-2.21)	0.978
DPB1	binned	40.7	36.4	0.83 (0.39-1.75)	0.608	binned	40.7	36.4	0.83 (0.39-1.75)	0.608	binned	52.0	47.7	0.84 (0.41-1.75)	0.618

Table 5 | HLA haplotype analysis for HLA-DRB1, -DQA1, and -DQB1 in 75 IgG-S weak decliners and 22 IgG-S strong decliners.
Abbreviations: OR, odds ratio; 95%CI, 95% confidence interval; binned, rare HLA alleles with expected count < 5 are combined into a common class.

Haplotype	DRB1~DQB1	IgG-S Weak decliner (2n=150) %	IgG-S Strong decliner (2n=44) %	OR (95%CI)	P-value
DRB1~DQB1	04:05:01~04:01:01	10.7	6.8	0.61 (0.17-2.20)	0.454
DRB1~DQB1	09:01:02~03:03:02	12.0	13.6	1.16 (0.35-3.32)	0.772
DRB1~DQB1	15:01:01~06:02:01	8.0	11.4	1.47 (0.49-4.44)	0.490
DRB1~DQB1	15:02:01~06:01:01	11.3	6.8	0.57 (0.16-2.05)	0.392
DRB1~DQB1	binned	58.0	61.4	1.15 (0.58-2.29)	0.690
DQA1~DQB1	01:03:01~06:01:01	21.3	6.8	0.27 (0.05-0.94)	0.028
DQA1~DQB1	03:01:01~03:02:01	11.3	13.6	1.24 (0.37-3.58)	0.678
DQA1~DQB1	03:02:01~03:03:02	13.3	15.9	1.23 (0.41-3.33)	0.664
DQA1~DQB1	03:03:01~04:01:01	10.7	6.8	0.61 (0.17-2.21)	0.454
DQA1~DQB1	01:02:01~06:02:01	8.0	11.4	1.47 (0.49-4.44)	0.490
DQA1~DQB1	binned	54.0	63.6	1.49 (0.75-2.98)	0.259
DRB1~DQA1~DQB1	04:05:01~03:03:01~04:01:01	10.7	6.8	0.61 (0.17-2.21)	0.454
DRB1~DQA1~DQB1	09:01:02~03:02:01~03:03:02	12.0	13.6	1.16 (0.35-3.32)	0.772
DRB1~DQA1~DQB1	15:01:01~01:02:01~06:02:01	8.0	11.4	1.47 (0.49-4.44)	0.490
DRB1~DQA1~DQB1	15:02:01~01:03:01~06:01:01	11.3	6.8	0.57 (0.16-2.05)	0.392
DRB1~DQA1~DQB1	binned	58.0	61.4	1.15 (0.58-2.29)	0.690