The urothelial transcriptomic response to interferon gamma predicts T1 recurrence-free and basal/squamous muscle-invasive bladder cancer survival and better targeted strategies for immune checkpoint blocking.

Simon C. Baker^{1,*}, Andrew S. Mason¹, Raphael G. Slip¹, Pontus Eriksson², Gottfrid Sjö Dahl², Ludwik K Trejdosiewicz¹ and Jennifer Southgate¹.

Affiliation

¹Jack Birch Unit of Molecular Carcinogenesis, Department of Biology and York Biomedical Research Institute, University of York, Heslington, York YO10 5DD, UK.

²Lund University Cancer Centre, Lunds Universitet, Box 117, 221 00 Lund, Sweden.

Contact Information for Corresponding Author* (Address as above)

Tel: +44 1904 328706

E-mail address: simon.baker@york.ac.uk

Keywords: urothelium, IFNg, non muscle invasive bladder cancer, APOBEC

Abstract

Intravesical Bacillus Calmette-Guérin vaccine (BCG) is an established immunotherapeutic in bladder cancer (BlCa), provoking inflammation leading to tumour-specific immunity. Immune checkpoint blockers such as anti-PD-L1 have potential for enhancing tumour-specific lymphocyte-mediated cytotoxicity in BCG-refractive or advanced disease. In both cases, Interferon-gamma (IFN γ) plays a central role. We investigated the transcriptomic response of normal human urothelium to IFN γ to disentangle mechanisms of BCG and anti-PD-L1 therapy failure.

Exposure of differentiated human urothelium to IFN γ resulted in upregulated MHC class I and class II and *de novo* expression of CXCL9-11 chemokine genes. Normal urothelium expressed only immuno-inhibitory B7 family members: PD-L1 expression was induced by IFN γ , whereas VISTA was expressed constitutively.

A urothelial IFN γ response gene set was derived and used for unsupervised clustering of tumours, which predicted longer recurrence-free survival in non-muscle invasive bladder cancer (NMIBC). In muscle invasive bladder cancer (MIBC), the IFN γ -signature split the basal/squamous consensus subtype, with significantly worse overall survival when weak/absent.

Normal urothelium has few resident lymphocytes. Tumour cell killing requires recruitment and activation of IFN γ -secreting pro-inflammatory/cytotoxic lymphocytes while surmounting both innate (VISTA) and upregulated (PD-L1) inhibitory mechanisms. This study offers supportive evidence for strategies to enhance immunotherapy *via* the IFN γ and VISTA/PD-L1 nexus.

39 **Patient Summary**

40 Immunotherapy brings promise of harnessing a patient's own immune system to seek and
41 destroy malignant cells, but it has yet to deliver widespread clinical benefit. We exposed
42 human urothelium to interferon gamma, a key messenger of the immune system and
43 identified a novel signature of 33 genes that predicted cancers with better outcomes. Our
44 study revealed alternative strategies for targeting checkpoint proteins to improve
45 immunotherapy in the future.

Manuscript

BCG is widely used in NMIBC therapy as a potent inducer of inflammation, provoking infiltration by IFN γ -secreting proinflammatory leukocytes which become exposed to tumour neoantigens, resulting in tumour-specific immunity [1]. In an orthotopic MB49 mouse BlCa model, IFN γ was critical for intrinsic tumour surveillance and an *Ifng*-knockout rendered BCG ineffective [2].

We report the results of exposing mitotically-quiescent (G0-arrested), *in vitro*-differentiated normal human urothelium from six independent donors to 200 U/mL IFN γ to generate a tissue-specific urothelial response transcriptome. mRNA-sequencing identified 107 genes significantly ($q < 0.05$) >2-fold increased by seven-day exposure to IFN γ (Fig. 1A and Supplementary Table 1).

The IFN γ receptor genes (*IFNGR1* and *IFNGR2*) were abundantly expressed by urothelium (Supplementary Fig. 1). In all six donors, irrespective of haplotype, IFN γ induced gain of human leukocyte antigen (HLA) gene expression associated with both major histocompatibility complex (MHC) class I and class II (Supplementary Fig. 2). Gains in expression of β 2-microglobulin (*B2M*), MHC class II invariant chain (*CD74*) and trans-activator (*CIITA*), as well as the antigen peptide transporters *TAP1* and *TAP2*, suggest an induced role in modulating antigen-presentation to CD4 T cells (Supplementary Fig. 2). However, urothelial cells are unlikely to act as professional antigen presenting cells, because the co-stimulatory B7 family CD80 and CD86 genes were absent and were not inducible by IFN γ (Fig. 1C).

Blocking antibodies against the PD-L1 (CD274) immune checkpoint have been trialled clinically to disinhibit tumour-specific cytotoxic lymphocytes. *CD274* (PD-L1) was strongly induced by IFN γ (Fig. 1C). By contrast, the inhibitory B7 family member *VSIR* (VISTA) was expressed

constitutively by normal human urothelium but has to-date been overlooked in the bladder [3]. This indicates that normal urothelium, and likely its malignant counterpart, is a locally immunosuppressive environment, with induction of PD-L1 by IFN γ acting as a negative feedback loop to prevent runaway inflammation.

IFN γ -treatment induced a shift in expression of chemokines and cytokines implicated in immune recruitment, with loss (*CCL20*, *CXCL8*, *CXCL1*, *CXCL6*, *IL23A*, *IL1B*, *IL17C*, *CXCL5* and *CXCL2*) and gain (*CXCL11*, *CXCL10*, *CXCL9*, *IL18BP*, *IL32*) of different signalling factors (Fig. 1B). Of particular note, the CXCL9/10/11-CXCR3 axis is a critical regulator of leukocyte migration, differentiation and activation [4], including recruitment of effector T cells. We have shown elsewhere that *CXCL10/11* are highly induced following exposure of normal urothelium to BK virus [5] and were the only genes to be further upregulated, rather than repressed, by treatment with exogenous IFN γ . We therefore suggest that attenuated BK virus could be used to generate a pro-inflammatory, IFN γ -rich local environment akin to that induced by BCG.

Cancer subtyping is usually performed by unsupervised clustering of most statistically informative genes detected as transcripts in extracted tumour tissue; the latter representing a heterogeneous mix of tumour, stromal and immune cells. Whilst able to categorise tumours into subsets by similarity, it is a blunt tool for informing therapy. We hypothesised that using signature gene sets representing tissue-specific pathway responses would reduce “noise” and provide greater therapeutic insight.

The *IFNG* transcript can be detected in cancer cohorts but gives weak sensitivity and, as a diffusible factor, IFN γ can elicit a response in the tumour even when originating from peritumoural leukocytes. We hypothesised that a urothelial IFN γ -response signature would provide a more informative means of detecting IFN γ -signalling in a tumour. We therefore

applied a curated urothelial *in vitro* IFN γ -response gene set (Supplementary Table 2) to the unsupervised clustering of BlCa cohorts, yielding an “IFN γ -signature” score.

The IFN γ -signature was significantly correlated with *IFNG* transcript in all four BlCa cohorts analysed (see figure legends for details). In NMIBC from the UROMOL2021 cohort [6], the IFN γ -signature was weak in Class 2a tumours (Supplementary Fig. 3). In T1 tumours (combined from UROMOL2021 [6] and Northwestern Memorial Hospital (NMH) [7] cohorts; Fig. 2A) dichotomised into IFN γ -signature high and low groups, recurrence was 50% more likely in patients whose tumours lacked the IFN γ -signature (Fig. 2B). A Cox proportional hazards regression analysis of the pseudo-continuous unit-length scaled IFN γ -signature scores in T1 tumours predicted a recurrence hazard ratio of 3.174 $p=0.0230$ (Supplementary Tables 3-4).

Analysis of TCGA-BLCA data [8] revealed that MIBC lacking the IFN γ -signature were most likely to be graded histologically as lymphocyte-negative and of luminal papillary sub-type according to the consensus classification [9] (Supplementary Fig. 5). By contrast, MIBC classed as basal/squamous split into two clear subgroups (Fig. 2C). To examine any difference in outcomes associated with the IFN γ -signature in basal/squamous tumours, the Lund MIBC cohort [10] was analysed in parallel (Supplementary Fig. 6). When dichotomised into IFN γ -signature high and low groups, overall survival for basal/squamous tumours (pooled from TCGA-BLCA and Lund cohorts) was significantly lower in tumours with a weak IFN γ -signature (Fig. 2D). A high survival hazard ratio of 4.846 $p=0.0006$ was predicted by Cox proportional hazards regression based on the pseudo-continuous IFN γ -signature scores from both cohorts (Supplementary Tables 5-7).

Bladder cancer is a disease characterised by high mutational loads derived by different processes, whose historic contributions can be assessed using mutational signatures. IFN γ -signature^{high} basal/squamous MIBC tumours had significantly enriched genomic damage from APOBEC enzymes (Supplementary Fig. 7). Widespread APOBEC damage leads to tumour suppressor loss and creates neoantigens which were significantly more abundant in IFN γ -signature^{high} basal/squamous tumours [8]. In an inflammatory context, neoantigens would induce antigen-specific T cell immunity. Conversely, our data suggest that normal and malignant urothelium would suppress T cell activation through PD-L1/VISTA checkpoint inhibition, thereby inhibiting type-1 anti-tumour immune responses. It seems most probable that a balance becomes established between activated tumoricidal effector cells and regulatory/immunosuppressive elements.

To conclude, this study suggests a reconsideration of the role for IFN γ in motivating immune clearance in non-BCG-responsive NMIBC. The IFN γ -signature^{high} basal/squamous MIBC tumours identified here appear to represent a target for immune checkpoint blockade. However, in the absence of IFN γ , expression of PD-L1 is likely to be too low to provide an effective target. We suggest that BCG therapy would be most effective when combined with blocking of both the innate (VISTA [3]) and inducible (PD-L1) immune checkpoints. We anticipate the urothelium-derived transcriptomic IFN γ -signature will prove a valuable tool for the design of more efficacious immunotherapy protocols.

CRediT Author Statement

Conceptualization – SCB, JS

Methodology – SCB

137 Software – ASM, SCB, PE, GS

138 Formal analysis – SCB, ASM

139 Investigation – SCB, RGS

140 Data Curation – ASM, SCB, PE, GS

141 Writing – SCB, LKT, JS

142 Writing - Review & Editing – SCB, ASM, RGS, PE, GS, LKT, JS

143 Visualization - SCB

144 Supervision – SCB, JS

145 Project administration - SCB

146 Funding acquisition - JS

147

148 **Acknowledgements**

149 This study was funded by York Against Cancer.

150 Figures

152 *Fig. 1 – (A) mRNA-sequencing volcano plot for the IFN γ -induced normal urothelial*
153 *transcriptome. Data is expressed as transcripts per million (TPM) values and significance was*
154 *assessed using a likelihood ratio test (LRT) to generate Benjamini-Hochberg corrected q-*
155 *values. The volcano highlights the potential for urothelial-immune signalling via (B) observed*
156 *shift in chemokine-related gene expression. Stars following gene names indicate >2-fold*
157 *changes with significance *=q<0.05, **=q<0.01 and ***=q<0.001. (C) Urothelial expression*
158 *of the family of B7 peripheral membrane protein genes. The MHC co-stimulatory molecules*
159 *CD80 and CD86 were not expressed. CD274 (“PD-L1”) was significantly induced by IFN γ*
160 *treatment (mean fold change = 5.2; q=0.0095). VSIR (“VISTA”) was abundantly expressed*
161 *and significantly induced by IFN γ treatment (mean fold change = 1.2; q=0.0322). For the x-*
162 *axis labels, the B7-family name and other common names are provided in normal font above*
163 *the italicised HGNC gene name. Data in all panels is derived from normal human urothelium*
164 *established from six independent donors.*

Figure 2 – (A) Heatmap and k means clustering based on expression of the IFN γ -signature genes in the T1 tumours of UROMOL2021 and Northwestern Memorial Hospital (NMH) cohorts (n=236). The IFN γ -signature shows a Spearman rank correlation of 0.66 ($p=4 \times 10^{-31}$) with the IFNG gene in T1 tumours. The full UROMOL2021 cohort is shown as Supplemental Fig. 3; [6]). The separate NMH T1 cohort (n=99; [7]) is shown as Supplemental Fig. 4. (B) Kaplan-Meier analysis of IFN γ -signature high and low T1 tumours with survival data from UROMOL2021 and NMH cohorts combined (n=201). The Kaplan-Meier curve is truncated but no events were recorded after the truncation. A Cox proportional hazards regression was also performed using the unit-length scaled IFN γ -signature values from T1 tumours which

predicted a recurrence hazard ratio of 3.174 $p=0.0230$ (Supplementary Table 4). (C)

Heatmap for expression of the $IFN\gamma$ -signature in the Basal/Squamous group of TCGA MIBC tumours ($n=151$; full cohort is shown as Supplemental Fig. 5; [8]; shown classified in red according to the consensus report [9]). The $IFN\gamma$ -signature shows a Spearman rank correlation of 0.87 ($p=3.85 \times 10^{-47}$) with the *IFNG* gene in Ba/Sq TCGA-BLCA tumours. Similar analysis for the Lund MIBC cohort ($n=88$; [10]) is shown as Supplemental Fig. 6. (D) Kaplan-Meier analysis of the $IFN\gamma$ -signature in Basal/Squamous tumours from TCGA and Lund MIBC cohorts combined ($n=232$). The Kaplan-Meier curve is truncated but no events were recorded after the truncation. A Cox proportional hazards regression was also performed using the unit-length scaled $IFN\gamma$ -signature values from both cohorts which predicted a hazard ratio of 4.846 $p=0.0006$ (Supplementary Table 7).

References

- [1] Antonelli AC, Binyamin A, Hohl TM, Glickman MS, Redelman-Sidi G. Bacterial immunotherapy for cancer induces CD4-dependent tumor-specific immunity through tumor-intrinsic interferon-gamma signaling. *Proc Natl Acad Sci U S A*. 2020;117:18627-37.
- [2] Riemensberger J, Bohle A, Brandau S. IFN-gamma and IL-12 but not IL-10 are required for local tumour surveillance in a syngeneic model of orthotopic bladder cancer. *Clin Exp Immunol*. 2002;127:20-6.
- [3] Yum JI, Hong YK. Terminating Cancer by Blocking VISTA as a Novel Immunotherapy: Hasta la vista, baby. *Front Oncol*. 2021;11:658488.
- [4] Vollmer T, Schlickeiser S, Amini L, et al. The intratumoral CXCR3 chemokine system is predictive of chemotherapy response in human bladder cancer. *Sci Transl Med*. 2021;13.
- [5] Baker SC, Mason AS, Slip RS, et al. BK polyoma virus (BKPv) is a risk factor for bladder cancer through its induction of APOBEC3-mediated genomic damage. *bioRxiv*. 2021;<https://doi.org/10.1101/2021.05.13.443803>.
- [6] Lindskrog SV, Prip F, Lamy P, et al. An integrated multi-omics analysis identifies prognostic molecular subtypes of non-muscle-invasive bladder cancer. *Nat Commun*. 2021;12:2301.
- [7] Robertson AG, Groeneveld CS, Jordan B, et al. Identification of Differential Tumor Subtypes of T1 Bladder Cancer. *Eur Urol*. 2020;78:533-7.
- [8] Robertson AG, Kim J, Al-Ahmadie H, et al. Comprehensive Molecular Characterization of Muscle-Invasive Bladder Cancer. *Cell*. 2017;171:540-56 e25.
- [9] Kamoun A, Reynies A, Allory Y, et al. A Consensus Molecular Classification of Muscle-invasive Bladder Cancer. *Eur Urol*. 2019.
- [10] Sjodahl G, Eriksson P, Liedberg F, Hoglund M. Molecular classification of urothelial carcinoma: global mRNA classification versus tumour-cell phenotype classification. *J Pathol*. 2017;242:113-25.