USA Winter 2021 CoVID-19 Resurgence Post-Christmas Update

Genghmun Eng

PhD Physics 1978, University of Illinois at Urbana-Champaign

February 4, 2022

Abstract

We have successfully modeled every USA CoVID-19 wave using:

$$N(t) \approx \max[N_o \exp\{+(t/[t_R(1 + \alpha_S t)]) \exp(-\delta_o t)\}],$$

where $N(t)$ is the total number of new CoVID-19 cases above a prior baseline, and t_R sets the doubling time $t_{dbl} = t_R (\ln 2)$. The new parameters $\{\alpha_S; \delta_o\}$ measure mitigation efforts among the *uninfected population*, with $\{\alpha_S > 0\}$ being associated with *Social Distancing* and *vaccinations*; while $\{\delta_o > 0\}$ is associated with *mask-wearing*, which gives a faster $\frac{d}{dt}N(t)$ post-peak *drop-off*. The predicted pandemic wave end is when $N(t)$ no longer increases.

Using data from 11/15/21-12/30/21, our prior *medrxiv.org* preprint* showed this initial *Omicron* CoVID-19 wave had values that matched the *initial* stage of the prior USA Winter 2020 resurgence $\{t_R \approx 8.05 \text{ days}; \alpha_S \approx 0.011 / \text{day}\}$, when practically no one was vaccinated. In addition, this *initial* Winter 2021 wave showed virtually no *mask-wearing* $\{\delta_o \approx 0.001 \times 10^{-3} / \text{day}\}$, making it capable of infecting virtually everyone. These parameter values indicated that the *Omicron* variant was likely evading the vaccines in people who thought they were protected.

As a result, stopping the *Omicron* CoVID-19 spread must once again rely on enhanced *Social Distancing* and *mask-wearing*, just like the initial pandemic wave in March 2020. Analyzing the USA follow-on data from 12/25/21-1/31/22 shows that people did exactly that after the *Christmas Holiday* season, resulting in the following model parameters and values:

$$\begin{aligned} N_o[12/25/21] &= 3,121,000 ; \\ N[t = 1/31/22] &\approx 26,870,400 ; N[t = 3/21/22] \approx 50,522,800 ; \\ t_R &= 7.636 \text{ days} ; t_{dbl} = 5.293 \text{ days} ; \\ \alpha_S &= 0.03168 / \text{day} ; \\ \delta_o &= 0.96 \times 10^{-3} / \text{day} ; \end{aligned}$$

for this wave by itself, with all prior waves subtracted out as a baseline. Combining all the USA CoVID-19 waves gives these updated totals:

$$\begin{aligned} N_{tot}(t = [1/31/22]) &\approx 77,122,000 \text{ vs } N_{data} = 75,399,650 ; \\ N_{tot}(t = [3/21/22]) &\approx 101,476,500 ; N_{tot}(t = [6/10/22]) \approx 111,531,800 \end{aligned}$$

assuming no future CoVID-19 Resurgence (*with 4 Figures*).

*(10.1101_2021.10.15.21265078)

1 Introduction

Each USA CoVID-19 wave¹⁻⁷, from the pandemic start (3/21/20) to the present day (1/31/22), has been successfully modeled using this basic $N(t)$ function for the total number of new CoVID-19 cases above a prior baseline:

$$N(t) \approx \max[N_o \exp\{+(t / [t_R(1 + \alpha_S t)]) \exp(-\delta_o t)\}], \quad [1.1]$$

with t_R setting the pandemic doubling time $t_{dbl} = t_R (\ln 2)$, as in standard **SEIR** (**S**usceptible, **E**xposed, **I**nfected, **R**ecovered or **R**emoved) epidemiology models. The new Eq. [1.1] parameters model mitigation efforts among the *uninfected population*, with $\{\alpha_S > 0\}$ associated with *Social Distancing* and *vaccinations*; and $\{\delta_o > 0\}$ associated with *mask-wearing*, which gives a faster $\frac{d}{dt}N(t)$ post-peak *drop-off*. The predicted pandemic wave ends when $N(t)$ no longer increases.

Given a total population of N_{ALL} , the *uninfected population* $U(t)$ is:

$$U(t) = [N_{ALL} - N(t)]. \quad [1.2]$$

Using Eq. [1.1] assumes $N(t) \ll N_{ALL}$, so that pandemic saturation effects can be ignored. Also, **SEIR** models do not generally include what the $U(t)$ *uninfected population* is doing in response to the pandemic. In contrast, Eq. [1.1] was developed as a *non-local* extension of **SEIR** models, to account for how the *uninfected population*, as a whole, is mitigating the pandemic spread.

Each new USA CoVID-19 wave starts with a sharp rise in the total number $N(t)$ of new cases, while the $t = 0$ point is chosen to be when the resurgence is first easily identified, with $N(t = 0) = N_o$ being the number of cases above baseline at that time. Since Eq. [1.1] is empirically based, it does not predict when each new CoVID-19 wave will start, or what biological and social circumstances are causing the new wave. But once the CoVID-19 wave becomes established, Eq. [1.1] appears to successfully predict its time evolution.

Since the same few parameters in Eq. [1.1] have successfully modeled the time evolution of each USA CoVID-19 wave¹⁻⁷, this result shows that the **response** of the $U(t)$ *uninfected population* has been similar for each wave, even if different dominating factors were driving the CoVID-19 resurgences.

All CoVID-19 data used here came from the open-source *bing.com* CoVID-Tracker⁸ database. The initial stage of the USA Winter 2021 CoVID-19 Resurgence started when the *Omicron CoVID-19* variant became ascendant. Our prior *medrxiv.org* preprint⁷ covering data from 11/15/21-12/30/21 showed that this stage was characterized by:

$$N_o(11/15/21) = 107,000; \quad [1.3a]$$

$$t_R(11/15/21 - 12/30/21) \approx 8.05 \text{ days}; \quad [1.3b]$$

$$\alpha_S(11/15/21 - 12/30/21) \approx 0.011 / \text{days}. \quad [1.3c]$$

These values are similar to the USA Winter 2020 CoVID-19 Resurgence, which only had a fully-vaccinated rate of ~0.3%. Since the USA now has a significantly vaccinated population, it makes it likely that the *Omicron CoVID-19* variant is evading the vaccines.

While the prior 2020 Winter Resurgence had a significant amount of *mask-wearing* [$\delta_o(2020) \approx 1.748 \times 10^{-3} / \text{day}$], the *initial* part of the 2021 Winter Resurgence was associated with virtually no *mask-wearing*:

$$\delta_o(11/15/21 - 12/30/21) \approx 0.001 \times 10^{-3} / \text{day}; \quad [1.4]$$

indicating that the *Omicron* variant was infecting people who thought they were protected. While *Omicron* vaccine evasion is now generally accepted, vaccination still offers protection against hospitalization from an *Omicron CoVID-19* infection. However, with virtually no *mask-wearing*, the Eqs. [1.3a]-[1.3c] values showed that this *initial* stage of the Winter 2021 CoVID-19 wave showed that it could infect virtually all susceptible people⁷.

This result similar to the *initial* stage of the USA Summer 2021 wave⁵. That commonality shows that resurgences may be driven by people letting their guard down with respect to *Social Distancing* and *mask-wearing*. As this *initial* Winter 2021 CoVID-19 Resurgence occurred between *Thanksgiving* and *Christmas*, festivities likely contributed to this CoVID-19 surge.

However, right after the *Christmas Holiday*, when the number of new cases reached $\sim 3,120,000$ cases above baseline, the *uninfected population* began practicing enhanced *Social Distancing* and *mask-wearing*. This change is similar to what occurred during the latter portion of the USA Summer 2021 CoVID-19 wave⁵, and it is modeled here in this update.

2 Post-*Christmas* USA Winter 2021 Resurgence

Using the USA Summer 2021 CoVID-19 Resurgence as a baseline, it showed that the USA Winter 2021 CoVID-19 Resurgence started around 11/15/21. The dots in *Figure 1* show the total number of new CoVID-19 cases above the Summer 2021 USA CoVID-19 baseline⁵⁻⁶. The *Figure 1* continuous line shows the original CoVID-19 projections for this *initial* stage of the USA Winter 2021 CoVID-19 Resurgence⁷, using data from 11/15/21-12/30/21.

Figure 2 includes follow-on data after this *initial* USA Winter 2021 CoVID-19 Resurgence, with data and models updates for 12/25/21-1/31/22, covering the post-*Christmas* CoVID-19 surge. Almost immediately after the *Christmas Holiday*, the *uninfected population* returned to enhanced *Social Distancing* and *mask-wearing*, creating a distinct change in the $N(t)$ function after 12/25/21.

Modeling the 12/25/21-1/31/22 period of the USA Winter 2021 wave in *Figure 2* shows that this CoVID-19 wave can subside prior to infection of the entire USA population. This *latter* portion has these model parameters:

$$N_o(12/25/21) = 3,121,000; \quad [2.1a]$$

$$t_R(12/25/21 - 1/31/22) \approx 7.636 \text{ days}; \quad [2.1b]$$

$$t_{dbl}(12/25/21 - 1/31/22) \approx 5.293 \text{ days}; \quad [2.1c]$$

$$\alpha_S(12/25/21 - 1/31/22) \approx 0.03168 / \text{day}; \quad [2.1d]$$

$$\delta_o(12/25/21 - 1/31/22) \approx 0.96 \times 10^{-3} / \text{day}. \quad [2.1e]$$

The $N_o(12/25/21) = 3,121,000$ starting point for the behavior change in $N(t)$ is similar to the prior Summer 2021 Resurgence, which also showed a behavior change in $N(t)$ at $N_o(8/13/21) = 3,200,000$. This commonality shows that the *uninfected population* altered their behavior at similar points, which is likely when some hospitals became overwhelmed.

Figure 3 shows the total number of CoVID-19 cases, $N_{tot}(t)$, resulting from combining all waves of the CoVID-19 pandemic, from March 2020 through January 2022. Whenever the pandemic appeared to be beaten down, people became more complacent, allowing new CoVID-19 variants to spread, and the pandemic rose up again, almost with every season.

The daily number of new cases $dN_{tot}(t)/dt$ in Figure 3 has peaks for the initial Spring 2020 pandemic; a Summer 2020 resurgence; the long Winter 2020 Resurgence; a small uptick in Spring 2021; the Summer 2021 Resurgence; and now the Winter 2021 Resurgence.

Figure 4 presents a tabulated summary of all the model parameters that were derived for each CoVID-19 wave. It shows these overall features:

- (a) The $t_{dbl} = t_R (\ln 2)$ doubling times for several CoVID-19 waves vary only from $\{5.3 - 5.6 \text{ days}\}$.
- (b) The α_S *Social Distancing* and *vaccination* parameter for this CoVID-19 wave is one of the largest values, except for the early 2020 pandemic start.
- (c) Given a fast-rising CoVID-19 pandemic Resurgence, the *uninfected population* generally takes notice of the severity of a new CoVID-19 wave when the new infections reach a level of $\sim 3,000,000$ above its baseline.
- (d) The *mask-wearing* parameter, δ_o , models the width and duration of the post-peak tail. Increased *mask-wearing* remains one the most powerful factors for hastening the CoVID-19 pandemic end.

3 Summary

Since our CoVID-19 modeling has been successful at predicting the time evolution of each USA CoVID-19 wave¹⁻⁷, using the same few parameters:

$$N(t) \approx \max[N_o \exp\{+(t/[t_R(1 + \alpha_S t)]) \exp(-\delta_o t)\}], \quad [3.1]$$

it indicates that the **response** of the $U(t)$ *uninfected population* was similar for each wave, even if different dominating factors drive each new resurgence.

The initial stage of the USA Winter 2021 CoVID-19 Resurgence started with the *Omicron CoVID-19* variant becoming dominant. Our prior *medrxiv.org* preprint⁷, covering data from 11/15/21-12/30/21, showed that this *initial* stage was characterized by:

$$N_o(11/15/21) = 107,000; \quad [3.2a]$$

$$t_R(11/15/21 - 12/30/21) = 8.05 \text{ days}; \quad [3.2b]$$

$$\alpha_S(11/15/21 - 12/30/21) = 0.011 / \text{days}; \quad [3.2c]$$

$$N(t = 12/25/21) \approx 3,372,300. \quad [3.2d]$$

These $\{N_o; t_R; \alpha_S\}_{11/15/21}$ model parameters are very similar to the prior USA Winter 2020 CoVID-19 Resurgence, when the USA only had a fully-vaccinated rate of $\sim 0.3\%$. Since the USA now has a significantly vaccinated population, this new Winter 2021 Resurgence *Omicron CoVID-19* variant is likely evading the vaccines. Since the $N_o(12/25/21)$ starting point of Eq. [2.1a] was nearly six weeks after the Eq. [3.2a] starting point, those two values should be similar, as they are.

Just after the *Christmas Holiday*, the *uninfected population* responded to this CoVID-19 surge with enhanced *Social Distancing* and *mask-wearing*. The 12/25/21-1/31/22 post-*Christmas* data gives Eq. [2.1a]-[2.1e] as model parameters for this *latter* portion of the USA Winter 2021 CoVID-19 Resurgence, along with these updated $N(t)$ predictions for this CoVID-19 wave by itself:

$$N[t = 1/31/22] \approx 26,870,400 ; \quad [3.4a]$$

$$N[t = 3/21/22] \approx 50,522,800 ; \quad [3.4b]$$

$$N[t = 6/10/22] \approx 60,329,500 ; \quad [3.4b]$$

with this wave no longer infecting practically everyone. Combining these results with the prior *medrxiv.org* preprint⁷ data, gives these updated USA CoVID-19 projections for the total number of USA CoVID-19 cases:

$$N_{tot}(t = [1/31/22]) \approx 77,122,000 \text{ vs } N_{data} = 75,399,650 ; \quad [3.5a]$$

$$N_{tot}(t = [3/21/22]) \approx 101,476,500 ; \quad [3.5b]$$

$$N_{tot}(t = [6/10/22]) \approx 111,531,800 ; \quad [3.5c]$$

from all USA CoVID-19 waves, assuming no new CoVID-19 resurgence intervenes. More *mask-wearing* would further mitigate CoVID-19 spread, and it could significantly reduce these projected future USA CoVID-19 totals.

4 List of Figures

Fig. 1: The Initial USA CoVID-19 Winter 2021 Resurgence, by Itself

Fig. 2: USA CoVID-19 Winter 2021 Resurgence Post-Christmas Update

Fig. 3: USA CoVID-19 Totals: 3/21/2020 through 1/31/2022

Fig. 4: Summary of CoVID-19 Model and Parameter Values

5 References

1. <https://medrxiv.org/cgi/content/short/2020.05.04.20091207v1>
"Initial Model for the Impact of Social Distancing on CoVID-19 Spread"
2. <https://medrxiv.org/cgi/content/short/2020.06.30.20143149v1>
"Orthogonal Functions for Evaluating
Social Distancing Impact on CoVID-19 Spread"
3. <https://medrxiv.org/cgi/content/short/2020.08.07.20169904>
"Model to Describe Fast Shutoff of CoVID-19 Pandemic Spread"
4. <https://medrxiv.org/cgi/content/short/2020.09.16.20196063>
"Initial Model for USA CoVID-19 Resurgence"
5. <https://medrxiv.org/cgi/content/short/2021.08.16.21262150>
"The *IHME* vs Me: Modeling USA CoVID-19 Spread,
Early Data to the *Fifth Wave*"
6. <https://medrxiv.org/cgi/content/short/2021.10.15.21265078>
"Updated Model for the USA Summer 2021 CoVID-19 Resurgence"
7. <https://medrxiv.org/cgi/content/short/2022.01.06.22268868>
"Modeling the USA Winter 2021 CoVID-19 Resurgence"
8. www.bing.com/covid/local/unitedstates?form=COVD07

Fig. 1: The Initial USA CoVID-19 Winter 2021 Resurgence, by Itself

Fig. 2: USA CoVID-19 Winter 2021 Resurgence Post-Christmas Update

Fig. 3: USA CoVID-19 Totals: 3/21/2020 through 1/31/2022

Fig. 4: Summary of CoVID-19 Model and Parameter Values

Index	U.S.A. CoVID-19 Stage	N_0 , (Initial $t=0$ Value)	$t=0$ Date	t/dbl days, at $t=0$	t_R value	α_S value	δ_o value
1	Ini. Model 6/7/20 Update	23,710	03/21/2020	1.99657	2.88044	0.06618	0.00000
2	1 st Resurgence Summer 2020 =>	15,650	06/07/2020	2.88000	4.15496	0.058__	0.0108_
3	Winter 2020 Resurgence =>	88,900	09/25/2020	5.57983	8.05__	0.011__	0.001748
4	Small Spring 2021 Resurgence =>	146,000	03/19/2021	5.57983	8.05__	0.00128	0.01365
5	Initial Portion, Summer 2021 Resurgence =>	41,000	06/07/2021	9.08924	13.113_	0.0030_	0.00000
6	Latter Portion, Summer 2021 Resurgence =>	3,200,000	08/13/2021	14.44339	20.83740	0.019999	0.000489
7	Initial Portion, Winter 2021 Resurgence =>	107,000	11/15/2021	5.57983	8.05__	0.011__	0.000001
8	Latter Portion, Winter 2021 Resurgence =>	3,121,000	12/25/2021	5.29287	7.636_	0.03168_	0.00096_