Estimating the risk of incident SARS-CoV-2 infection among healthcare workers in quarantine hospitals: the Egyptian example

Sofía Jijón^{1,2†} (PhD), Ahmad Al Shafie^{3†} (MSc), Essam Hassan⁴ (MD), EMAE-MESuRS working group on nosocomial SARS-CoV-2 modeling, Laura Temime^{1,2} (PhD), Kévin Jean^{1,2,5*} (PhD), Mohamed El-Kassas^{3*} (MD)

¹ Laboratoire Modélisation, épidémiologie et surveillance des risques sanitaires (MESuRS), Conservatoire national des Arts et Métiers, Paris, France

² Unité PACRI, Institut Pasteur, Conservatoire national des Arts et Métiers, Paris, France

³ Endemic Medicine Department, Faculty of Medicine, Helwan University, Cairo, Egypt

⁴ Tropical Medicine Department, Faculty of Medicine, Fayoum University, Fayoum, Egypt

⁵ MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London, United Kingdom

[†] Contributed equally to the work

*Contributed equally to the work

Corresponding author:

Sofía Jijón, sofia.jijon_alban@sorbonne-universite.fr

Current address : Sorbonne Université, Campus Pierre et Marie Curie, 4 Place Jussieu, 75005 Paris, France.

Running title: SARS-CoV-2 infection risk among HCW in quarantine hospitals

Word count. Abstract: 296 words; Main text: 3219 words (3932 including captions and tables)

26 ABSTRACT

27 In response to the COVID-19 epidemic, Egypt established a unique care model based on
28 quarantine hospitals where only externally-referred confirmed COVID-19 patients were
29 admitted, and healthcare workers resided continuously over 1- to 2-week working shifts.
30 While the COVID-19 risk for HCWs has been widely reported in standard healthcare settings,
31 it has not been evaluated yet in quarantine hospitals.

32 Here, we relied on longitudinal data, including results of routine RT-PCR tests, collected
33 within three quarantine hospitals located in Cairo and Fayoum, Egypt. Using a model-based
34 approach that accounts for the time-since-exposure variation in false-negative rates of RT-
35 PCR tests, we computed the incidence of SARS-CoV-2 infection among HCWs. Over a total
36 follow-up of 6,064 person-days (PD), we estimated an incidence rate (per 100 PD) of 1.05
37 (95% CrI: 0.58–1.65) at Hospital 1, 1.92 (95% CrI: 0.93–3.28) at Hospital 2 and 7.62 (95%
38 CrI: 3.47–13.70) at Hospital 3. The probability for an HCW to be infected at the end of a
39 shift was 13.7% (95% CrI: 7.8%–20.8%) and 23.8% (95% CrI: 12.2%–37.3%) for a 2-week
40 shift at Hospital 1 and Hospital 2, respectively, which lies within the range of risk levels
41 previously documented in standard healthcare settings, whereas it was >3-fold higher for a 7-
42 day shift at Hospital 2 (42.6%, 95%CrI: 21.9%–64.4%). Our model-based estimates unveil a
43 proportion of undiagnosed infections among HCWs of 46.4% (95% CrI: 18.8%–66.7%),
44 45.0% (95% CrI: 5.6%–70.8%) and 59.2% (95% CrI: 34.8%–78.8%), for Hosp1 to 3,
45 respectively.

46 The large variation in SARS-CoV-2 incidence we document here suggests that HCWs from
47 quarantine hospitals may face a high occupational risk of infection, but that, with sufficient
48 anticipation and infection control measures, this risk can be brought down to levels similar to
49 those observed in standard healthcare settings.

50
51 **Keywords.** COVID-19; SARS-CoV-2; Nosocomial infection; Healthcare workers;
52 Quarantine hospitals.

53 **WHAT THIS PAPER ADDS**

54 **What is already known on this topic**

55 Previous studies conducted in standard care setting documented that frontline healthcare
56 workers (HCWs) face a high COVID-19 risk. Whether risk levels differ in alternative care
57 models, such as the Egyptian COVID-19 care organization based on quarantine hospitals
58 where HCWs resided days and nights for various length, is unknown.

59 **What this study adds**

60 The COVID-19 risk for HCWs in quarantine hospitals varies substantially between facilities,
61 from risk levels that are in the range of those documented in standard healthcare settings to
62 levels that were approximatively 3 times higher.

63 **How this study might affect research, practice or policy**

64 With sufficient anticipation and infection control measures, occupational COVID-19 risk for
65 HCWs working in quarantine hospitals can be brought down to levels similar to those
66 observed in standard healthcare settings.

67 INTRODUCTION

68 Healthcare settings have faced multiple challenges related to the currently ongoing severe
69 acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic. They have notably had
70 to deal with the influx of infected patients across successive epidemics waves while
71 controlling the risk of nosocomial spread to other patients and staff. As a result, healthcare
72 workers (HCWs) have been a population of interest in terms of risk assessment and control
73 measures implementations [1].

74 At early stages of the pandemic, Egypt was identified as one of the African countries most
75 vulnerable to a SARS-CoV-2 importation [2]. In February 14, 2020, Egypt reported the first
76 confirmed case of coronavirus disease (COVID-19) in Africa and remained among the five
77 African countries most affected by the COVID-19 epidemic up to the end of 2020 [3]. From
78 February 14 to August 31, 2020 (period to which we refer to as the first wave), the COVID-
79 19 epidemic in Egypt resulted in about 9,700 confirmed infections and 5,500 deaths reported
80 nationally [4]. These numbers most certainly represent an underreporting of the real number
81 of infections, due to the high proportion of asymptomatic SARS-CoV-2 infections [5,6] and
82 limited testing.

83 To mitigate the high risk of SARS-CoV-2 spread within the population, Egypt established a
84 unique care model under the World Health Organization supervision, whereby specific
85 hospitals were assigned as quarantine hospitals for patients with COVID-19, and where
86 dedicated medical teams stayed in the hospital days and nights during working shifts of
87 various lengths [7].

88 While the quarantine-hospital strategy has the potential to be highly efficient in terms of
89 patient care, as well as in limiting the potential spread of the virus from hospitals into the
90 community, its impact on the infection risk for HCWs remains understudied. One theoretical
91 modelling study assessing healthcare working force organization found that alternating HCW
92 teams by 1-week periods may reduce the overall number of infected HCWs [8]. On the other
93 hand, to the best of our knowledge, previous epidemiological studies focusing on the infection
94 risk faced by Egyptian HCWs have been conducted in non-quarantine settings exclusively [9–
95 11]. Yet, the quarantine strategy was adopted again to face the second wave of the COVID-19
96 epidemic at the national level, and could be adopted for future epidemics waves in Egypt
97 and/or in other countries [12].

98 Here, we aimed at using mathematical modelling to estimate the risk of SARS-CoV-2
99 infection among HCWs participating in quarantine-hospital interventions, relying on detailed
100 longitudinal data collected in three Egyptian healthcare facilities, during the first wave of the
101 COVID-19 epidemic that hit the country.

METHODS

Study settings

Data was collected within three Egyptian hospitals (hereafter denoted by Hosp1, Hosp2 and Hosp3) located in Cairo (Hosp1 and Hosp3), and Fayoum (middle Egypt, Hosp2), that were temporarily transformed into quarantine hospitals during the first COVID-19 wave. During the quarantine-organization period (Hosp1: March 14th to August 1st, 2020; Hosp2: April 1st to July 31th; Hosp3: June 6th to July 11th, 2020), only externally-referred COVID-19 confirmed patients were admitted to these hospitals for medical care. Multidisciplinary medical teams fully dedicated to COVID-19 patients management worked in total isolation, organized by shifts (Hosp1: 2-week shifts; Hosp2: 1 to 2-week shifts; Hosp3: 1-week shifts). During their shift, HCWs were affected either to intensive care units (ICU) or non-ICU. HCWs were screened for SARS-CoV-2 infection before starting a working shift using rapid serological IgM/IgG antibody tests (Artron laboratories Burnaby, Canada; sensitivity: 83.3%, specificity: 100% [13]). Only HCWs with no SARS-CoV-2 antibodies were allowed to start working in the hospital; except for the last shift in Hosp3, where staff recruitment relied exclusively on HCWs who had been previously infected (positive serological tests), and were additionally tested using reverse transcription-polymerase chain reaction (RT-PCR) on nasopharyngeal swabs, obtaining negative results. During the working shifts, HCWs were tested for SARS-CoV-2 infection using RT-PCR tests: i) routinely at the end of the shift, ii) upon symptoms, and iii) in case of outbreak suspicion (>2 positive tests among HCWs). HCWs testing negative at the end of the shift were then released for self-isolation at home for two weeks. HCWs testing positive before or during their shift self-isolated at home in the case of presenting no or mild symptoms, or were admitted to the same quarantine hospital for medical care in the case of presenting moderate to severe symptoms.

Observed risk of SARS-CoV-2 infection

Crude incidence rates were obtained from the number of incident SARS-Cov-2 infections observed (ie, diagnosed during or at the end of a shift) among HCWs screened as negative for SARS-CoV-2 before their working shift. Incidence rate ratios were obtained through a Poisson regression adjusted by hospital and by affectation (ICU/ non-ICU). In addition, we computed the attack rates over each working shift, based on the observed infections among HCWs having a negative screening test at the beginning of the working shift. Observed per-

shift attack rates of $\geq 20\%$ were considered to be healthcare-associated outbreaks among HCWs.

Model-based estimates of the incidence rate of SARS-CoV-2 infection

An important feature arising from the COVID-19 surveillance within quarantine hospitals is the right-truncation of data: HCWs infected a short time before the end of a shift are likely to remain undiagnosed despite the systematic testing at the end of shifts, especially if the test is performed early in the incubation period. We thus developed the following mathematical model to estimate the incidence rate of both diagnosed and undiagnosed SARS-CoV-2 infections.

We simulated the daily number of incident infections among HCWs, $X_{k,i}^A$ based on an unobserved binomial process:

$$X_{k,i}^A \sim \text{Bin}(n = S_{k,i}^A, p = \lambda)$$

where k denotes the working shift, i denotes the day within the shift, A is the affectation of HCWs ($A \in \{ICU, Non-ICU\}$), $S_{k,i}^A$ denotes the number of susceptible HCWs in affectation A , at the beginning of the i -th day of shift k and λ denotes the constant daily probability of SARS-CoV-2 infection faced by HCWs. The number of susceptible HCWs was updated on a daily basis:

$$S_{k,i+1}^A = S_{k,i}^A - X_{k,i}^A$$

To model right-truncation in the surveillance data, we simulated an observation process representing the systematic RT-PCR testing of HCWs at the end of each shift, accounting for documented changes in the test sensitivity with the time since infection [14]. The period of time since infection was computed as $j_k - i$, where j_k denotes the day at the end of shift k (i.e., the day where testing was performed) and i denotes the day of infection. Then, the daily number of incident infections eventually diagnosed at the end of the k -th shift was given by

$$I_{k,i}^A = X_{k,i}^A \theta_{j_k - i}$$

where θ_{j_k-i} denotes the probability of testing positive to RT-PCR, $j_k - i$ days after infection.

We used a Bayesian Markov Chain Monte-Carlo (MCMC) approach to estimate the parameter λ that best fitted the observed number of infections. We assumed that λ was constant over the study period. We considered a non-informative uniform prior distribution $\sim U(0,1)$ for λ and the following binomial likelihood function:

$$L(\lambda|data) = \prod_A \left(\frac{\sum_{k,i} N_{k,i}^{S,A}}{\sum_{k,i} N_{k,i}^{I,A}} \right) \left(\frac{\sum_{k,i} I_{k,i}^A}{\sum_{k,i} S_{k,i}^A} \right)^{\sum_{k,i} N_{k,i}^{I,A}} \left(1 - \frac{\sum_{k,i} I_{k,i}^A}{\sum_{k,i} S_{k,i}^A} \right)^{\sum_{k,i} N_{k,i}^{S,A} - \sum_{k,i} N_{k,i}^{I,A}},$$

where $\sum_{k,i} N_{k,i}^{S,A}$ and $\sum_{k,i} N_{k,i}^{I,A}$ denote the total number of susceptible HCWs (denoted by the superscripts S) and the total number of HCW infected and diagnosed with COVID-19 (denoted by the superscript I), observed in affectionation A , during the study period, respectively. The model was thus fitted to the total number of observed infections in each affectionation of the hospital.

Medians and 95% credibility intervals (CrI) for λ were computed from posterior samples obtained after 10,000 model runs. The chains were visually inspected for convergence. For each hospital, we further estimated the probability for a susceptible HCW to be infected at the end of a shift as

$$P(InI) = 1 - (1 - \lambda)^d,$$

where d is the length of the working shift, in days.

Finally, to assess the proportion of undetected infections among HCWs, we ran our model on each hospital, using the posterior distributions for λ , to obtain an estimate for the real number of SARS-CoV-2 infections in each hospital (both diagnosed and undiagnosed), over the study period.

The numeric implementation of the model was built in R 4.0.3 [15], using the FME package [16] for parameter estimation.

Ethical approval

This study was approved by the Research Ethics Committee (REC) of the Central Directorate of Research and Health Development and Reviews at the Egyptian Ministry of Health and Population (Serial: 25-2020/16), and by REC for human subject research at the Faculty of Medicine, Helwan University (Serial: 50-2020).

RESULTS

The study period covered ten 2-week shifts in Hosp1, nine ~2-week shifts in Hosp2 and five 1-week shifts in Hosp3. The mean (min–max) number of HCWs per shift was 46 (34–63) in Hosp1, 15 (5–26) in Hosp2 and 19 (16–20) in Hosp3 (Table 1). Over a total follow-up of 8,733 person-days (PD), 54 SARS-CoV-2 infections were observed (ie., diagnosed) among 722 HCWs showing no evidence of SARS-CoV-2 antibodies at the beginning of their shifts, within the three hospitals. This represented an overall incidence rate of 0.62 (95% CI: 0.45–0.78) diagnosed SARS-Cov-2 infections per 100 PD (Table 2). A significantly higher incidence rate was observed for HCWs working at Hosp3 as compared to the other two hospitals; and HCWs working in non-ICU units tended to be more at risk than HCWs working in ICU, though this difference was not statistically significant (Table 2).

Table 1. Hospital characteristics.

Quarantine organization period, location and mean number of HCWs and patients by hospital (Hosp1, Hosp2 and Hosp3). Abbreviations: HCWs = Healthcare workers, PD = person-days.

	Hosp1	Hosp2	Hosp3
Quarantine organization period	March 14 – August 1	April 1 – July 31	June 6 – July 11
Location	Cairo	Fayoum	Cairo
Mean daily number of patients (min–max)	62 (0–108)	37 (0–103)	8 (0–20)
Mean number of HCWs per shift (min–max)	46 (34–63)	15 (5–26)	19 (16–20)
Shift duration	14 days	7–14 days	7 days

Table 2. Observed risk of SARS-CoV-2 infection among HCWs, by hospital and by affectation.

A total of 54 infections were observed (ie., diagnosed before or at the end of a shift) over 8,733 person-days, in the three hospitals (Hosp1, Hosp1 and Hosp3). Crude rates of SARS-CoV-2 infection among HCWs by hospital and by affectation: intensive and non-intensive care units and 95% confidence intervals. A Poisson regression adjusted by hospital and by affectation was performed on the observed infections to obtain the incidence rate ratios and 95% confidence intervals. Abbreviations: CI = confidence interval, ICU = intensive care unit, IRR = incidence rate ratio, PD = person-days.

		Observations		Crude rates		Adjusted Poisson Regression		
		Events	PD	Rate per 100 PD	95% CI	IRR	95% CI	p
Hospital	Hosp1	28	6258	0.45	0.28–0.61	1	ref	–
	Hosp2	11	1808	0.61	0.25–0.97	1.49	0.74–2.8745	0.28
	Hosp3	15	667	2.25	1.1–3.39	5.59	3.13–10.01	<0.001
Affectation	ICU	15	2628	0.57	0.28–0.86	1	ref	–
	Non-ICU	39	6105	0.64	0.44–0.84	1.49	0.85–2.59	0.16

Healthcare-associated outbreaks among HCWs (observed attack rates of $\geq 20\%$) represented $\sim 70\%$ (38/54) of all SARS-CoV-2 infections observed over the study period (Figure 1). Two of these outbreaks occurred in Hosp2 with 30% (3/10) and 24% (5/21) of HCWs being infected over two different working shifts. An outbreak where 36% of HCWs were infected (16/44) occurred in Hosp1 around the same time period. Two outbreaks took place later on in Hosp3, leading to, respectively, 30% (6/20) and 40% (8/20) of susceptible HCWs being infected over two different shifts. Of note, each outbreak consisted in infections in both ICU and non-ICU affectations. For the last shift in Hosp3, staff recruitment relied exclusively on HCWs who had been infected during the first or the second shifts. This explains why this last shift remained free of infections among HCWs in Hosp3 (cf. Figure 1).

Figure 1. Per-week number of infections by affection.

For a given working shift, healthcare workers were affected to either the intensive care unit (ICU, yellow) or a non-ICU (blue). Dates are given in weeks, denoted by the prefix W followed by the week number of year 2020. Hosp1 was established as a quarantine hospital from March 14th to August 1st (W11 to W31), Hosp2 from April 1st to July 31th (W14 to W31) and Hosp3 from June 6th to July 11th (W23 to W28). In Hosp1 and Hosp2, most

To estimate the mean incidence rate of both observed and unobserved infections in each

hospital, we ran our model considering a constant risk, , over the whole study period, in each

affection of each hospital. The chains for λ , for the three hospitals, are depicted in the

Supplementary Figure 1. We estimated significantly different risk levels between the three

hospitals (Kolmogorov-Smirnov test; p-value < 0.001), with an incidence rate of 1.05 (95%

CrI: 0.58–1.65) per 100 PD in Hosp1, 1.92 (95% CrI: 0.93–3.28) per 100 PD in Hosp2 and

7.62 (95% CrI: 3.47–13.70) per 100 PD in Hosp3.

The probability for a HCW to be infected at the end of a working shift was estimated at

13.7% (95% CrI: 7.8%–20.8%) and 23.8% (95% CrI: 12.2%–37.3%) for 2-week shifts at

Hosp1 and Hosp2, respectively, whereas a much higher probability of 42.6% (95% CrI:

21.9%–64.4%) was found for a 7-day shift at Hosp3 (see Table 3). These probabilities

become 7.1% (95% CrI: 4.0%–11.0%) and 12.7% (95% CrI: 6.3%–20.8%), for Hosp1 and

Hosp2, respectively, if 7-day shifts are considered as well, for easier comparison (Table 3).

Given the notable differences between hospitals (Figure 1), we did not assess infection risk by

affection across hospitals, but within each hospital.

The posterior distributions for the incidence rate λ yield the following model-based median

numbers of total SARS-CoV-2 infections: 59 (95% CrI: 45–74) at Hosp1, 31 (95% CrI: 14–

52) at Hosp2 and 41 (95% CrI: 19–63) at Hosp3 (Table 3), which unveils a proportion of

undiagnosed infections among HCWs of 46.4% (95% CrI: 18.8%–66.7%), 45.0% (95% CrI:

5.6%–70.8%) and 59.2% (95% CrI: 34.8%–78.8%), respectively.

We further ran our model on each affectation on each hospital, which confirmed a notable difference between Hosp1–2 and Hosp3: our results suggest that HCWs working in Hosp1 and Hosp2 face a higher risk when affected to non-ICU, as compared to those affected to the ICU; whereas HCWs in the ICU of Hosp3 face the higher risk (see Fig. 1 and Table 3).

Table 3. Model-based estimates of all SARS-CoV-2 infections

The model-based estimates of the incidence rate, the probability of infection for a 7-day working shift and the estimated number of SARS-CoV-2 infections in each hospital, over the study period. Running our model for each affectation (ICU and non-ICU) in each hospital unveils the difference in the risk faced by HCWs between the three hospitals of our study. Indeed, HCWs working in Hosp1 and Hosp2 face a higher risk when affected to non-ICU, as compared with those working in the ICU; whereas HCWs in Hosp3 face a slightly higher risk when affected to the ICU. All estimates concern both diagnosed and undiagnosed infections (all), unless otherwise stated (diagnosed). Abbreviations: CrI = credibility interval, ICU = intensive care unit, PD = person-days.

	Median estimated incidence rate per 100 PD (95% CrI)			Probability of infection for a 7- day shift (95% CrI)	Median number of SARS-CoV-2 infections (95% CrI)		
	ICU	Non-ICU	Overall		All	Diagnosed	Proportion of undiagnosed infections
Hosp1	0.59 (0.11–1.61)	1.27 (0.68–2.01)	1.05 (0.58–1.65)	7.1% (4.0%–11.0%)	59 (45–74)	31 (15–51)	46.4% (18.8%–66.7%)
Hosp2	1.33 (0.33–3.33)	2.75 (1.12–5.02)	1.92 (0.93–3.28)	12.7% (6.3%–20.8%)	31 (14–52)	17 (7–31)	45.0% (5.6%–70.8%)
Hosp3	9.37 (3.48–19.66)	7.12 (2.06–18.71)	7.62 (3.47–13.70)	42.6% (21.9%–64.4%)	41 (19–63)	16 (6–30)	59.2% (34.8%–78.8%)

DISCUSSION

Here, we studied the risk of incident SARS-CoV-2 infection among HCWs residing days and nights in quarantine hospitals, relying on detailed longitudinal data collected in three Egyptian quarantine hospitals. We found that most diagnosed infections (70%) occurred during what we considered as healthcare-associated outbreaks (as compared to isolated spill-over infections). We observed a high variability in the incidence, ranging from 0.45 to 2.25 per 100 PD across the three hospitals. Relying on a model-based approach, we estimated the risk of both diagnosed and undiagnosed SARS-CoV-2 infections among HCWs, by hospital and by affectation (ICU and non-ICU). We further estimated that a substantial proportion of infections may remain undetected, ranging from 45.0% to 59.2% across the three hospitals.

Our study encounters several limitations. First, screening HCWs using serological tests before starting their working shifts may not detect infections at the time of testing. Indeed, antibodies may be detected by serological tests in less than 40% of infected individuals within 7 days since symptoms onset, reaching the higher detection levels at day 16 after symptom onset [17]. Hence, serological screening may allow recently infected HCWs –who are probably infectious– to start a working shift; especially in Hosp3, where the working shifts lasted 7 days. However, the Egyptian public health authorities’ choice of a protocol relying on serological tests before working shifts was mainly driven by material constraints. Indeed, the use of rapid serological tests as a diagnostic tool was and remains of particular interest in limited-resources contexts due to their lower cost, the minimal equipment required, and faster results, as compared with RT-PCR tests [18]. Second, the assumption of a constant overall risk of infection among HCWs conducted to disregard dynamics driven by variations in the number of HCWs and/or patients in the hospitals, as well as the frequency and nature of HCWs’ contacts during their working shifts. This simplification was made in order to estimate average transmission rate first; fine reproduction of transmission dynamics were left for future work. Third, despite the quarantine strategy being adopted nationally, the differences between the organizations of each of the three hospitals (working shifts lengths, quarantine-organization period) resulted in some difficulties in comparing the risk estimations across the three hospitals. Data on other hospitals adopting the quarantine strategy may have been beneficial to better understand its performance of the quarantine-organization strategy, with regard to standard settings.

We found a ~5-fold higher observed risk for Hosp3 as compared with the risk found in Hosp1; whereas the risk of infection at Hosp2 was just slightly higher. The higher risk of SARS-CoV-2 infection we report for Hosp3 and the higher risk faced by HCWs affected to the ICU of Hosp3 may be partly explained by the short period over which this hospital adopted the quarantine organization, which coincided with the country's highest epidemic activity of the first wave [4]. This may have led to a high proportion of severe and thus highly contagious COVID-19 patients referred to quarantine hospitals together with a higher workload for HCWs. Contrarily, Hosp1 adopted the quarantine organization in mid-March 2020, and thus experienced some weeks of lower epidemic intensity that may have improved preparedness for intense COVID-19 activity, as well as implementing control measures for invasive procedures (ie., the ICU).

Our model-based approach, accounting for false-negativity rates of testing, as well as the right-truncation of our data, allowed us to estimate the risk of both diagnosed and undiagnosed SARS-CoV-2 infections among HCWs, by hospital and by affectation. Of note, the higher proportion of infections that remained undetected in Hosp3 (59.2%, versus 46.4% and 45.0% in Hosp1 and Hosp2, respectively) may be explained by the shorter duration of the working shifts (7 days), which may result in false-negative tests for HCWs infected a short time before the end of their shift. We estimated that a rigorous quarantine organization with systematic testing of HCWs at the end of their working shifts could potentially capture only about half of all HCWs infections (depending on the length of the working shift). Moreover, these results suggest that, in the absence of the quarantine-hospital organization and, more specifically, in the absence of systematic testing at the end of the working shifts, a probably larger proportion of infections among HCWs may remain undetected, thus putting HCWs' close contacts at risk of infection [19] and, consequently, putting themselves at risk for adverse psychological symptoms [20].

The model-based estimates we found for the SARS-CoV-2 infection risk in Hosp1 and Hosp2 are consistent with the infection point-prevalence reported in earlier studies performed in non-quarantine Egyptian hospitals (varying from 4.2% to 14.3%) or with the incidence reported in front-line HCWs in the UK (13% infection rate after one month) [10,11,21]. A summary of previous results obtained early in the Covid-19 pandemic and/or specifically in the Egyptian context is presented in Table 4, to ease the comparison. Moreover, our results suggest that HCWs affected to non-ICU face a higher risk than those affected to the ICU, in Hosp1 and

Hosp2, which is in line with what was observed in previous studies addressing the occupational risk of SARS-CoV-2 infection for HCWs in non-quarantine settings [22]. Overall, our findings on Hosp1 and Hosp2 suggest that, providing sufficient preparedness, HCWs working in quarantine hospitals may not face a higher infection risk, and thus highlight the benefits of implementing a quarantine-hospital strategy.

Table 4. Our results in a context.

Our estimation of the per-shift probability of infection in Hosp1–3, along with estimates of the point-prevalence in other Egyptian studies and the observed cumulative incidence rate among HCWs in an English study.

Indicator	Estimate (%)	Context	Study
Per-shift probability of infection (95% CrI)	12.8 (7.6–19.5)	Hosp1 (14-days shift)	Our results
	17.3 (7.5–30.7)	Hosp2 (14-days shift)	
	48.2 (23.8–74.5)	Hosp3 (7-days shift)	
Point-prevalence	4.2	Symptomatic and asymptomatic HCWs in 12 healthcare facilities (N=4040).	[23]
Point-prevalence	14.3	Asymptomatic HCWs in emergency departments of tertiary care facility (N=203).	[10]
Incidence rate	13.0	Front-line HCWs testing negative at enrollment.	[21]

A notable strength of our study lies in the specific nature of the quarantine hospital set-up: because HCWs resided continuously in the hospitals over their entire working shifts, we were able to specifically quantify the nosocomial risk for HCWs. Conversely, results reported from non-quarantine hospitals worldwide are generally unable to distinguish between the nosocomial vs. community risk of infection. However, as most of studies conducted in healthcare settings, we were unable to distinguish between patient-to-HCW and HCW-to-HCW routes of transmission. A previous study on several nosocomial Covid-19 healthcare-associated outbreaks in Germany reported that HCW-to-HCW transmission could represent an outweighing risk as compared to the one due to infected patients [24]. Still, assessing the relative contribution between patient-to-HCW versus HCW-to-HCW SARS-Cov-2 transmission was left for future work. However, it is worth to note that during the outbreaks observed in our study, infections occur in different affectations, which may prime HCW-to-HCW transmission, rather than simultaneous independent events of patient-to-HCW

transmission. Personal protective equipment (PPE) places a focus on the risk induced by patients, but could contribute to underestimate the risk of infection from colleagues. This may be especially true for quarantine hospitals in which HCWs share longer resting times. Adapting the use of PPE and social distancing between colleagues while ensuring sufficient social interaction and support to maintain HCWs mental health thus constitutes a specific challenge for quarantine hospitals [20].

Preventing nosocomial SARS-CoV-2 infection remains an urgent need, especially in settings where preparedness does not match the risk faced by HCWs. The large variation in the infection risk we found between the hospitals we studied suggests that HCWs from quarantine hospitals may face a high risk of infection, but that, with sufficient anticipation and infection control measures, this risk can be brought down to levels similar to those observed in standard healthcare settings. A comprehensive assessment of the quarantine hospitals care model should also include its impact on HCWs' mental health as well as the benefits of an early diagnosis of those infected, which is likely to reduce further household and community transmission [20].

EMAEE-MESuRS working group on nosocomial SARS-CoV-2 modeling

Audrey Duval^{6,7,8}, Kenza Hamzi¹, Niels Hendrickx⁶, Kévin Jean^{1,2,5}, Sofía Jijón^{1,2}, Ajmal Oodally^{6,7,1}, Lulla Opatowski^{6,7}, George Shirreff^{6,7,8}, David RM Smith^{6,7,1}, Cynthia Tamandjou⁷ and Laura Temime^{1,2}

⁶ Institut Pasteur, Epidemiology and Modelling of Antibiotic Evasion (EMAEE), Paris, France

⁷ Université Paris-Saclay, UVSQ, Inserm, CESP, Anti-infective evasion and pharmacoepidemiology team, Montigny-Le-Bretonneux, France

⁸ IAME, UMR 1137, Université Paris 13, Sorbonne Paris Cité, France

Declaration of interest

The authors declare no conflicts of interest.

Funding

This study was supported by INSERM-ANRS (France Recherche Nord and Sud Sida-HIV Hépatites, grant number ANRS-COV-19).

Acknowledgments

The authors thank all the dedicated HCWs who contributed to this study.

REFERENCES

1. Lai X, Wang M, Qin C, et al. Coronavirus Disease 2019 (COVID-2019) Infection Among Health Care Workers and Implications for Prevention Measures in a Tertiary Hospital in Wuhan, China. *JAMA Netw Open* **2020**; 3:e209666. Available at: <https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2766227>.
2. Gilbert M, Pullano G, Pinotti F, et al. Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study. *Lancet* **2020**; 395:871–877.
3. Salyer SJ, Maeda J, Sembuche S, et al. The first and second waves of the COVID-19 pandemic in Africa: a cross-sectional study. *Lancet* **2021**; 397:1265–1275. Available at: [http://dx.doi.org/10.1016/S0140-6736\(21\)00632-2](http://dx.doi.org/10.1016/S0140-6736(21)00632-2).
4. WHO. Coronavirus Disease (COVID-19) Dashboard. Egypt. 2020; Available at: <https://covid19.who.int/region/emro/country/eg>.
5. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, et al. Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLOS Med* **2020**; 17:e1003346. Available at: <https://dx.plos.org/10.1371/journal.pmed.1003346>.
6. Yanes-Lane M, Winters N, Fregonese F, et al. Proportion of asymptomatic infection among COVID-19 positive persons and their transmission potential: A systematic review and meta-analysis. *PLoS One* **2020**; 15:e0241536. Available at: <https://dx.plos.org/10.1371/journal.pone.0241536>.
7. Medhat MA, El Kassas M. COVID-19 in Egypt: Uncovered figures or a different situation? *J Glob Health* **2020**; 10. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7303805>.
8. Sánchez-Taltavull D, Castelo-Szekely V, Candinas D, Roldán E, Beldi G. Modelling strategies to organize healthcare workforce during pandemics: Application to COVID-19. *J Theor Biol* **2021**; 523.
9. Meguid A, Talaat H, Shawky S, et al. SARS-CoV-2 infection among healthcare workers of a gastroenterological service in a tertiary care facility. *Arab J Gastroenterol* **2020**; 2020. Available at: <https://doi.org/10.1016/j.ajg.2020.07.005>.
10. Abdelmoniem R, Fouad R, Shawky S, et al. SARS-CoV-2 infection among asymptomatic healthcare workers of the emergency department in a tertiary care facility. *J Clin Virol* **2021**; 134:104710. Available at: <https://doi.org/10.1016/j.jcv.2020.104710>.
11. Mostafa A, Kandil S, El-Sayed MH, et al. Universal COVID-19 screening of 4040 health care workers in a resource-limited setting: an Egyptian pilot model in a university with 12 public hospitals and medical centers. *Int J Epidemiol* **2020**; :1–12. Available at: <https://academic.oup.com/ije/advance-article/doi/10.1093/ije/dyaa173/5936053>.
12. Egypt reopens 21 coronavirus quarantine hospitals. 2020. Available at:

- 415 [https://www.egyptindependent.com/egypt-reopens-21-coronavirus-quarantine-](https://www.egyptindependent.com/egypt-reopens-21-coronavirus-quarantine-hospitals/)
416 [hospitals/](https://www.egyptindependent.com/egypt-reopens-21-coronavirus-quarantine-hospitals/). Accessed 21 November 2021.
- 417 13. Lassaunière R, Frische A, Harboe ZB, et al. Evaluation of nine commercial SARS-
418 CoV-2 immunoassays. *medRxiv* **2020**; :2020.04.09.20056325. Available at:
419 <http://medrxiv.org/content/early/2020/04/10/2020.04.09.20056325.abstract>.
- 420 14. Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-
421 Negative Rate of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-
422 CoV-2 Tests by Time Since Exposure. *Ann Intern Med* **2020**; 173:262–267.
- 423 15. R Core Team. R: A language and environment for statistical computing. 2020;
424 Available at: <https://www.r-project.org/>.
- 425 16. Soetaert K, Petzoldt T. Inverse modelling, sensitivity and monte carlo analysis in R
426 using package FME. *J Stat Softw* **2010**; 33:1–28.
- 427 17. Zhao J, Yuan Q, Wang H, et al. Antibody Responses to SARS-CoV-2 in Patients With
428 Novel Coronavirus Disease 2019. *Clin Infect Dis* **2020**; 71:2027–2034. Available at:
429 <https://academic.oup.com/cid/article/71/16/2027/5812996>.
- 430 18. Jiang J-C, Zhang Y. Serological antibody testing in the COVID-19 pandemic: their
431 molecular basis and applications. *Biochem Soc Trans* **2020**; 48:2851–2863. Available
432 at: [https://portlandpress.com/biochemsoctrans/article/48/6/2851/226924/Serological-](https://portlandpress.com/biochemsoctrans/article/48/6/2851/226924/Serological-antibody-testing-in-the-COVID-19)
433 [antibody-testing-in-the-COVID-19](https://portlandpress.com/biochemsoctrans/article/48/6/2851/226924/Serological-antibody-testing-in-the-COVID-19).
- 434 19. Krastinova E, Garrait V, Lecam M-T, et al. Household transmission and incidence of
435 positive SARS-CoV-2 RT-PCR in symptomatic healthcare workers, clinical course and
436 outcome: a French hospital experience. *Occup Environ Med* **2020**; :oemed-2020-
437 106866. Available at: <https://oem.bmj.com/lookup/doi/10.1136/oemed-2020-106866>.
- 438 20. Youssef N, Mostafa A, Ezzat R, Yosef M, El Kassas M. Mental health status of health-
439 care professionals working in quarantine and non-quarantine Egyptian hospitals during
440 the COVID-19 pandemic. *East Mediterr Heal J* **2020**; 26:1155–1164. Available at:
441 [https://applications.emro.who.int/emhj/v26/10/1020-3397-2020-2610-1155-1164-](https://applications.emro.who.int/emhj/v26/10/1020-3397-2020-2610-1155-1164-eng.pdf)
442 [eng.pdf](https://applications.emro.who.int/emhj/v26/10/1020-3397-2020-2610-1155-1164-eng.pdf).
- 443 21. Houlihan CF, Vora N, Byrne T, et al. Pandemic peak SARS-CoV-2 infection and
444 seroconversion rates in London frontline health-care workers. *Lancet* **2020**; 0.
445 Available at: [https://www.thelancet.com/journals/lancet/article/PIIS0140-](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31484-7/abstract)
446 [6736\(20\)31484-7/abstract](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31484-7/abstract).
- 447 22. Eyre DW, Lumley SF, O'Donnell D, et al. Differential occupational risks to healthcare
448 workers from SARS-CoV-2 observed during a prospective observational study. *Elife*
449 **2020**; 9. Available at: <https://elifesciences.org/articles/60675>.
- 450 23. Mostafa A, Kandil S, El-Sayed MH, et al. SARS-CoV-2 seroconversion among 4040
451 Egyptian healthcare workers in 12 resource-limited healthcare facilities: A prospective
452 cohort study. *Int J Infect Dis* **2021**; Available at:
453 <https://doi.org/10.1016/j.ijid.2021.01.037>.

24. Schneider S, Piening B, Nouri-Pasovsky PA, Krüger AC, Gastmeier P, Aghdassi SJS. SARS-Coronavirus-2 cases in healthcare workers may not regularly originate from patient care: lessons from a university hospital on the underestimated risk of healthcare worker to healthcare worker transmission. *Antimicrob Resist Infect Control* **2020**; 9:192. Available at: <https://aricjournal.biomedcentral.com/articles/10.1186/s13756-020-00848-w>.